

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 November 2001 (29.11.2001)

PCT

(10) International Publication Number
WO 01/90358 A2

(51) International Patent Classification⁷: C12N 15/12,
C07K 14/715, 16/18, G01N 33/53, C12N 5/10

(21) International Application Number: PCT/US01/16767

(22) International Filing Date: 23 May 2001 (23.05.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/206,862 24 May 2000 (24.05.2000) US

(71) Applicant: SCHERING CORPORATION [US/US];
2000 Galloping Hill Road, Kenilworth, NJ 07033-0530
(US).

(72) Inventor: GORMAN, Daniel, M.; 6371 Central Avenue,
Newark, CA 94560 (US).

(74) Agent: ZARADIC, Sandy; Schering-Plough Corpora-
tion, Patent Department, K-6-1, 1990, 2000 Galloping Hill
Road, Kenilworth, NJ 07033-0530 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CZ,
DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, HR, HU,
ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LU, LV,
MA, MD, MG, MK, MN, MX, MZ, NO, NZ, PL, PT, RO,
RU, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UZ, VN,
YU, ZA.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— *as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii)) for all designations*

Published:

— *without international search report and to be republished
upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*



WO 01/90358 A2

(54) Title: MAMMALIAN RECEPTOR PROTEINS; RELATED REAGENTS AND METHODS

(57) Abstract: Nucleic acids encoding mammalian, e.g., primate, receptors, purified receptor proteins and fragments thereof. Anti-
bodies, both polyclonal and monoclonal, are also provided. Methods of using the compositions for both diagnostic and therapeutic
utilities are described.

5

MAMMALIAN RECEPTOR PROTEINS; RELATED REAGENTS AND METHODS

10

FIELD OF THE INVENTION

The present invention relates to compositions and methods for affecting mammalian physiology, including immune system function. In particular, it provides methods to regulate development and/or the immune system. Diagnostic and therapeutic uses of these materials are also disclosed.

15

BACKGROUND OF THE INVENTION

Recombinant DNA technology refers generally to techniques of integrating genetic information from a donor source into vectors for subsequent processing, such as through introduction into a host, whereby the transferred genetic information is copied and/or expressed in the new environment. Commonly, the genetic information exists in the form of complementary DNA (cDNA) derived from messenger RNA (mRNA) coding for a desired protein product. The carrier is frequently a plasmid having the capacity to incorporate cDNA for later replication in a host and, in some cases, actually to control expression of the cDNA and thereby direct synthesis of the encoded product in the host. See, e.g., Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual, (2d ed.) vols. 1-3, CSH Press, NY.

For some time, it has been known that the mammalian immune response is based on a series of complex cellular interactions, called the "immune network". Recent research has provided new insights into the inner workings of this network. While it remains clear that much of the immune response does, in fact, revolve around the network-like interactions of lymphocytes, macrophages, granulocytes, and other cells, immunologists now generally hold the opinion that soluble proteins, known as lymphokines, cytokines, or monokines, play critical roles in controlling these cellular interactions. Thus, there is considerable interest in the isolation, characterization, and mechanisms of action of cell modulatory factors, an understanding of which will lead to significant advancements in the diagnosis and therapy of numerous medical abnormalities, e.g., immune system disorders.

The immune system of vertebrates consists of a number of organs and several different cell types. Two major cell types include the myeloid and lymphoid lineages. Among the lymphoid cell lineage are B cells, which were originally characterized as differentiating in fetal liver or adult bone marrow, and T cells, which were originally
5 characterized as differentiating in the thymus. See, e.g., Paul (ed. 1998) Fundamental Immunology (4th ed.) Raven Press, New York; and Thomson (ed. 1994) The Cytokine Handbook 2d ed., Academic Press, San Diego. Lymphokines apparently mediate cellular activities in a variety of ways. They have been shown to support the proliferation, growth, and/or differentiation of cells, e.g., pluripotential hematopoietic stem cells, into
10 vast numbers of progenitors comprising diverse cellular lineages which make up a complex immune system. Proper and balanced interactions between the cellular components are necessary for a healthy immune response. The different cellular lineages often respond in a different manner when lymphokines are administered in conjunction with other agents.

15 Cell lineages especially important to the immune response include two classes of lymphocytes: B-cells, which can produce and secrete immunoglobulins (proteins with the capability of recognizing and binding to foreign matter to effect its removal), and T-cells of various subsets that secrete lymphokines and induce or suppress the B-cells and various other cells (including other T-cells) making up the immune network. These
20 lymphocytes interact with many other cell types.

Research to better understand and treat various immune disorders has been hampered by the general inability to maintain cells of the immune system in vitro. Immunologists have discovered that culturing many of these cells can be accomplished through the use of T-cell and other cell supernatants, which contain various growth
25 factors, including many of the lymphokines.

Various growth and regulatory factors exist which modulate morphogenetic development. And many receptors for cytokines are also known. Often there are at least two critical subunits in the functional receptor. See, e.g., Gonda and D'Andrea (1997) Blood 89:355-369; Presky, et al. (1996) Proc. Nat'l Acad. Sci. USA 93:14002-14007;
30 Drachman and Kaushansky (1995) Curr. Opin. Hematol. 2:22-28; Theze (1994) Eur. Cytokine Netw. 5:353-368; and Lemmon and Schlessinger (1994) Trends Biochem. Sci. 19:459-463.

From the foregoing, it is evident that the discovery and development of new soluble proteins and their receptors, including ones similar to lymphokines, should
35 contribute to new therapies for a wide range of degenerative or abnormal conditions which directly or indirectly involve development, differentiation, or function, e.g., of the

immune system and/or hematopoietic cells. In particular, the discovery and understanding of novel receptors for lymphokine-like molecules which enhance or potentiate the beneficial activities of other lymphokines would be highly advantageous. However, the lack of understanding of how the immune system is regulated or differentiates has blocked the ability to advantageously modulate the normal defensive mechanisms to biological challenges. Medical conditions characterized by abnormal or inappropriate regulation of the development or physiology of relevant cells thus remain unmanageable. The discovery and characterization of specific cytokines and their receptors will contribute to the development of therapies for a broad range of degenerative or other conditions which affect the immune system, hematopoietic cells, as well as other cell types. The present invention provides new receptors for ligands exhibiting similarity to cytokine like compositions and related compounds, and methods for their use.

SUMMARY OF THE INVENTION

The present invention is directed to novel receptors related to cytokine receptors, e.g., primate, cytokine receptor like molecular structures, designated DNAX Cytokine Receptor Subunits (DCRS), and their biological activities. In particular, it provides description of various subunits, designated DCRS6, DCRS7, DCRS8, DCRS9, and DCRS10. Primate, e.g, human, and rodent, e.g., mouse, embodiments of the various subunits are provided. It includes nucleic acids coding for the polypeptides themselves and methods for their production and use. The nucleic acids of the invention are characterized, in part, by their homology to cloned complementary DNA (cDNA) sequences enclosed herein.

The present invention provides a composition of matter selected from: a substantially pure or recombinant polypeptide comprising at least three distinct nonoverlapping segments of at least four amino acids identical to segments of SEQ ID NO: 2, 5, 8, 11, 23, or 26; a substantially pure or recombinant polypeptide comprising at least three distinct nonoverlapping segments of at least four amino acids identical to segments of SEQ ID NO: 14; a substantially pure or recombinant polypeptide comprising at least two distinct nonoverlapping segments of at least five amino acids identical to segments of SEQ ID NO: 14; a natural sequence DCRS8 comprising mature SEQ ID NO: 14; a fusion polypeptide comprising DCRS8 sequence; a substantially pure or recombinant polypeptide comprising at least three distinct nonoverlapping segments of at least four amino acids identical to segments of SEQ ID NO: 17 or 20; a substantially pure or recombinant polypeptide comprising at least two distinct nonoverlapping segments of at least five amino acids identical to segments of SEQ ID NO: 17 or 20; a natural

sequence DCRS9 comprising mature SEQ ID NO: 17 or 20; or a fusion polypeptide comprising DCRS9 sequence. Preferably, wherein the distinct nonoverlapping segments of identity include: one of at least eight amino acids; one of at least four amino acids and a second of at least five amino acids; at least three segments of at least four, five, and six amino acids, or one of at least twelve amino acids. In other embodiments, the:
5 polypeptide: comprises a mature sequence of Tables 1, 2, 3, 4, or 5; is an unglycosylated form of DCRS8 or DCRS9; is from a primate, such as a human; comprises at least seventeen amino acids of SEQ ID NO: 14 or 17; exhibits at least four nonoverlapping segments of at least seven amino acids of SEQ ID NO: 14 or 17; is a natural allelic
10 variant of DCRS8 or DCRS9; has a length at least about 30 amino acids; exhibits at least two non-overlapping epitopes which are specific for a primate DCRS8 or DCRS9; is glycosylated; has a molecular weight of at least 30 kD with natural glycosylation; is a synthetic polypeptide; is attached to a solid substrate; is conjugated to another chemical moiety; is a 5-fold or less substitution from natural sequence; or is a deletion or insertion
15 variant from a natural sequence.

The invention further embraces a composition comprising: a substantially pure DCRS8 or DCRS9 and another cytokine receptor family member; a sterile DCRS8 or DCRS9 polypeptide; the DCRS8 or DCRS9 polypeptide and a carrier, wherein the carrier is: an aqueous compound, including water, saline, and/or buffer; and/or formulated for
20 oral, rectal, nasal, topical, or parenteral administration. Additional embodiments include a polypeptide comprising: mature protein sequence of Tables 1, 2, 3, 4, or 5; a detection or purification tag, including a FLAG, His6, or Ig sequence; or sequence of another cytokine receptor protein. Kit embodiments include ones comprising a described polypeptide, and: a compartment comprising the protein or polypeptide; or instructions
25 for use or disposal of reagents in the kit.

Binding compositions are provided, e.g., comprising an antigen binding site from an antibody, which specifically binds to a natural DCRS8 or DCRS9 polypeptide, wherein: the binding compound is in a container; the DCRS8 or DCRS9 polypeptide is from a human; the binding compound is an Fv, Fab, or Fab2 fragment; the binding
30 compound is conjugated to another chemical moiety; or the antibody: is raised against a peptide sequence of a mature polypeptide of Table 3 or 4; is raised against a mature DCRS8 or DCRS9; is raised to a purified human DCRS8 or DCRS9; is immunoselected; is a polyclonal antibody; binds to a denatured DCRS8 or DCRS9; exhibits a K_d to antigen of at least 30 μM; is attached to a solid substrate, including a bead or plastic membrane; is
35 in a sterile composition; or is detectably labeled, including a radioactive or fluorescent label. Kits include ones comprising such a binding compound, and: a compartment

comprising the binding compound; or instructions for use or disposal of reagents in the kit.

The invention also provides methods of producing an antigen:antibody complex, comprising contacting under appropriate conditions a primate DCRS8 or DCRS9 polypeptide with a described antibody, thereby allowing the complex to form. Preferred methods include ones wherein: the complex is purified from other cytokine receptors; the complex is purified from other antibody; the contacting is with a sample comprising an interferon; the contacting allows quantitative detection of the antigen; the contacting is with a sample comprising the antibody; or the contacting allows quantitative detection of the antibody. Further compositions include those comprising: a sterile binding compound, as described, or the binding compound and a carrier, wherein the carrier is: an aqueous compound, including water, saline, and/or buffer; and/or formulated for oral, rectal, nasal, topical, or parenteral administration.

Nucleic acid compositions include an isolated or recombinant nucleic acid encoding a described polypeptide wherein the: DCRS8 or DCRS9 is from a human; or the nucleic acid: encodes an antigenic peptide sequence of Table 3 or 4; encodes a plurality of antigenic peptide sequences of Table 3 or 4; exhibits identity over at least thirteen nucleotides to a natural cDNA encoding the segment; is an expression vector; further comprises an origin of replication; is from a natural source; comprises a detectable label; comprises synthetic nucleotide sequence; is less than 6 kb, preferably less than 3 kb; is from a primate; comprises a natural full length coding sequence; is a hybridization probe for a gene encoding the DCRS8 or DCRS9; or is a PCR primer, PCR product, or mutagenesis primer. Also provided are a cell or tissue comprising such a recombinant nucleic acid, e.g., where the cell is: a prokaryotic cell; a eukaryotic cell; a bacterial cell; a yeast cell; an insect cell; a mammalian cell; a mouse cell; a primate cell; or a human cell.

Kit embodiments include those comprising a described nucleic acid and: a compartment comprising the nucleic acid; a compartment further comprising a primate DCRS8 or DCRS9 polypeptide; or instructions for use or disposal of reagents in the kit.

Other nucleic acids provided include ones which: hybridize under wash conditions of 30 minutes at 30° C and less than 2M salt to the coding portion of SEQ ID NO: 13 or 16; or exhibit identity over a stretch of at least about 30 nucleotides to a primate DCRS8 or DCRS9. Preferably, such will be nucleic acids where: the wash conditions are: at 45° C and/or 500 mM salt; at 55° C and/or 150 mM salt; or the stretch is at least 55 or 75 nucleotides.

Also provided are methods of modulating physiology or development of a cell or tissue culture cells comprising contacting the cell with an agonist or antagonist of a

mammalian DCRS8 or DCRS9. Preferably, the cell is transformed with a nucleic acid encoding the DCRS8 or DCRS9 and another cytokine receptor subunit.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

5

OUTLINE

- I. General
- II. Activities
- III. Nucleic acids
 - 10 A. encoding fragments, sequence, probes
 - B. mutations, chimeras, fusions
 - C. making nucleic acids
 - D. vectors, cells comprising
- IV. Proteins, Peptides
 - 15 A. fragments, sequence, immunogens, antigens
 - B. muteins
 - C. agonists/antagonists, functional equivalents
 - D. making proteins
- V. Making nucleic acids, proteins
 - 20 A. synthetic
 - B. recombinant
 - C. natural sources
- VI. Antibodies
 - 25 A. polyclonals
 - B. monoclonal
 - C. fragments; Kd
 - D. anti-idiotypic antibodies
 - E. hybridoma cell lines
- VII. Kits and Methods to quantify DCRSs
 - 30 A. ELISA
 - B. assay mRNA encoding
 - C. qualitative/quantitative
 - D. kits
- VIII. Therapeutic compositions, methods
 - 35 A. combination compositions
 - B. unit dose
 - C. administration
- IX. Screening
- X. Ligands

40

I. General

The present invention provides the amino acid sequence and DNA sequence of mammalian, herein primate, cytokine receptor-like subunit molecules, these designated DNAX Cytokine Receptor Subunits 6 (DCRS6), 7 (DCRS7), 8 (DCRS8), 9 (DCRS9),
 45 and 10 (DCRS10) having particular defined properties, both structural and biological.

Various cDNAs encoding these molecules were obtained from primate, e.g., human, and/or rodent, e.g., mouse, cDNA sequence libraries. Other primate or other mammalian counterparts would also be desired.

Some of the standard methods applicable are described or referenced, e.g., in
 5 Maniatis, et al. (1982) Molecular Cloning. A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor Press; Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual, (2d ed.), vols. 1-3, CSH Press, NY; Ausubel, et al., Biology, Greene Publishing Associates, Brooklyn, NY; or Ausubel, et al. (1987 and periodic supplements) Current Protocols in Molecular Biology, Greene/Wiley, New York; each of which is
 10 incorporated herein by reference.

Nucleotide (SEQ ID NO: 1) and corresponding amino acid sequence (SEQ ID NO: 2) of a primate, e.g., human, DCRS6 coding segment is shown in Table 1 along with reverse translation (SEQ ID NO: 3). Rodent, e.g., mouse, counterpart sequences are provided, e.g., SEQ ID NO: 4-6.

Similarly, nucleotide (SEQ ID NO: 7) and corresponding amino acid sequence (SEQ ID NO: 8) of a primate, e.g., human, DCRS7 coding segment is shown in Table 2 along with reverse translation (SEQ ID NO: 9). Rodent, e.g., mouse, counterpart sequences are provided, e.g., SEQ ID NO: 10-12. Nucleotide (SEQ ID NO: 13) and corresponding amino acid sequence (SEQ ID NO: 14) of a primate, e.g., human, DCRS8
 20 coding segment is shown in Table 3 along with reverse translation (SEQ ID NO: 15).

Nucleotide (SEQ ID NO: 16) and corresponding amino acid sequence (SEQ ID NO: 17) of a primate, e.g., human, DCRS9 coding segment is shown in Table 4 along with reverse translation (SEQ ID NO: 18). Rodent, e.g., mouse, counterpart sequences are provided, e.g., SEQ ID NO: 19-21. Nucleotide (SEQ ID NO: 22) and corresponding
 25 amino acid sequence (SEQ ID NO: 23) of a primate, e.g., human, DCRS10 coding segment is shown in Table 5 along with reverse translation (SEQ ID NO: 24). Rodent, e.g., mouse, counterpart sequences are provided, e.g., SEQ ID NO: 26-27.

30 Table 1: Nucleotide and polypeptide sequences of DNAX Cytokine Receptor Subunit like embodiments (DCRS6). Primate, e.g., human, embodiment (see SEQ ID NO: 1 and 2). Predicted signal sequence indicated, but may vary by a few positions and depending upon cell type.

35	gcg atg tcg ctc gtg ctg cta agc ctg gcc gcg ctg tgc agg agc gcc	48
	Met Ser Leu Val Leu Leu Ser Leu Ala Ala Leu Cys Arg Ser Ala	
	-10 -5 -1 1	
40	gta ccc cga gag ccg acc gtt caa tgt ggc tct gaa act ggg cca tct	96
	Val Pro Arg Glu Pro Thr Val Gln Cys Gly Ser Glu Thr Gly Pro Ser	
	5 10 15	

	cca gag tgg atg cta caa cat gat cta atc ccg gga gac ttg agg gac	144
	Pro Glu Trp Met Leu Gln His Asp Leu Ile Pro Gly Asp Leu Arg Asp	
	20 25 30	
5	ctc cga gta gaa cct gtt aca act agt gtt gca aca ggg gac tat tca	192
	Leu Arg Val Glu Pro Val Thr Thr Ser Val Ala Thr Gly Asp Tyr Ser	
	35 40 45	
10	att ttg atg aat gta agc tgg gta ctc cgg gca gat gcc agc atc cgc	240
	Ile Leu Met Asn Val Ser Trp Val Leu Arg Ala Asp Ala Ser Ile Arg	
	50 55 60 65	
15	ttg ttg aag gcc acc aag att tgt gtg acg ggc aaa agc aac ttc cag	288
	Leu Leu Lys Ala Thr Lys Ile Cys Val Thr Gly Lys Ser Asn Phe Gln	
	70 75 80	
20	tcc tac agc tgt gtg agg tgc aat tac aca gag gcc ttc cag act cag	336
	Ser Tyr Ser Cys Val Arg Cys Asn Tyr Thr Glu Ala Phe Gln Thr Gln	
	85 90 95	
25	acc aga ccc tct ggt ggt aaa tgg aca ttt tcc tat atc ggc ttc cct	384
	Thr Arg Pro Ser Gly Gly Lys Trp Thr Phe Ser Tyr Ile Gly Phe Pro	
	100 105 110	
30	gta gag ctg aac aca gtc tat ttc att ggg gcc cat aat att cct aat	432
	Val Glu Leu Asn Thr Val Tyr Phe Ile Gly Ala His Asn Ile Pro Asn	
	115 120 125	
35	gca aat atg aat gaa gat ggc cct tcc atg tct gtg aat ttc acc tca	480
	Ala Asn Met Asn Glu Asp Gly Pro Ser Met Ser Val Asn Phe Thr Ser	
	130 135 140 145	
40	cca ggc tgc cta gac cac ata atg aaa tat aaa aaa aag tgt gtc aag	528
	Pro Gly Cys Leu Asp His Ile Met Lys Tyr Lys Lys Lys Cys Val Lys	
	150 155 160	
45	gcc gga agc ctg tgg gat ccg aac atc act gct tgt aag aag aat gag	576
	Ala Gly Ser Leu Trp Asp Pro Asn Ile Thr Ala Cys Lys Lys Asn Glu	
	165 170 175	
50	gag aca gta gaa gtg aac ttc aca acc act ccc ctg gga aac aga tac	624
	Glu Thr Val Glu Val Asn Phe Thr Thr Thr Pro Leu Gly Asn Arg Tyr	
	180 185 190	
55	atg gct ctt atc caa cac agc act atc atc ggg ttt tct cag gtg ttt	672
	Met Ala Leu Ile Gln His Ser Thr Ile Ile Gly Phe Ser Gln Val Phe	
	195 200 205	
60	gag cca cac cag aag aaa caa acg cga gct tca gtg gtg att cca gtg	720
	Glu Pro His Gln Lys Lys Gln Thr Arg Ala Ser Val Val Ile Pro Val	
	210 215 220 225	
65	act ggg gat agt gaa ggt gct acg gtg cag ctg act cca tat ttt cct	768
	Thr Gly Asp Ser Glu Gly Ala Thr Val Gln Leu Thr Pro Tyr Phe Pro	
	230 235 240	

	act tgt ggc agc gac tgc atc cga cat aaa gga aca gtt gtg ctc tgc	816
	Thr Cys Gly Ser Asp Cys Ile Arg His Lys Gly Thr Val Val Leu Cys	
	245 250 255	
5	cca caa aca ggc gtc cct ttc cct ctg gat aac aac aaa agc aag ccg	864
	Pro Gln Thr Gly Val Pro Phe Pro Leu Asp Asn Asn Lys Ser Lys Pro	
	260 265 270	
10	gga ggc tgg ctg cct ctc ctc ctg ctg tct ctg ctg gtg gcc aca tgg	912
	Gly Gly Trp Leu Pro Leu Leu Leu Leu Ser Leu Leu Val Ala Thr Trp	
	275 280 285	
15	gtg ctg gtg gca ggg atc tat cta atg tgg agg cac gaa agg atc aag	960
	Val Leu Val Ala Gly Ile Tyr Leu Met Trp Arg His Glu Arg Ile Lys	
	290 295 300 305	
20	aag act tcc ttt tct acc acc aca cta ctg ccc ccc att aag gtt ctt	1008
	Lys Thr Ser Phe Ser Thr Thr Thr Leu Leu Pro Pro Ile Lys Val Leu	
	310 315 320	
	gtg gtt tac cca tct gaa ata tgt ttc cat cac aca att tgt tac ttc	1056
	Val Val Tyr Pro Ser Glu Ile Cys Phe His His Thr Ile Cys Tyr Phe	
	325 330 335	
25	act gaa ttt ctt caa aac cat tgc aga agt gag gtc atc ctt gaa aag	1104
	Thr Glu Phe Leu Gln Asn His Cys Arg Ser Glu Val Ile Leu Glu Lys	
	340 345 350	
30	tgg cag aaa aag aaa ata gca gag atg ggt cca gtg cag tgg ctt gcc	1152
	Trp Gln Lys Lys Lys Ile Ala Glu Met Gly Pro Val Gln Trp Leu Ala	
	355 360 365	
35	act caa aag aag gca gca gac aaa gtc gtc ttc ctt ctt tcc aat gac	1200
	Thr Gln Lys Lys Ala Ala Asp Lys Val Val Phe Leu Leu Ser Asn Asp	
	370 375 380 385	
40	gtc aac agt gtg tgc gat ggt acc tgt ggc aag agc gag ggc agt ccc	1248
	Val Asn Ser Val Cys Asp Gly Thr Cys Gly Lys Ser Glu Gly Ser Pro	
	390 395 400	
	agt gag aac tct caa gac ctc ttc ccc ctt gcc ttt aac ctt ttc tgc	1296
	Ser Glu Asn Ser Gln Asp Leu Phe Pro Leu Ala Phe Asn Leu Phe Cys	
	405 410 415	
45	agt gat cta aga agc cag att cat ctg cac aaa tac gtg gtg gtc tac	1344
	Ser Asp Leu Arg Ser Gln Ile His Leu His Lys Tyr Val Val Val Tyr	
	420 425 430	
50	ttt aga gag att gat aca aaa gac gat tac aat gct ctc agt gtc tgc	1392
	Phe Arg Glu Ile Asp Thr Lys Asp Asp Tyr Asn Ala Leu Ser Val Cys	
	435 440 445	
55	ccc aag tac cac ctc atg aag gat gcc act gct ttc tgt gca gaa ctt	1440
	Pro Lys Tyr His Leu Met Lys Asp Ala Thr Ala Phe Cys Ala Glu Leu	
	450 455 460 465	

ctc cat gtc aag cag cag gtg tca gca gga aaa aga tca caa gcc tgc 1488
 Leu His Val Lys Gln Gln Val Ser Ala Gly Lys Arg Ser Gln Ala Cys
 470 475 480

5 cac gat ggc tgc tgc tcc ttg tagccaccc atgagaagca agagacctta 1539
 His Asp Gly Cys Cys Ser Leu
 485

10 aaggcttcct atcccaccaa ttacagggaa aaaacgtgtg atgatcctga agcttactat 1599
 gcagcctaca aacagcctta gtaattaaaa ctttttatac caataaaatt ttcaaattat 1659
 gctaactaat gtagcattaa ctaacgattg gaaactacat ttacaacttc aaagctgttt 1719

15 tatacataga aatcaattac agctttaatt gaaaactgta accattttga taatgcaaca 1779
 ataaagcatc ttcagcc 1796

20 MSLVLLSLAALCRSAVPREPTVQCGSETGPSPEWMLQHDLPGLRDLRVEPVTTSVATGDYSILMNVSWSVL
 RADASIRLLKATKICVTGKSNFQSYSCVRCNYTEAFQTQTRPSGGKWTFSYIGFPVELNTVYFIGAHNIPNA
 NMNEDGPSMSVNFTSPGCLDHIMKYKKKCVKAGSLWDPNITACKKNEETVEVNFTTTPLGNRYMALIQHSTI
 IGFSQVFEPHQKKQTRASVVIPTGDSGATVQLTPYFPTCGSDCIRHKGTVVLCPTGVPFPLDNNKSKPG
 GWLPLLLLSLLVATWVLVAGIYLMWRHERIKKTSFSTTTLLPPIKVLVVPSEICFHHTICYFTEFLQNHCR
 SEVILEKWQKKKIAEMGPVQWLATQKKAADKVFLSNDVNSVCDGTCGKSEGSPSENSQDLFPLAFNLFCS
 25 DLRSQIHLHKYVVVYFREIDTKDDYNALSVC PKYHLMKDATAFCAELLHV KQVVSAGKRSQACHDGCCSL.

Reverse translation of primate, e.g., human, DCRS6 (SEQ ID NO: 3):

30 atgwsnytnng tnytnytnws nytnngcngcn ytntgymgnw sngcngtncc nmngngarcen 60
 acngtncart gyggwnwsnga racngngccn wsncngart ggatgytnca rcaygayytn 120

35 athccnggng ayytnmngnga yytnmngnngn garccngtna cnacnwsngt ngcnacnggn 180
 gaytaywsna thytnatgaa ygtwnsntgg gtnytnmngn cngaygcnws nathmngnytn 240
 ytnaargcna cnaarathtg ygtnacnggn aarwsnaayt tycarwsnta ywsntgygtn 300

40 mgntgyaayt ayacngargc nttycaracn caracnmngn cnwsnggngg naartggacn 360
 ttywsntaya thggnttycc ngtngarytn aayacngtnt aytyathgg ngcnacayaay 420
 athccnaayg cnaayatgaa ygargayggn ccnwsnatgw sngtnaaytt yacnwsnccn 480

45 ggntgyytng aycayathat gaartayaar aaraartgyg tnaargcngg nwsnytnngg 540
 gayccnaaya thacngcntg yaaraaraay gargaracng tngargtnaa yttyacnacn 600

50 acncnytnng gnaaymgnta yatggcnytn athcarcayw snacnathat hggnttywsn 660
 cargtnnttyg arccncayca raaraarcac acnmngngcnw sngtnngtnat hccngtnacn 720
 ggngaywsng arggngcnac ngtnrcarytn acncntayt tyccnacntg yggnwsngay 780

55 tgyathmgnc ayaarggnac ngtnngtnytn tgyccncara cngngngtncc nttyccnytn 840
 gayaayaaya arwsnaarcc ngngngntgg ytnccnytny tnytnytnws nytnytnngtn 900

gcnacntggg tnytngtngc nggnathtay ytnatgtggm gncaygarmg nathaaraar 960
 acnwsnttyw snacnacnac nytnytncn ccnathaarg tnytngtngt ntayccnwsn 1020
 5 garathtgyt tycaycayac nathtgytay ttyacngart tyytncaraa ycaytgymgn 1080
 wsgargttna thytngaraa rtggcaraar aaraarathg cngaratggg nccngtncar 1140
 10 tggytngcna cncaraaraa rgcngcngay aargtngtnt tyytnytnws naaygaygtn 1200
 aaywsngtnt gygaygnac ntgyggnaar wsgarggnw snccnwsnga raaywsncar 1260
 gayytnttyc cnytngcntt yaayytntty tgywsngayy tnmgnwsnca rathcayytn 1320
 15 cayaartayg tngtngtnta yttymngar athgayacna argaygayta yaaygcnytn 1380
 wsngtntgyc cnaartayca yytnatgaar gaygcnacng cnttytgygc ngarytnytn 1440
 20 caygtnaarc arcargtnws ngcnggnaar mgnwsncarg cntgycayga yggntgytgy 1500
 wsnytn 1506

25 Rodent, e.g., mouse embodiment (see SEQ ID NO: 4 and 5).

gat ttc agc agc cag acg cat ctg cac aaa tac ctg gag gtc tat ctt 48
 Asp Phe Ser Ser Gln Thr His Leu His Lys Tyr Leu Glu Val Tyr Leu
 1 5 10 15
 30 ggg gga gca gac ctc aaa ggc gac tat aat gcc ctg agt gtc tgc ccg 96
 Gly Gly Ala Asp Leu Lys Gly Asp Tyr Asn Ala Leu Ser Val Cys Pro
 20 25 30
 35 caa tat cat ctc atg aag gac gcc aca gct ttc cac aca gaa ctt ctc 144
 Gln Tyr His Leu Met Lys Asp Ala Thr Ala Phe His Thr Glu Leu Leu
 35 40 45
 40 aag gct acg cag agc atg tca gtg aag aaa cgc tca caa gcc tgc cat 192
 Lys Ala Thr Gln Ser Met Ser Val Lys Lys Arg Ser Gln Ala Cys His
 50 55 60
 gat agc tgt tca ccc ttg tagtccaccc gggggaatag agactctgaa 240
 Asp Ser Cys Ser Pro Leu
 45 65 70
 gccttcctac tctcccttcc agtgacaaat gctgtgtgac gactctgaaa tgtgtgggag 300
 aggctgtgtg gaggtagtgc tatgtacaaa cttgctttaa aactggagtt tgcaaagtca 360
 50 acctgagcat acacgcctga ggctagtcac tggctggatt tatgaagaca acacagttac 420
 agacaataat gagtgggacc tacatttggg atatacccaa agctgggtaa tgattatcac 480
 55 tgagaaccac gcaactctggc catgaggtaa tacggcactt ccctgtcagg ctgtctgtca 540
 gggtgggtct gtcttgact gccatgctc tatgctgcac gtagaccgtt ttgtaacatt 600
 ttaatctgtt aatgaataat ccgtttggga ggctctc 637

DFSSQTHLHKYLEVYLGGADLKG DYNALSVCPQYHLMKDATAFHTELLKATQSMSVKKRSQACHDSCSPL.

5 Reverse translation of rodent, e.g., mouse, DCRS6 (SEQ ID NO: 6):

gayttywsnw sncaracnca yytncayaar tayytnngarg tntayytnng ngngngcngay 60
 ytnaarggng aytayaaygc nytnwsngtn tgyccncart aycayytnat gaargaygcn 120
 10 acngcnttyc ayacngaryt nytnaargcn acncarwsna tgwsngtnaa raarmgnwsn 180
 cargcntgyc aygaywsntg ywsnccnytn 210

15

Table 2: Nucleotide and polypeptide sequences of DNAX Cytokine Receptor Subunit like
 embodiments (DCRS7). Primate, e.g., human, embodiment (see SEQ ID NO: 7 and 8).
 Predicted signal sequence indicated, but may vary by a few positions and depending upon cell
 type.

20

gagtcaggac tcccaggaca gagagtgcac aaactaccca gcacagcccc ctccgcccc 60
 tctggaggct gaagagggat tccagcccct gccaccaca gacacgggct gactgggggtg 120

25

tctgcccccc ttgggggcan ccacagggcc tcaggcctgg gtgccacctg gcactagaag 180

atg cct gtg ccc tgg ttc ttg ctg tcc ttg gca ctg ggc cga agc cag 228
 Met Pro Val Pro Trp Phe Leu Leu Ser Leu Ala Leu Gly Arg Ser Gln
 -20 -15 -10 -5

30

tgg atc ctt tct ctg gag agg ctt gtg ggg cct cag gac gct acc cac 276
 Trp Ile Leu Ser Leu Glu Arg Leu Val Gly Pro Gln Asp Ala Thr His
 -1 1 5 10

35

tgc tct ccg ggc ctc tcc tgc cgc ctc tgg gac agt gac ata ctc tgc 324
 Cys Ser Pro Gly Leu Ser Cys Arg Leu Trp Asp Ser Asp Ile Leu Cys
 15 20 25

40

ctg cct ggg gac atc gtg cct gct ccg ggc ccc gtg ctg gcg cct acg 372
 Leu Pro Gly Asp Ile Val Pro Ala Pro Gly Pro Val Leu Ala Pro Thr
 30 35 40

45

cac ctg cag aca gag ctg gtg ctg agg tgc cag aag gag acc gac tgt 420
 His Leu Gln Thr Glu Leu Val Leu Arg Cys Gln Lys Glu Thr Asp Cys
 45 50 55 60

50

gac ctc tgt ctg cgt gtg gct gtc cac ttg gcc gtg cat ggg cac tgg 468
 Asp Leu Cys Leu Arg Val Ala Val His Leu Ala Val His Gly His Trp
 65 70 75

gaa gag cct gaa gat gag gaa aag ttt gga gga gca gct gac tta ggg 516
 Glu Glu Pro Glu Asp Glu Glu Lys Phe Gly Gly Ala Ala Asp Leu Gly
 80 85 90

55

gtg gag gag cct agg aat gcc tct ctc cag gcc caa gtc gtg ctc tcc 564
 Val Glu Glu Pro Arg Asn Ala Ser Leu Gln Ala Gln Val Val Leu Ser
 95 100 105

	ttc cag gcc tac cct act gcc cgc tgc gtc ctg ctg gag gtg caa gtg	612
	Phe Gln Ala Tyr Pro Thr Ala Arg Cys Val Leu Leu Glu Val Gln Val	
	110 115 120	
5	cct gct gcc ctt gtg cag ttt ggt cag tct gtg ggc tct gtg gta tat	660
	Pro Ala Ala Leu Val Gln Phe Gly Gln Ser Val Gly Ser Val Val Tyr	
	125 130 135 140	
10	gac tgc ttc gag gct gcc cta ggg agt gag gta cga atc tgg tcc tat	708
	Asp Cys Phe Glu Ala Ala Leu Gly Ser Glu Val Arg Ile Trp Ser Tyr	
	145 150 155	
15	act cag ccc agg tac gag aag gaa ctc aac cac aca cag cag ctg cct	756
	Thr Gln Pro Arg Tyr Glu Lys Glu Leu Asn His Thr Gln Gln Leu Pro	
	160 165 170	
20	gac tgc agg ggg ctc gaa gtc tgg aac agc atc ccg agc tgc tgg gcc	804
	Asp Cys Arg Gly Leu Glu Val Trp Asn Ser Ile Pro Ser Cys Trp Ala	
	175 180 185	
25	ctg ccc tgg ctc aac gtg tca gca gat ggt gac aac gtg cat ctg gtt	852
	Leu Pro Trp Leu Asn Val Ser Ala Asp Gly Asp Asn Val His Leu Val	
	190 195 200	
30	ctg aat gtc tct gag gag cag cac ttc ggc ctc tcc ctg tac tgg aat	900
	Leu Asn Val Ser Glu Glu Gln His Phe Gly Leu Ser Leu Tyr Trp Asn	
	205 210 215 220	
35	cag gtc cag ggc ccc cca aaa ccc cgg tgg cac aaa aac ctg act gga	948
	Gln Val Gln Gly Pro Pro Lys Pro Arg Trp His Lys Asn Leu Thr Gly	
	225 230 235	
40	ccg cag atc att acc ttg aac cac aca gac ctg gtt ccc tgc ctc tgt	996
	Pro Gln Ile Ile Thr Leu Asn His Thr Asp Leu Val Pro Cys Leu Cys	
	240 245 250	
45	att cag gtg tgg cct ctg gaa cct gac tcc gtt agg acg aac atc tgc	1044
	Ile Gln Val Trp Pro Leu Glu Pro Asp Ser Val Arg Thr Asn Ile Cys	
	255 260 265	
50	ccc ttc agg gag gac ccc cgc gca cac cag aac ctc tgg caa gcc gcc	1092
	Pro Phe Arg Glu Asp Pro Arg Ala His Gln Asn Leu Trp Gln Ala Ala	
	270 275 280	
55	cga ctg cga ctg ctg acc ctg cag agc tgg ctg ctg gac gca ccg tgc	1140
	Arg Leu Arg Leu Leu Thr Leu Gln Ser Trp Leu Leu Asp Ala Pro Cys	
	285 290 295 300	
55	tcg ctg ccc gca gaa gcg gca ctg tgc tgg cgg gct ccg ggt ggg gac	1188
	Ser Leu Pro Ala Glu Ala Ala Leu Cys Trp Arg Ala Pro Gly Gly Asp	
	305 310 315	
55	ccc tgc cag cca ctg gtc cca ccg ctt tcc tgg gag aat gtc act gtg	1236
	Pro Cys Gln Pro Leu Val Pro Pro Leu Ser Trp Glu Asn Val Thr Val	
	320 325 330	
55	gac gtg aac agc tcg gag aag ctg cag ctg cag gag tgc ttg tgg gct	1284
	Asp Val Asn Ser Ser Glu Lys Leu Gln Leu Gln Glu Cys Leu Trp Ala	
	335 340 345	

5	gac tcc ctg ggg cct ctc aaa gac gat gtg cta ctg ttg gag aca cga 1332														
	Asp Ser Leu Gly Pro Leu Lys Asp Asp Val Leu Leu Leu Glu Thr Arg	350				355				360					
10	ggc ccc cag gac aac aga tcc ctc tgt gcc ttg gaa ccc agt ggc tgt 1380														
	Gly Pro Gln Asp Asn Arg Ser Leu Cys Ala Leu Glu Pro Ser Gly Cys	365			370				375					380	
15	act tca cta ccc agc aaa gcc tcc acg agg gca gct cgc ctt gga gag 1428														
	Thr Ser Leu Pro Ser Lys Ala Ser Thr Arg Ala Ala Arg Leu Gly Glu			385				390					395		
20	tac tta cta caa gac ctg cag tca ggc cag tgt ctg cag cta tgg gac 1476														
	Tyr Leu Leu Gln Asp Leu Gln Ser Gly Gln Cys Leu Gln Leu Trp Asp		400				405				410				
25	gat gac ttg gga gcg cta tgg gcc tgc ccc atg gac aaa tac atc cac 1524														
	Asp Asp Leu Gly Ala Leu Trp Ala Cys Pro Met Asp Lys Tyr Ile His		415				420				425				
30	aag cgc tgg gcc ctc gtg tgg ctg gcc tgc cta ctc ttt gcc gct gcg 1572														
	Lys Arg Trp Ala Leu Val Trp Leu Ala Cys Leu Leu Phe Ala Ala Ala		430			435				440					
35	ctt tcc ctc atc ctc ctt ctc aaa aag gat cac gcg aaa ggg tgg ctg 1620														
	Leu Ser Leu Ile Leu Leu Leu Lys Lys Asp His Ala Lys Gly Trp Leu	445			450				455					460	
40	agg ctc ttg aaa cag gac gtc cgc tcg ggg gcg gcc gcc agg ggc cgc 1668														
	Arg Leu Leu Lys Gln Asp Val Arg Ser Gly Ala Ala Ala Arg Gly Arg			465				470					475		
45	gcg gct ctg ctc ctc tac tca gcc gat gac tcg ggt ttc gag cgc ctg 1716														
	Ala Ala Leu Leu Leu Tyr Ser Ala Asp Asp Ser Gly Phe Glu Arg Leu		480				485				490				
50	gtg ggc gcc ctg gcg tcg gcc ctg tgc cag ctg ccg ctg cgc gtg gcc 1764														
	Val Gly Ala Leu Ala Ser Ala Leu Cys Gln Leu Pro Leu Arg Val Ala		495				500				505				
55	gta gac ctg tgg agc cgt cgt gaa ctg agc gcg cag ggg ccc gtg gct 1812														
	Val Asp Leu Trp Ser Arg Arg Glu Leu Ser Ala Gln Gly Pro Val Ala		510			515				520					
60	tgg ttt cac gcg cag cgg cgc cag acc ctg cag gag ggc ggc gtg gtg 1860														
	Trp Phe His Ala Gln Arg Arg Gln Thr Leu Gln Glu Gly Gly Val Val	525			530				535					540	
65	gtc ttg ctc ttc tct ccc ggt gcg gtg gcg ctg tgc agc gag tgg cta 1908														
	Val Leu Leu Phe Ser Pro Gly Ala Val Ala Leu Cys Ser Glu Trp Leu			545				550					555		
70	cag gat ggg gtg tcc ggg ccc ggg gcg cac ggc ccg cac gac gcc ttc 1956														
	Gln Asp Gly Val Ser Gly Pro Gly Ala His Gly Pro His Asp Ala Phe		560				565				570				

	cgc gcc tcg ctc agc tgc gtg ctg ccc gac ttc ttg cag ggc cgg gcg	2004
	Arg Ala Ser Leu Ser Cys Val Leu Pro Asp Phe Leu Gln Gly Arg Ala	
	575 580 585	
5	ccc ggc agc tac gtg ggg gcc tgc ttc gac agg ctg ctc cac ccg gac	2052
	Pro Gly Ser Tyr Val Gly Ala Cys Phe Asp Arg Leu Leu His Pro Asp	
	590 595 600	
10	gcc gta ccc gcc ctt ttc cgc acc gtg ccc gtc ttc aca ctg ccc tcc	2100
	Ala Val Pro Ala Leu Phe Arg Thr Val Pro Val Phe Thr Leu Pro Ser	
	605 610 615 620	
15	caa ctg cca gac ttc ctg ggg gcc ctg cag cag cct cgc gcc ccg cgt	2148
	Gln Leu Pro Asp Phe Leu Gly Ala Leu Gln Gln Pro Arg Ala Pro Arg	
	625 630 635	
20	tcc ggg cgg ctc caa gag aga gcg gag caa gtg tcc cgg gcc ctt cag	2196
	Ser Gly Arg Leu Gln Glu Arg Ala Glu Gln Val Ser Arg Ala Leu Gln	
	640 645 650	
25	cca gcc ctg gat agc tac ttc cat ccc ccg ggg acn tcc gcg ccg gga	2244
	Pro Ala Leu Asp Ser Tyr Phe His Pro Pro Gly Xaa Ser Ala Pro Gly	
	655 660 665	
30	cgc ggg gtg gga cca ggg gcg gga cct ggg gcg ggg gac ggg act	2289
	Arg Gly Val Gly Pro Gly Ala Gly Pro Gly Ala Gly Asp Gly Thr	
	670 675 680	
35	taaataaagg cagacgctg	2308
40	MPVPWFLLSLALGRSQWILSLERLVGPQDATHCSPGLSCLWSDILCLPGDIVPAPGPVLAPTHLQTELVL RCQKETDCDLCLRVAVHLAVHGHWEPEDEEKFGGAADLGVEEPRNASLQAQVVLVSFQAYPTARCVLLELVQV PAALVQFGQSVGSVVYDCFEAALGSEVRIWSYTQPRYEKELNHTQQLPDCRGLEVWNSIPSCWALPWLNVSA DGDNVHLVLNVSEEQHFGLSLYWNQVQGPCKPRWHKNLTGPGQIITLNHTDLVPCLCIQVWPLEPDSVRTNIC PFREDPRAHQNLWQAARLRLTLQSWLLDAPCSLPAAALCWRAAGGDPQCPLVPPLSWENVTVDVNSSEKL QLQECWLWADSLGPKDDVLLLETRGPQDNRLCALEPSGCTSLPSKASTRAARLGEYLLQDLQSGQCLQLWD DDLGALWACPMCKYIHKRWALVWLACLLFAAALSLILLKKDHAKGWLRLKQDVRSGAAARGRAALLLYSA DDSGFERLVGALASALCQLPLRVAVDLWSRRELSAQGPVAVFWFAHQRRQTLEGGVVVLLFSFGAVALCSEWL QDGVSGPGAHGPHDAFRASLSCVLPDFLQGRAPGSYVGACFDRLHLPDAVPALFRTVPVFTLPSQLPDLFLGA LQQPRAPRSGRLQERAEQVSRALQPALDSYFHPGTSAPGRGVGPGAGPGAGDGT.	

Reverse translation of primate, e.g., human, DCRS7 (SEQ ID NO: 9):

45	atgccngtnc cntgggttyt nytnwsnytn gcnytnggnm gnwsncartg gathytnwsn	60
	ytngarmgny tngtnggncc ncargaygcn acncaytgyw snccnggnytn nwsntgymgn	120
50	ytntggggyw sngayathyt ntgyytnccn ggngayathg tncngcncncc nggncngtn	180
	ytngcncnca cncayytnc racngarytn gtnytnmgnt gycaraarga racngaytgy	240
	gayytnntgyy tnmngntngc ngtncayytn gcngtncayg gncaytggga rgarccngar	300
55	gaygargara arttyggngg ngcngcngay ytngnggtng argarccnmg naaygcnwsn	360
	ytncargcnc argtngtnyt nwsnttycar gcntayccna cngcnmgntg ygtnytnytn	420
	gargtncarg tncngcngc nytngtncar ttyggncarw sngtnggnws ngtngtntay	480

5 gaytgyttyg argcngcnyt nggnwsngar gtnmgnatht ggwsntayac ncarccnmgn 540
 taygaraarg arytnaayca yacncarcar ytncngayt gymngggnyt ngargtntgg 600
 aaywsnathc cnwsntgytg ggcnytnccn tggytnaayg tnwsngcnga yggngayaay 660
 gtncayytng tnytnaaygt nwsngargar carcayttyg gnytnwsnyt ntaytggay 720
 10 cargtncarg gncncnccnaa rccnmgttg cayaaraayy tnacnggncc ncarathath 780
 acnytnaayc ayacngayyt ngtnccntgy ytntgyathc argtntggcc nytngarccn 840
 15 gaywsngtnm gnacnaayat htgyccntty mgngargayc cnmgngcnca ycaraayytn 900
 tggcargcng cnmgnytnmg nytnytnacn ytncarwsnt ggytnytnga ygcncntgy 960
 wsnytnccng cngargcngc nytntgytgg mgngcncng gngnggaycc ntgycarccn 1020
 20 ytngtnccnc cnytnwsntg ggaraaygtn acngtngayg tnaaywsnws ngaraarytn 1080
 carytncarg artgyytntg ggcngaywsn ytnggncny tnaargayga ygtnytnytn 1140
 ytngaracnm gnggncnca rgayaaymgn wsnytnthyg cnytngarcc nwsnggntgy 1200
 25 acnwsnytn cwnsnaargc nwsnacnmgn gcngcnmgny tngngarta yytnytnear 1260
 gayytnearw snggncartg yytnarytn tgggaygayg ayytnggngc nytntgggcn 1320
 30 tgyccnatgg ayaartayat hcayaarmgn tgggcnytn tntggytngc ntgyytnytn 1380
 ttygcngcng cnytnwsnyt nathytnytn ytnaaraarg aycaygcnaa rggntggytn 1440
 35 mgnytnytna arcargaygt nmgnwsnggn gcngcngcnm gnggnmgngc ngcnytnytn 1500
 ytntaywsng cngaygayws nggnttygar mgnytngtng gngcnytnge nwsngcnytn 1560
 tgycarytn cnytnmgnt ngcngtngay ytntggwsnm gnmngaryt nwsngcncar 1620
 40 ggnccngtng cntggtyca ygcncarmgn mgncaracny tncargargg ngngtngtn 1680
 gtnytnytn tywsncngg ngcngtngcn ytntgywsng artggytnca rgayggngtn 1740
 wsnggncng gngcncaygg nccncaygay gcnttymng cwnsnytnws ntgygtnytn 1800
 45 ccngayttyy tncarggnmg ngcncnggn wsntaygtng gngcntgytt ygaymgnytn 1860
 ytncaycng aygngtncc ngcnytnntty mgnacngtn cngtnnttyac nytnccnwsn 1920
 50 carytnccng aytyytngg ngcnytnear carccnmng cncnmgng nsgnmgnyn 1980
 cargarmng cngarcargt nwsnmngcn ytncarccng cnytngayws ntayttycay 2040
 ccncnggna cwnsgncnc nggnmgnggn gtnggncng gngcnggnc ngngcnggn 2100
 55 gayggnacn 2109

Rodent, e.g., mouse, embodiment (see SEQ ID NO: 10 and 11). Predicted signal sequence indicated, but may vary by a few positions and depending upon cell type.

5	ccaaatcgaa agcacgggag ctgatactgg gcctggagtc caggctcact ggagtgggga	60
	agcatggctg gagaggaatt ctagcccttg ctctctccca gggacacggg gctgattgtc	120
	agcaggggag aggggtctgc ccccccttgg gggggcagga cggggcctca ggcctgggtg	180
10	ctgtccggca cctggaag atg cct gtg tcc tgg ttc ctg ctg tcc ttg gca	231
	Met Pro Val Ser Trp Phe Leu Leu Ser Leu Ala	
	-20 -15 -10	
15	ctg ggc cga aac cct gtg gtc gtc tct ctg gag aga ctg atg gag cct	279
	Leu Gly Arg Asn Pro Val Val Val Ser Leu Glu Arg Leu Met Glu Pro	
	-5 -1 1 5	
20	cag gac act gca cgc tgc tct cta ggc ctc tcc tgc cac ctc tgg gat	327
	Gln Asp Thr Ala Arg Cys Ser Leu Gly Leu Ser Cys His Leu Trp Asp	
	10 15 20	
25	ggt gac gtg ctc tgc ctg cct gga agc ctc cag tct gcc cca ggc cct	375
	Gly Asp Val Leu Cys Leu Pro Gly Ser Leu Gln Ser Ala Pro Gly Pro	
	25 30 35	
30	gtg cta gtg cct acc cgc ctg cag acg gag ctg gtg ctg agg tgt cca	423
	Val Leu Val Pro Thr Arg Leu Gln Thr Glu Leu Val Leu Arg Cys Pro	
	40 45 50 55	
35	cag aag aca gat tgc gcc ctc tgt gtc cgt gtg gtg gtc cac ttg gcc	471
	Gln Lys Thr Asp Cys Ala Leu Cys Val Arg Val Val Val His Leu Ala	
	60 65 70	
40	gtg cat ggg cac tgg gca gag cct gaa gaa gct gga aag tct gat tca	519
	Val His Gly His Trp Ala Glu Pro Glu Glu Ala Gly Lys Ser Asp Ser	
	75 80 85	
45	gaa ctc cag gag tct agg aac gcc tct ctc cag gcc cag gtg gtg ctc	567
	Glu Leu Gln Glu Ser Arg Asn Ala Ser Leu Gln Ala Gln Val Val Leu	
	90 95 100	
50	tcc ttc cag gcc tac ccc atc gcc cgc tgt gcc ctg ctg gag gtc cag	615
	Ser Phe Gln Ala Tyr Pro Ile Ala Arg Cys Ala Leu Leu Glu Val Gln	
	105 110 115	
55	gtg ccc gct gac ctg gtg cag cct ggt cag tcc gtg ggt tct gcg gta	663
	Val Pro Ala Asp Leu Val Gln Pro Gly Gln Ser Val Gly Ser Ala Val	
	120 125 130 135	
55	ttt gac tgt ttc gag gct agt ctt ggg gct gag gta cag atc tgg tcc	711
	Phe Asp Cys Phe Glu Ala Ser Leu Gly Ala Glu Val Gln Ile Trp Ser	
	140 145 150	
55	tac acg aag ccc agg tac cag aaa gag ctc aac ctc aca cag cag ctg	759
	Tyr Thr Lys Pro Arg Tyr Gln Lys Glu Leu Asn Leu Thr Gln Gln Leu	
	155 160 165	

	cct gac tgc agg ggt ctt gaa gtc cgg gac agc atc cag agc tgc tgg	807
	Pro Asp Cys Arg Gly Leu Glu Val Arg Asp Ser Ile Gln Ser Cys Trp	
	170 175 180	
5	gtc ctg ccc tgg ctc aat gtg tct aca gat ggt gac aat gtc ctt ctg	855
	Val Leu Pro Trp Leu Asn Val Ser Thr Asp Gly Asp Asn Val Leu Leu	
	185 190 195	
10	aca ctg gat gtc tct gag gag cag gac ttt agc ttc tta ctg tac ctg	903
	Thr Leu Asp Val Ser Glu Glu Gln Asp Phe Ser Phe Leu Leu Tyr Leu	
	200 205 210 215	
15	cgt cca gtc ccg gat gct ctc aaa tcc ttg tgg tac aaa aac ctg act	951
	Arg Pro Val Pro Asp Ala Leu Lys Ser Leu Trp Tyr Lys Asn Leu Thr	
	220 225 230	
20	gga cct cag aac att act tta aac cac aca gac ctg gtt ccc tgc ctc	999
	Gly Pro Gln Asn Ile Thr Leu Asn His Thr Asp Leu Val Pro Cys Leu	
	235 240 245	
25	tgc att cag gtg tgg tcg cta gag cca gac tct gag agg gtc gaa ttc	1047
	Cys Ile Gln Val Trp Ser Leu Glu Pro Asp Ser Glu Arg Val Glu Phe	
	250 255 260	
30	tgc ccc ttc cgg gaa gat ccc ggt gca cac agg aac ctc tgg cac ata	1095
	Cys Pro Phe Arg Glu Asp Pro Gly Ala His Arg Asn Leu Trp His Ile	
	265 270 275	
35	gcc agg ctg cgg gta ctg tcc cca ggg gta tgg cag cta gat gcg cct	1143
	Ala Arg Leu Arg Val Leu Ser Pro Gly Val Trp Gln Leu Asp Ala Pro	
	280 285 290 295	
40	tgc tgt ctg ccg ggc aag gta aca ctg tgc tgg cag gca cca gac cag	1191
	Cys Cys Leu Pro Gly Lys Val Thr Leu Cys Trp Gln Ala Pro Asp Gln	
	300 305 310	
45	agt ccc tgc cag cca ctt gtg cca cca gtg ccc cag aag aac gcc act	1239
	Ser Pro Cys Gln Pro Leu Val Pro Pro Val Pro Gln Lys Asn Ala Thr	
	315 320 325	
50	gtg aat gag cca caa gat ttc cag ttg gtg gca ggc cac ccc aac ctc	1287
	Val Asn Glu Pro Gln Asp Phe Gln Leu Val Ala Gly His Pro Asn Leu	
	330 335 340	
55	tgt gtc cag gtg agc acc tgg gag aag gtt cag ctg caa gcg tgc ttg	1335
	Cys Val Gln Val Ser Thr Trp Glu Lys Val Gln Leu Gln Ala Cys Leu	
	345 350 355	
60	tgg gct gac tcc ttg ggg ccc ttc aag gat gat atg ctg tta gtg gag	1383
	Trp Ala Asp Ser Leu Gly Pro Phe Lys Asp Asp Met Leu Leu Val Glu	
	360 365 370 375	
65	atg aaa acc ggc ctc aac aac aca tca gtc tgt gcc ttg gaa ccc agt	1431
	Met Lys Thr Gly Leu Asn Asn Thr Ser Val Cys Ala Leu Glu Pro Ser	
	380 385 390	
70	ggc tgt aca cca ctg ccc agc atg gcc tcc acg aga gct gct cgc ctg	1479
	Gly Cys Thr Pro Leu Pro Ser Met Ala Ser Thr Arg Ala Ala Arg Leu	
	395 400 405	

5	gga gag gag ttg ctg caa gac ttc cga tca cac cag tgt atg cag ctg 1527													
	Gly Glu Glu Leu Leu Gln Asp Phe Arg Ser His Gln Cys Met Gln Leu	410				415					420			
10	tgg aac gat gac aac atg gga tgc cta tgg gcc tgc ccc atg gac aag 1575													
	Trp Asn Asp Asp Asn Met Gly Ser Leu Trp Ala Cys Pro Met Asp Lys	425				430				435				
15	tac atc cac agg cgc tgg gtc cta gta tgg ctg gcc tgc cta ctc ttg 1623													
	Tyr Ile His Arg Arg Trp Val Leu Val Trp Leu Ala Cys Leu Leu Leu	440				445				450				455
20	gct gcg gcg ctt ttc ttc ttc ctc ctt cta aaa aag gac cgc agg aaa 1671													
	Ala Ala Ala Leu Phe Phe Phe Leu Leu Leu Lys Lys Asp Arg Arg Lys		460					465					470	
25	gcg gcc cgt ggc tcc cgc acg gcc ttg ctc ctc cac tcc gcc gac gga 1719													
	Ala Ala Arg Gly Ser Arg Thr Ala Leu Leu Leu His Ser Ala Asp Gly		475					480				485		
30	gcg ggc tac gag cgc ctg gtg gga gca ctg gcg tcc gcg ttg agc cag 1767													
	Ala Gly Tyr Glu Arg Leu Val Gly Ala Leu Ala Ser Ala Leu Ser Gln		490				495					500		
35	atg cca ctg cgc gtg gcc gtg gac ctg tgg agc cgc cgc gag ctg agc 1815													
	Met Pro Leu Arg Val Ala Val Asp Leu Trp Ser Arg Arg Glu Leu Ser	505				510				515				
40	gcg cac gga gcc cta gcc tgg ttc cac cac cag cga cgc cgt atc ctg 1863													
	Ala His Gly Ala Leu Ala Trp Phe His His Gln Arg Arg Arg Ile Leu	520				525				530				535
45	cag gag ggt ggc gtg gta atc ctt ctc ttc tgc ccc gcg gcc gtg gcg 1911													
	Gln Glu Gly Gly Val Val Ile Leu Leu Phe Ser Pro Ala Ala Val Ala		540					545					550	
50	cag tgt cag cag tgg ctg cag ctc cag aca gtg gag ccc ggg ccg cat 1959													
	Gln Cys Gln Gln Trp Leu Gln Leu Gln Thr Val Glu Pro Gly Pro His		555					560				565		
55	gac gcc ctc gcc gcc tgg ctc agc tgc gtg cta ccc gat ttc ctg caa 2007													
	Asp Ala Leu Ala Ala Trp Leu Ser Cys Val Leu Pro Asp Phe Leu Gln		570				575				580			
60	ggc cgg gcg acc ggc cgc tac gtc ggg gtc tac ttc gac ggg ctg ctg 2055													
	Gly Arg Ala Thr Gly Arg Tyr Val Gly Val Tyr Phe Asp Gly Leu Leu	585				590				595				
65	cac cca gac tct gtg ccc tcc ccg ttc cgc gtc gcc ccg ctc ttc tcc 2103													
	His Pro Asp Ser Val Pro Ser Pro Phe Arg Val Ala Pro Leu Phe Ser	600				605				610				615
70	ctg ccc tgc cag ctg ccg gct ttc ctg gat gca ctg cag gga ggc tgc 2151													
	Leu Pro Ser Gln Leu Pro Ala Phe Leu Asp Ala Leu Gln Gly Gly Cys		620					625					630	

tcc act tcc gcg ggg cga ccc gcg gac cgg gtg gaa cga gtg acc cag 2199
 Ser Thr Ser Ala Gly Arg Pro Ala Asp Arg Val Glu Arg Val Thr Gln
 635 640 645
 5 gcg ctg cgg tcc gcc ctg gac agc tgt act tct agc tcg gaa gcc cca 2247
 Ala Leu Arg Ser Ala Leu Asp Ser Cys Thr Ser Ser Ser Glu Ala Pro
 650 655 660
 10 ggc tgc tgc gag gaa tgg gac ctg gga ccc tgc act aca cta gaa 2292
 Gly Cys Cys Glu Glu Trp Asp Leu Gly Pro Cys Thr Thr Leu Glu
 665 670 675
 taaaagccga tacagtattc ct 2314
 15 MPVSWFLLSLALGRNPVVVSLERLMEPQD TARCSLGLSCHLWDGDVLCPLPGSLQSA
 RCPQKTDCALCVRVVHLAVHGHWAEPPEEAGKSDSELQESRNASLQAQVVLSFQAYPIAR
 CALLEVQVPADLVQPGQSVGSAVFDCFEASLGAEVQIWSYTKPRYQKELNLTQQLPDCRGLE
 VRDSIQSCWVLPWLVNSTDGDNVLLTLDVSEEQDFSLLYLRPVPDALKSLWYKNLTGPQNI
 TLNHTDLVPCLCIQVWSLEPDSEVEFCPFREDPGAHRNLWHIARLRVLSPGVWQLDAP
 CCLPGKVTLWCQAPDQSPCQPLVPPVPQKNATVNEPQDFQLVAGHPNL CVQVSTWEKVQ
 LQACLWADSLGPFKDDMLLVEMKTGLNNTSVCALPSGCTPLPSMASTRAARLGEELL
 20 QDFRSHQCMQLWDDNMGSLWACPMDKYIHRRWVLVWLACLLAAALFFFLLLKKDRRKA
 ARGSR TALLLHSA DGAGYERLVGALASALSQMPLRVAVDLWSRRELSAHGALAWFHHQ
 RRRILQEGGVVILLFSPA AVAQCCQWLQLQTVEPGPHDALAAWLSCVLPDFLQGRATGRY
 VGVYFDGLLHPDSVPSPFRVAPLPSLPSQLPAFLDALQGGCSTSAGR PADRVERVTQAL
 RSALDSCTSSSEAPGCCEWDLGPCTTLE. 25

Reverse translation of rodent, e.g., mouse, DCRS7 (SEQ ID NO: 12):

atgccngtnw sntgggttyt nytnwsnytn gcnytnggm gnaayccngt ngtnngtnwsn 60
 30 ytnngarmgny tnatggarcc ncargayacn gcnmgntgyw snytnggny nwsntgycay 120
 ytnntgggag gngaygtnyt ntgyytnccn ggnwsnytn arwsngcnc nggncngtn 180
 35 ytngtncena cnmgnytnca racngarytn gtnytnmgnt gyccncaraa racngaytgy 240
 gcnytntgyg tnmngntngt ngtncayytn gcngtncayg gncaytgggc ngarccngar 300
 gargcnggna arwsngayws ngarytnear garwsnmgna aygnwsnytn ncargcncar 360
 40 gtngtnytnw snttycargc ntayccnath gcnmgntgyg cnytnytna rgtncargtn 420
 ccngcngayy tngtnearcc nggncarwsn gtnggnwsng cngtnttyga ytgyttygar 480
 45 gcnwsnytn gngcngargt ncarathtg wstnayacna arccnmngnta ycaraargar 540
 ytnaayytna cncarcaryt nccngaytgy mgnggnytn argtnmgnga ywsnathcar 600
 wstngytggg tnytnccntg gytnaaygtn wsnacngayg gngayaaygt nytnytnacn 660
 50 ytngaygtnw sngargarca rgayttywsn tyytnytn ayytnmgnc ngtnccngay 720
 gcnynaarw snyntggta yaaraayytn acnggncnc araayathac nytnaaycay 780
 55 acngayytn tncntgyt ntgyathcar gtntggwsny tngarccnga ywsngarmgn 840
 gtngarttyt gycenttymg ngargayccn ggngcncaym gnaayytn gcnayathgc 900
 mgnytnmgng tnytnwsncc ngngntntgg carytngayg cncntgyt ytnccnggn 960

aargtnacny tntgytggca rgcnccngay carwncnt gycarccnyt ngtnccnccn 1020
 5 gtnccncara araaygcna ngtnaaygar ccncargayt tycarytngt ngcnggncay 1080
 ccnaayytnt gygtncargt nwsnacntgg garaargtnc arytnccargc ntgyytntgg 1140
 gcngaywsny tnggncntt yaargaygay atgytntng tngaratgaa racnggnytn 1200
 10 aayaayacnw sngtntgygc nytngarccn wsggntgya cncnytncc nwsnatggcn 1260
 wsnacnmng cngcnmngyt nggngargar ytnytnccarg ayttymgnws ncaycartgy 1320
 atgcarytnt ggaaygayga yaayatgggn wsnynttggg cntgyccnat ggayaartay 1380
 15 athcaymngm gntgggtnyt ngtnnggytn gcntgyytny tnytnccngc ngcnytnntt 1440
 ttyttytny tnytnaaraa rgaymngmgn aargcngcnm gnggnwsnmg nacngcnytn 1500
 20 ytnytnccayw sngcngaygg ngcnggntay garmgnytn tnggngcnytn ngcnwsngcn 1560
 ytnwsncara tgccnytnmg ngtnccngtn gayytnngw smngnmngna rytwnsngcn 1620
 cayggngcny tngcntggtt ycaycaycar mngnmngmna thytncarga rggnggngtn 1680
 25 gtnathytny tnttywsncc ngcngcngtn gcncartgyc arcartggyt ncarytnccar 1740
 acngtnccarc cnggncncca ygaygcnytn gcngcntggy tnwsntgygt nytncngay 1800
 30 ttytnccarg gnmngcncac nggngntay gtnngngntnt ayttgyaygg nytnytnccay 1860
 ccngaywsng tncnwsncc nttymgngtn gcncnytn tywsnytncc nwsncarytn 1920
 ccngcnttyy tngaygcnytn ncargggngn tgywsnacnw sngcnggngm nccngcngay 1980
 35 mngntngarm gngtnacnca rgcnytnmgn wsgcnytn aywsntgyac nwsnwsnwsn 2040
 gargcncng gntgytgyga rgartgggay ytnngncnt gyacnacnytn ngar 2094

40 Table 3: Nucleotide and polypeptide sequences of DNAX Cytokine Receptor Subunit like
 embodiments (DCRS8). Primate, e.g., human, embodiment (see SEQ ID NO: 13 and 14).
 Predicted signal sequence indicated, but may vary by a few positions and depending upon cell
 type.

45 cccacgcntc cgggccagca gcgggcggcc ggggcgcaga gaacggcctg gctgggagag 60
 cgcacggcc atg gcc ccg tgg ctg cag ctc tgc tcc gtc ttc ttt acg gtc 111
 Met Ala Pro Trp Leu Gln Leu Cys Ser Val Phe Phe Thr Val
 50 -15 -10 -5
 aac gcc tgc ctc aac ggc tcg cag ctg gct gtn gcc gct ggc ggg tcc 159
 Asn Ala Cys Leu Asn Gly Ser Gln Leu Ala Xaa Ala Ala Gly Gly Ser
 -1 1 5 10
 55 ggc cgc gcg cng ggc gcc gac acc tgt agc tgg ang gga gtg ggg cca 207
 Gly Arg Ala Xaa Gly Ala Asp Thr Cys Ser Trp Xaa Gly Val Gly Pro
 15 20 25 30

5	gcc agc aga aac agt ggg ctg tac aac atc acc ttc aaa tat gac aat 255	
	Ala Ser Arg Asn Ser Gly Leu Tyr Asn Ile Thr Phe Lys Tyr Asp Asn	
	35 40 45	
10	tgt acc acc tac ttg aat cca gtg ggg aag cat gtg att gct gac gcc 303	
	Cys Thr Thr Tyr Leu Asn Pro Val Gly Lys His Val Ile Ala Asp Ala	
	50 55 60	
15	cag aat atc acc atc agc cag tat gct tgc cat gac caa gtg gca gtc 351	
	Gln Asn Ile Thr Ile Ser Gln Tyr Ala Cys His Asp Gln Val Ala Val	
	65 70 75	
20	acc att ctt tgg tcc cca ggg gcc ctc ggc atc gaa ttc ctg aaa gga 399	
	Thr Ile Leu Trp Ser Pro Gly Ala Leu Gly Ile Glu Phe Leu Lys Gly	
	80 85 90	
25	ttt cgg gta ata ctg gag gag ctg aag tcg gag gga aga cag ngc caa 447	
	Phe Arg Val Ile Leu Glu Glu Leu Lys Ser Glu Gly Arg Gln Xaa Gln	
	95 100 105 110	
30	caa ctg att cta aag gat ccg aag cag ntc aac agt agc ttc aaa aga 495	
	Gln Leu Ile Leu Lys Asp Pro Lys Gln Xaa Asn Ser Ser Phe Lys Arg	
	115 120 125	
35	act gga atg gaa tct caa cct ttn ctg aat atg aaa ttt gaa acg gat 543	
	Thr Gly Met Ser Gln Pro Xaa Leu Asn Met Lys Phe Glu Thr Asp	
	130 135 140	
40	tat ttc gta agg ttg tcc ttt tcc ttc att aaa aac gaa agc aat tac 591	
	Tyr Phe Val Arg Leu Ser Phe Ser Phe Ile Lys Asn Glu Ser Asn Tyr	
	145 150 155	
45	cac cct ttc ttc ttt aga acc cga gcc tgt gac ctg ttg tta cag ccg 639	
	His Pro Phe Phe Phe Arg Thr Arg Ala Cys Asp Leu Leu Leu Gln Pro	
	160 165 170	
50	gac aat cta gct tgt aaa ccc ttc tgg aag cct cgg aac ctg aac atc 687	
	Asp Asn Leu Ala Cys Lys Pro Phe Trp Lys Pro Arg Asn Leu Asn Ile	
	175 180 185 190	
55	agc cag cat ggc tcg gac atg cag gtg tcc ttc gac cac gca ccg cac 735	
	Ser Gln His Gly Ser Asp Met Gln Val Ser Phe Asp His Ala Pro His	
	195 200 205	
60	aac ttc ggc ttc cgt ttc ttc tat ctt cac tac aag ctc aag cac gaa 783	
	Asn Phe Gly Phe Arg Phe Phe Tyr Leu His Tyr Lys Leu Lys His Glu	
	210 215 220	
65	gga cct ttc aag cga aag acc tgt aag cag gag caa act aca gag atg 831	
	Gly Pro Phe Lys Arg Lys Thr Cys Lys Gln Glu Gln Thr Thr Glu Met	
	225 230 235	
70	acc agc tgc ctc ctt caa aat gtt tct cca ggg gat tat ata att gag 879	
	Thr Ser Cys Leu Leu Gln Asn Val Ser Pro Gly Asp Tyr Ile Ile Glu	
	240 245 250	

	ctg	gtg	gat	gac	act	aac	aca	aca	aga	aaa	gtg	atg	cat	tat	gcc	tta	927
	Leu	Val	Asp	Asp	Thr	Asn	Thr	Thr	Arg	Lys	Val	Met	His	Tyr	Ala	Leu	
	255					260					265					270	
5	aag	cca	gtg	cac	tcc	ccg	tgg	gcc	ggg	ccc	atc	aga	gcc	gtg	gcc	atc	975
	Lys	Pro	Val	His	Ser	Pro	Trp	Ala	Gly	Pro	Ile	Arg	Ala	Val	Ala	Ile	
					275					280					285		
10	aca	gtg	cca	ctg	gta	gtc	ata	tcg	gca	ttc	gcg	acg	ctc	ttc	act	gtg	1023
	Thr	Val	Pro	Leu	Val	Val	Ile	Ser	Ala	Phe	Ala	Thr	Leu	Phe	Thr	Val	
				290					295					300			
15	atg	tgc	cgc	aag	aag	caa	caa	gaa	aat	ata	tat	tca	cat	tta	gat	gaa	1071
	Met	Cys	Arg	Lys	Lys	Gln	Gln	Glu	Asn	Ile	Tyr	Ser	His	Leu	Asp	Glu	
			305					310					315				
20	gag	agc	tct	gag	tct	tcc	aca	tac	act	gca	gca	ctc	cca	aga	gag	agg	1119
	Glu	Ser	Ser	Glu	Ser	Ser	Thr	Tyr	Thr	Ala	Ala	Leu	Pro	Arg	Glu	Arg	
		320					325					330					
	ctc	cgg	ccg	cgg	ccg	aag	gtc	ttt	ctc	tgc	tat	tcc	agt	aaa	gat	ggc	1167
	Leu	Arg	Pro	Arg	Pro	Lys	Val	Phe	Leu	Cys	Tyr	Ser	Ser	Lys	Asp	Gly	
	335					340					345				350		
25	cag	aat	cac	atg	aat	gtc	gtc	cag	tgt	ttc	gcc	tac	ttc	ctc	cag	gac	1215
	Gln	Asn	His	Met	Asn	Val	Val	Gln	Cys	Phe	Ala	Tyr	Phe	Leu	Gln	Asp	
					355					360					365		
30	ttc	tgt	ggc	tgt	gag	gtg	gct	ctg	gac	ctg	tgg	gaa	gac	ttc	agc	ctc	1263
	Phe	Cys	Gly	Cys	Glu	Val	Ala	Leu	Asp	Leu	Trp	Glu	Asp	Phe	Ser	Leu	
				370				375						380			
35	tgt	aga	gaa	ggg	cag	aga	gaa	tgg	gtc	atc	cag	aag	atc	cac	gag	tcc	1311
	Cys	Arg	Glu	Gly	Gln	Arg	Glu	Trp	Val	Ile	Gln	Lys	Ile	His	Glu	Ser	
			385				390						395				
40	cag	ttc	atc	att	gtg	gtt	tgt	tcc	aaa	ggg	atg	aag	tac	ttt	gtg	gac	1359
	Gln	Phe	Ile	Ile	Val	Val	Cys	Ser	Lys	Gly	Met	Lys	Tyr	Phe	Val	Asp	
		400					405					410					
	aag	aag	aac	tac	aaa	cac	aaa	gga	ggg	ggc	cga	ggc	tcg	ggg	aaa	gga	1407
	Lys	Lys	Asn	Tyr	Lys	His	Lys	Gly	Gly	Gly	Arg	Gly	Ser	Gly	Lys	Gly	
	415					420					425				430		
45	gag	ctc	ttc	ctg	gtg	gcg	gtg	tca	gcc	att	gcc	gaa	aag	ctc	cgc	cag	1455
	Glu	Leu	Phe	Leu	Val	Ala	Val	Ser	Ala	Ile	Ala	Glu	Lys	Leu	Arg	Gln	
					435				440						445		
50	gcc	aag	cag	agt	tcg	tcc	gcg	gcg	ctc	agc	aag	ttt	atc	gcc	gtc	tac	1503
	Ala	Lys	Gln	Ser	Ser	Ser	Ala	Ala	Leu	Ser	Lys	Phe	Ile	Ala	Val	Tyr	
				450					455					460			
	ttt	gat	tat	tcc	tcg	gag	gga	gac	gtc	ccc	ggg	atc	cta	gac	ctg	agt	1551
	Phe	Asp	Tyr	Ser	Cys	Glu	Gly	Asp	Val	Pro	Gly	Ile	Leu	Asp	Leu	Ser	
			465					470					475				
55	acc	aag	tac	aga	ctc	atg	gac	aat	ctt	cct	cag	ctc	tgt	tcc	cac	ctg	1599
	Thr	Lys	Tyr	Arg	Leu	Met	Asp	Asn	Leu	Pro	Gln	Leu	Cys	Ser	His	Leu	
		480					485					490					

	cac tcc cga gac cac ggc ctc cag gag ccg ggg cag cac acg cga cag	1647
	His Ser Arg Asp His Gly Leu Gln Glu Pro Gly Gln His Thr Arg Gln	
	495 500 505 510	
5	ggc agc aga agg aac tac ttc cgg agc aag tca ggc cgg tcc cta tac	1695
	Gly Ser Arg Arg Asn Tyr Phe Arg Ser Lys Ser Gly Arg Ser Leu Tyr	
	515 520 525	
10	gtc gcc att tgc aac atg cac cag ttt att gac gag gag ccc gac tgg	1743
	Val Ala Ile Cys Asn Met His Gln Phe Ile Asp Glu Glu Pro Asp Trp	
	530 535 540	
15	ttc gaa aag cag ttc gtt ccc ttc cat cct cct cca ctg cgc tac cgg	1791
	Phe Glu Lys Gln Phe Val Pro Phe His Pro Pro Pro Leu Arg Tyr Arg	
	545 550 555	
20	gag cca gtc ttg gag aaa ttt gat tcg ggc ttg gtt tta aat gat gtc	1839
	Glu Pro Val Leu Glu Lys Phe Asp Ser Gly Leu Val Leu Asn Asp Val	
	560 565 570	
25	atg tgc aaa cca ggg cct gag agt gac ttc tgc cta aag gta gag gcg	1887
	Met Cys Lys Pro Gly Pro Glu Ser Asp Phe Cys Leu Lys Val Glu Ala	
	575 580 585 590	
30	gct gtt ctt ggg gca acc gga cca gcc gac tcc cag cac gag agt cag	1935
	Ala Val Leu Gly Ala Thr Gly Pro Ala Asp Ser Gln His Glu Ser Gln	
	595 600 605	
35	cat ggg ggc ctg gac caa gac ggg gag gcc cgg cct gcc ctt gac ggt	1983
	His Gly Gly Leu Asp Gln Asp Gly Glu Ala Arg Pro Ala Leu Asp Gly	
	610 615 620	
40	agc gcc gcc ctg caa ccc ctg ctg cac acg gtg aaa gcc ggc agc ccc	2031
	Ser Ala Ala Leu Gln Pro Leu Leu His Thr Val Lys Ala Gly Ser Pro	
	625 630 635	
45	tcg gac atg ccg cgg gac tca ggc atc tat gac tcg tct gtg ccc tca	2079
	Ser Asp Met Pro Arg Asp Ser Gly Ile Tyr Asp Ser Ser Val Pro Ser	
	640 645 650	
50	tcc gag ctg tct ctg cca ctg atg gaa gga ctc tcg acg gac cag aca	2127
	Ser Glu Leu Ser Leu Pro Leu Met Glu Gly Leu Ser Thr Asp Gln Thr	
	655 660 665 670	
55	gaa acg tct tcc ctg acg gag agc gtg tcc tcc tct tca ggc ctg ggt	2175
	Glu Thr Ser Ser Leu Thr Glu Ser Val Ser Ser Ser Ser Gly Leu Gly	
	675 680 685	
60	gag gag gaa cct cct gcc ctt cct tcc aag ctc ctc tct tct ggg tca	2223
	Glu Glu Glu Pro Pro Ala Leu Pro Ser Lys Leu Leu Ser Ser Gly Ser	
	690 695 700	
65	tgc aaa gca gat ctt ggt tgc cgc agc tac act gat gaa ctc cac gcg	2271
	Cys Lys Ala Asp Leu Gly Cys Arg Ser Tyr Thr Asp Glu Leu His Ala	
	705 710 715	
70	gtc gcc cct ttg taacaaaacg aaagagtcta agcattgccca ctttagctgc	2323
	Val Ala Pro Leu	
	720	

5 tgcctccctc tgattcccca gctcatctcc ctggttgcat ggcccacttg gagctgaggt 2383
 ctcatacaag gatatttgga gtgaaatgct ggccagtact tgttctccct tgccccaacc 2443
 ctttaccgga tatcttgaca aactctccaa ttttctaaaa tgatatggag ctctgaaagg 2503
 catgtccata aggtctgaca acagcttgcc aaatttggtt agtccttgga tcagagcctg 2563
 10 ttgtgggagg tagggaggaa atatgtaaaag aaaaacagga agatacctgc actaatcatt 2623
 cagacttcat tgagctctgc aaactttgcc tgtttgctat tggctacctt gatttgaaat 2683
 gctttgtgaa aaaaggcact tttaacatca tagccacaga aatcaagtgc cagtctatct 2743
 15 ggaatccatg ttgtattgca gataatgttc tcatttattt ttg 2786

MAPWLQLCSVFFTVNACLNGSQLAVAAGGSGRAXGADTCSWXGVGPASRNSGLYNITPKYDNCTTYLNPVGK
 20 HVIADAQNITISQYACHDQVAVTILWSPGALGIEFLKGFRVILEELKSEGRQXQQLILKDPKQXNSSFKRTG
 MESQPxLNMKFETDYFVRLSFSFIKNESNYHPPFFRTRACDLLLPDNLACKPFWKPRNLNISQHGSDMQVS
 FDHAPHNFGFRFFYLHYKLKHEGPFKRKTCKQEQTTEMTSCLLQNVSPGDYIIELVDDTNTTRKVMHYALKP
 VHSPWAGPIRAVAITVPLVVISAFATLFTVMCRKKQENIYSHLDEESSESSTYTAALPRERLRPRPKVFLC
 YSSKDGQNMNVVQCFAFLQDFCGCEVALDLWEDFSLCREGQREWVIQKIHSQFIIIVVCSKGMKYFVDK
 25 NYKHKGGGRSGSGKELFLVAVSAIAEKLQAKQSSSAALSKFIAVYFDYSCGDVPGILDSTKYRLMDNLP
 QLCSHLSRDRHGLQEPGQHTROGSRNRYFRSKSGRSLYVAICNMHQFIDEEDWFEKQVFPFHPPPLRYREP
 VLEKFDGLVNDVMCKPGPESDFCLKVEAAVLGATGPADSQHESQHGGLDQDGEARPALDGSAAALQPLLHT
 VKAGSPDMPRDSGIYDSSVPSSLSLPLMEGLSTDQTETSSLTESVSSSSGLGEEPPALPSKLLSSGSCK
 ADLGCRSYTDELHAVAPL.

30 Reverse translation of primate, e.g., human, DCRS8 (SEQ ID NO: 15):
 atggcnccnt ggytncaryt ntgywsngtn ttyttyacng tnaaygcntg yytnaayggn 60
 35 wsnrcarytng cngtngcngc ngngngnwns ggnmgngcnn nngngngnga yacntgywsn 120
 tggnnngngng tngngcngc nwsnmgnaay wsngngnynt ayaayathac nttyaartay 180
 gayaaytgga cnacntayyt naayccngtn ggnaarcayg tnathgnga ygcncaraay 240
 40 athacnathw sncartaygc ntgycaygay cargtngcng tnacnathyt ntggwsnccn 300
 ggngcnytn gnatggartt yytnaarggn ttmngngtna thytngarga rytnaarwsn 360
 45 gargngmgnc arnnncarca rytnathytn aargayccna arcarnnnaa ywsnwsntty 420
 aarmgnacng gnatggarws ncarccnnnn ytnaayatga arttygarac ngaytaytty 480
 gtnmgnytnw snttywsntt yathaaraay garwsnaayt aycayccntt yttyttymgn 540
 50 acnmngngent gygayytnyt nytnrcarccn gayaayytn cngtyaarcc nttytggaar 600
 ccnmgnaayy tnaayathws ncarcayggn wsngayatgc argtnwsntt ygaycaygn 660
 55 ccncayaayt tyggnttymg nttyttytay ytncaytaya arytnaarca ygargngccn 720
 ttyaarmgna aracntgyaa rcargarcar acnacngara tgacnwsntg yytnytnar 780
 aaygtnwsnc cngnggayta yathathgar ytngtngayg ayacnaayac nacnmgnaar 840

gtnatgcayt aygcnytnaa rccngtncay wsnccntggg cnggnccnat hmgngcngtn 900
 5 gcnathacng tncnnytngt ngtnathwsn gcnttygcna cnytnntyac ngtnatgtgy 960
 mgnaaraarc arcargaraa yathtaywsn cayytn gayg argarwsnws ngarwsnwsn 1020
 acntayacng cngcnytncc nmngngarmgn ytnmgncnm gncnaargt nttyytnatgy 1080
 10 taywsnwsna argayggna raaycaytg aaygtngtnc artgytgyt ntayttyytn 1140
 cargayttyt gyggntgyga rgtngcnytn gayytn tggg argaytgyws nytnngymgn 1200
 garggncarm gngartgggt nathcaraar athcaygarw sncarttyat hathgtngtn 1260
 15 tgywsnaarg gnatgaarta yttygtngay aaraaraayt ayaarcayaa rggngggngn 1320
 mgnggnwsng gnaarggnga rytnttyytn gtngcngtnw sngcnathgc ngaraarytn 1380
 20 mgncargcna arcarwsnws nwsngcngcn ytnwsnaart tyathgcngt ntaytgyay 1440
 taywsntgyg arggngaygt nccnggnath ytngayytnw snacnaarta ymgnytnatg 1500
 gayaayytn cncarytn tgywsncayytn caywsnmngn aycayggnyt ncargarccn 1560
 25 ggncarcaya cnmngncargg nwsnmngmgn aaytaytym gnwsnaarws nggnmgwnwsn 1620
 ytntaygtng cnathgyaa yatgcaycar ttyathgayg argarccnga ytggttygar 1680
 30 aarcarttyg tncnttyca yccnccnccn ytnmgntaym gngarccngt nytnagaraar 1740
 ttygaywsng gnytngtnyt naaygaygt atgtgyaarc cnggnccnga rwsngaytty 1800
 35 tgyytnaarg tngargcngc ngtnytnngn gcnacnggnc cngcngayws ncarcaygar 1860
 wsnarcayg gnggnytna ycargaygg gargcnmgnc cngcnytna yggwnsngcn 1920
 gcnytnarc cnytnytnca yacngtnaar gcnggnwsnc cnwsngayat gccnmngay 1980
 40 wsggnatht aygaywsnws ngtnccnwsn wsgarytnw snytnccnyt natggarggn 2040
 ytnwsnacng aycaracnga racnwsnwsn ytnacngarw sngtnwsnws nwsnwsngn 2100
 ytnggngarg argarccncc ngcnytnccn wsnaarytny tnwsnwsngg nwsntgyaar 2160
 45 gcngayytn gntgymgnws ntayacngay garytn cayg cngtngcncc nytn 2214

50 Table 4: Nucleotide and polypeptide sequences of DNAX Cytokine Receptor Subunit like
 embodiments (DCRS9). Primate, e.g., human, embodiment (see SEQ ID NO: 16 and 17).
 Predicted signal sequence indicated, but may vary by a few positions and depending upon cell
 type.

55 atg ggg agc tcc aga ctg gca gcc ctg ctc ctg cct ctc ctc ctc ata 48
 Met Gly Ser Ser Arg Leu Ala Ala Leu Leu Leu Pro Leu Leu Leu Ile
 -20 -15 -10

27.

	gtc atc gac ctc tct gac tct gct ggg att ggc ttt cgc cac ctg ccc	96
	Val Ile Asp Leu Ser Asp Ser Ala Gly Ile Gly Phe Arg His Leu Pro	
	-5 -1 1 5	
5	cac tgg aac acc cgc tgt cct ctg gcc tcc cac acg gaa gtt ctg cct	144
	His Trp Asn Thr Arg Cys Pro Leu Ala Ser His Thr Glu Val Leu Pro	
	10 15 20 25	
10	ata tcc ctt gcc gca cct ggt ggg ccc tct tct cca caa agc ctt ggt	192
	Ile Ser Leu Ala Ala Pro Gly Gly Pro Ser Ser Pro Gln Ser Leu Gly	
	30 35 40	
15	gtg tgc gag tct ggc act gtt ccc gct gtt tgt gcc agc atc tgc tgt	240
	Val Cys Glu Ser Gly Thr Val Pro Ala Val Cys Ala Ser Ile Cys Cys	
	45 50 55	
20	cag gtg gct cag gtc ttc aac ggg gcc tct tcc acc tcc tgg tgc aga	288
	Gln Val Ala Gln Val Phe Asn Gly Ala Ser Ser Thr Ser Trp Cys Arg	
	60 65 70	
25	aat cca aaa agt ctt cca cat tca agt tct ata gga gac aca aga tgc	336
	Asn Pro Lys Ser Leu Pro His Ser Ser Ser Ile Gly Asp Thr Arg Cys	
	75 80 85	
30	cag cac ctg ctc aga gga agc tgc tgc ctc gtc gtc acc tgt ctg aga	384
	Gln His Leu Leu Arg Gly Ser Cys Cys Leu Val Val Thr Cys Leu Arg	
	90 95 100 105	
35	aga gcc atc aca ttt cca tcc cct ccc cag aca tct ccc aca agg gac	432
	Arg Ala Ile Thr Phe Pro Ser Pro Pro Gln Thr Ser Pro Thr Arg Asp	
	110 115 120	
40	ttc gct cta aaa gga ccc aac ctt cgg atc cag aga cat ggg aaa gtc	480
	Phe Ala Leu Lys Gly Pro Asn Leu Arg Ile Gln Arg His Gly Lys Val	
	125 130 135	
45	ttc cca gat tgg act cac aaa ggc atg gag gtg ggc act ggg tac aac	528
	Phe Pro Asp Trp Thr His Lys Gly Met Glu Val Gly Thr Gly Tyr Asn	
	140 145 150	
50	agg aga tgg gtt cag ctg agt ggt gga ccc gag ttc tcc ttt gat ttg	576
	Arg Arg Trp Val Gln Leu Ser Gly Gly Pro Glu Phe Ser Phe Asp Leu	
	155 160 165	
55	ctg cct gag gcc cgg gct att cgg gtg acc ata tct tca ggc cct gag	624
	Leu Pro Glu Ala Arg Ala Ile Arg Val Thr Ile Ser Ser Gly Pro Glu	
	170 175 180 185	
60	gtc agc gtg cgt ctt tgt cac cag tgg gca ctg gag tgt gaa gag ctg	672
	Val Ser Val Arg Leu Cys His Gln Trp Ala Leu Glu Cys Glu Glu Leu	
	190 195 200	
65	agc agt ccc tat gat gtc cag aaa att gtg tct ggg ggc cac act gta	720
	Ser Ser Pro Tyr Asp Val Gln Lys Ile Val Ser Gly Gly His Thr Val	
	205 210 215	
70	gag ctg cct tat gaa ttc ctt ctg ccc tgt ctg tgc ata gag gca tcc	768
	Glu Leu Pro Tyr Glu Phe Leu Leu Pro Cys Leu Cys Ile Glu Ala Ser	
	220 225 230	

5	tac	ctg	caa	gag	gac	act	gtg	agg	cgc	aaa	aaa	tgt	ccc	ttc	cag	agc	816
	Tyr	Leu	Gln	Glu	Asp	Thr	Val	Arg	Arg	Lys	Lys	Cys	Pro	Phe	Gln	Ser	
	235						240					245					
10	tgg	cca	gaa	gcc	tat	ggc	tcg	gac	ttc	tgg	aag	tca	gtg	cac	ttc	act	864
	Trp	Pro	Glu	Ala	Tyr	Gly	Ser	Asp	Phe	Trp	Lys	Ser	Val	His	Phe	Thr	
	250					255					260					265	
15	gac	tac	agc	cag	cac	act	cag	atg	gtc	atg	gcc	ctg	aca	ctc	cgc	tgc	912
	Asp	Tyr	Ser	Gln	His	Thr	Gln	Met	Val	Met	Ala	Leu	Thr	Leu	Arg	Cys	
					270					275					280		
20	cca	ctg	aag	ctg	gaa	gct	gcc	ctc	tgc	cag	agg	cac	gac	tgg	cat	acc	960
	Pro	Leu	Lys	Leu	Glu	Ala	Ala	Leu	Cys	Gln	Arg	His	Asp	Trp	His	Thr	
				285					290					295			
25	ctt	tgc	aaa	gac	ctc	ccg	aat	gcc	acg	gct	cga	gag	tca	gat	ggg	tgg	1008
	Leu	Cys	Lys	Asp	Leu	Pro	Asn	Ala	Thr	Ala	Arg	Glu	Ser	Asp	Gly	Trp	
			300					305					310				
30	tat	gtt	ttg	gag	aag	gtg	gac	ctg	cac	ccc	cag	ctc	tgc	ttc	aag	gta	1056
	Tyr	Val	Leu	Glu	Lys	Val	Asp	Leu	His	Pro	Gln	Leu	Cys	Phe	Lys	Val	
	315						320					325					
35	caa	cca	tgg	ttc	tct	ttt	gga	aac	agc	agc	cat	gtt	gaa	tgc	ccc	cac	1104
	Gln	Pro	Trp	Phe	Ser	Phe	Gly	Asn	Ser	Ser	His	Val	Glu	Cys	Pro	His	
	330					335					340				345		
40	cag	act	ggg	tct	ctc	aca	tcc	tgg	aat	gta	agc	atg	gat	acc	caa	gcc	1152
	Gln	Thr	Gly	Ser	Leu	Thr	Ser	Trp	Asn	Val	Ser	Met	Asp	Thr	Gln	Ala	
					350				355						360		
45	cag	cag	ctg	att	ctt	cac	ttc	tcc	tca	aga	atg	cat	gcc	acc	ttc	agt	1200
	Gln	Gln	Leu	Ile	Leu	His	Phe	Ser	Ser	Arg	Met	His	Ala	Thr	Phe	Ser	
				365					370					375			
50	gct	gcc	tgg	agc	ctc	cca	ggc	ttg	ggg	cag	gac	act	ttg	gtg	ccc	ccc	1248
	Ala	Ala	Trp	Ser	Leu	Pro	Gly	Leu	Gly	Gln	Asp	Thr	Leu	Val	Pro	Pro	
			380				385						390				
55	gtg	tac	act	gtc	agc	cag	gtg	tgg	cgg	tca	gat	gtc	cag	ttt	gcc	tgg	1296
	Val	Tyr	Thr	Val	Ser	Gln	Val	Trp	Arg	Ser	Asp	Val	Gln	Phe	Ala	Trp	
	395						400					405					
60	aag	cac	ctc	ttg	tgt	cca	gat	gtc	tct	tac	aga	cac	ctg	ggg	ctc	ttg	1344
	Lys	His	Leu	Leu	Cys	Pro	Asp	Val	Ser	Tyr	Arg	His	Leu	Gly	Leu	Leu	
	410					415					420				425		
65	atc	ctg	gca	ctg	ctg	gcc	ctc	ctc	acc	cta	ctg	ggg	gtt	gtt	ctg	gcc	1392
	Ile	Leu	Ala	Leu	Leu	Ala	Leu	Leu	Thr	Leu	Leu	Gly	Val	Val	Leu	Ala	
					430					435					440		
70	ctc	acc	tgc	cgg	cgc	cca	cag	tca	ggc	ccg	ggc	cca	gcg	cgg	cca	gtg	1440
	Leu	Thr	Cys	Arg	Arg	Pro	Gln	Ser	Gly	Pro	Gly	Pro	Ala	Arg	Pro	Val	
				445					450					455			

	ctc ctc ctg cac gcg gcg gac tcg gag gcg cag cgg cgc ctg gtg gga	1488
	Leu Leu Leu His Ala Ala Asp Ser Glu Ala Gln Arg Arg Leu Val Gly	
	460 465 470	
5	gcg ctg gct gaa ctg cta cgg gca gcg ctg ggc ggc ggg cgc gac gtg	1536
	Ala Leu Ala Glu Leu Leu Arg Ala Ala Leu Gly Gly Gly Arg Asp Val	
	475 480 485	
10	atc gtg gac ctg tgg gag ggg agg cac gtg gcg cgc gtg ggc ccg ctg	1584
	Ile Val Asp Leu Trp Glu Gly Arg His Val Ala Arg Val Gly Pro Leu	
	490 495 500 505	
15	ccg tgg ctc tgg gcg gcg cgg acg cgc gta gcg cgg gag cag ggc act	1632
	Pro Trp Leu Trp Ala Ala Arg Thr Arg Val Ala Arg Glu Gln Gly Thr	
	510 515 520	
20	gtg ctg ctg ctg tgg agc ggc gcc gac ctt cgc ccg gtc agc ggc ccc	1680
	Val Leu Leu Leu Trp Ser Gly Ala Asp Leu Arg Pro Val Ser Gly Pro	
	525 530 535	
	gac ccc cgc gcc gcg ccc ctg ctc gcc ctg ctc cac gct gcc ccg cgc	1728
	Asp Pro Arg Ala Ala Pro Leu Leu Ala Leu Leu His Ala Ala Pro Arg	
	540 545 550	
25	ccg ctg ctg ctg ctc gct tac ttc agt cgc ctc tgc gcc aag ggc gac	1776
	Pro Leu Leu Leu Leu Ala Tyr Phe Ser Arg Leu Cys Ala Lys Gly Asp	
	555 560 565	
30	atc ccc ccg ccg ctg cgc gcc ctg ccg cgc tac cgc ctg ctg cgc gac	1824
	Ile Pro Pro Pro Leu Arg Ala Leu Pro Arg Tyr Arg Leu Leu Arg Asp	
	570 575 580 585	
35	ctg ccg cgt ctg ctg cgg gcg ctg gac gcg cgg cct ttc gca gag gcc	1872
	Leu Pro Arg Leu Leu Arg Ala Leu Asp Ala Arg Pro Phe Ala Glu Ala	
	590 595 600	
40	acc agc tgg ggc cgc ctt ggg gcg cgg cag cgc agg cag agc cgc cta	1920
	Thr Ser Trp Gly Arg Leu Gly Ala Arg Gln Arg Arg Gln Ser Arg Leu	
	605 610 615	
	gag ctg tgc agc cgg ctc gaa cga gag gcc gcc cga ctt gca gac cta	1968
	Glu Leu Cys Ser Arg Leu Glu Arg Glu Ala Ala Arg Leu Ala Asp Leu	
	620 625 630	
45	ggt tgagcagagc tccaccgcag tcccgggtgt ctgcggccgc t	2012
	Gly	
50	MGSSRLAALLPLLLIVIDLSDSAGIGFRHLPHWNTRCPLASHTEVLPISLAAPGGPSSPQSLGVCESGTVF	
	AVCASICQVAQVFNGASSTSWCRNPKSLPHSSSIGDTRCQHLLRGSCCLVVTCLRRAITFPSPPTSPTRD	
	FALKGPNLRIQRHGKVFDPDWITHKGMEVGTGYNRRWVQLSGGPEFSFDLLPEARAIRVTISSGPEVSVRLCHQ	
	WALECEELSSPYDVQKIVSGGHTVELPYEFLLPCLCIEASYLQEDTVRRKKCPFSWPEAYGSDFWKSVHFT	
	DYSQHTQMVMALTLRCPLKLEAALCQRHDWHTLCKDLPNATARESDGWYVLEKVDLHPQLCFKVQPWFSFGN	
	SSHVECPHQTGSLSWNVSMQTQAQQLILHFSSRMHATFSAAWSLPGLGQDTLVPPVYTVSVQVWRSVDQFAW	
	KHLLCPDVSYRHLGLLILALLALLTLTGVLALTCCRPGSGPGPARPVLLLHAADSEAQRRLVGALAEELLRA	
55	ALGGGRDVIVDLWEGRHVARVGPLPWLWAARTRVAREQGTVLLWLGADLRPVSGPDPRAPLLALLHAAPR	
	PLLLLAYFSRLCAKGDIPPLRALPRYRLRLDLPRLLRALDARPF AEATSWGRLGARQRRQSRLELCSRLER	
	EAARLADLG.	

Reverse translation of primate, e.g., human, DCRS9 (SEQ ID NO: 18):

5 atgggnwsnw snmgnytn gc ngenytnytn ytnccnytny tnytnathgt nathgayytn 60
 wsgaywsng cnggnathgg nttymgncay ytnccncayt ggaayacnmg ntgyccnytn 120
 gcnwsncaya cngargtnyt nccnathwsn ytn gcngcnc cnggnggncc nwsnwsnccn 180
 10 carwsnytn gngtntgyga rwsnggnacn gtncgcngcng tntgygcnws nathtgytgy 240
 cargtn gcnc argtnttyaa yggngcnwsn wsnacnwsnt ggtgymgnaa yccnaarwsn 300
 ytnccncayw snwsnwsnat hggngayacn mgntgy carc ayytnytnmg nggnwsntgy 360
 15 tgyytn gtn tnaentgyt nmgnmgngcn athacnttyc cnwsnccncc ncaracnwsn 420
 ccnacnmng aytygcny naarggnccn ayytnmgna thcarmgna yggnaargtn 480
 20 ttyccngayt ggacncayaa rggnatggar gtnggnacng gntayaaymg nmngtgggt 540
 carytnwsng gnggnccnga rtywsntty gayytnytn cngargcnmg ngcnathmgn 600
 gtnacnathw snwsnggncc ngargtnwsn gtnmgnytn gycaycartg ggcnytn gar 660
 25 tgygargary tnwsnwsncc ntaygaygn caraarathg tnwsngngng ncayacngtn 720
 garytnccnt aygarttyt nytnccntgy ytn tgyathg argcnwsnta yytn cargar 780
 30 gayacngtnm gnmgnaaraa rtgyccntty carwsntggc cngargenta yggnwsngay 840
 ttytggaarw sngtn cayt yacngaytay wsnarcaya cncaratggt natggcnytn 900
 acnytnmgnt gyccnytnaa rytngargcn gcnytn tgyc armgncayga ytggcayacn 960
 35 ytn tgyaarg ayytnccnaa ygcncngcn mgngarwsng ayggntggta ygtnytn gar 1020
 aargtn gayy tncayccna rytntgytty aargtn carc cntggtyws nttyggnaay 1080
 40 wsnwsncayg tngartgycc ncaycaracn ggnwsnytna cnwsntggaa ygtnwsnatg 1140
 gayacncarg cncarcaryt nathytn cay ttwsnwsnm gnatgcaygc nacnttywsn 1200
 gcngcntggw snytnccngg nytn ggn car gayacnytn tncncngt ntayacngtn 1260
 45 wsn cargtn ggmgnwsnga ygtncartty gcntggaarc ayytnytn tgy ccngaygn 1320
 wsntaymgnc ayytn gny nytnathytn gcnytnytn cnytnytnac nytnytn ggn 1380
 50 gtngtnytn gnytnacntg ymgngmgnccn carwsnggnc cnggnccngc nmgnccngtn 1440
 ytnytnytn aygcngcnga ywsngargcn carmgmgn ytn gngngc nytn gngar 1500
 ytnytnmgng cngcnytn ggnggngmgn gaygtnathg tngayytn tgy ggarggngmgn 1560
 55 caygtngcnm gngtnggncc nytnccntgg ytn tggcng cnmgncnm ngtn gcnmgn 1620
 garcarggna cngtnytn nytn tggwsn ggngcngayy tnmgnccngt nwsnggnccn 1680

gayccnmngng cngcncnyt nytnngcnytn ytncaygeng cncnmgnc nytnytnytn 1740
 ytnngcntayt tywsnmgnyt ntgygcnaar ggngayathc cncncncnyt nmngncnytn 1800
 5 ccnmgntaym gnytnytnmg ngayytnccn mgnytnytnm gngcnytnga ygcnmgnccn 1860
 ttygcngarg cnacnwsntg gggnmgnytn ggngcngmc armgnmgnc rwsnmgnytn 1920
 garytntgyw snmgnytnga rmngngargcn gcnmgnytn cngayytnng n 1971
 10

Rodent, e.g., mouse, embodiment (see SEQ ID NO: 19 and 20). Predicted signal sequence indicated, but may vary by a few positions and depending upon cell type.

15 cagctccggg ccaggccctg ctgccctctt gcagacagga aagacatggt ctctgcgccc 60
 tgatcctaca gaagctc atg ggg agc ccc aga ctg gca gcc ttg ctc ctg 110
 Met Gly Ser Pro Arg Leu Ala Ala Leu Leu Leu
 -20 -15
 20 tct ctc ccg cta ctg ctc atc ggc ctc gct gtg tct gct cgg gtt gcc 158
 Ser Leu Pro Leu Leu Leu Ile Gly Leu Ala Val Ser Ala Arg Val Ala
 -10 -5 -1 1
 25 tgc ccc tgc ctg cgg agt tgg acc agc cac tgt ctc ctg gcc tac cgt 206
 Cys Pro Cys Leu Arg Ser Trp Thr Ser His Cys Leu Leu Ala Tyr Arg
 5 10 15 20
 30 gtg gat aaa cgt ttt gct ggc ctt cag tgg ggc tgg ttc cct ctc ttg 254
 Val Asp Lys Arg Phe Ala Gly Leu Gln Trp Gly Trp Phe Pro Leu Leu
 25 30 35
 35 gtg agg aaa tct aaa agt cct cct aaa ttt gaa gac tat tgg agg cac 302
 Val Arg Lys Ser Lys Ser Pro Pro Lys Phe Glu Asp Tyr Trp Arg His
 40 45 50
 40 agg aca cca gca tcc ttc cag agg aag ctg cta ggc agc cct tcc ctg 350
 Arg Thr Pro Ala Ser Phe Gln Arg Lys Leu Leu Gly Ser Pro Ser Leu
 55 60 65
 45 tct gag gaa agc cat cga att tcc atc ccc tcc tca gcc atc tcc cac 398
 Ser Glu Glu Ser His Arg Ile Ser Ile Pro Ser Ser Ala Ile Ser His
 70 75 80
 50 aga ggc caa cgc acc aaa agg gcc cag cct tca gct gca gaa gga aga 446
 Arg Gly Gln Arg Thr Lys Arg Ala Gln Pro Ser Ala Ala Glu Gly Arg
 85 90 95 100
 55 gaa cat ctc cct gaa gca ggg tca caa aag tgt gga gga cct gaa ttc 494
 Glu His Leu Pro Glu Ala Gly Ser Gln Lys Cys Gly Gly Pro Glu Phe
 105 110 115
 tcc ttt gat ttg ctg ccc gag gtg cag gct gtt cgg gtg act att cct 542
 Ser Phe Asp Leu Leu Pro Glu Val Gln Ala Val Arg Val Thr Ile Pro
 120 125 130

	gca ggc ccc aag gca cgt gtg cgc ctt tgt tat cag tgg gca ctg gaa	590
	Ala Gly Pro Lys Ala Arg Val Arg Leu Cys Tyr Gln Trp Ala Leu Glu	
	135 140 145	
5	tgt gaa gac ttg agt agc cct ttt gat acc cag aaa att gtg tct gga	638
	Cys Glu Asp Leu Ser Ser Pro Phe Asp Thr Gln Lys Ile Val Ser Gly	
	150 155 160	
10	ggg cac act gta gac ctg cct tat gaa ttc ctt ctg ccc tgc atg tgc	686
	Gly His Thr Val Asp Leu Pro Tyr Glu Phe Leu Leu Pro Cys Met Cys	
	165 170 175 180	
15	ata gag gcc tcc tac ctg caa gag gac act gtg agg cgc aaa agt gtc	734
	Ile Glu Ala Ser Tyr Leu Gln Glu Asp Thr Val Arg Arg Lys Ser Val	
	185 190 195	
20	cct tcc aga gct ggc ctg aag ctt atg gct cag act tct ggc agt caa	782
	Pro Ser Arg Ala Gly Leu Lys Leu Met Ala Gln Thr Ser Gly Ser Gln	
	200 205 210	
	tac gct tca ctg act aca gcc agc ac	808
	Tyr Ala Ser Leu Thr Thr Ala Ser	
	215 220	
25	MGSPRLAALLLSLPLLLIGLAVSARVACPCLRSWTSCHLLAYRVDKRFAGLQGWGFPLLVRKSKSPPKFEDY WRHRTPASFQRKLLGSPSLSEESHRIIPSSAISHRGQRTKRAQPSAAEGREHLPEAGSQKCGGPEFSFDLL PEVQAVRVTIPAGPKARVRLCYQWALECEDLSSPFDTKIVSGGHTVDLPYEFLLPCMCIEASYLQEDTVRR KSVPSRAGLKLMAQTSGSQYASLTTAS	
30	Reverse translation of rodent, e.g., mouse, DCRS9 (SEQ ID NO: 21):	
	atgggnwsnc cnmgnytngc ngcnynytyn ytnwsnytnc cnytnytnyt nathggnytn	60
35	gcngtnwsng cnmgngtngc ntgyccntgy ytnmgwsnt ggacnwsnca ytgyytnytyn	120
	gcntaymgng tngayaarmg nttygcnggn ytncartggg gntgggttycc nytnytnngtn	180
40	mgnaarwsna arwsncchcc naarttygar gaytaytggm gncaymgna nccngcnwsn	240
	ttycarmgna arytnytnng nwsnccnwsn ytnwsngarg arwsncaymg nathwsnath	300
	ccnwsnwsng cnathwsnca ymgnggncar mgnaacnaarm gngcncarcc nwsngcngcn	360
45	garggnmgng arcayytnc ngargcnggn wscaraart gyggnggngcc ngarttywsn	420
	ttygayytyn tncngargt ncargcngtn mgngtnacna thcngcngg nccnaargcn	480
50	mgngtnmgny tntgytayca rtgggcnytn gartgygarg ayytnwsnws nccnttygay	540
	acncaraara thgtnwsngg nggncayacn gtngayytnc cntaygartt yytnytnccn	600
	tgyatgtgya thgargcnws ntayytncar gargayacng tnmgnmgnaa rwsngtnccn	660
55	wsnmgngcng gnytnaaryt natggcncar acnwsnggnw sncartaygc nwsnytnacn	720
	acngcnwsn	729

Table 5: Nucleotide and polypeptide sequences of DNAX Cytokine Receptor Subunit like embodiments (DCRS10). Primate, e.g., human, embodiment (see SEQ ID NO: 22 and 23).

5	ttttgagcag aggcttccta ggctccgtag aaatttgcat acagcttcca cttcctgctt 60	
	cagagcctgt tcttctactt acctgggccc ggagaagggtg gagggagacg agaagccgcc 120	
10	gagagccgac taccctccgg gccagctctg tctgtccgtg gtggatctaa gaaactaga 179	
	atg aac cga agc att cct gtg gag gtt gat gaa tca gaa cca tac cca 227	
	Met Asn Arg Ser Ile Pro Val Glu Val Asp Glu Ser Glu Pro Tyr Pro	
	1 5 10 15	
15	agt cag ttg ctg aaa cca atc cca gaa tat tcc ccg gaa gag gaa tca 275	
	Ser Gln Leu Leu Lys Pro Ile Pro Glu Tyr Ser Pro Glu Glu Glu Ser	
	20 25 30	
20	gaa cca cct gct cca aat ata agg aac atg gca ccc aac agc ttg tct 323	
	Glu Pro Pro Ala Pro Asn Ile Arg Asn Met Ala Pro Asn Ser Leu Ser	
	35 40 45	
25	gca ccc aca atg ctt cac aat tcc tcc gga gac ttt tct caa gct cac 371	
	Ala Pro Thr Met Leu His Asn Ser Ser Gly Asp Phe Ser Gln Ala His	
	50 55 60	
30	tca acc ctg aaa ctt gca aat cac cag cgg cct gta tcc cgg cag gtc 419	
	Ser Thr Leu Lys Leu Ala Asn His Gln Arg Pro Val Ser Arg Gln Val	
	65 70 75 80	
35	acc tgc ctg cgc act caa gtt ctg gag gac agt gaa gac agt ttc tgc 467	
	Thr Cys Leu Arg Thr Gln Val Leu Glu Asp Ser Glu Asp Ser Phe Cys	
	85 90 95	
40	agg aga cac cca ggc ctg ggc aaa gct ttc cct tct ggg tgc tct gca 515	
	Arg Arg His Pro Gly Leu Gly Lys Ala Phe Pro Ser Gly Cys Ser Ala	
	100 105 110	
45	gtc agc gag cct gcg tct gag tct gtg gtt gga gcc ctc cct gca gag 563	
	Val Ser Glu Pro Ala Ser Glu Ser Val Val Gly Ala Leu Pro Ala Glu	
	115 120 125	
50	cat cag ttt tca ttt atg gaa aaa cgt aat caa tgg ctg gta tct cag 611	
	His Gln Phe Ser Phe Met Glu Lys Arg Asn Gln Trp Leu Val Ser Gln	
	130 135 140	
55	ctt tca gcg gct tct cct gac act ggc cat gac tca gac aaa tca gac 659	
	Leu Ser Ala Ala Ser Pro Asp Thr Gly His Asp Ser Asp Lys Ser Asp	
	145 150 155 160	
60	caa agt tta cct aat gcc tca gca gac tcc ttg ggc ggt agc cag gag 707	
	Gln Ser Leu Pro Asn Ala Ser Ala Asp Ser Leu Gly Gly Ser Gln Glu	
	165 170 175	
65	atg gtg caa cgg ccc cag cct cac agg aac cga gca ggc ctg gat ctg 755	
	Met Val Gln Arg Pro Gln Pro His Arg Asn Arg Ala Gly Leu Asp Leu	
	180 185 190	

	cca acc ata gac acg gga tat gat tcc cag ccc cag gat gtc ctg ggc	803
	Pro Thr Ile Asp Thr Gly Tyr Asp Ser Gln Pro Gln Asp Val Leu Gly	
	195 200 205	
5	atc agg cag ctg gaa agg ccc ctg ccc ctc acc tcc gtg tgt tac ccc	851
	Ile Arg Gln Leu Glu Arg Pro Leu Pro Leu Thr Ser Val Cys Tyr Pro	
	210 215 220	
10	cag gac ctc ccc aga cct ctc agg tcc agg gag ttc cct cag ttt gaa	899
	Gln Asp Leu Pro Arg Pro Leu Arg Ser Arg Glu Phe Pro Gln Phe Glu	
	225 230 235 240	
15	cct cag agg tat cca gca tgt gca cag atg ctg cct ccc aat ctt tcc	947
	Pro Gln Arg Tyr Pro Ala Cys Ala Gln Met Leu Pro Pro Asn Leu Ser	
	245 250 255	
20	cca cat gct cca tgg aac tat cat tac cat tgt cct gga agt ccc gat	995
	Pro His Ala Pro Trp Asn Tyr His Tyr His Cys Pro Gly Ser Pro Asp	
	260 265 270	
	cac cag gtg cca tat ggc cat gac tac cct cga gca gcc tac cag caa	1043
	His Gln Val Pro Tyr Gly His Asp Tyr Pro Arg Ala Ala Tyr Gln Gln	
	275 280 285	
25	gtg atc cag ccg gct ctg cct ggg cag ccc ctg cct gga gcc agt gtg	1091
	Val Ile Gln Pro Ala Leu Pro Gly Gln Pro Leu Pro Gly Ala Ser Val	
	290 295 300	
30	aga ggc ctg cac cct gtg cag aag gtt atc ctg aat tat ccc agc ccc	1139
	Arg Gly Leu His Pro Val Gln Lys Val Ile Leu Asn Tyr Pro Ser Pro	
	305 310 315 320	
35	tgg gac caa gaa gag agg ccc gca cag aga gac tgc tcc ttt ccg ggg	1187
	Trp Asp Gln Glu Arg Pro Ala Gln Arg Asp Cys Ser Phe Pro Gly	
	325 330 335	
40	ctt cca agg cac cag gac cag cca cat cac cag cca cct aat aga gct	1235
	Leu Pro Arg His Gln Asp Gln Pro His His Gln Pro Pro Asn Arg Ala	
	340 345 350	
	ggg gct cct ggg gag tcc ttg gag tgc cct gca gag ctg aga cca cag	1283
	Gly Ala Pro Gly Glu Ser Leu Glu Cys Pro Ala Glu Leu Arg Pro Gln	
	355 360 365	
45	gtt ccc cag cct ccg tcc cca gct gct gtg cct aga ccc cct agc aac	1331
	Val Pro Gln Pro Pro Ser Pro Ala Ala Val Pro Arg Pro Pro Ser Asn	
	370 375 380	
50	cct cca gcc aga gga act cta aaa aca agc aat ttg cca gaa gaa ttg	1379
	Pro Pro Ala Arg Gly Thr Leu Lys Thr Ser Asn Leu Pro Glu Glu Leu	
	385 390 395 400	
55	cgg aaa gtc ttt atc act tat tgc atg gac aca gct atg gag gtg gtg	1427
	Arg Lys Val Phe Ile Thr Tyr Ser Met Asp Thr Ala Met Glu Val Val	
	405 410 415	
	aaa ttc gtg aac ttt ttg ttg gta aat ggc ttc caa act gca att gac	1475
	Lys Phe Val Asn Phe Leu Leu Val Asn Gly Phe Gln Thr Ala Ile Asp	
	420 425 430	

ata ttt gag gat aga atc cga ggc att gat atc att aaa tgg atg gag 1523
 Ile Phe Glu Asp Arg Ile Arg Gly Ile Asp Ile Ile Lys Trp Met Glu
 435 440 445

5

cgc tac ctt agg gat aag acc gtg atg ata atc gta gca atc agc ccc 1571
 Arg Tyr Leu Arg Asp Lys Thr Val Met Ile Ile Val Ala Ile Ser Pro
 450 455 460

10

aaa tac aaa cag gac gtg gaa ggc gct gag tgc cag ctg gac gag gat 1619
 Lys Tyr Lys Gln Asp Val Glu Gly Ala Glu Ser Gln Leu Asp Glu Asp
 465 470 475 480

15

gag cat ggc tta cat act aag tac att cat cga atg atg cag att gag 1667
 Glu His Gly Leu His Thr Lys Tyr Ile His Arg Met Met Gln Ile Glu
 485 490 495

20

ttc ata aaa caa gga agc atg aat ttc aga ttc atc cct gtg ctc ttc 1715
 Phe Ile Lys Gln Gly Ser Met Asn Phe Arg Phe Ile Pro Val Leu Phe
 500 505 510

25

cca aat gct aag aag gag cat gtg ccc acc tgg ctt cag aac act cat 1763
 Pro Asn Ala Lys Lys Glu His Val Pro Thr Trp Leu Gln Asn Thr His
 515 520 525

gtc tac agc tgg ccc aag aat aaa aaa aac atc ctg ctg cgg ctg ctg 1811
 Val Tyr Ser Trp Pro Lys Asn Lys Lys Asn Ile Leu Leu Arg Leu Leu
 530 535 540

30

aga gag gaa gag tat gtg gct cct cca cgg ggg cct ctg ccc acc ctt 1859
 Arg Glu Glu Glu Tyr Val Ala Pro Pro Arg Gly Pro Leu Pro Thr Leu
 545 550 555 560

35

cag gtg gtt ccc ttg tgacaccgtt catccccaga tcaactgaggc caggccatgt 1914
 Gln Val Val Pro Leu
 565

40

ttggggcctt gttctgacag cattctggct gaggctggtc ggtagcactc ctggctgggtt 1974
 tttttctgtt cctccccgag aggccctctg gccccagga aacctgttgt gcagagctct 2034
 tccccggaga cctccacaca ccctggcttt gaagtggagt ctgtgactgc tctgcattct 2094

45

ctgcttttaa aaaaaccatt gcagggtcca gtgtcccata tgttcctcct gacagtttga 2154
 tgtgtccatt ctgggcctct cagtgttag caagtagata atgtaaggga tgtggcagca 2214
 aatggaaatg actacaaaca ctctcctatc aatcacttca ggctactttt atgagtttagc 2274

50

cagatgcttg tgtatcctca gaccaaactg attcatgtac aaataataaa atgtttactc 2334
 ttttgtaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaa 2377

5 MNR SIPVEVDESEFPYPSQLLKPIPEYSPEESEEPPAPNIRNMAPNSLSAPTMLHNSSGDFSQAHSTLKLANH
 QRPVSRQVTCLRTQVLEDSFSCRRHPGLGKAPPSGCSAVSEPASES VVGALPAEHQFSFMEKRNQWLVSQ
 LSAASPDGTGSDSDQSLPNASADSLGGSQEMVQRPPQPHNRAGLDLPTIDTGYDSQPQDVLGIRQLERPL
 10 PLTSVCYPQDLPRPLRSREFPQFEPQRYPACAQMLPPNLS PHAPWNYHYHCPGSPDHQVPYGHDPRAAYQQ
 VIQPALPGQPLPGASVRGLHPVQKVILNYPSPWDQBERPAQRDCSFPGLPRHQDQPHHQPNNRAGAPGESLE
 CPABLRPQVPQPPSPAAPVRPPSNPPARGTLKTSNLPEELRKVFITYSMDTAMEVVKFVNFLLVNGFQTAID
 IFEDRIRGIDI IKWMERYLRDKTVMIIIVAI SPKYKQDVEGAESQLDEDEHGLHTKYIHRMMQIEFIKQGS MN
 FRFIPVLPNAPNAKEHVPTWLQNTHVYSWPKNKNILLRLLREEEYVAPPRGPLPTLQVVPL

Reverse translation of primate, e.g., human, DCRS10 (SEQ ID NO: 24):

15 atgaaymgw snathccngt ngargtngay garwsngarc cntayccnws ncarytnytn 60
 aarccnathc cngartayws nccngargar garwsngarc cncngcnc naayathmgn 120
 aayatggcnc cnaaywsnyt nwsngcncn acnatgytnc ayaaywsnws nggngaytty 180
 20 wsnccargcnc aywsnacnyt naarytngcn aaycaycarm gncngtnws nmgnccargtn 240
 acntgyytnm gnacncargt nytngargay wsnccargayw snttytgymg nmgnccayccn 300
 ggnytnngna argcnttycc nwsnggntgy wsnccngtnw sngarccngc nwsngarwsn 360
 25 gtngtnngng cnytnccngc ngarcaycar ttywsnttya tggaraarmg naaycartgg 420
 ytngtnwsnc arytnwsngc ngcnwsncn gayacnggnc aygaywsnga yaarwsngay 480
 30 carwsnytn cnaaygcnc nngcngaywsn ytngngngnw sncargarat ggtncarmgn 540
 ccncarccnc aymgnaaymg ngcnggnytn gayytnccna cnathgayac nggntaygay 600
 wsnccarccnc argaygtnyt nggnathmgn carytngarm gncnytncc nytnacnwsn 660
 35 gtntgytayc cncargayt nccnmgnccn ytnmgnwsnm gngarttycc ncarttygar 720
 ccncarmgt aycngcngtg ygcncaratg ytnccncna ayytnwsncc ncaygcncn 780
 40 tggaytayc aytaycaytg yccngnwsn ccngaycayc argtnccnta yggncaygay 840
 tayccnmng cngcntayca rcargtnath carccngcny tncnggnc rccnytnccn 900
 ggngcnwsng tnmnggnytn ncayccngtn caraargtna thytnaayta yccnwsncn 960
 45 tggaycarg argarmgnc ngncarmgn gaytgywsnt tyccnggnytn nccnmgnccay 1020
 cargaycarg cncaycayca rccncnaay mgngcnggng cncnggnga rwsnytngar 1080
 50 tgyccngcng arytnmgncc ncargtnccn carccncnw sncngcngc ngtnccnmgn 1140
 ccncnwsna aycncngc nmnggnacn ytnaaracnw snaaytncc ngargarytn 1200
 mgnaargtn tyathacnta ywsnatggay acngcnatgg argtngtana rtyygtana 1260
 55 tyytnytn tnaaygntt ycaracngcn athgayatht tygargaymg nathmnggng 1320
 athgayatha thaartggat ggarmgntay ytnmnggaya aracngtnat gathathgtn 1380

genathwsnc cnaartayaa rcargaygtn gargngcng arwsncaryt ngaygargay 1440
 garcayggny tncayacnaa rtayathcay mgnatgatgc arathgartt yathaarc 1500
 5 ggnwsnatga ayttymgntt yathccngtn ytnttyccna aygcnaaraa rgarcaygtn 1560
 ccnactggy tncaraayac ncaygtntay wsntggccna araayaaraa raayathytn 1620
 10 ytnmgnytny tnmngngarga rgartaygtn gcncncncnm gnggnccnyt nccnacnytn 1680
 cargtngtnc cnytn 1695

Rodent, e.g., mouse, embodiment (see SEQ ID NO: 25 and 26).

15 cag gac ctc cct ggg cct ctg agg tcc agg gaa ttg cca cct cag ttt 48
 Gln Asp Leu Pro Gly Pro Leu Arg Ser Arg Glu Leu Pro Pro Gln Phe
 1 5 10 15
 20 gaa ctt gag agg tat cca atg aac gcc cag ctg ctg ccg ccc cat cct 96
 Glu Leu Glu Arg Tyr Pro Met Asn Ala Gln Leu Leu Pro Pro His Pro
 20 25 30
 25 tcc cca cag gcc cca tgg aac tgt cag tac tac tgc ccc gga ggg ccc 144
 Ser Pro Gln Ala Pro Trp Asn Cys Gln Tyr Tyr Cys Pro Gly Gly Pro
 35 40 45
 30 tac cac cac cag gtg cca cac ggc cat ggc tac cct cca gca gca gcc 192
 Tyr His His Gln Val Pro His Gly His Gly Tyr Pro Pro Ala Ala Ala
 50 55 60
 35 tac cag caa gta ctc cag cct gct ctg cct ggg cag gtc ctt cct ggg 240
 Tyr Gln Gln Val Leu Gln Pro Ala Leu Pro Gly Gln Val Leu Pro Gly
 65 70 75 80
 40 gca agg gca aga ggc cca cgc cct gtg cag aag gtc atc ctg aat gac 288
 Ala Arg Ala Arg Gly Pro Arg Pro Val Gln Lys Val Ile Leu Asn Asp
 85 90 95
 45 tcc agc ccc caa gac caa gaa gag aga cct gca cag aga gac ttc tct 336
 Ser Ser Pro Gln Asp Gln Glu Glu Arg Pro Ala Gln Arg Asp Phe Ser
 100 105 110
 50 ttc ccg agg ctc ccg agg gac cag ctc tac cgc cca cca tct aat gga 384
 Phe Pro Arg Leu Pro Arg Asp Gln Leu Tyr Arg Pro Pro Ser Asn Gly
 115 120 125
 55 gtg gaa gcc cct gag gag tcc ttg gac ctt cct gca gag ctg aga cca 432
 Val Glu Ala Pro Glu Glu Ser Leu Asp Leu Pro Ala Glu Leu Arg Pro
 130 135 140
 cat ggt ccc cag gct cca tcc cta gct gcc gtg cct aga ccc cct agc 480
 His Gly Pro Gln Ala Pro Ser Leu Ala Ala Val Pro Arg Pro Pro Ser
 145 150 155 160
 55 aac ccc tta gcc cga gga act cta aga acc agc aat ttg cca gaa gaa 528
 Asn Pro Leu Ala Arg Gly Thr Leu Arg Thr Ser Asn Leu Pro Glu Glu
 165 170 175

	tta cgg aaa gtc ttt atc act tat tct atg gac aca gcc atg gag gtg	576
	Leu Arg Lys Val Phe Ile Thr Tyr Ser Met Asp Thr Ala Met Glu Val	
	180 185 190	
5	gtg aaa ttt gtg aac ttt ctg ttg gtg aac ggc ttc caa act gcg att	624
	Val Lys Phe Val Asn Phe Leu Leu Val Asn Gly Phe Gln Thr Ala Ile	
	195 200 205	
10	gac ata ttt gag gat aga atc cgg ggt att gat atc att aaa tgg atg	672
	Asp Ile Phe Glu Asp Arg Ile Arg Gly Ile Asp Ile Ile Lys Trp Met	
	210 215 220	
15	gag cgc tat ctt cga gat aag aca gtg atg ata atc gta gca atc agc	720
	Glu Arg Tyr Leu Arg Asp Lys Thr Val Met Ile Ile Val Ala Ile Ser	
	225 230 235 240	
20	ccc aaa tac aaa cag gat gtg gaa ggc gct gag tgc cag ctg gac gag	768
	Pro Lys Tyr Lys Gln Asp Val Glu Gly Ala Glu Ser Gln Leu Asp Glu	
	245 250 255	
	gac gag cat ggc tta cat act aag tac att cat cgg atg atg cag att	816
	Asp Glu His Gly Leu His Thr Lys Tyr Ile His Arg Met Met Gln Ile	
	260 265 270	
25	gag ttc ata agt cag gga agc atg aac ttc aga ttc atc cct gtg ctc	864
	Glu Phe Ile Ser Gln Gly Ser Met Asn Phe Arg Phe Ile Pro Val Leu	
	275 280 285	
30	ttc cca aat gcc aag aag gag cat gtg ccg acc tgg ctt cag aac act	912
	Phe Pro Asn Ala Lys Lys Glu His Val Pro Thr Trp Leu Gln Asn Thr	
	290 295 300	
35	cat gtt tac agc tgg ccc aag aat aag aaa aac atc ctg ctg cgg ctg	960
	His Val Tyr Ser Trp Pro Lys Asn Lys Lys Asn Ile Leu Leu Arg Leu	
	305 310 315 320	
40	ctc agg gag gaa gag tat gtg gct cct ccc cga ggc cct ctg ccc acc	1008
	Leu Arg Glu Glu Glu Tyr Val Ala Pro Pro Arg Gly Pro Leu Pro Thr	
	325 330 335	
	ctt cag gtg gta ccc ttg tgacgatggc cactccagct cagtgccagc	1056
	Leu Gln Val Val Pro Leu	
	340	
45	ctgtttctcac agcattcttc tagcggagct ggctggtggc acccaggccc tggaacacct	1116
	cttctacaga gtcctctgtc tcttgagtct gagttgtcct cgctgggctt ccagagcttc	1176
50	agtgcctgga tgctgcaggt gacagaaaca aacatctatg accacaaaaa ctctcatcac	1236
	ttcagctact tttatgagtc ggtcagatgc tctgtgtcct tagaccagtc taaatcatgc	1296
	tcaaataata aaatgattat tctttgt	1323
55	QDLPGPLRSRELPPQFELERYPMNAQLLPPHPSPQAPWNCQYYCPGGPYHHQVPHGHGYPPAAAYQQVLQPA LPGQVLPGARARGPRPVQKVLNDSSPDQEERPAQRDFSFPRLPRDQLYRPPSNGVEAPEESLDLPAELRP HGPQAPSLAAVPRPPSNPLARGTLRTSNLPEELRKVFITYSMDTAMEVVKFVNFLLVNGFQTATDIFEDRIR GIDIWKWMERYLRDKTVMIIVAISPKYKQDVEGAESQLDEDEHGLHTKYIHRMMQIEFISQGSMMNFRFIPVL FPNAKKEHVPTWLQNTHVYSWPKNKNILLRLLREEYVAPPRGPLPTLQVVPL.	

Reverse translation of rodent, e.g., mouse, DCRS6 (SEQ ID NO: 27):

5 cargayytnc cnggnccnyt nmgnwnsmgn garytnccnc cncarttyga rytngarmgn 60
 tayccnatga aygcncaryt nytnccnccn cayccnwsnc cncargcncc ntggaaaytgy 120
 10 cartaytayt gyccngggngg nccntaycay caycargtnc cncayggnga yggntayccn 180
 ccngcngcng cntaycarca rgtnytnear ccngcnytn cnggncargt nytnccnggn 240
 gcnmgngcnm gnggnccnmg nccngtnear aargtnathy tnaaygayws nwsnccncar 300
 15 gaycargarg armgnccngc ncarmnggay ttywsnttyc cnmgnytncc nmngaycar 360
 ytntaymgnc cncnwsnaa yggngtngar gcncngarg arwsnytna yytnccngcn 420
 garytnmgnc cncayggnc nccargcnccn wsnytnngc cngtnccnmg nccnccnwsn 480
 20 aayccnytn cnmngngnac nytnmgnaen wsnaayytnc cngargaryt nmgnaargtn 540
 ttyathacnt aywsnatgga yacngcnatg gargtngtna arttygtnaa yttyytnytn 600
 25 gtnaayggnt tyacaracngc nathgayath ttygargaym gnathmgngg nathgayath 660
 athaartgga tggarmgnta yytnmgngay aaracngtna tgathathgt ngcnathwsn 720
 ccnaartaya arcargaygt ngarggngcn garwsncary tngaygarga ygarayggg 780
 30 ytncayacna artayathca ymgntatgat carathgart tyathwsnca rggwnsnatg 840
 aayttnmgnt tyathccngt nytnnttyccn aaygcnaara argarcaygt nccnacntgg 900
 35 ytncaraaya cncaygtnta ywsntggccn aaraayaara araayathyt nytnmgnytn 960
 ytnmgngarg argartaygt ngcnccnccn mgnggncny tncnacnytn ncargtngtn 1020
 ccnytn 1026
 40

Table 6: Alignment of the cytoplasmic portions of various cytokine receptor subunits. The IL-17R_Hu (SEQ ID NO: 28) is GenBank AAB99730.1(U58917), gi|7657230; the IL-17R_Mu (SEQ ID NO: 29) is GenBank AAC52357.1(U31993), gi|6680411; the IL-17R_Ce (SEQ ID NO: 30) is GenBank AAA811100.1(U39997), gi|1353171; and the DCRS6_Ce (SEQ ID NO: 31) is EMBCAA90543.1(Z50177), gi|7503597. Of particular interest are motifs or features corresponding, in primate DCRS8 to: R/K at 339/340; D/E at 348/349; alpha helical regions from H353-Q365, C370-S381, E389-H396, K410-D414, and D485-H495; beta sheet regions correspond to F400-V404 and F458-Y462; E at 431; E/D at 442/443; Y/F at 458; D/E at 468-470; Y/F at 481; and Q/R/F at 523.

45

50

5	DCRS7_Mu DCRS7_Hu IL-17R_Hu IL-17R_Mu DCRS10 DCRS10_Mu DCRS9_Hu DCRS8_Hu IL-17R_Ce DCRS6_Hu DCRS6_Ce	RTALLLSADG-AGYERLVGALASALSQMP---LRVAVDLWSRRE-LSAHGALAWFHHQR RAALLLYSADD-SGFERLVGALASALCQLP---LRVAVDLWSRRE-LSAQGPVAVFHAQR RKVWIIYSADH-PLYVDVVLKFAQFLLTACG--TEVALDLLLEEQA-ISEAGVMTWVGRQK RKVWIVYSADH-PLYVEVVVLKFAQFLITACG--TEVALDLLEEQV-ISEVGVMTWVSRQK RKVFITYSMD---TAMEVVVKFVNFLVNG---FQTAIDIFEDR--IRGIDI IKWMERYL RKVFITYSMD---TAMEVVVKFVNFLVNG---FQTAIDIFEDR--IRGIDI IKWMERYL RPVLLLHAADS-EAQRRLVGALAE LLRAALGGGRDVI VDLWEGRH-VARVGPLPWLWAAR PKVFLCYSSKDGQNMNVVQCFAYFLQDFCG--CEVALDLWEDFS-LCREGQREWVIQKI VKVMIVYADDN-DLHTDCVKKLVENLRNCAS--CDPVFDLEKLI--TAEIVPSRWLVQDI IKVLVVPSEI--CFHHTICYFTEFLQNHCR--SEVILEKWQKKK-IAEMGPVQWLATQK FKVMLVCPEVS-GRDEDFMMRIADALKKSN--NKVVCDRWFEFSKNAEENMLHWVYEQT . : . : : *
15	DCRS7_Mu DCRS7_Hu IL-17R_Hu IL-17R_Mu DCRS10 DCRS10_Mu DCRS9_Hu DCRS8_Hu IL-17R_Ce DCRS6_Hu DCRS6_Ce	RRILQEGGVVILLFSPAAVAQCQ---QWLQLQTVEP---GP---HDLAALWLSCVLPDFL RQTLQEGGVVLLFSPGAVALCS---EWLQDGVSGPGAHP---HDAFRASLSCVLPDFL QEMVESNSKIIIVLCSRGTRAKWQALLGRGAP-VRLRCDHGKPV-GDLFTAAMNMILPDFK QEMVESNSKIIILCSRGTOAKWKAILGWAEPVQLRCDHWKPA-GDLFTAAMNMILPDFK R---DKTVMIIIVAI SPKYQDVE---GAESQLDED-EHGL---HTKYIHRM-MQIEFIK R---DKTVMIIIVAI SPKYQDVE---GAESQLDED-EHGL---HTKYIHRM-MQIEFIS TRVAREQGTVLLWLSGADLRPV S---GPD P-RAAP-----LLA---LLHAAP H---ESQFIIIVCSKGMKYFVD---KKNYKHKGGRGSGK---GELFLVAVSAIAEKL R S---SLKKFIIIVSDCAEKILD---TEASETHQLVQARP--FADLFGPAMEMIIRDAT K---AADKVVFLLSNDVNSVCD---GTCGKSEGSPSENS---QDLFPLAFNLFCSDLR K---IAEKIIVFHSAYYHPRCG---IYDVINNFPCDTPR-----LAHIALT---PEAQ . . . *
30	DCRS7_Mu DCRS7_Hu IL-17R_Hu IL-17R_Mu DCRS10 DCRS10_Mu DCRS9_Hu DCRS8_Hu IL-17R_Ce DCRS6_Hu DCRS6_Ce	QGRATGR-----YGVVYFDGLLHPDSVPSFPRVAPLFSLP-SQLPAFLDALQ--GGCSTS QGRAPGS-----YVGACFDRLLHPDAVPALFRTVPVFTLP-SQLPDLFGALQ--QPRAPR RPACFGT-----YVVCYFSEVSCDGDVPDLFGAAPRYPLM-DRFEEVYFRIQ--DLEMFO RPACFGT-----YVVCYFSGICSERDVPDLFNITSRYPLM-DRFEEVYFRIQ--DLEMFE QGS MNFR-----FIPVLFPNAK-KEHVPTWLQNTHVYSWP-KNKKNILLRLL-REEEYVA QGS MNFR-----FIPVLFPNAK-KEHVPTWLQNTHVYSWP-KNKKNILLRLL-REEEYVA RPL-----LLLAYFSRLCAKGDIPPLRALPRYRL- RDLPRLLRALD--ARPFAE QAKQSSSAALS KFI AVYFDYSC-EGDVP GILD LSTKYRLM-DNLPQLCSHLHSRDHGLQE HNFPEAR---KKYAVVRFNYS P--HVPPNLAILNLPTFIPEQFAQLTAF LHN-VEHTER SQIHLHK----YVVVYFREID-TKDDYNALSVC PKYHLM-KDATAFCAELL---HVKQQ RSVPKEV----EYVLP RDQKLL--EDAFDITIADPLVIDIPIEDVAIPENVP--IH HES C
40	DCRS7_Mu DCRS7_Hu IL-17R_Hu IL-17R_Mu DCRS10 DCRS10_Mu DCRS9_Hu DCRS8_Hu IL-17R_Ce DCRS6_Hu DCRS6_Ce	AGRPAD RVER-----VT-----QALRSALDSCTS----- SGRLQERAEQ-----VS-----RALQPALDSYFHPP----- PGRMHRVGELSGDNYLRS---PGRQLRAALDRFRDWQVRCPDW PGRMHVREL TGDNYLQS---PSGRQLKEAVLR FQEWQTQCPDW P----PRGPL-----PTLQVVPL----- P----PRGPL-----PTLQVVPL----- ATSWGRLGAR-----QRRQSRLELCSR----- PGQHTRQGS R-----RNYFRSKSGRSLYVAICNMHQFIDE EPDW ANVTQNISEA-----Q-----IHEWNL CASRMMSFFVRNPW VS----AGKR-----SQACHDGCCSL----- DSIDSRNNSK-----THSTDSGVSSLSS---NS--

Table 6 shows comparison of the available sequences of primate, rodent, and various other receptors. Various conserved residues are aligned and indicated. The structurally homologous cytoplasmic domains most likely signal through pathways like IL-17, e.g., through NFkB. Similar to IL-1 signalling, it is likely that these receptors are involved in innate immunity and/or development.

As used herein, the term DCRS shall be used to describe a protein comprising amino acid sequences shown in Tables 1-5, respectively. In many cases, a substantial fragment thereof will be functionally or structurally equivalent, including, e.g., an extracellular or intracellular domain. The invention also includes a protein variation of the respective DCRS allele whose sequence is provided, e.g., a mutein or soluble extracellular construct. Typically, such agonists or antagonists will exhibit less than about 10% sequence differences, and thus will often have between 1 and 11 substitutions, e.g., 2-, 3-, 5-, 7-fold, and others. It also encompasses allelic and other variants, e.g., natural polymorphic, of the protein described. Typically, it will bind to its corresponding biological ligand, perhaps in a dimerized state with an alpha receptor subunit, with high affinity, e.g., at least about 100 nM, usually better than about 30 nM, preferably better than about 10 nM, and more preferably at better than about 3 nM. The term shall also be used herein to refer to related naturally occurring forms, e.g., alleles, polymorphic variants, and metabolic variants of the mammalian protein. Preferred forms of the receptor complexes will bind the appropriate ligand with an affinity and selectivity appropriate for a ligand-receptor interaction.

This invention also encompasses combinations of proteins or peptides having substantial amino acid sequence identity with an amino acid sequence in Tables 1-5. It will include sequence variants with relatively few residue substitutions, e.g., preferably less than about 3-5.

A substantial polypeptide "fragment", or "segment", is a stretch of amino acid residues of at least about 8 amino acids, generally at least 10 amino acids, more generally at least 12 amino acids, often at least 14 amino acids, more often at least 16 amino acids, typically at least 18 amino acids, more typically at least 20 amino acids, usually at least 22 amino acids, more usually at least 24 amino acids, preferably at least 26 amino acids, more preferably at least 28 amino acids, and, in particularly preferred embodiments, at least about 30 or more amino acids. This includes, e.g., 40, 50, 60, 70, 85, 100, 115, 130, 150, and other lengths. Sequences of segments of different proteins can be compared to one another over appropriate length stretches, typically between conserved motifs. In many situations, fragments may exhibit functional properties of the intact subunits, e.g., the extracellular domain of the transmembrane receptor may retain the ligand binding features, and may be used to prepare a soluble receptor-like complex.

Amino acid sequence homology, or sequence identity, is determined by optimizing residue matches. In some comparisons, gaps may be introduced, as required. See, e.g., Needleham, et al., (1970) *J. Mol. Biol.* 48:443-453; Sankoff, et al., (1983) chapter one in Time Warps, String Edits, and Macromolecules: The Theory and Practice of Sequence Comparison, Addison-Wesley, Reading, MA; and software packages from IntelliGenetics, Mountain View, CA; and the University of Wisconsin Genetics Computer Group (GCG), Madison, WI; each of which is incorporated herein by reference. This changes when considering conservative substitutions as matches. Conservative substitutions typically include substitutions within the following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid; asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine. Homologous amino acid sequences are intended to include natural allelic and interspecies variations in the cytokine sequence. Typical homologous proteins or peptides will have from 50-100% homology (if gaps can be introduced), to 60-100% homology (if conservative substitutions are included) with an amino acid sequence segment of, e.g., Table 3 or 4. Homology measures will be at least about 70%, generally at least 76%, more generally at least 81%, often at least 85%, more often at least 88%, typically at least 90%, more typically at least 92%, usually at least 94%, more usually at least 95%, preferably at least 96%, and more preferably at least 97%, and in particularly preferred embodiments, at least 98% or more. The degree of homology will vary with the length of the compared segments. Homologous proteins or peptides, such as the allelic variants, will share most biological activities with the embodiments described in Tables 1-5.

As used herein, the term "biological activity" is used to describe, without limitation, effects on inflammatory responses, innate immunity, and/or morphogenic development by cytokine-like ligands. For example, these receptors should mediate phosphatase or phosphorylase activities, which activities are easily measured by standard procedures. See, e.g., Hardie, et al. (eds. 1995) The Protein Kinase FactBook vols. I and II, Academic Press, San Diego, CA; Hanks, et al. (1991) Meth. Enzymol. 200:38-62; Hunter, et al. (1992) Cell 70:375-388; Lewin (1990) Cell 61:743-752; Pines, et al. (1991) Cold Spring Harbor Symp. Quant. Biol. 56:449-463; and Parker, et al. (1993) Nature 363:736-738. The receptors, or portions thereof, may be useful as phosphate labeling enzymes to label general or specific substrates. The subunits may also be functional immunogens to elicit recognizing antibodies, or antigens capable of binding antibodies.

The terms ligand, agonist, antagonist, and analog of, e.g., a DCRS8 or DCRS9, include molecules that modulate the characteristic cellular responses to cytokine ligand proteins, as well as molecules possessing the more standard structural binding competition features of ligand-receptor interactions, e.g., where the receptor is a natural

receptor or an antibody. The cellular responses likely are typically mediated through receptor tyrosine kinase pathways.

Also, a ligand is a molecule which serves either as a natural ligand to which said receptor, or an analog thereof, binds, or a molecule which is a functional analog of the natural ligand. The functional analog may be a ligand with structural modifications, or may be a wholly unrelated molecule which has a molecular shape which interacts with the appropriate ligand binding determinants. The ligands may serve as agonists or antagonists, see, e.g., Goodman, et al. (eds. 1990) Goodman & Gilman's: The Pharmacological Bases of Therapeutics, Pergamon Press, New York.

Rational drug design may also be based upon structural studies of the molecular shapes of a receptor or antibody and other effectors or ligands. See, e.g., Herz, et al. (1997) J. Recept. Signal Transduct. Res. 17:671-776; and Chaiken, et al. (1996) Trends Biotechnol. 14:369-375. Effectors may be other proteins which mediate other functions in response to ligand binding, or other proteins which normally interact with the receptor. One means for determining which sites interact with specific other proteins is a physical structure determination, e.g., x-ray crystallography or 2 dimensional NMR techniques. These will provide guidance as to which amino acid residues form molecular contact regions. For a detailed description of protein structural determination, see, e.g., Blundell and Johnson (1976) Protein Crystallography, Academic Press, New York, which is hereby incorporated herein by reference.

II. Activities

The cytokine receptor-like proteins will have a number of different biological activities, e.g., modulating cell proliferation, or in phosphate metabolism, being added to or removed from specific substrates, typically proteins. Such will generally result in modulation of an inflammatory function, other innate immunity response, or a morphological effect. The subunit will probably have a specific low affinity binding to the ligand.

The DCRS8 and DCRS9 have characteristic motifs of receptors signaling through the JAK pathway. See, e.g., Ihle, et al. (1997) Stem Cells 15(suppl. 1):105-111; Silvennoinen, et al. (1997) APMIS 105:497-509; Levy (1997) Cytokine Growth Factor Review 8:81-90; Winston and Hunter (1996) Current Biol. 6:668-671; Barrett (1996) Baillieres Clin. Gastroenterol. 10:1-15; and Briscoe, et al. (1996) Philos. Trans. R. Soc. Lond. B. Biol. Sci. 351:167-171.

The biological activities of the cytokine receptor subunits will be related to addition or removal of phosphate moieties to substrates, typically in a specific manner, but occasionally in a non specific manner. Substrates may be identified, or conditions for

enzymatic activity may be assayed by standard methods, e.g., as described in Hardie, et al. (eds. 1995) The Protein Kinase FactBook vols. I and II, Academic Press, San Diego, CA; Hanks, et al. (1991) Meth. Enzymol. 200:38-62; Hunter, et al. (1992) Cell 70:375-388; Lewin (1990) Cell 61:743-752; Pines, et al. (1991) Cold Spring Harbor Symp. Quant. Biol. 56:449-463; and Parker, et al. (1993) Nature 363:736-738.

The receptor subunits may combine to form functional complexes, e.g., which may be useful for binding ligand or preparing antibodies. These will have substantial diagnostic uses, including detection or quantitation.

10 III. Nucleic Acids

This invention contemplates use of isolated nucleic acid or fragments, e.g., which encode these or closely related proteins, or fragments thereof, e.g., to encode a corresponding polypeptide, preferably one which is biologically active. In addition, this invention covers isolated or recombinant DNAs which encode combinations of such
15 proteins or polypeptides having characteristic sequences, e.g., of the DCRSs. Typically, the nucleic acid is capable of hybridizing, under appropriate conditions, with a nucleic acid sequence segment shown in Tables 1-5, but preferably not with a corresponding segment of other receptors described in Table 6. Said biologically active protein or polypeptide can be a full length protein, or fragment, and will typically have a segment of
20 amino acid sequence highly homologous, e.g., exhibiting significant stretches of identity, to one shown in Tables 1-5. Further, this invention covers the use of isolated or recombinant nucleic acid, or fragments thereof, which encode proteins having fragments which are equivalent to the DCRS8 or DCRS9 proteins. The isolated nucleic acids can have the respective regulatory sequences in the 5' and 3' flanks, e.g., promoters,
25 enhancers, poly-A addition signals, and others from the natural gene. Combinations, as described, are also provided.

An "isolated" nucleic acid is a nucleic acid, e.g., an RNA, DNA, or a mixed polymer, which is substantially pure, e.g., separated from other components which naturally accompany a native sequence, such as ribosomes, polymerases, and flanking
30 genomic sequences from the originating species. The term embraces a nucleic acid sequence which has been removed from its naturally occurring environment, and includes recombinant or cloned DNA isolates, which are thereby distinguishable from naturally occurring compositions, and chemically synthesized analogs or analogs biologically synthesized by heterologous systems. A substantially pure molecule includes isolated
35 forms of the molecule, either completely or substantially pure.

An isolated nucleic acid will generally be a homogeneous composition of molecules, but will, in some embodiments, contain heterogeneity, preferably minor. This

heterogeneity is typically found at the polymer ends or portions not critical to a desired biological function or activity.

A "recombinant" nucleic acid is typically defined either by its method of production or its structure. In reference to its method of production, e.g., a product made
5 by a process, the process is use of recombinant nucleic acid techniques, e.g., involving human intervention in the nucleotide sequence. Typically this intervention involves in vitro manipulation, although under certain circumstances it may involve more classical animal breeding techniques. Alternatively, it can be a nucleic acid made by generating a sequence comprising fusion of two fragments which are not naturally contiguous to each
10 other, but is meant to exclude products of nature, e.g., naturally occurring mutants as found in their natural state. Thus, for example, products made by transforming cells with an unnaturally occurring vector is encompassed, as are nucleic acids comprising sequence derived using any synthetic oligonucleotide process. Such a process is often done to replace a codon with a redundant codon encoding the same or a conservative amino acid,
15 while typically introducing or removing a restriction enzyme sequence recognition site. Alternatively, the process is performed to join together nucleic acid segments of desired functions to generate a single genetic entity comprising a desired combination of functions not found in the commonly available natural forms, e.g., encoding a fusion protein. Restriction enzyme recognition sites are often the target of such artificial
20 manipulations, but other site specific targets, e.g., promoters, DNA replication sites, regulation sequences, control sequences, or other useful features may be incorporated by design. A similar concept is intended for a recombinant, e.g., fusion, polypeptide. This will include a dimeric repeat. Specifically included are synthetic nucleic acids which, by genetic code redundancy, encode equivalent polypeptides to fragments of DCRSs and
25 fusions of sequences from various different related molecules, e.g., other cytokine receptor family members.

A "fragment" in a nucleic acid context is a contiguous segment of at least about 17 nucleotides, generally at least 21 nucleotides, more generally at least 25 nucleotides, ordinarily at least 30 nucleotides, more ordinarily at least 35 nucleotides, often at least 39
30 nucleotides, more often at least 45 nucleotides, typically at least 50 nucleotides, more typically at least 55 nucleotides, usually at least 60 nucleotides, more usually at least 66 nucleotides, preferably at least 72 nucleotides, more preferably at least 79 nucleotides, and in particularly preferred embodiments will be at least 85 or more nucleotides. Typically, fragments of different genetic sequences can be compared to one another over
35 appropriate length stretches, particularly defined segments such as the domains described below.

A nucleic acid which codes for the DCRS8 or DCRS9 will be particularly useful to identify genes, mRNA, and cDNA species which code for itself or closely related proteins, as well as DNAs which code for polymorphic, allelic, or other genetic variants, e.g., from different individuals or related species. Preferred probes for such screens are those regions of the interleukin which are conserved between different polymorphic variants or which contain nucleotides which lack specificity, and will preferably be full length or nearly so. In other situations, polymorphic variant specific sequences will be more useful.

This invention further covers recombinant nucleic acid molecules and fragments having a nucleic acid sequence identical to or highly homologous to the isolated DNA set forth herein. In particular, the sequences will often be operably linked to DNA segments which control transcription, translation, and DNA replication. These additional segments typically assist in expression of the desired nucleic acid segment.

Homologous, or highly identical, nucleic acid sequences, when compared to one another, e.g., DCRS8 sequences, exhibit significant similarity. The standards for homology in nucleic acids are either measures for homology generally used in the art by sequence comparison or based upon hybridization conditions. Comparative hybridization conditions are described in greater detail below.

Substantial identity in the nucleic acid sequence comparison context means either that the segments, or their complementary strands, when compared, are identical when optimally aligned, with appropriate nucleotide insertions or deletions, in at least about 60% of the nucleotides, generally at least 66%, ordinarily at least 71%, often at least 76%, more often at least 80%, usually at least 84%, more usually at least 88%, typically at least 91%, more typically at least about 93%, preferably at least about 95%, more preferably at least about 96 to 98% or more, and in particular embodiments, as high as about 99% or more of the nucleotides, including, e.g., segments encoding structural domains such as the segments described below. Alternatively, substantial identity will exist when the segments will hybridize under selective hybridization conditions, to a strand or its complement, typically using a sequence derived from Tables 1-5. Typically, selective hybridization will occur when there is at least about 55% homology over a stretch of at least about 14 nucleotides, more typically at least about 65%, preferably at least about 75%, and more preferably at least about 90%. See, Kanehisa (1984) Nucl. Acids Res. 12:203-213, which is incorporated herein by reference. The length of homology comparison, as described, may be over longer stretches, and in certain embodiments will be over a stretch of at least about 17 nucleotides, generally at least about 20 nucleotides, ordinarily at least about 24 nucleotides, usually at least about 28 nucleotides, typically at least about 32 nucleotides, more typically at least about 40 nucleotides, preferably at least

about 50 nucleotides, and more preferably at least about 75 to 100 or more nucleotides. This includes, e.g., 125, 150, 175, 200, 225, 246, 273, and other lengths.

Stringent conditions, in referring to homology in the hybridization context, will be stringent combined conditions of salt, temperature, organic solvents, and other parameters typically controlled in hybridization reactions. Stringent temperature conditions will usually include temperatures in excess of about 30 C, more usually in excess of about 37 C, typically in excess of about 45 C, more typically in excess of about 55 C, preferably in excess of about 65 C, and more preferably in excess of about 70 C. Stringent salt conditions will ordinarily be less than about 500 mM, usually less than about 400 mM, more usually less than about 300 mM, typically less than about 200 mM, preferably less than about 100 mM, and more preferably less than about 80 mM, even down to less than about 20 mM. However, the combination of parameters is much more important than the measure of any single parameter. See, e.g., Wetmur and Davidson (1968) J. Mol. Biol. 31:349-370, which is hereby incorporated herein by reference.

The isolated DNA can be readily modified by nucleotide substitutions, nucleotide deletions, nucleotide insertions, and inversions of nucleotide stretches. These modifications result in novel DNA sequences which encode this protein or its derivatives. These modified sequences can be used to produce mutant proteins (muteins) or to enhance the expression of variant species. Enhanced expression may involve gene amplification, increased transcription, increased translation, and other mechanisms. Such mutant DCRS8-like derivatives include predetermined or site-specific mutations of the protein or its fragments, including silent mutations using genetic code degeneracy. "Mutant DCRS8" as used herein encompasses a polypeptide otherwise falling within the homology definition of the DCRS8 as set forth above, but having an amino acid sequence which differs from that of other cytokine receptor-like proteins as found in nature, whether by way of deletion, substitution, or insertion. In particular, "site specific mutant DCRS8" encompasses a protein having substantial sequence identity with a protein of Table 3, and typically shares most of the biological activities or effects of the forms disclosed herein.

Although site specific mutation sites are predetermined, mutants need not be site specific. Mammalian DCRS8 mutagenesis can be achieved by making amino acid insertions or deletions in the gene, coupled with expression. Substitutions, deletions, insertions, or many combinations may be generated to arrive at a final construct. Insertions include amino- or carboxy- terminal fusions. Random mutagenesis can be conducted at a target codon and the expressed mammalian DCRS mutants can then be screened for the desired activity, providing some aspect of a structure-activity relationship. Methods for making substitution mutations at predetermined sites in DNA

having a known sequence are well known in the art, e.g., by M13 primer mutagenesis. See also Sambrook, et al. (1989) and Ausubel, et al. (1987 and periodic Supplements).

5 The mutations in the DNA normally should not place coding sequences out of reading frames and preferably will not create complementary regions that could hybridize to produce secondary mRNA structure such as loops or hairpins.

The phosphoramidite method described by Beaucage and Carruthers (1981) Tetra. Letts. 22:1859-1862, will produce suitable synthetic DNA fragments. A double stranded fragment will often be obtained either by synthesizing the complementary strand and annealing the strand together under appropriate conditions or by adding the
10 complementary strand using DNA polymerase with an appropriate primer sequence.

Polymerase chain reaction (PCR) techniques can often be applied in mutagenesis. Alternatively, mutagenesis primers are commonly used methods for generating defined mutations at predetermined sites. See, e.g., Innis, et al. (eds. 1990) PCR Protocols: A Guide to Methods and Applications Academic Press, San Diego, CA; and Dieffenbach
15 and Dveksler (1995; eds.) PCR Primer: A Laboratory Manual Cold Spring Harbor Press, CSH, NY.

Certain embodiments of the invention are directed to combination compositions comprising the receptor or ligand sequences described. In other embodiments, functional portions of the sequences may be joined to encode fusion proteins. In other forms,
20 variants of the described sequences may be substituted.

IV. Proteins, Peptides

As described above, the present invention encompasses primate DCRS6-10, e.g., whose sequences are disclosed in Tables 1-5, and described above. Allelic and other
25 variants are also contemplated, including, e.g., fusion proteins combining portions of such sequences with others, including, e.g., epitope tags and functional domains.

The present invention also provides recombinant proteins, e.g., heterologous fusion proteins using segments from these primate or rodent proteins. A heterologous fusion protein is a fusion of proteins or segments which are naturally not normally fused
30 in the same manner. Thus, the fusion product of, e.g., a DCRS8 with another cytokine receptor is a continuous protein molecule having sequences fused in a typical peptide linkage, typically made as a single translation product and exhibiting properties, e.g., sequence or antigenicity, derived from each source peptide. A similar concept applies to heterologous nucleic acid sequences. Combinations of various designated proteins into
35 complexes are also provided.

In addition, new constructs may be made from combining similar functional or structural domains from other related proteins, e.g., cytokine receptors or Toll-like

receptors, including species variants. For example, ligand-binding or other segments may be "swapped" between different new fusion polypeptides or fragments. See, e.g., Cunningham, et al. (1989) Science 243:1330-1336; and O'Dowd, et al. (1988) J. Biol. Chem. 263:15985-15992, each of which is incorporated herein by reference. Thus, new chimeric polypeptides exhibiting new combinations of specificities will result from the functional linkage of receptor-binding specificities. For example, the ligand binding domains from other related receptor molecules may be added or substituted for other domains of this or related proteins. The resulting protein will often have hybrid function and properties. For example, a fusion protein may include a targeting domain which may serve to provide sequestering of the fusion protein to a particular subcellular organelle.

Candidate fusion partners and sequences can be selected from various sequence data bases, e.g., GenBank, c/o IntelliGenetics, Mountain View, CA; and BCG, University of Wisconsin Biotechnology Computing Group, Madison, WI, which are each incorporated herein by reference. In particular, combinations of polypeptide sequences provided in Tables 1-5 are particularly preferred. Variant forms of the proteins may be substituted in the described combinations.

The present invention particularly provides muteins which bind cytokine-like ligands, and/or which are affected in signal transduction. Structural alignment of human DCRSs with other members of the cytokine receptor family show conserved features/residues. See Table 6. Alignment of the human DCRS8 sequence with other members of the cytokine receptor family indicates various structural and functionally shared features. See also, Bazan, et al. (1996) Nature 379:591; Lodi, et al. (1994) Science 263:1762-1766; Sayle and Milner-White (1995) TIBS 20:374-376; and Gronenberg, et al. (1991) Protein Engineering 4:263-269.

Substitutions with either mouse sequences or human sequences are particularly preferred. Conversely, conservative substitutions away from the ligand binding interaction regions will probably preserve most signaling activities; and conservative substitutions away from the intracellular domains will probably preserve most ligand binding properties.

"Derivatives" of the primate DCRS8 include amino acid sequence mutants, glycosylation variants, metabolic derivatives and covalent or aggregative conjugates with other chemical moieties. Covalent derivatives can be prepared by linkage of functionalities to groups which are found in the DCRS8 amino acid side chains or at the N- or C- termini, e.g., by means which are well known in the art. These derivatives can include, without limitation, aliphatic esters or amides of the carboxyl terminus, or of residues containing carboxyl side chains, O-acyl derivatives of hydroxyl group-containing residues, and N-acyl derivatives of the amino terminal amino acid or amino-group

containing residues, e.g., lysine or arginine. Acyl groups are selected from the group of alkyl-moieties, including C3 to C18 normal alkyl, thereby forming alkanoyl aroyl species.

5 In particular, glycosylation alterations are included, e.g., made by modifying the glycosylation patterns of a polypeptide during its synthesis and processing, or in further processing steps. Particularly preferred means for accomplishing this are by exposing the polypeptide to glycosylating enzymes derived from cells which normally provide such processing, e.g., mammalian glycosylation enzymes. Deglycosylation enzymes are also contemplated. Also embraced are versions of the same primary amino acid sequence which have other minor modifications, including phosphorylated amino acid residues,
10 e.g., phosphotyrosine, phosphoserine, or phosphothreonine.

A major group of derivatives are covalent conjugates of the receptors or fragments thereof with other proteins of polypeptides. These derivatives can be synthesized in recombinant culture such as N- or C-terminal fusions or by the use of agents known in the art for their usefulness in cross-linking proteins through reactive side groups. Preferred
15 derivatization sites with cross-linking agents are at free amino groups, carbohydrate moieties, and cysteine residues.

Fusion polypeptides between the receptors and other homologous or heterologous proteins are also provided. Homologous polypeptides may be fusions between different receptors, resulting in, for instance, a hybrid protein exhibiting binding specificity for
20 multiple different cytokine ligands, or a receptor which may have broadened or weakened specificity of substrate effect. Likewise, heterologous fusions may be constructed which would exhibit a combination of properties or activities of the derivative proteins. Typical examples are fusions of a reporter polypeptide, e.g., luciferase, with a segment or domain of a receptor, e.g., a ligand-binding segment, so that the presence or location of a desired
25 ligand may be easily determined. See, e.g., Dull, et al., U.S. Patent No. 4,859,609, which is hereby incorporated herein by reference. Other gene fusion partners include glutathione-S-transferase (GST), bacterial β -galactosidase, trpE, Protein A, β -lactamase, alpha amylase, alcohol dehydrogenase, and yeast alpha mating factor. See, e.g., Godowski, et al. (1988) Science 241:812-816. Labeled proteins will often be substituted
30 in the described combinations of proteins.

The phosphoramidite method described by Beaucage and Carruthers (1981) Tetra. Letts. 22:1859-1862, will produce suitable synthetic DNA fragments. A double stranded fragment will often be obtained either by synthesizing the complementary strand and annealing the strand together under appropriate conditions or by adding the
35 complementary strand using DNA polymerase with an appropriate primer sequence.

Such polypeptides may also have amino acid residues which have been chemically modified by phosphorylation, sulfonation, biotinylation, or the addition or removal of

other moieties, particularly those which have molecular shapes similar to phosphate groups. In some embodiments, the modifications will be useful labeling reagents, or serve as purification targets, e.g., affinity ligands.

5 Fusion proteins will typically be made by either recombinant nucleic acid methods or by synthetic polypeptide methods. Techniques for nucleic acid manipulation and expression are described generally, for example, in Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual (2d ed.), Vols. 1-3, Cold Spring Harbor Laboratory, and Ausubel, et al. (eds. 1987 and periodic supplements) Current Protocols in Molecular Biology, Greene/Wiley, New York, which are each incorporated herein by reference.

10 Techniques for synthesis of polypeptides are described, for example, in Merrifield (1963) J. Amer. Chem. Soc. 85:2149-2156; Merrifield (1986) Science 232: 341-347; and Atherton, et al. (1989) Solid Phase Peptide Synthesis: A Practical Approach, IRL Press, Oxford; each of which is incorporated herein by reference. See also Dawson, et al. (1994) Science 266:776-779 for methods to make larger polypeptides.

15 This invention also contemplates the use of derivatives of a DCRS8 other than variations in amino acid sequence or glycosylation. Such derivatives may involve covalent or aggregative association with chemical moieties. These derivatives generally fall into three classes: (1) salts, (2) side chain and terminal residue covalent modifications, and (3) adsorption complexes, for example with cell membranes. Such covalent or

20 aggregative derivatives are useful as immunogens, as reagents in immunoassays, or in purification methods such as for affinity purification of a receptor or other binding molecule, e.g., an antibody. For example, a cytokine ligand can be immobilized by covalent bonding to a solid support such as cyanogen bromide-activated Sepharose, by methods which are well known in the art, or adsorbed onto polyolefin surfaces, with or

25 without glutaraldehyde cross-linking, for use in the assay or purification of a cytokine receptor, antibodies, or other similar molecules. The ligand can also be labeled with a detectable group, for example radioiodinated by the chloramine T procedure, covalently bound to rare earth chelates, or conjugated to another fluorescent moiety for use in diagnostic assays.

30 A combination, e.g., including a DCRS8, of this invention can be used as an immunogen for the production of antisera or antibodies specific, e.g., capable of distinguishing between other cytokine receptor family members, for the combinations described. The complexes can be used to screen monoclonal antibodies or antigen-binding fragments prepared by immunization with various forms of impure preparations

35 containing the protein. In particular, the term "antibodies" also encompasses antigen binding fragments of natural antibodies, e.g., Fab, Fab2, Fv, etc. The purified DCRS8 can also be used as a reagent to detect antibodies generated in response to the presence of

elevated levels of expression, or immunological disorders which lead to antibody production to the endogenous receptor. Additionally, DCRS8 fragments may also serve as immunogens to produce the antibodies of the present invention, as described immediately below. For example, this invention contemplates antibodies having binding affinity to or being raised against the amino acid sequences shown in Tables 1-5, fragments thereof, or various homologous peptides. In particular, this invention contemplates antibodies having binding affinity to, or having been raised against, specific fragments which are predicted to be, or actually are, exposed at the exterior protein surface of the native DCRS8 or DCRS9. Complexes of combinations of proteins will also be useful, and antibody preparations thereto can be made.

The blocking of physiological response to the receptor ligands may result from the inhibition of binding of the ligand to the receptor, likely through competitive inhibition. Thus, in vitro assays of the present invention will often use antibodies or antigen binding segments of these antibodies, or fragments attached to solid phase substrates. These assays will also allow for the diagnostic determination of the effects of either ligand binding region mutations and modifications, or other mutations and modifications, e.g., which affect signaling or enzymatic function.

This invention also contemplates the use of competitive drug screening assays, e.g., where neutralizing antibodies to the receptor complexes or fragments compete with a test compound for binding to a ligand or other antibody. In this manner, the neutralizing antibodies or fragments can be used to detect the presence of a polypeptide which shares one or more binding sites to a receptor and can also be used to occupy binding sites on a receptor that might otherwise bind a ligand.

V. Making Nucleic Acids and Protein

DNA which encodes the protein or fragments thereof can be obtained by chemical synthesis, screening cDNA libraries, or by screening genomic libraries prepared from a wide variety of cell lines or tissue samples. Natural sequences can be isolated using standard methods and the sequences provided herein, e.g., in Tables 1-5. Other species counterparts can be identified by hybridization techniques, or by various PCR techniques, combined with or by searching in sequence databases, e.g., GenBank.

This DNA can be expressed in a wide variety of host cells for the synthesis of a full-length receptor or fragments which can in turn, for example, be used to generate polyclonal or monoclonal antibodies; for binding studies; for construction and expression of modified ligand binding or kinase/phosphatase domains; and for structure/function studies. Variants or fragments can be expressed in host cells that are transformed or transfected with appropriate expression vectors. These molecules can be substantially

free of protein or cellular contaminants, other than those derived from the recombinant host, and therefore are particularly useful in pharmaceutical compositions when combined with a pharmaceutically acceptable carrier and/or diluent. The protein, or portions thereof, may be expressed as fusions with other proteins. Combinations of the described
5 proteins, or nucleic acids encoding them, are particularly interesting.

Expression vectors are typically self-replicating DNA or RNA constructs containing the desired receptor gene or its fragments, usually operably linked to suitable genetic control elements that are recognized in a suitable host cell. These control
10 elements are capable of effecting expression within a suitable host. The multiple genes may be coordinately expressed, and may be on a polycistronic message. The specific type of control elements necessary to effect expression will depend upon the eventual host cell used. Generally, the genetic control elements can include a prokaryotic promoter system or a eukaryotic promoter expression control system, and typically include a transcriptional promoter, an optional operator to control the onset of transcription,
15 transcription enhancers to elevate the level of mRNA expression, a sequence that encodes a suitable ribosome binding site, and sequences that terminate transcription and translation. Expression vectors also usually contain an origin of replication that allows the vector to replicate independently of the host cell.

The vectors of this invention include those which contain DNA which encodes a
20 combination of proteins, as described, or a biologically active equivalent polypeptide. The DNA can be under the control of a viral promoter and can encode a selection marker. This invention further contemplates use of such expression vectors which are capable of expressing eukaryotic cDNAs coding for such proteins in a prokaryotic or eukaryotic host, where the vector is compatible with the host and where the eukaryotic cDNAs are
25 inserted into the vector such that growth of the host containing the vector expresses the cDNAs in question. Usually, expression vectors are designed for stable replication in their host cells or for amplification to greatly increase the total number of copies of the desirable gene per cell. It is not always necessary to require that an expression vector replicate in a host cell, e.g., it is possible to effect transient expression of the protein or its
30 fragments in various hosts using vectors that do not contain a replication origin that is recognized by the host cell. It is also possible to use vectors that cause integration of the protein encoding portions into the host DNA by recombination.

Vectors, as used herein, comprise plasmids, viruses, bacteriophage, integratable DNA fragments, and other vehicles which enable the integration of DNA fragments into
35 the genome of the host. Expression vectors are specialized vectors which contain genetic control elements that effect expression of operably linked genes. Plasmids are the most commonly used form of vector but all other forms of vectors which serve an equivalent

function and which are, or become, known in the art are suitable for use herein. See, e.g., Pouwels, et al. (1985 and Supplements) Cloning Vectors: A Laboratory Manual, Elsevier, N.Y., and Rodriguez, et al. (eds. 1988) Vectors: A Survey of Molecular Cloning Vectors and Their Uses, Butterworth, Boston, which are incorporated herein by reference.

5 Transformed cells are cells, preferably mammalian, that have been transformed or transfected with vectors constructed using recombinant DNA techniques. Transformed host cells usually express the desired proteins, but for purposes of cloning, amplifying, and manipulating its DNA, do not need to express the subject proteins. This invention further contemplates culturing transformed cells in a nutrient medium, thus permitting the
10 proteins to accumulate. The proteins can be recovered, either from the culture or, in certain instances, from the culture medium.

For purposes of this invention, nucleic sequences are operably linked when they are functionally related to each other. For example, DNA for a presequence or secretory leader is operably linked to a polypeptide if it is expressed as a preprotein or participates
15 in directing the polypeptide to the cell membrane or in secretion of the polypeptide. A promoter is operably linked to a coding sequence if it controls the transcription of the polypeptide; a ribosome binding site is operably linked to a coding sequence if it is positioned to permit translation. Usually, operably linked means contiguous and in reading frame, however, certain genetic elements such as repressor genes are not
20 contiguously linked but still bind to operator sequences that in turn control expression.

Suitable host cells include prokaryotes, lower eukaryotes, and higher eukaryotes. Prokaryotes include both gram negative and gram positive organisms, e.g., E. coli and B. subtilis. Lower eukaryotes include yeasts, e.g., S. cerevisiae and Pichia, and species of the genus Dictyostelium. Higher eukaryotes include established tissue culture cell lines
25 from animal cells, both of non-mammalian origin, e.g., insect cells, and birds, and of mammalian origin, e.g., human, primates, and rodents.

Prokaryotic host-vector systems include a wide variety of vectors for many different species. As used herein, E. coli and its vectors will be used generically to include equivalent vectors used in other prokaryotes. A representative vector for
30 amplifying DNA is pBR322 or many of its derivatives. Vectors that can be used to express the receptor or its fragments include, but are not limited to, such vectors as those containing the lac promoter (pUC-series); trp promoter (pBR322-trp); Ipp promoter (the pIN-series); lambda-pP or pR promoters (pOTS); or hybrid promoters such as ptac (pDR540). See Brosius, et al. (1988) "Expression Vectors Employing Lambda-, trp-, lac-,
35 and Ipp-derived Promoters", in Vectors: A Survey of Molecular Cloning Vectors and Their Uses, (eds. Rodriguez and Denhardt), Butterworth, Boston, Chapter 10, pp. 205-236, which is incorporated herein by reference.

Lower eukaryotes, e.g., yeasts and Dictyostelium, may be transformed with DCRS8 sequence containing vectors. For purposes of this invention, the most common lower eukaryotic host is the baker's yeast, Saccharomyces cerevisiae. It will be used to generically represent lower eukaryotes although a number of other strains and species are also available. Yeast vectors typically consist of a replication origin (unless of the integrating type), a selection gene, a promoter, DNA encoding the receptor or its fragments, and sequences for translation termination, polyadenylation, and transcription termination. Suitable expression vectors for yeast include such constitutive promoters as 3-phosphoglycerate kinase and various other glycolytic enzyme gene promoters or such inducible promoters as the alcohol dehydrogenase 2 promoter or metallothionein promoter. Suitable vectors include derivatives of the following types: self-replicating low copy number (such as the YRp-series), self-replicating high copy number (such as the YE_p-series); integrating types (such as the YIp-series), or mini-chromosomes (such as the YCp-series).

Higher eukaryotic tissue culture cells are normally the preferred host cells for expression of the functionally active interleukin or receptor proteins. In principle, many higher eukaryotic tissue culture cell lines are workable, e.g., insect baculovirus expression systems, whether from an invertebrate or vertebrate source. However, mammalian cells are preferred. Transformation or transfection and propagation of such cells has become a routine procedure. Examples of useful cell lines include HeLa cells, Chinese hamster ovary (CHO) cell lines, baby rat kidney (BRK) cell lines, insect cell lines, bird cell lines, and monkey (COS) cell lines. Expression vectors for such cell lines usually include an origin of replication, a promoter, a translation initiation site, RNA splice sites (if genomic DNA is used), a polyadenylation site, and a transcription termination site. These vectors also usually contain a selection gene or amplification gene. Suitable expression vectors may be plasmids, viruses, or retroviruses carrying promoters derived, e.g., from such sources as from adenovirus, SV40, parvoviruses, vaccinia virus, or cytomegalovirus. Representative examples of suitable expression vectors include pCDNA1; pCD, see Okayama, et al. (1985) Mol. Cell Biol. 5:1136-1142; pMC1neo PolyA, see Thomas, et al. (1987) Cell 51:503-512; and a baculovirus vector such as pAC 373 or pAC 610.

For secreted proteins and some membrane proteins, an open reading frame usually encodes a polypeptide that consists of a mature or secreted product covalently linked at its N-terminus to a signal peptide. The signal peptide is cleaved prior to secretion of the mature, or active, polypeptide. The cleavage site can be predicted with a high degree of accuracy from empirical rules, e.g., von-Heijne (1986) Nucleic Acids Research 14:4683-4690; and Nielsen, et al. (1997) Protein Eng. 10:1-12, and the precise amino acid composition of the signal peptide often does not appear to be critical to its function, e.g.,

Randall, et al. (1989) Science 243:1156-1159; and Kaiser, et al. (1987) Science 235:312-317. The mature proteins of the invention can be readily determined using standard methods.

5 It will often be desired to express these polypeptides in a system which provides a specific or defined glycosylation pattern. In this case, the usual pattern will be that provided naturally by the expression system. However, the pattern will be modifiable by exposing the polypeptide, e.g., an unglycosylated form, to appropriate glycosylating proteins introduced into a heterologous expression system. For example, the receptor gene may be co-transformed with one or more genes encoding mammalian or other
10 glycosylating enzymes. Using this approach, certain mammalian glycosylation patterns will be achievable in prokaryote or other cells. Expression in prokaryote cells will typically lead to unglycosylated forms of protein.

The source of DCRS8 can be a eukaryotic or prokaryotic host expressing recombinant DCRS8, such as is described above. The source can also be a cell line, but
15 other mammalian cell lines are also contemplated by this invention, with the preferred cell line being from the human species.

Now that the sequences are known, the primate DCRS8 or DCRS9, fragments, or derivatives thereof can be prepared by conventional processes for synthesizing peptides. These include processes such as are described in Stewart and Young (1984) Solid Phase Peptide Synthesis, Pierce Chemical Co., Rockford, IL; Bodanszky and Bodanszky (1984) The Practice of Peptide Synthesis, Springer-Verlag, New York; and Bodanszky (1984) The Principles of Peptide Synthesis, Springer-Verlag, New York; all of each which are incorporated herein by reference. For example, an azide process, an acid chloride process, an acid anhydride process, a mixed anhydride process, an active ester process
25 (for example, p-nitrophenyl ester, N-hydroxysuccinimide ester, or cyanomethyl ester), a carbodiimidazole process, an oxidative-reductive process, or a dicyclohexylcarbodiimide (DCCD)/additive process can be used. Solid phase and solution phase syntheses are both applicable to the foregoing processes. Similar techniques can be used with partial DCRS8 or DCRS9 sequences.

30 The DCRS8 proteins, fragments, or derivatives are suitably prepared in accordance with the above processes as typically employed in peptide synthesis, generally either by a so-called stepwise process which comprises condensing an amino acid to the terminal amino acid, one by one in sequence, or by coupling peptide fragments to the terminal amino acid. Amino groups that are not being used in the coupling reaction
35 typically must be protected to prevent coupling at an incorrect location.

If a solid phase synthesis is adopted, the C-terminal amino acid is bound to an insoluble carrier or support through its carboxyl group. The insoluble carrier is not

particularly limited as long as it has a binding capability to a reactive carboxyl group. Examples of such insoluble carriers include halomethyl resins, such as chloromethyl resin or bromomethyl resin, hydroxymethyl resins, phenol resins, tert-alkyloxycarbonylhydrazidated resins, and the like.

5 An amino group-protected amino acid is bound in sequence through condensation of its activated carboxyl group and the reactive amino group of the previously formed peptide or chain, to synthesize the peptide step by step. After synthesizing the complete sequence, the peptide is split off from the insoluble carrier to produce the peptide. This solid-phase approach is generally described by Merrifield, et al. (1963) in J. Am. Chem.
10 Soc. 85:2149-2156, which is incorporated herein by reference.

 The prepared protein and fragments thereof can be isolated and purified from the reaction mixture by means of peptide separation, e.g., by extraction, precipitation, electrophoresis, various forms of chromatography, and the like. The receptors of this invention can be obtained in varying degrees of purity depending upon desired uses.
15 Purification can be accomplished by use of the protein purification techniques disclosed herein, see below, or by the use of the antibodies herein described in methods of immunoabsorbant affinity chromatography. This immunoabsorbant affinity chromatography is carried out by first linking the antibodies to a solid support and then contacting the linked antibodies with solubilized lysates of appropriate cells, lysates of
20 other cells expressing the receptor, or lysates or supernatants of cells producing the protein as a result of DNA techniques, see below.

 Generally, the purified protein will be at least about 40% pure, ordinarily at least about 50% pure, usually at least about 60% pure, typically at least about 70% pure, more typically at least about 80% pure, preferable at least about 90% pure and more preferably
25 at least about 95% pure, and in particular embodiments, 97%-99% or more. Purity will usually be on a weight basis, but can also be on a molar basis. Different assays will be applied as appropriate. Individual proteins may be purified and thereafter combined.

VI. Antibodies

30 Antibodies can be raised to the various mammalian, e.g., primate DCRS8 or DCRS9 proteins and fragments thereof, both in naturally occurring native forms and in their recombinant forms, the difference being that antibodies to the active receptor are more likely to recognize epitopes which are only present in the native conformations. Denatured antigen detection can also be useful in, e.g., Western analysis. Anti-idiotypic
35 antibodies are also contemplated, which would be useful as agonists or antagonists of a natural receptor or an antibody.

Antibodies, including binding fragments and single chain versions, against predetermined fragments of the protein can be raised by immunization of animals with conjugates of the fragments with immunogenic proteins. Monoclonal antibodies are prepared from cells secreting the desired antibody. These antibodies can be screened for binding to normal or defective protein, or screened for agonistic or antagonistic activity. These monoclonal antibodies will usually bind with at least a K_D of about 1 mM, more usually at least about 300 μ M, typically at least about 100 μ M, more typically at least about 30 μ M, preferably at least about 10 μ M, and more preferably at least about 3 μ M or better.

The antibodies, including antigen binding fragments, of this invention can have significant diagnostic or therapeutic value. They can be potent antagonists that bind to the receptor and inhibit binding to ligand or inhibit the ability of the receptor to elicit a biological response, e.g., act on its substrate. They also can be useful as non-neutralizing antibodies and can be coupled to toxins or radionuclides to bind producing cells, or cells localized to the source of the interleukin. Further, these antibodies can be conjugated to drugs or other therapeutic agents, either directly or indirectly by means of a linker.

The antibodies of this invention can also be useful in diagnostic applications. As capture or non-neutralizing antibodies, they might bind to the receptor without inhibiting ligand or substrate binding. As neutralizing antibodies, they can be useful in competitive binding assays. They will also be useful in detecting or quantifying ligand. They may be used as reagents for Western blot analysis, or for immunoprecipitation or immunopurification of the respective protein. Likewise, nucleic acids and proteins may be immobilized to solid substrates for affinity purification or detection methods. The substrates may be, e.g., solid resin beads or sheets of plastic.

Protein fragments may be joined to other materials, particularly polypeptides, as fused or covalently joined polypeptides to be used as immunogens. Mammalian cytokine receptors and fragments may be fused or covalently linked to a variety of immunogens, such as keyhole limpet hemocyanin, bovine serum albumin, tetanus toxoid, etc. See (1969) Microbiology, Hoeber Medical Division, Harper and Row; Landsteiner (1962) Specificity of Serological Reactions, Dover Publications, New York; and Williams, et al. (1967) Methods in Immunology and Immunochemistry, Vol. 1, Academic Press, New York; each of which is incorporated herein by reference, for descriptions of methods of preparing polyclonal antisera. A typical method involves hyperimmunization of an animal with an antigen. The blood of the animal is then collected shortly after the repeated immunizations and the gamma globulin is isolated.

In some instances, it is desirable to prepare monoclonal antibodies from various mammalian hosts, such as mice, rodents, primates, humans, etc. Description of

techniques for preparing such monoclonal antibodies may be found in, e.g., Stites, et al. (eds.) Basic and Clinical Immunology (4th ed.), Lange Medical Publications, Los Altos, CA, and references cited therein; Harlow and Lane (1988) Antibodies: A Laboratory Manual, CSH Press; Goding (1986) Monoclonal Antibodies: Principles and Practice (2d ed.) Academic Press, New York; and particularly in Kohler and Milstein (1975) Nature 256:495-497, which discusses one method of generating monoclonal antibodies. Each of these references is incorporated herein by reference. Summarized briefly, this method involves injecting an animal with an immunogen. The animal is then sacrificed and cells taken from its spleen, which are then fused with myeloma cells. The result is a hybrid cell or "hybridoma" that is capable of reproducing in vitro. The population of hybridomas is then screened to isolate individual clones, each of which secrete a single antibody species to the immunogen. In this manner, the individual antibody species obtained are the products of immortalized and cloned single B cells from the immune animal generated in response to a specific site recognized on the immunogenic substance.

Other suitable techniques involve in vitro exposure of lymphocytes to the antigenic polypeptides or alternatively to selection of libraries of antibodies in phage or similar vectors. See, Huse, et al. (1989) "Generation of a Large Combinatorial Library of the Immunoglobulin Repertoire in Phage Lambda," Science 246:1275-1281; and Ward, et al. (1989) Nature 341:544-546, each of which is incorporated herein by reference. The polypeptides and antibodies of the present invention may be used with or without modification, including chimeric or humanized antibodies. Frequently, the polypeptides and antibodies will be labeled by joining, either covalently or non-covalently, a substance which provides for a detectable signal. A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent moieties, chemiluminescent moieties, magnetic particles, and the like. Patents, teaching the use of such labels include U.S. Patent Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241. Also, recombinant or chimeric immunoglobulins may be produced, see Cabilly, U.S. Patent No. 4,816,567; or made in transgenic mice, see Mendez, et al. (1997) Nature Genetics 15:146-156; Abgenix; and Medarex. These references are incorporated herein by reference.

The antibodies of this invention can also be used for affinity chromatography in isolating the DCRS8 proteins or peptides. Columns can be prepared where the antibodies are linked to a solid support, e.g., particles, such as agarose, Sephadex, or the like, where a cell lysate may be passed through the column, the column washed, followed by increasing concentrations of a mild denaturant, whereby the purified protein will be

released. Alternatively, the protein may be used to purify antibody. Appropriate cross absorptions or depletions may be applied.

5 The antibodies may also be used to screen expression libraries for particular expression products. Usually the antibodies used in such a procedure will be labeled with a moiety allowing easy detection of presence of antigen by antibody binding.

Antibodies raised against a cytokine receptor will also be used to raise anti-idiotypic antibodies. These will be useful in detecting or diagnosing various immunological conditions related to expression of the protein or cells which express the protein. They also will be useful as agonists or antagonists of the ligand, which may be competitive inhibitors or substitutes for naturally occurring ligands.

10 A cytokine receptor protein that specifically binds to or that is specifically immunoreactive with an antibody generated against a defined immunogen, such as an immunogen consisting of the amino acid sequence of SEQ ID NO: 14, is typically determined in an immunoassay. The immunoassay typically uses a polyclonal antiserum which was raised, e.g., to a protein of SEQ ID NO: 14. This antiserum is selected to have low crossreactivity against other cytokine receptor family members, preferably from the same species, and any such crossreactivity is removed by immunoabsorption prior to use in the immunoassay.

15 In order to produce antisera for use in an immunoassay, the protein, e.g., of SEQ ID NO: 14, is isolated as described herein. For example, recombinant protein may be produced in a mammalian cell line. An appropriate host, e.g., an inbred strain of mice such as Balb/c, is immunized with the selected protein, typically using a standard adjuvant, such as Freund's adjuvant, and a standard mouse immunization protocol (see Harlow and Lane, supra). Alternatively, a synthetic peptide derived from the sequences disclosed herein and conjugated to a carrier protein can be used as an immunogen.

25 Polyclonal sera are collected and titered against the immunogen protein in an immunoassay, e.g., a solid phase immunoassay with the immunogen immobilized on a solid support. Polyclonal antisera with a titer of 10^4 or greater are selected and tested for their cross reactivity against other cytokine receptor family members using a competitive binding immunoassay such as the one described in Harlow and Lane, supra, at pages 570-573. Preferably at least two cytokine receptor family members are used in this determination. These cytokine receptor family members can be produced as recombinant proteins and isolated using standard molecular biology and protein chemistry techniques as described herein.

30 Immunoassays in the competitive binding format can be used for the crossreactivity determinations. For example, the protein of SEQ ID NO: 14 can be immobilized to a solid support. Proteins added to the assay compete with the binding of

the antisera to the immobilized antigen. The ability of the above proteins to compete with the binding of the antisera to the immobilized protein is compared to the other proteins. The percent crossreactivity for the above proteins is calculated, using standard calculations. Those antisera with less than 10% crossreactivity with each of the proteins listed above are selected and pooled. The cross-reacting antibodies are then removed from the pooled antisera by immunoabsorption with the above-listed proteins.

The immunoabsorbed and pooled antisera are then used in a competitive binding immunoassay as described above to compare a second protein to the immunogen protein (e.g., the DCRS8 like protein of SEQ ID NO: 14). In order to make this comparison, the two proteins are each assayed at a wide range of concentrations and the amount of each protein required to inhibit 50% of the binding of the antisera to the immobilized protein is determined. If the amount of the second protein required is less than twice the amount of the protein of the selected protein or proteins that is required, then the second protein is said to specifically bind to an antibody generated to the immunogen.

It is understood that these cytokine receptor proteins are members of a family of homologous proteins that comprise at least 9 so far identified members, 6 mammalian and 3 worm embodiments. For a particular gene product, such as the DCRS8, the term refers not only to the amino acid sequences disclosed herein, but also to other proteins that are allelic, non-allelic, or species variants. It is also understood that the terms include nonnatural mutations introduced by deliberate mutation using conventional recombinant technology such as single site mutation, or by excising short sections of DNA encoding the respective proteins, or by substituting new amino acids, or adding new amino acids. Such minor alterations typically will substantially maintain the immunoidentity of the original molecule and/or its biological activity. Thus, these alterations include proteins that are specifically immunoreactive with a designated naturally occurring DCRS8 protein. The biological properties of the altered proteins can be determined by expressing the protein in an appropriate cell line and measuring the appropriate effect, e.g., upon transfected lymphocytes. Particular protein modifications considered minor would include conservative substitution of amino acids with similar chemical properties, as described above for the cytokine receptor family as a whole. By aligning a protein optimally with the protein of the cytokine receptors and by using the conventional immunoassays described herein to determine immunoidentity, one can determine the protein compositions of the invention.

VII. Kits and quantitation

Both naturally occurring and recombinant forms of the cytokine receptor like molecules of this invention are particularly useful in kits and assay methods. For

example, these methods would also be applied to screening for binding activity, e.g., ligands for these proteins. Several methods of automating assays have been developed in recent years so as to permit screening of tens of thousands of compounds per year. See, e.g., a BIOMEK automated workstation, Beckman Instruments, Palo Alto, California, and
5 Fodor, et al. (1991) Science 251:767-773, which is incorporated herein by reference. The latter describes means for testing binding by a plurality of defined polymers synthesized on a solid substrate. The development of suitable assays to screen for a ligand or agonist/antagonist homologous proteins can be greatly facilitated by the availability of large amounts of purified, soluble cytokine receptors in an active state such as is provided
10 by this invention.

Purified protein can be coated directly onto plates for use in the aforementioned ligand screening techniques. However, non-neutralizing antibodies to these proteins can be used as capture antibodies to immobilize the respective receptor on the solid phase, useful, e.g., in diagnostic uses.

15 This invention also contemplates use of receptor subunit, fragments thereof, peptides, and their fusion products in a variety of diagnostic kits and methods for detecting the presence of the protein or its ligand. Alternatively, or additionally, antibodies against the molecules may be incorporated into the kits and methods. Typically the kit will have a compartment containing, e.g., a DCRS8 peptide or gene
20 segment or a reagent which recognizes one or the other. Typically, recognition reagents, in the case of peptide, would be a receptor or antibody, or in the case of a gene segment, would usually be a hybridization probe.

A preferred kit for determining the concentration of DCRS8 in a sample would typically comprise a labeled compound, e.g., ligand or antibody, having known binding
25 affinity for DCRS8, a source of DCRS8 (naturally occurring or recombinant) as a positive control, and a means for separating the bound from free labeled compound, e.g., a solid phase for immobilizing the DCRS8 in the test sample. Compartments containing reagents, and instructions, will normally be provided. Appropriate nucleic acid or protein containing kits are also provided.

30 Antibodies, including antigen binding fragments, specific for mammalian DCRS8 or a peptide fragment, or receptor fragments are useful in diagnostic applications to detect the presence of elevated levels of ligand and/or its fragments. Diagnostic assays may be homogeneous (without a separation step between free reagent and antibody-antigen complex) or heterogeneous (with a separation step). Various commercial assays exist,
35 such as radioimmunoassay (RIA), enzyme-linked immunosorbent assay (ELISA), enzyme immunoassay (EIA), enzyme-multiplied immunoassay technique (EMIT), substrate-labeled fluorescent immunoassay (SLFIA) and the like. For example, unlabeled

antibodies can be employed by using a second antibody which is labeled and which recognizes the antibody to a cytokine receptor or to a particular fragment thereof. These assays have also been extensively discussed in the literature. See, e.g., Harlow and Lane (1988) Antibodies: A Laboratory Manual, CSH, and Coligan (ed. 1991 and periodic supplements) Current Protocols In Immunology Greene/Wiley, New York.

Anti-idiotypic antibodies may have similar use to serve as agonists or antagonists of cytokine receptors. These should be useful as therapeutic reagents under appropriate circumstances.

Frequently, the reagents for diagnostic assays are supplied in kits, so as to optimize the sensitivity of the assay. For the subject invention, depending upon the nature of the assay, the protocol, and the label, either labeled or unlabeled antibody, or labeled ligand is provided. This is usually in conjunction with other additives, such as buffers, stabilizers, materials necessary for signal production such as substrates for enzymes, and the like. Preferably, the kit will also contain instructions for proper use and disposal of the contents after use. Typically the kit has compartments for each useful reagent, and will contain instructions for proper use and disposal of reagents. Desirably, the reagents are provided as a dry lyophilized powder, where the reagents may be reconstituted in an aqueous medium having appropriate concentrations for performing the assay.

The aforementioned constituents of the diagnostic assays may be used without modification or may be modified in a variety of ways. For example, labeling may be achieved by covalently or non-covalently joining a moiety which directly or indirectly provides a detectable signal. In many of these assays, a test compound, cytokine receptor, or antibodies thereto can be labeled either directly or indirectly. Possibilities for direct labeling include label groups: radiolabels such as ^{125}I , enzymes (U.S. Pat. No. 3,645,090) such as peroxidase and alkaline phosphatase, and fluorescent labels (U.S. Pat. No. 3,940,475) capable of monitoring the change in fluorescence intensity, wavelength shift, or fluorescence polarization. Both of the patents are incorporated herein by reference. Possibilities for indirect labeling include biotinylation of one constituent followed by binding to avidin coupled to one of the above label groups.

There are also numerous methods of separating the bound from the free ligand, or alternatively the bound from the free test compound. The cytokine receptor can be immobilized on various matrixes followed by washing. Suitable matrices include plastic such as an ELISA plate, filters, and beads. Methods of immobilizing the receptor to a matrix include, without limitation, direct adhesion to plastic, use of a capture antibody, chemical coupling, and biotin-avidin. The last step in this approach involves the precipitation of antibody/antigen complex by any of several methods including those

utilizing, e.g., an organic solvent such as polyethylene glycol or a salt such as ammonium sulfate. Other suitable separation techniques include, without limitation, the fluorescein antibody magnetizable particle method described in Rattle, et al. (1984) Clin. Chem. 30(9):1457-1461, and the double antibody magnetic particle separation as described in
5 U.S. Pat. No. 4,659,678, each of which is incorporated herein by reference.

The methods for linking protein or fragments to various labels have been extensively reported in the literature and do not require detailed discussion here. Many of the techniques involve the use of activated carboxyl groups either through the use of carbodiimide or active esters to form peptide bonds, the formation of thioethers by
10 reaction of a mercapto group with an activated halogen such as chloroacetyl, or an activated olefin such as maleimide, for linkage, or the like. Fusion proteins will also find use in these applications.

Another diagnostic aspect of this invention involves use of oligonucleotide or polynucleotide sequences taken from the sequence of an cytokine receptor. These
15 sequences can be used as probes for detecting levels of the respective cytokine receptor in patients suspected of having an immunological disorder. The preparation of both RNA and DNA nucleotide sequences, the labeling of the sequences, and the preferred size of the sequences has received ample description and discussion in the literature. Normally an oligonucleotide probe should have at least about 14 nucleotides, usually at least about
20 18 nucleotides, and the polynucleotide probes may be up to several kilobases. Various labels may be employed, most commonly radionuclides, particularly ^{32}P . However, other techniques may also be employed, such as using biotin modified nucleotides for introduction into a polynucleotide. The biotin then serves as the site for binding to avidin or antibodies, which may be labeled with a wide variety of labels, such as radionuclides,
25 fluorescers, enzymes, or the like. Alternatively, antibodies may be employed which can recognize specific duplexes, including DNA duplexes, RNA duplexes, DNA-RNA hybrid duplexes, or DNA-protein duplexes. The antibodies in turn may be labeled and the assay carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected. The use of
30 probes to the novel RNA may be carried out in conventional techniques such as nucleic acid hybridization, plus and minus screening, recombinational probing, hybrid released translation (HRT), and hybrid arrested translation (HART). Antisense nucleic acids, which may be used to block protein expression, are also provided. See, e.g., Isis Pharmaceuticals, Sequitur, Inc., or Hybridon. This also includes amplification techniques
35 such as polymerase chain reaction (PCR).

Diagnostic kits which also test for the qualitative or quantitative presence of other markers are also contemplated. Diagnosis or prognosis may depend on the combination

of multiple indications used as markers. Thus, kits may test for combinations of markers. See, e.g., Viallet, et al. (1989) Progress in Growth Factor Res. 1:89-97.

VIII. Therapeutic Utility

5 This invention provides reagents with significant therapeutic value. See, e.g., Levitzki (1996) Curr. Opin. Cell Biol. 8:239-244. The cytokine receptors (naturally occurring or recombinant), fragments thereof, mutein receptors, and antibodies, along with compounds identified as having binding affinity to the receptors or antibodies, should be useful in the treatment of conditions exhibiting abnormal expression of the
10 receptors of their ligands. Such abnormality will typically be manifested by immunological disorders, e.g., innate immunity, or developmentally. Additionally, this invention should provide therapeutic value in various diseases or disorders associated with abnormal expression or abnormal triggering of response to the ligand. For example, the IL-1 ligands have been suggested to be involved in morphologic development, e.g.,
15 dorso-ventral polarity determination, and immune responses, particularly the primitive innate responses. See, e.g., Sun, et al. (1991) Eur. J. Biochem. 196:247-254; and Hultmark (1994) Nature 367:116-117.

 Recombinant cytokine receptors, muteins, agonist or antagonist antibodies thereto, or antibodies can be purified and then administered to a patient. These reagents can be
20 combined for therapeutic use with additional active ingredients, e.g., in conventional pharmaceutically acceptable carriers or diluents, along with physiologically innocuous stabilizers and excipients. These combinations can be sterile, e.g., filtered, and placed into dosage forms as by lyophilization in dosage vials or storage in stabilized aqueous preparations. This invention also contemplates use of antibodies or binding fragments
25 thereof which are not complement binding.

 Ligand screening using cytokine receptor or fragments thereof can be performed to identify molecules having binding affinity to the receptors. Subsequent biological assays can then be utilized to determine if a putative ligand can provide competitive binding, which can block intrinsic stimulating activity. Receptor fragments can be used
30 as a blocker or antagonist in that it blocks the activity of ligand. Likewise, a compound having intrinsic stimulating activity can activate the receptor and is thus an agonist in that it simulates the activity of ligand, e.g., inducing signaling. This invention further contemplates the therapeutic use of antibodies to cytokine receptors as antagonists.

 The quantities of reagents necessary for effective therapy will depend upon many
35 different factors, including means of administration, target site, reagent physiological life, pharmacological life, physiological state of the patient, and other medicants administered. Thus, treatment dosages should be titrated to optimize safety and efficacy. Typically,

dosages used in vitro may provide useful guidance in the amounts useful for in situ administration of these reagents. Animal testing of effective doses for treatment of particular disorders will provide further predictive indication of human dosage. Various considerations are described, e.g., in Gilman, et al. (eds. 1990) Goodman and Gilman's: The Pharmacological Bases of Therapeutics, 8th Ed., Pergamon Press; and Remington's Pharmaceutical Sciences, 17th ed. (1990), Mack Publishing Co., Easton, Penn.; each of which is hereby incorporated herein by reference. Methods for administration are discussed therein and below, e.g., for oral, intravenous, intraperitoneal, or intramuscular administration, transdermal diffusion, and others. Pharmaceutically acceptable carriers will include water, saline, buffers, and other compounds described, e.g., in the Merck Index, Merck & Co., Rahway, New Jersey. Because of the likely high affinity binding, or turnover numbers, between a putative ligand and its receptors, low dosages of these reagents would be initially expected to be effective. And the signaling pathway suggests extremely low amounts of ligand may have effect. Thus, dosage ranges would ordinarily be expected to be in amounts lower than 1 mM concentrations, typically less than about 10 μ M concentrations, usually less than about 100 nM, preferably less than about 10 pM (picomolar), and most preferably less than about 1 fM (femtomolar), with an appropriate carrier. Slow release formulations, or slow release apparatus will often be utilized for continuous administration.

Cytokine receptors, fragments thereof, and antibodies or its fragments, antagonists, and agonists, may be administered directly to the host to be treated or, depending on the size of the compounds, it may be desirable to conjugate them to carrier proteins such as ovalbumin or serum albumin prior to their administration. Therapeutic formulations may be administered in many conventional dosage formulations. While it is possible for the active ingredient to be administered alone, it is preferable to present it as a pharmaceutical formulation. Formulations comprise at least one active ingredient, as defined above, together with one or more acceptable carriers thereof. Each carrier must be both pharmaceutically and physiologically acceptable in the sense of being compatible with the other ingredients and not injurious to the patient. Formulations include those suitable for oral, rectal, nasal, or parenteral (including subcutaneous, intramuscular, intravenous and intradermal) administration. The formulations may conveniently be presented in unit dosage form and may be prepared by methods well known in the art of pharmacy. See, e.g., Gilman, et al. (eds. 1990) Goodman and Gilman's: The Pharmacological Bases of Therapeutics, 8th Ed., Pergamon Press; and Remington's Pharmaceutical Sciences, 17th ed. (1990), Mack Publishing Co., Easton, Penn.; Avis, et al. (eds. 1993) Pharmaceutical Dosage Forms: Parenteral Medications Dekker, NY; Lieberman, et al. (eds. 1990) Pharmaceutical Dosage Forms: Tablets Dekker, NY; and

Lieberman, et al. (eds. 1990) Pharmaceutical Dosage Forms: Disperse Systems Dekker, NY. The therapy of this invention may be combined with or used in association with other therapeutic agents, particularly agonists or antagonists of other cytokine receptor family members.

5

IX. Screening

Drug screening using DCRS8 or fragments thereof can be performed to identify compounds having binding affinity to the receptor subunit, including isolation of associated components. Subsequent biological assays can then be utilized to determine if the compound has intrinsic stimulating activity and is therefore a blocker or antagonist in that it blocks the activity of the ligand. Likewise, a compound having intrinsic stimulating activity can activate the receptor and is thus an agonist in that it simulates the activity of a cytokine ligand. This invention further contemplates the therapeutic use of antibodies to the receptor as cytokine agonists or antagonists.

10

Similarly, complexes comprising multiple proteins may be used to screen for ligands or reagents capable of recognizing the complex. Most cytokine receptors comprise at least two subunits, which may be the same, or distinct. Alternatively, the transmembrane receptor may bind to a complex comprising a cytokine-like ligand associated with another soluble protein serving, e.g., as a second receptor subunit.

15

One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant DNA molecules expressing the DCRS8 in combination with another cytokine receptor subunit. Cells may be isolated which express a receptor in isolation from other functional receptors. Such cells, either in viable or fixed form, can be used for standard antibody/antigen or ligand/receptor binding assays. See also, Parce, et al. (1989) Science 246:243-247; and Owicki, et al. (1990) Proc. Nat'l Acad. Sci. USA 87:4007-4011, which describe sensitive methods to detect cellular responses. Competitive assays are particularly useful, where the cells (source of putative ligand) are contacted and incubated with a labeled receptor or antibody having known binding affinity to the ligand, such as ¹²⁵I-antibody, and a test sample whose binding affinity to the binding composition is being measured. The bound and free labeled binding compositions are then separated to assess the degree of ligand binding. The amount of test compound bound is inversely proportional to the amount of labeled receptor binding to the known source. Many techniques can be used to separate bound from free ligand to assess the degree of ligand binding. This separation step could typically involve a procedure such as adhesion to filters followed by washing, adhesion to plastic followed by washing, or centrifugation of the cell membranes. Viable cells could also be used to screen for the effects of drugs on cytokine mediated functions, e.g., second messenger

20

25

30

35

levels, e.g., Ca^{++} ; cell proliferation; inositol phosphate pool changes; and others. Some detection methods allow for elimination of a separation step, e.g., a proximity sensitive detection system. Calcium sensitive dyes will be useful for detecting Ca^{++} levels, with a fluorimeter or a fluorescence cell sorting apparatus.

5

X. Ligands

The descriptions of the DCRS8 herein provides means to identify ligands, as described above. Such ligand should bind specifically to the respective receptor with reasonably high affinity. Various constructs are made available which allow either labeling of the receptor to detect its ligand. For example, directly labeling cytokine receptor, fusing onto it markers for secondary labeling, e.g., FLAG or other epitope tags, etc., will allow detection of receptor. This can be histological, as an affinity method for biochemical purification, or labeling or selection in an expression cloning approach. A two-hybrid selection system may also be applied making appropriate constructs with the available cytokine receptor sequences. See, e.g., Fields and Song (1989) Nature 340:245-246.

10

15

Most likely candidates will be structurally related to members of the IL-17 family. See, e.g., USSN 09/480,287.

The broad scope of this invention is best understood with reference to the following examples, which are not intended to limit the inventions to the specific embodiments.

20

EXAMPLES

I. General Methods

25

Some of the standard methods are described or referenced, e.g., in Maniatis, et al. (1982) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor Press; Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual, (2d ed.), vols. 1-3, CSH Press, NY; or Ausubel, et al. (1987 and Supplements) Current Protocols in Molecular Biology, Greene/Wiley, New York. Methods for protein purification include such methods as ammonium sulfate precipitation, column chromatography, electrophoresis, centrifugation, crystallization, and others. See, e.g., Ausubel, et al. (1987 and periodic supplements); Coligan, et al. (ed. 1996) and periodic supplements, Current Protocols In Protein Science Greene/Wiley, New York; Deutscher (1990) "Guide to Protein Purification" in Methods in Enzymology, vol. 182, and other volumes in this series; and manufacturer's literature on use of protein purification products, e.g., Pharmacia, Piscataway, N.J., or Bio-Rad, Richmond, CA. Combination

30

35

with recombinant techniques allow fusion to appropriate segments, e.g., to a FLAG sequence or an equivalent which can be fused via a protease-removable sequence. See, e.g., Hochuli (1990) "Purification of Recombinant Proteins with Metal Chelate Absorbent" in Setlow (ed.) Genetic Engineering, Principle and Methods 12:87-98, Plenum Press, N.Y.; and Crowe, et al. (1992) QIAexpress: The High Level Expression & Protein Purification System QUIAGEN, Inc., Chatsworth, CA.

Computer sequence analysis is performed, e.g., using available software programs, including those from the GCG (U. Wisconsin) and GenBank sources. Public sequence databases were also used, e.g., from GenBank and others.

Many techniques applicable to IL-10 receptors may be applied to the DCRSs, as described, e.g., in USSN 08/110,683 (IL-10 receptor), which is incorporated herein by reference.

II. Computational Analysis

Human sequences related to cytokine receptors were identified from genomic sequence database using, e.g., the BLAST server (Altschul, et al. (1994) Nature Genet. 6:119-129). Standard analysis programs may be used to evaluate structure, e.g., PHD (Rost and Sander (1994) Proteins 19:55-72) and DSC (King and Sternberg (1996) Protein Sci. 5:2298-2310). Standard comparison software includes, e.g., Altschul, et al. (1990) J. Mol. Biol. 215:403-10; Waterman (1995) Introduction to Computational Biology: Maps, Sequences, and Genomes Chapman & Hall; Lander and Waterman (eds. 1995) Calculating the Secrets of Life: Applications of the Mathematical Sciences in Molecular Biology National Academy Press; and Speed and Waterman (eds. 1996) Genetic Mapping and DNA Sequencing (IMA Volumes in Mathematics and Its Applications, Vol 81) Springer Verlag. Each reference is incorporate herein by reference.

III. Cloning of full-length cDNAs; Chromosomal localization

PCR primers derived from the sequences are used to probe a human cDNA library. Sequences may be derived, e.g., from Tables 1-5, preferably those adjacent the ends of sequences. Full length cDNAs for primate, rodent, or other species DCRS8 are cloned, e.g., by DNA hybridization screening of λ gt10 phage. PCR reactions are conducted using T. aquaticus Taqplus DNA polymerase (Stratagene) under appropriate conditions. Extending partial length cDNA clones is typically routine.

Chromosome spreads are prepared. In situ hybridization is performed on chromosome preparations obtained from phytohemagglutinin-stimulated human lymphocytes cultured for 72 h. 5-bromodeoxyuridine was added for the final seven hours

of culture (60 µg/ml of medium), to ensure a posthybridization chromosomal banding of good quality.

A PCR fragment, amplified with the help of primers, is cloned into an appropriate vector. The vector is labeled by nick-translation with ^3H . The radiolabeled probe is
5 hybridized to metaphase spreads at final concentration of 200 ng/ml of hybridization solution as described, e.g., in Mattei, et al. (1985) Hum. Genet. 69:327-331.

After coating with nuclear track emulsion (KODAK NTB₂), slides are exposed. To avoid any slipping of silver grains during the banding procedure, chromosome spreads are first stained with buffered Giemsa solution and metaphase photographed. R-banding
10 is then performed by the fluorochrome-photolysis-Giemsa (FPG) method and metaphases rephotographed before analysis.

Similar appropriate methods are used for other species.

IV. Localization of mRNA

15 Human multiple tissue (Cat# 1, 2) and cancer cell line blots (Cat# 7757-1), containing approximately 2 µg of poly(A)⁺ RNA per lane, are purchased from Clontech (Palo Alto, CA). Probes are radiolabeled with [α - ^{32}P] dATP, e.g., using the Amersham Rediprime random primer labeling kit (RPN1633). Prehybridization and hybridizations are performed, e.g., at 65° C in 0.5 M Na₂HPO₄, 7% SDS, 0.5 M EDTA (pH 8.0). High
20 stringency washes are conducted, e.g., at 65° C with two initial washes in 2 x SSC, 0.1% SDS for 40 min followed by a subsequent wash in 0.1 x SSC, 0.1% SDS for 20 min. Membranes are then exposed at -70° C to X-Ray film (Kodak) in the presence of intensifying screens. More detailed studies by cDNA library Southern blots are performed with selected appropriate human DCRS clones to examine their expression in
25 hemopoietic or other cell subsets.

Alternatively, two appropriate primers are selected from Tables 1-5. RT-PCR is used on an appropriate mRNA sample selected for the presence of message to produce a cDNA, e.g., a sample which expresses the gene.

30 Full length clones may be isolated by hybridization of cDNA libraries from appropriate tissues pre-selected by PCR signal. Northern blots can be performed.

Message for genes encoding DCRS will be assayed by appropriate technology, e.g., PCR, immunoassay, hybridization, or otherwise. Tissue and organ cDNA preparations are available, e.g., from Clontech, Mountain View, CA. Identification of sources of natural expression are useful, as described. And the identification of functional
35 receptor subunit pairings will allow for prediction of what cells express the combination of receptor subunits which will result in a physiological responsiveness to each of the cytokine ligands.

For mouse counterpart distribution, e.g., Southern Analysis can be performed: DNA (5 µg) from a primary amplified cDNA library was digested with appropriate restriction enzymes to release the inserts, run on a 1% agarose gel and transferred to a nylon membrane (Schleicher and Schuell, Keene, NH).

- 5 Samples for mouse mRNA isolation may include: resting mouse fibroblastic L cell line (C200); Braf:ER (Braf fusion to estrogen receptor) transfected cells, control (C201); T cells, TH1 polarized (Mel14 bright, CD4+ cells from spleen, polarized for 7 days with IFN-γ and anti IL-4; T200); T cells, TH2 polarized (Mel14 bright, CD4+ cells from spleen, polarized for 7 days with IL-4 and anti-IFN-γ; T201); T cells, highly TH1 polarized (see Openshaw, et al. (1995) J. Exp. Med. 182:1357-1367; activated with anti-CD3 for 2, 6, 16 h pooled; T202); T cells, highly TH2 polarized (see Openshaw, et al. (1995) J. Exp. Med. 182:1357-1367; activated with anti-CD3 for 2, 6, 16 h pooled; T203); CD44- CD25+ pre T cells, sorted from thymus (T204); TH1 T cell clone D1.1, resting for 3 weeks after last stimulation with antigen (T205); TH1 T cell clone D1.1, 10
10 µg/ml ConA stimulated 15 h (T206); TH2 T cell clone CDC35, resting for 3 weeks after last stimulation with antigen (T207); TH2 T cell clone CDC35, 10 µg/ml ConA stimulated 15 h (T208); Mel14+ naive T cells from spleen, resting (T209); Mel14+ T cells, polarized to Th1 with IFN-γ/IL-12/anti-IL-4 for 6, 12, 24 h pooled (T210); Mel14+ T cells, polarized to Th2 with IL-4/anti-IFN-γ for 6, 13, 24 h pooled (T211); unstimulated
15 mature B cell leukemia cell line A20 (B200); unstimulated B cell line CH12 (B201); unstimulated large B cells from spleen (B202); B cells from total spleen, LPS activated (B203); metrizamide enriched dendritic cells from spleen, resting (D200); dendritic cells from bone marrow, resting (D201); monocyte cell line RAW 264.7 activated with LPS 4 h (M200); bone-marrow macrophages derived with GM and M-CSF (M201); macrophage
20 cell line J774, resting (M202); macrophage cell line J774 + LPS + anti-IL-10 at 0.5, 1, 3, 6, 12 h pooled (M203); macrophage cell line J774 + LPS + IL-10 at 0.5, 1, 3, 5, 12 h pooled (M204); aerosol challenged mouse lung tissue, Th2 primers, aerosol OVA challenge 7, 14, 23 h pooled (see Garlisi, et al. (1995) Clinical Immunology and Immunopathology 75:75-83; X206); Nippostrongylus-infected lung tissue (see Coffman, et al. (1989) Science 245:308-310; X200); total adult lung, normal (O200); total lung, rag-1 (see Schwarz, et al. (1993) Immunodeficiency 4:249-252; O205); IL-10 K.O. spleen (see Kuhn, et al. (1991) Cell 75:263-274; X201); total adult spleen, normal (O201); total spleen, rag-1 (O207); IL-10 K.O. Peyer's patches (O202); total Peyer's patches, normal (O210); IL-10 K.O. mesenteric lymph nodes (X203); total mesenteric lymph nodes,
25 normal (O211); IL-10 K.O. colon (X203); total colon, normal (O212); NOD mouse pancreas (see Makino, et al. (1980) Jikken Dobutsu 29:1-13; X205); total thymus, rag-1 (O208); total kidney, rag-1 (O209); total heart, rag-1 (O202); total brain, rag-1 (O203);
30
35

total testes, rag-1 (O204); total liver, rag-1 (O206); rat normal joint tissue (O300); and rat arthritic joint tissue (X300).

Samples for human mRNA isolation may include, e.g.: peripheral blood mononuclear cells (monocytes, T cells, NK cells, granulocytes, B cells), resting (T100);

5 peripheral blood mononuclear cells, activated with anti-CD3 for 2, 6, 12 h pooled (T101); T cell, TH0 clone Mot 72, resting (T102); T cell, TH0 clone Mot 72, activated with anti-CD28 and anti-CD3 for 3, 6, 12 h pooled (T103); T cell, TH0 clone Mot 72, anergic treated with specific peptide for 2, 7, 12 h pooled (T104); T cell, TH1 clone HY06, resting (T107); T cell, TH1 clone HY06, activated with anti-CD28 and anti-CD3 for 3, 6,

10 12 h pooled (T108); T cell, TH1 clone HY06, anergic treated with specific peptide for 2, 6, 12 h pooled (T109); T cell, TH2 clone HY935, resting (T110); T cell, TH2 clone HY935, activated with anti-CD28 and anti-CD3 for 2, 7, 12 h pooled (T111); T cells CD4+CD45RO- T cells polarized 27 days in anti-CD28, IL-4, and anti IFN- γ , TH2 polarized, activated with anti-CD3 and anti-CD28 4 h (T116); T cell tumor lines Jurkat and Hut78, resting (T117); T cell clones, pooled AD130.2, Tc783.12, Tc783.13,

15 Tc783.58, Tc782.69, resting (T118); T cell random $\gamma\delta$ T cell clones, resting (T119); Splenocytes, resting (B100); Splenocytes, activated with anti-CD40 and IL-4 (B101); B cell EBV lines pooled WT49, RSB, JY, CVIR, 721.221, RM3, HSY, resting (B102); B cell line JY, activated with PMA and ionomycin for 1, 6 h pooled (B103); NK 20 clones pooled, resting (K100); NK 20 clones pooled, activated with PMA and ionomycin for 6 h (K101); NKL clone, derived from peripheral blood of LGL leukemia patient, IL-2 treated (K106); NK cytotoxic clone 640-A30-1, resting (K107); hematopoietic precursor line TF1, activated with PMA and ionomycin for 1, 6 h pooled (C100); U937 premonocytic line, resting (M100); U937 premonocytic line, activated with PMA and ionomycin for 1,

20 6 h pooled (M101); elutriated monocytes, activated with LPS, IFN γ , anti-IL-10 for 1, 2, 6, 12, 24 h pooled (M102); elutriated monocytes, activated with LPS, IFN γ , IL-10 for 1, 2, 6, 12, 24 h pooled (M103); elutriated monocytes, activated with LPS, IFN γ , anti-IL-10 for 4, 16 h pooled (M106); elutriated monocytes, activated with LPS, IFN γ , IL-10 for 4, 16 h pooled (M107); elutriated monocytes, activated LPS for 1 h (M108); elutriated

30 monocytes, activated LPS for 6 h (M109); DC 70% CD1a+, from CD34+ GM-CSF, TNF α 12 days, resting (D101); DC 70% CD1a+, from CD34+ GM-CSF, TNF α 12 days, activated with PMA and ionomycin for 1 hr (D102); DC 70% CD1a+, from CD34+ GM-CSF, TNF α 12 days, activated with PMA and ionomycin for 6 hr (D103); DC 95% CD1a+, from CD34+ GM-CSF, TNF α 12 days FACS sorted, activated with PMA and

35 ionomycin for 1, 6 h pooled (D104); DC 95% CD14+, ex CD34+ GM-CSF, TNF α 12 days FACS sorted, activated with PMA and ionomycin 1, 6 hr pooled (D105); DC CD1a+ CD86+, from CD34+ GM-CSF, TNF α 12 days FACS sorted, activated with PMA and

ionomycin for 1, 6 h pooled (D106); DC from monocytes GM-CSF, IL-4 5 days, resting (D107); DC from monocytes GM-CSF, IL-4 5 days, resting (D108); DC from monocytes GM-CSF, IL-4 5 days, activated LPS 4, 16 h pooled (D109); DC from monocytes GM-CSF, IL-4 5 days, activated TNF α , monocyte supe for 4, 16 h pooled (D110); leiomyoma L11 benign tumor (X101); normal myometrium M5 (O115); malignant leiomyosarcoma GS1 (X103); lung fibroblast sarcoma line MRC5, activated with PMA and ionomycin for 1, 6 h pooled (C101); kidney epithelial carcinoma cell line CHA, activated with PMA and ionomycin for 1, 6 h pooled (C102); kidney fetal 28 wk male (O100); lung fetal 28 wk male (O101); liver fetal 28 wk male (O102); heart fetal 28 wk male (O103); brain fetal 28 wk male (O104); gallbladder fetal 28 wk male (O106); small intestine fetal 28 wk male (O107); adipose tissue fetal 28 wk male (O108); ovary fetal 25 wk female (O109); uterus fetal 25 wk female (O110); testes fetal 28 wk male (O111); spleen fetal 28 wk male (O112); adult placenta 28 wk (O113); and tonsil inflamed, from 12 year old (X100).

TaqMan quantitative PCR techniques have shown the DCRS6, in both mouse and human, to be expressed on T cells, including thymocytes and CD4+ naive and differentiated (hDCRS6 is also expressed on dendritic cells), in gastrointestinal tissue, including stomach, intestine, colon and associated lymphoid tissue, e.g., Peyer's patches and mesenteric lymph nodes, and upregulated in inflammatory models of bowel disease, e.g., IL-10 KO mice. The hDCRS7 was detected in both resting and activated dendritic cells, epithelial cells, and mucosal tissues, including GI and reproductive tracts. These data suggest that family members are expressed in mucosal tissues and immune system cell types, and/or in gastrointestinal, airway, and reproductive tract development.

As such, therapeutic indications include, e.g., short bowel syndrome, post chemo/radio-therapy or alcoholic recovery, combinations with ulcer treatments or arthritis medication, Th2 pregnancy skewing, stomach lining/tissue regeneration, loss of adsorptive surface conditions, etc. See, e.g., Yamada, et al. (eds. 1999) Textbook of Gastroenterology; Yamada, et al. (eds. 1999) Textbook and Atlas of Gastroenterology; Gore and Levine (2000) Textbook of Gastrointestinal Radiology; and (1987) Textbook of Pediatric Gastroenterology.

Similar samples may isolated in other species for evaluation.

Primers specific for IL-17RA were designed and used in Taqman quantative PCR against various human libraries. IL-17RA is highly expressed in innate immune myeloid cells including dendritic cells and monocytes. Expression is also detected in T-cell libraries. These data demonstrate the receptor is expressed in immune cell types and may be regulated by activation conditions.

**Table for IL-17RA
library description**

	CT for IL- 17RA_H
DC ex monocytes GM-CSF, IL-4, resting	16.97
U937 premonocytic line, activated	17.14
DC ex monocytes GM-CSF, IL-4, resting	17.53
DC 70% CD1a+, ex CD34+ GM-CSF, TNFa, resting	18.17
monocytes, LPS, gIFN, anti-IL-10	18.27
DC ex monocytes GM-CSF, IL-4, LPS activated 4+16 hr	18.51
DC ex monocytes GM-CSF, IL-4, monokine activated 4+16 hr	18.68
kidney epithelial carcinoma cell line CHA, activated	18.69
monocytes, LPS, 1 hr	18.72
monocytes, LPS, 6 hr	18.72
DC 70% CD1a+, ex CD34+ GM-CSF, TNFa, activated 1 hr	18.91
DC 70% CD1a+, ex CD34+ GM-CSF, TNFa, activated 6 hr	18.94
T cell, TH1 clone HY06, activated	18.99
lung fetal	19.15
T cell, TH1 clone HY06, resting	19.18
T cell, TH1 clone HY06, anergic	19.23
monocytes, LPS, gIFN, IL-10, 4+16 hr	19.3
spleen fetal	19.51
testes fetal	19.7
T cell, TH0 clone Mot 72, resting	19.71
T cell, TH0 clone Mot 72, resting	19.84
DC CD1a+ CD86+, ex CD34+ GM-CSF, TNFa, activated 1+6 hr	19.94
peripheral blood mononuclear cells, activated	20.01
hematopoietic precursor line TF1, activated	20.07
lung fibroblast sarcoma line MRC5, activated	20.18
Splenocytes, activated	20.21
T cell gd clones, resting	20.27
ovary fetal	20.45
T cells CD4+, TH2 polarized, activated	20.57
Splenocytes, resting	20.6
uterus fetal	20.62
DC 95% CD1a+, ex CD34+ GM-CSF, TNFa, activated 1+6 hr	20.94
epithelial cells, unstimulated	20.96
peripheral blood mononuclear cells, resting	20.97
adipose tissue fetal	21.13

B cell line JY, activated	21.28
monocytes, LPS, gIFN, IL-10	21.37
placenta 28 wk	21.38
NK 20 clones pooled, activated	21.55
pool of two normal human lung samples	21.63
normal human thyroid	21.65
epithelial cells, IL-1b activated	21.72
normal human skin	21.84
T cell, TH0 clone Mot 72, anergic	21.87
small intestine fetal	22.01
CD28- T cell clone in pME	22.08
T cell, TH2 clone HY935, activated	22.09
T cell clones, pooled, resting	22.29
Hashimoto's thyroiditis thyroid sample	22.3
NK 20 clones pooled, resting	22.4
B cell EBV lines, resting	22.45
T cell, TH2 clone HY935, resting	22.86
T cell, TH0 clone Mot 72, activated	23.3
monocytes, LPS, gIFN, anti-IL-10, 4+16 hr	23.39
T cell lines Jurkat and Hut78, resting	23.4
T cell, TH0 clone Mot 72, activated	23.56
<i>Pneumocystis carinii</i> pneumonia lung sample	24.05
U937 premonocytic line, resting	25.01
pool of rheumatoid arthritis samples, human	25.85
pool of three heavy smoker human lung samples	26.1
DC 95% CD14+, ex CD34+ GM-CSF, TNFa, activated 1+6 hr	32.69
kidney fetal	33.7
liver fetal	34.4
NK cytotoxic clone, resting	34.49
tonsil inflamed	35.02
normal w.t. monkey lung	35.45
gallbladder fetal	35.84
TR1 T cell clone	35.86
allergic lung sample	36.39
Psoriasis patient skin sample	36.44
normal human colon	37.34
brain fetal	37.35
<i>Ascaris</i> -challenged monkey lung, 4 hr.	37.75
<i>Ascaris</i> -challenged monkey lung, 24 hr.	40
heart fetal	40
normal w.t. monkey colon	40
ulcerative colitis human colon sample	40

Primers specific for DCRS6_H were designed and used in Taqman quantitative PCR against various human libraries. DCRS6_H is expressed in innate immune myeloid cells including dendritic cells and monocytes. Expression is also detected in T-cell libraries. These data demonstrate the receptor is expressed in immune cell types and may be regulated by activation conditions.

5

Table for DCRS6_H library description	CT for DCRS6_H
T cell, TH0 clone Mot 72, resting	15.54
T cell, TH0 clone Mot 72, resting	15.7
DC ex monocytes GM-CSF, IL-4, resting	17.84
DC ex monocytes GM-CSF, IL-4, resting	18.19
DC ex monocytes GM-CSF, IL-4, LPS activated 4+16 hr	18.3
DC ex monocytes GM-CSF, IL-4, monokine activated 4+16 hr	18.3
T cell, TH1 clone HY06, resting	18.43
NK cytotoxic clone, resting	18.53
T cell clones, pooled, resting	18.8
T cell, TH1 clone HY06, activated	19.03
T cell, TH2 clone HY935, activated	19.1
TR1 T cell clone	19.12
T cells CD4+, TH2 polarized, activated	20.06
B cell EBV lines, resting	20.3
T cell, TH2 clone HY935, resting	20.48
kidney epithelial carcinoma cell line CHA, activated	21.07
T cell, TH1 clone HY06, anergic	21.14
normal human colon	21.29
NK 20 clones pooled, resting	21.49
T cell gd clones, resting	21.58
gallbladder fetal	22.21
kidney fetal	22.79
liver fetal	22.8
<i>Pneumocystic carinii</i> pneumonia lung sample	23.06
CD28- T cell clone in pME	23.18
T cell, TH0 clone Mot 72, anergic	23.2
ovary fetal	23.51
normal human thyroid	24.03
small intestine fetal	24.13
testes fetal	24.82
epithelial cells, IL-1b activated	26.08
pool of three heavy smoker human lung samples	26.49
placenta 28 wk	26.56
normal w.t. monkey lung	28.65
peripheral blood mononuclear cells,	33.39

activated	
<i>Ascaris</i> -challenged monkey lung, 4 hr.	36.59
spleen fetal	38.43
peripheral blood mononuclear cells, resting	40
T cell, TH0 clone Mot 72, activated	40
T cell lines Jurkat and Hut78, resting	40
Splenocytes, resting	40
Splenocytes, activated	40
B cell line JY, activated	40
NK 20 clones pooled, activated	40
hematopoietic precursor line TF1, activated	40
U937 premonocytic line, resting	40
U937 premonocytic line, activated	40
monocytes, LPS, gIFN, anti-IL-10	40
monocytes, LPS, gIFN, IL-10	40
monocytes, LPS, gIFN, anti-IL-10, 4+16 hr	40
monocytes, LPS, gIFN, IL-10, 4+16 hr	40
monocytes, LPS, 1 hr	40
monocytes, LPS, 6 hr	40
DC 70% CD1a+, ex CD34+ GM-CSF, TNFa, resting	40
DC 70% CD1a+, ex CD34+ GM-CSF, TNFa, activated 1 hr	40
DC 70% CD1a+, ex CD34+ GM-CSF, TNFa, activated 6 hr	40
DC 95% CD1a+, ex CD34+ GM-CSF, TNFa, activated 1+6 hr	40
DC 95% CD14+, ex CD34+ GM-CSF, TNFa, activated 1+6 hr	40
DC CD1a+ CD86+, ex CD34+ GM-CSF, TNFa, activated 1+6 hr	40
epithelial cells, unstimulated	40
lung fibroblast sarcoma line MRC5, activated	40
<i>Ascaris</i> -challenged monkey lung, 24 hr.	40
pool of two normal human lung samples	40
allergic lung sample	40
normal w.t. monkey colon	40
ulcerative colitis human colon sample	40
Hashimoto's thyroiditis thyroid sample	40
pool of rheumatoid arthritis samples, human	40
normal human skin	40
Psoriasis patient skin sample	40
tonsil inflamed	40
lung fetal	40
heart fetal	40
brain fetal	40
adipose tissue fetal	40
uterus fetal	40

T cell, TH0 clone Mot 72, activated 40

- 5 Primers specific for DCRS7_H were designed and used in Taqman quantitative PCR against various human libraries. DCRS7_H is expressed in innate immune myeloid cells including dendritic cells and monocytes. Expression is also detected in fetal libraries. These data demonstrate the receptor is expressed in immune cell types and may be regulated by activation conditions.

Table for DCRS7_H library description	CT for DCRS7_H
fetal uterus	19.05
DC mix	19.34
fetal small intestine	19.46
fetal ovary	19.68
fetal testes	19.75
fetal lung	20.04
CHA	20.24
normal thyroid	20.32
DC/GM/IL-4	20.52
fetal spleen	20.86
normal lung	20.94
TF1	21
allergic lung #19	21.02
Psoriasis skin	21.07
fetal liver	21.15
MRC5	21.15
24 hr. Ascaris lung	21.17
hi dose IL-4 lung	21.23
CD1a+ 95%	21.32
Hashimotos thyroiditis	21.35
Crohns colon 4003197A	21.35
normal lung pool	21.36
70% DC resting	21.42
fetal kidney	21.58
adult placenta	21.68
lung 121897-1	21.8
Pneumocystis carinii lung	21.81
#20	
A549 unstim.	21.89
normal colon #22	21.94
18 hr. Ascaris lung	22.09
normal skin	22.1
Crohns colon 9609C144	22.13
fetal adipose tissue	22.35
D6	22.39

DC resting CD34-derived	22.45
DC TNF/TGFb act CD34-der.	22.54
fetal brain	22.9
DC CD40L activ. mono-deriv.	22.91
Crohns colon 403242A	22.91
ulcerative colitis colon #26	23
RA synovium pool	23.06
A549 activated	23.06
mono + IL-10	23.42
DC LPS	23.49
Mot 72 activated	23.66
CD1a+ CD86+	23.86
HY06 resting	23.87
U937 activated	23.97
inflamed tonsil	23.97
D1	24.06
M1	24.17
CD14+ 95%	24.21
lung 080698-2	24.28
4 hr. Ascaris lung	24.37
Jurkat activated pSPORT	24.42
DC resting mono-derived	24.48
HY06 activated	24.54
C+	24.64
Splenocytes resting	24.65
U937/CD004 resting	24.96
PBMC resting	25.8
Mot 72 resting	25.91
mono + anti-IL-10	26.14
NK pool	26.99
HY06 anti-peptide	27.34
mast cell pME	27.38
Tc gamma delta	28.14
TC1080 CD28- pMET7	31.05
PBMC activated	31.89
NK non cytotox.	32.3
RV-C30 TR1 pMET7	32.5
Bc	33.72
C-	33.8
Splenocytes activated	34.7
JY	35.05
NK cytotox.	36.44
NKL/IL-2	37.59
HY935 resting	37.6
NK pool activated	38.15
Mot 72 anti-peptide	38.87
fetal heart	40.92

B21 resting	42.05
Jurkat resting pSPORT	42.8
B21 activated	43.09
NKA6 pSPORT	44.85
HY935 activated	45
M6	45

- 5 Primers specific for DCRS9_H were designed and used in Taqman quantative PCR against various human libraries. DCRS9_H is expressed T-cells, fetal lung, and resting monocytes. These data demonstrate the receptor is expressed in immune cell types and may be regulated by activation conditions.

Table for DCRS9_H
library description CT for
DCRS9_H

HY06 resting	22.35
fetal lung	22.63
HY06 anti-peptide	22.72
HY06 activated	22.96
U937/CD004 resting	24.16
fetal small intestine	24.94
JY	25.04
Mot 72 resting	25.12
Jurkat activated pSPORT	25.2
RV-C30 TR1 pMET7	26.51
fetal kidney	26.76
MRC5	27.2
Psoriasis skin	27.3
Tc gamma delta	27.37
Crohns colon 4003197A	27.44
fetal spleen	27.72
normal lung	27.83
Hashimotos thyroiditis	28.03
B21 resting	28.32
TF1	28.39
NK cytotox.	28.44
TC1080 CD28- pMET7	28.61
Pneumocystis carinii lung #20	29.05
U937 activated	29.06
HY935 resting	29.09
CD1a+ 95%	29.13

B21 activated	29.2
Mot 72 activated	29.21
fetal testes	29.27
lung 080698-2	29.32
Jurkat resting	29.38
pSPORT	
CD14+ 95%	29.38
normal thyroid	29.53
Mot 72 anti-peptide	29.65
Splenocytes resting	29.85
Crohns colon 9609C144	30.28
lung 121897-1	30.37
24 hr. Ascaris lung	30.59
hi dose IL-4 lung	30.8
CD1a+ CD86+	31.42
normal skin	31.73
fetal uterus	31.79
PBMC activated	31.82
inflammed tonsil	31.98
fetal brain	32.21
RA synovium pool	32.77
allergic lung #19	33.18
18 hr. Ascaris lung	33.42
adult placenta	33.43
normal lung pool	33.45
Crohns colon 403242A	33.52
NK pool	33.72
HY935 activated	33.75
DC/GM/IL-4	34.28
DC resting mono-derived	34.57
fetal ovary	35.06
fetal adipose tissue	35.07
CHA	35.2
PBMC resting	35.95
Bc	36.19
A549 unstim.	36.4
fetal heart	36.87
ulcerative colitis colon #26	37.83
C-	38.32
4 hr. Ascaris lung	40.2
D6	40.62
C+	44.38

A549 activated	44.58
Splenocytes	45
activated	
NK pool activated	45
NKA6 pSPORT	45
NKL/IL-2	45
NK non cytotox.	45
mono + anti-IL-10	45
mono + IL-10	45
M1	45
M6	45
70% DC resting	45
D1	45
DC LPS	45
DC mix	45
fetal liver	45
mast cell pME	45
DC CD40L activ.	45
mono-deriv.	
DC resting CD34- derived	45
DC TNF/TGFb act	45
CD34-der.	
normal colon #22	45

V. Cloning of species counterparts

Various strategies are used to obtain species counterparts of the DCRSs, preferably from other primates or rodents. One method is by cross hybridization using closely related species DNA probes. It may be useful to go into evolutionarily similar species as intermediate steps. Another method is by using specific PCR primers based on the identification of blocks of similarity or difference between genes, e.g., areas of highly conserved or nonconserved polypeptide or nucleotide sequence. Sequence database searches may identify species counterparts.

10 VI. Production of mammalian protein

An appropriate, e.g., GST, fusion construct is engineered for expression, e.g., in *E. coli*. For example, a mouse IGIF pGex plasmid is constructed and transformed into *E. coli*. Freshly transformed cells are grown, e.g., in LB medium containing 50 µg/ml ampicillin and induced with IPTG (Sigma, St. Louis, MO). After overnight induction, the bacteria are harvested and the pellets containing the appropriate protein are isolated. The pellets are homogenized, e.g., in TE buffer (50 mM Tris-base pH 8.0, 10 mM EDTA and 2 mM pefabloc) in 2 liters. This material is passed through a microfluidizer (Microfluidics, Newton, MA) three times. The fluidized supernatant is spun down on a Sorvall GS-3 rotor for 1 h at 13,000 rpm. The resulting supernatant containing the cytokine receptor protein is filtered and passed over a glutathione-SEPHAROSE column equilibrated in 50 mM Tris-base pH 8.0. Fractions containing the DCRS8-GST fusion protein are pooled and cleaved, e.g., with thrombin (Enzyme Research Laboratories, Inc., South Bend, IN). The cleaved pool is then passed over a Q-SEPHAROSE column equilibrated in 50 mM Tris-base. Fractions containing DCRS8 are pooled and diluted in cold distilled H₂O, to lower the conductivity, and passed back over a fresh Q-Sepharose column, alone or in succession with an immunoaffinity antibody column. Fractions containing the DCRS8 protein are pooled, aliquoted, and stored in the -70° C freezer.

Comparison of the CD spectrum with cytokine receptor protein may suggest that the protein is correctly folded. See Hazuda, et al. (1969) J. Biol. Chem. 264:1689-1693.

30 VII. Preparation of specific antibodies

Inbred Balb/c mice are immunized intraperitoneally with recombinant forms of the protein, e.g., purified DCRS8 or stable transfected NIH-3T3 cells. Animals are boosted at appropriate time points with protein, with or without additional adjuvant, to further stimulate antibody production. Serum is collected, or hybridomas produced with harvested spleens.

Alternatively, Balb/c mice are immunized with cells transformed with the gene or fragments thereof, either endogenous or exogenous cells, or with isolated membranes enriched for expression of the antigen. Serum is collected at the appropriate time, typically after numerous further administrations. Various gene therapy techniques may be useful, e.g., in producing protein in situ, for generating an immune response. Serum or antibody preparations may be cross-absorbed or immunoselected to prepare substantially purified antibodies of defined specificity and high affinity.

Monoclonal antibodies may be made. For example, splenocytes are fused with an appropriate fusion partner and hybridomas are selected in growth medium by standard procedures. Hybridoma supernatants are screened for the presence of antibodies which bind to the DCRS8, e.g., by ELISA or other assay. Antibodies which specifically recognize specific DCRS8 embodiments may also be selected or prepared.

In another method, synthetic peptides or purified protein are presented to an immune system to generate monoclonal or polyclonal antibodies. See, e.g., Coligan (ed. 1991) Current Protocols in Immunology Wiley/Greene; and Harlow and Lane (1989) Antibodies: A Laboratory Manual Cold Spring Harbor Press. In appropriate situations, the binding reagent is either labeled as described above, e.g., fluorescence or otherwise, or immobilized to a substrate for panning methods. Nucleic acids may also be introduced into cells in an animal to produce the antigen, which serves to elicit an immune response. See, e.g., Wang, et al. (1993) Proc. Nat'l. Acad. Sci. 90:4156-4160; Barry, et al. (1994) BioTechniques 16:616-619; and Xiang, et al. (1995) Immunity 2: 129-135.

VIII. Production of fusion proteins

Various fusion constructs are made with DCRS8 or DCRS9. A portion of the appropriate gene is fused to an epitope tag, e.g., a FLAG tag, or to a two hybrid system construct. See, e.g., Fields and Song (1989) Nature 340:245-246.

The epitope tag may be used in an expression cloning procedure with detection with anti-FLAG antibodies to detect a binding partner, e.g., ligand for the respective cytokine receptor. The two hybrid system may also be used to isolate proteins which specifically bind to the receptor subunit.

IX. Structure activity relationship

Information on the criticality of particular residues is determined using standard procedures and analysis. Standard mutagenesis analysis is performed, e.g., by generating many different variants at determined positions, e.g., at the positions identified above, and evaluating biological activities of the variants. This may be performed to the extent of determining positions which modify activity, or to focus on specific positions to

determine the residues which can be substituted to either retain, block, or modulate biological activity.

Alternatively, analysis of natural variants can indicate what positions tolerate natural mutations. This may result from populational analysis of variation among
5 individuals, or across strains or species. Samples from selected individuals are analyzed, e.g., by PCR analysis and sequencing. This allows evaluation of population polymorphisms.

X. Isolation of a ligand

10 A cytokine receptor can be used as a specific binding reagent to identify its binding partner, by taking advantage of its specificity of binding, much like an antibody would be used. The binding receptor may be a heterodimer of receptor subunits; or may involve, e.g., a complex of the DCRS8 with another cytokine receptor subunit. A binding reagent is either labeled as described above, e.g., fluorescence or otherwise, or
15 immobilized to a substrate for panning methods.

The binding composition is used to screen an expression library made from a cell line which expresses a binding partner, i.e., ligand, preferably membrane associated. Standard staining techniques are used to detect or sort surface expressed ligand, or surface expressing transformed cells are screened by panning. Screening of intracellular
20 expression is performed by various staining or immunofluorescence procedures. See also McMahan, et al. (1991) EMBO J. 10:2821-2832.

For example, on day 0, precoat 2-chamber permanox slides with 1 ml per chamber of fibronectin, 10 ng/ml in PBS, for 30 min at room temperature. Rinse once with PBS. Then plate COS cells at $2-3 \times 10^5$ cells per chamber in 1.5 ml of growth media. Incubate
25 overnight at 37 C.

On day 1 for each sample, prepare 0.5 ml of a solution of 66 μ g/ml DEAE-dextran, 66 μ M chloroquine, and 4 μ g DNA in serum free DME. For each set, a positive control is prepared, e.g., of DCRS8-FLAG cDNA at 1 and 1/200 dilution, and a negative mock. Rinse cells with serum free DME. Add the DNA solution and incubate 5 hr at 37
30 C. Remove the medium and add 0.5 ml 10% DMSO in DME for 2.5 min. Remove and wash once with DME. Add 1.5 ml growth medium and incubate overnight.

On day 2, change the medium. On days 3 or 4, the cells are fixed and stained. Rinse the cells twice with Hank's Buffered Saline Solution (HBSS) and fix in 4% paraformaldehyde (PFA)/glucose for 5 min. Wash 3X with HBSS. The slides may be
35 stored at -80 C after all liquid is removed. For each chamber, 0.5 ml incubations are performed as follows. Add HBSS/saponin (0.1%) with 32 μ l/ml of 1 M NaN₃ for 20 min. Cells are then washed with HBSS/saponin 1X. Add appropriate DCRS8 or

DCRS8/antibody complex to cells and incubate for 30 min. Wash cells twice with HBSS/saponin. If appropriate, add first antibody for 30 min. Add second antibody, e.g., Vector anti-mouse antibody, at 1/200 dilution, and incubate for 30 min. Prepare ELISA solution, e.g., Vector Elite ABC horseradish peroxidase solution, and preincubate for 30 min. Use, e.g., 1 drop of solution A (avidin) and 1 drop solution B (biotin) per 2.5 ml HBSS/saponin. Wash cells twice with HBSS/saponin. Add ABC HRP solution and incubate for 30 min. Wash cells twice with HBSS, second wash for 2 min, which closes cells. Then add Vector diaminobenzoic acid (DAB) for 5 to 10 min. Use 2 drops of buffer plus 4 drops DAB plus 2 drops of H₂O₂ per 5 ml of glass distilled water.

Carefully remove chamber and rinse slide in water. Air dry for a few minutes, then add 1 drop of Crystal Mount and a cover slip. Bake for 5 min at 85-90 C.

Evaluate positive staining of pools and progressively subclone to isolation of single genes responsible for the binding.

Alternatively, receptor reagents are used to affinity purify or sort out cells expressing a putative ligand. See, e.g., Sambrook, et al. or Ausubel, et al.

Another strategy is to screen for a membrane bound receptor by panning. The receptor cDNA is constructed as described above. Immobilization may be achieved by use of appropriate antibodies which recognize, e.g., a FLAG sequence of a DCRS8 fusion construct, or by use of antibodies raised against the first antibodies. Recursive cycles of selection and amplification lead to enrichment of appropriate clones and eventual isolation of receptor expressing clones.

Phage expression libraries can be screened by mammalian DCRS8. Appropriate label techniques, e.g., anti-FLAG antibodies, will allow specific labeling of appropriate clones.

We tested the ability of DCRS receptors to specifically bind IL-17 family cytokines. Recombinant FLAG-hIL-17 family cytokines were used in binding experiments on Baf/3 DCRS receptor transfected expressing recombinant IL-17R_H, DCRS6_H, DCRS7_H, DCRS8_H and DCRS9_H and analyzed by FACS. We can demonstrate specific binding of IL-17 family member IL-74 to DCRS6 expressing Baf/3 cells. In additional experiments we have shown IL-17 specific binding to IL-17R_H, DCRS7_H, DCRS8_H. Further experiments show IL-71 binding to DCRS8_Hu transfectants. These experiments demonstrate the sequence homology among IL-17 related cytokine receptors confers functional binding to IL-17 cytokines.

All citations herein are incorporated herein by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

Many modifications and variations of this invention can be made without departing from its spirit and scope, as will be apparent to those skilled in the art. The specific embodiments described herein are offered by way of example only, and the invention is to be limited by the terms of the appended claims, along with the full scope of equivalents to which such claims are entitled; and the invention is not to be limited by the specific embodiments that have been presented herein by way of example.

WHAT IS CLAIMED IS:

1. A composition of matter selected from:
 - a) a substantially pure or recombinant polypeptide comprising at least three distinct nonoverlapping segments of at least four amino acids identical to segments of SEQ ID NO: 14;
 - b) a substantially pure or recombinant polypeptide comprising at least two distinct nonoverlapping segments of at least five amino acids identical to segments of SEQ ID NO: 14;
 - c) a natural sequence DCRS8 comprising mature SEQ ID NO: 14;
 - d) a fusion polypeptide comprising DCRS8 sequence;
 - e) a substantially pure or recombinant polypeptide comprising at least three distinct nonoverlapping segments of at least four amino acids identical to segments of SEQ ID NO: 17 or 20;
 - f) a substantially pure or recombinant polypeptide comprising at least two distinct nonoverlapping segments of at least five amino acids identical to segments of SEQ ID NO: 17 or 20;
 - g) a natural sequence DCRS9 comprising mature SEQ ID NO: 17 or 20; or
 - h) a fusion polypeptide comprising DCRS9 sequence.
2. The substantially pure or isolated antigenic polypeptide of Claim 1, wherein said distinct nonoverlapping segments of identity include:
 - a) one of at least eight amino acids;
 - b) one of at least four amino acids and a second of at least five amino acids;
 - c) at least three segments of at least four, five, and six amino acids, or
 - d) one of at least twelve amino acids.
3. The composition of matter of Claim 1, wherein said:
 - a) polypeptide:
 - i) comprises a mature sequence of Table 3 or 4;
 - ii) is an unglycosylated form of DCRS8 or DCRS9;
 - iii) is from a primate, such as a human;
 - iv) comprises at least seventeen amino acids of SEQ ID NO: 14 or 17;
 - v) exhibits at least four nonoverlapping segments of at least seven amino acids of SEQ ID NO: 14 or 17;
 - vi) is a natural allelic variant of DCRS8 or DCRS9;
 - vii) has a length at least about 30 amino acids;

- viii) exhibits at least two non-overlapping epitopes which are specific for a primate DCRS8 or DCRS9;
- ix) is glycosylated;
- x) has a molecular weight of at least 30 kD with natural glycosylation;
- xi) is a synthetic polypeptide;
- xii) is attached to a solid substrate;
- xiii) is conjugated to another chemical moiety;
- xiv) is a 5-fold or less substitution from natural sequence; or
- xv) is a deletion or insertion variant from a natural sequence.
4. A composition comprising:
- a) a substantially pure DCRS8 or DCRS9 and another cytokine receptor family member;
- b) a sterile DCRS8 or DCRS9 polypeptide of Claim 1;
- c) said DCRS8 or DCRS9 polypeptide of Claim 1 and a carrier, wherein said carrier is:
- i) an aqueous compound, including water, saline, and/or buffer; and/or
- ii) formulated for oral, rectal, nasal, topical, or parenteral administration.
5. The fusion polypeptide of Claim 1, comprising:
- a) mature protein sequence of Table 3 or 4;
- b) a detection or purification tag, including a FLAG, His6, or Ig sequence; or
- c) sequence of another cytokine receptor protein.
6. A kit comprising a polypeptide of Claim 1, and:
- a) a compartment comprising said protein or polypeptide; or
- b) instructions for use or disposal of reagents in said kit.
7. A binding compound comprising an antigen binding site from an antibody, which specifically binds to a natural DCRS8 or DCRS9 polypeptide of Claim 1, wherein:
- a) said binding compound is in a container;
- b) said DCRS8 or DCRS9 polypeptide is from a human;
- c) said binding compound is an Fv, Fab, or Fab2 fragment;
- d) said binding compound is conjugated to another chemical moiety; or
- e) said antibody:
- i) is raised against a peptide sequence of a mature polypeptide of Table 3 or 4;

- ii) is raised against a mature DCRS8 or DCRS9;
iii) is raised to a purified human DCRS8 or DCRS9;
iv) is immunoselected;
v) is a polyclonal antibody;
5 vi) binds to a denatured DCRS8 or DCRS9;
vii) exhibits a Kd to antigen of at least 30 μ M;
viii) is attached to a solid substrate, including a bead or plastic membrane;
ix) is in a sterile composition; or
x) is detectably labeled, including a radioactive or fluorescent label.
- 10 8. A kit comprising said binding compound of Claim 7, and:
a) a compartment comprising said binding compound; or
b) instructions for use or disposal of reagents in said kit.
- 15 9. A method of producing an antigen:antibody complex, comprising
contacting under appropriate conditions a primate DCRS8 or DCRS9 polypeptide with an
antibody of Claim 7, thereby allowing said complex to form.
10. The method of Claim 9, wherein:
20 a) said complex is purified from other cytokine receptors;
b) said complex is purified from other antibody;
c) said contacting is with a sample comprising an interferon;
d) said contacting allows quantitative detection of said antigen;
e) said contacting is with a sample comprising said antibody; or
25 f) said contacting allows quantitative detection of said antibody.
11. A composition comprising:
a) a sterile binding compound of Claim 7, or
b) said binding compound of Claim 7 and a carrier, wherein said carrier is:
30 i) an aqueous compound, including water, saline, and/or buffer; and/or
ii) formulated for oral, rectal, nasal, topical, or parenteral administration.
12. An isolated or recombinant nucleic acid encoding said polypeptide of
Claim 1, wherein said:
35 a) DCRS8 or DCRS9 is from a human; or
b) said nucleic acid:
i) encodes an antigenic peptide sequence of Table 3 or 4;

- ii) encodes a plurality of antigenic peptide sequences of Table 3 or 4;
iii) exhibits identity over at least thirteen nucleotides to a natural cDNA
encoding said segment;
iv) is an expression vector;
5 v) further comprises an origin of replication;
vi) is from a natural source;
vii) comprises a detectable label;
viii) comprises synthetic nucleotide sequence;
ix) is less than 6 kb, preferably less than 3 kb;
10 x) is from a primate;
xi) comprises a natural full length coding sequence;
xii) is a hybridization probe for a gene encoding said DCRS8 or DCRS9;
or
xiii) is a PCR primer, PCR product, or mutagenesis primer.
- 15 13. A cell or tissue comprising said recombinant nucleic acid of Claim 12.
14. The cell of Claim 13, wherein said cell is:
a) a prokaryotic cell;
20 b) a eukaryotic cell;
c) a bacterial cell;
d) a yeast cell;
e) an insect cell;
f) a mammalian cell;
25 g) a mouse cell;
h) a primate cell; or
i) a human cell.
15. A kit comprising said nucleic acid of Claim 12, and:
30 a) a compartment comprising said nucleic acid;
b) a compartment further comprising a primate DCRS8 or DCRS9 polypeptide;
or
c) instructions for use or disposal of reagents in said kit.
- 35 16. A nucleic acid which:
a) hybridizes under wash conditions of 30 minutes at 30° C and less than 2M salt
to the coding portion of SEQ ID NO: 13 or 16; or

- b) exhibits identity over a stretch of at least about 30 nucleotides to a primate DCRS8 or DCRS9.

17. The nucleic acid of Claim 16, wherein:

- 5 a) said wash conditions are at 45° C and/or 500 mM salt; or
b) said stretch is at least 55 nucleotides.

18. The nucleic acid of Claim 16, wherein:

- 10 a) said wash conditions are at 55° C and/or 150 mM salt; or
b) said stretch is at least 75 nucleotides.

19. A method of modulating physiology or development of a cell or tissue culture cells comprising contacting said cell with an agonist or antagonist of a mammalian DCRS8 or DCRS9.

15

20. The method of Claim 19, wherein said cell is transformed with a nucleic acid encoding said DCRS8 or DCRS9 and another cytokine receptor subunit.

1/2

DCRS7_Mu RTALLHSADG-AGYERLVGALASALSQMP---LRVAVDLWSRRE-LSAHGALAWFHHQR
 DCRS7_Hu RAALLYSADD-SGFERLVGALASALCQLP---LRVAVDLWSRRE-LSAQGPVAFWFAQR
 IL-17R_Hu RKVWIYYSADH-PLYVDVVLKFAQFLLTACG--TEVALDLLEEQA-ISEAGVMTWVGRQK
 IL-17R_Mu RKVWIVYSADH-PLYVEVVLKFAQFLITACG--TEVALDLLEEQV-ISEVGVMTWVSRQK
 DCRS10 RKVFITYSMD-----TAMEVVKFVNFLLVNG---FQTAIDIFEDR--IRGIDIIKWMERYL
 DCRS10_Mu RKVFITYSMD-----TAMEVVKFVNFLLVNG---FQTAIDIFEDR--IRGIDIIKWMERYL
 DCRS9_Hu RPVLLLHAADS-EAQRRLVGALAEELLRAALGGGRDVIVDLWEGRH-VARVGPLPWLWAAR
 DCRS8_Hu PKVFLCYSSKDGQNHMNVVQCFAFLQDFCG--CEVALDLWEDFS-LCREGQREWVIQKI
 IL-17R_Ce VKVMIVYADDN-DLHTDCVKKLVENLRNCAS--CDPVFDLEKLI--TAEIVPSRWLVDQI
 DCRS6_Hu IKVLVVPSEI--CFHHTICYFTEFLQNHCR--SEVILEKWQKKK-IAEMGPVQWLATQK
 DCRS6_Ce FKVMLVCPEVS-GRDEDFMMRIADALKKSN---NKVVCDRWFEDSKNAEENMLHWVYEQT
 . : . : . : * *

DCRS7_Mu RRILQEGGVILLFSPAAVAQCQ---QWLQLQTVEP--GP---HDALAAMLSCVLPDFL
 DCRS7_Hu RQTLQEGGVVLLFSPGAVALCS---EWLQDGVSGPGAHGP---HDAFRASLSCVLPDFL
 IL-17R_Hu QEMVESNSKIIVLCSRGTRAKWQALLGRGAP-VRLRCDHGKPV-GDLFTAAMNMILPDFK
 IL-17R_Mu QEMVESNSKIIILCSRGTOAKWKAILGWAEPVQLRCDHWKPA-GDLFTAAMNMILPDFK
 DCRS10 R---DKTVMIIVAISP KYKQDVE-----GAESQDDED-EHGL---HTKYIHRM-MQIEFIK
 DCRS10_Mu R---DKTVMIIVAISP KYKQDVE-----GAESQDDED-EHGL---HTKYIHRM-MQIEFIS
 DCRS9_Hu TRVAREQGTVLLWSGADLRPVS---GPD P-RAAP-----LLA----LLHAAP
 DCRS8_Hu H---ESQFIIVVCSKGMKYFVD---KKNYKHKGGRSGK---GELFLVAVSAIAEKL R
 IL-17R_Ce S---SLKKFIIVVS DCAEKILD-----TEASETHQLVQARP--FADLFGPAMEMIIRDAT
 DCRS6_Hu K---AADKVVFLLSNDVNSVCD-----GTCGKSEGSPSENS---QDLFPLAFNLFCSDLR
 DCRS6_Ce K---IAEKII VFHSAYYHPRCG---IYDVINFFPCTDPR-----LAHIALT---PEAQ
 . : . : . : *

FIG. 1A

DCRS7_Mu	QGRATGR-----YVGVFYDGLLHPDSVPSFVRVAPLFSLP-SQLPAFLDALQ--GGCSTS
DCRS7_Hu	QGRAPGS-----YVGACFDRLLHPDAVPALFRTVPVFTLP-SQLPDFLGALQ--QPRAPR
IL-17R_Hu	RPACFGT-----YVVCYFSEVSCDGDVPDLFGAAPRYPLM-DRFEEVYFRIQ--DLEMFO
IL-17R_Mu	RPACFGT-----YVVCYFSGICSERDVPDLFNITSRYPLM-DRFEEVYFRIQ--DLEMFE
DCRS10	QGS MNFR-----FIPVLF PNAK-KEHVPTWLQNT HVYSWP-KNKKNI LLRL-REEEYVA
DCRS10_Mu	QGS MNFR-----FIPVLF PNAK-KEHVPTWLQNT HVYSWP-KNKKNI LLRL-REEEYVA
DCRS9_Hu	RPL-----LLLAYFSRLCAKGDIPPLRALPRYRL- RDLPRLLRALD--ARPPAE
DCRS8_Hu	QAKQSSAALSKFIAVYFDYSC-EGDVPGILDLSTKYRLM-DNLPQLCSHLHSRDHGLQE
IL-17R_Ce	HNFPEAR---KKYAVVRFNYS P---HVPPNLA I LNLP TF IPEQFAQLTAF LHN-VEHTER
DCRS6_Hu	SQIHLHK-----YVVVYFREID-TKDDYNALSVCPKYHLM-KDATAFCAELL---HVKQQ
DCRS6_Ce	RSVPKEV---EYVLPRDQKLL--EDAFDITIADPLVIDIPIEDVAIPENVP--IHHE SC

: . :

DCRS7_Mu	AGRPAD RVER-----VT-----QALRSALDSCTS-----
DCRS7_Hu	SGRLQERAEQ-----VS-----RALQPALDSYFHPP-----
IL-17R_Hu	PGRMHRV GELSGDNYLRS---PGRQLRAALDRFRDWQVRC PDW
IL-17R_Mu	PGRMHHVREL TGDNYLQS---P SGRQLKEAVLRFQEWQTQCPDW
DCRS10	P-----PRG PL-----PTLQVVPL-----
DCRS10_Mu	P-----PRG PL-----PTLQVVPL-----
DCRS9_Hu	ATSWGRLGAR-----QRRQSRLELCSR-----
DCRS8_Hu	PGQHTRQGS R-----RNYFRSKSGRSLYVAICNMHQFIDE EPDW
IL-17R_Ce	ANVTQNISEA-----Q-----IHEWNLCASRMMSFFVRNP NW
DCRS6_Hu	VS-----AGKR-----SQACHDGCCSL-----
DCRS6_Ce	DSIDSRNNSK-----THSTD SGVSSLS S-----NS--

:

FIG. 1B

SEQUENCE SUBMISSION

SEQ ID NO: 1 is primate DCRS6 nucleotide sequence.
SEQ ID NO: 2 is primate DCRS6 polypeptide sequence.
SEQ ID NO: 3 is primate DCRS6 reverse translation.
SEQ ID NO: 4 is rodent DCRS6 nucleotide sequence.
SEQ ID NO: 5 is rodent DCRS6 polypeptide sequence.
SEQ ID NO: 6 is rodent DCRS6 reverse translation.
SEQ ID NO: 7 is primate DCRS7 nucleotide sequence.
SEQ ID NO: 8 is primate DCRS7 polypeptide sequence.
SEQ ID NO: 9 is primate DCRS7 reverse translation.
SEQ ID NO: 10 is rodent DCRS7 nucleotide sequence.
SEQ ID NO: 11 is rodent DCRS7 polypeptide sequence.
SEQ ID NO: 12 is rodent DCRS7 reverse translation.
SEQ ID NO: 13 is primate DCRS8 nucleotide sequence.
SEQ ID NO: 14 is primate DCRS8 polypeptide sequence.
SEQ ID NO: 15 is primate DCRS8 reverse translation.
SEQ ID NO: 16 is primate DCRS9 nucleotide sequence.
SEQ ID NO: 17 is primate DCRS9 polypeptide sequence.
SEQ ID NO: 18 is primate DCRS9 reverse translation.
SEQ ID NO: 19 is rodent DCRS9 nucleotide sequence.
SEQ ID NO: 20 is rodent DCRS9 polypeptide sequence.
SEQ ID NO: 21 is rodent DCRS9 reverse translation.
SEQ ID NO: 22 is primate DCRS10 nucleotide sequence.
SEQ ID NO: 23 is primate DCRS10 polypeptide sequence.
SEQ ID NO: 24 is primate DCRS10 reverse translation.
SEQ ID NO: 25 is rodent DCRS10 nucleotide sequence.
SEQ ID NO: 26 is rodent DCRS10 polypeptide sequence.
SEQ ID NO: 27 is rodent DCRS10 reverse translation.
SEQ ID NO: 28 is primate IL-17 receptor peptide sequence.
SEQ ID NO: 29 is rodent IL-17 receptor peptide sequence.
SEQ ID NO: 30 is worm IL-17 receptor peptide sequence.
SEQ ID NO: 31 is worm DCRS6 nucleotide sequence.

<110> Schering Corporation

<120> Mammalian Receptor Proteins; Related Reagents and
Methods

<130> DX01170K PCT

<140>

<141>

<150> US 60/206,862

<151> 2000-05-24

<160> 31

<170> PatentIn Ver. 2.0

<210> 1

<211> 1796

<212> DNA

<213> Unknown

<220>

<223> Description of Unknown Organism: primate; surmised
Homo sapiens

<220>
 <221> CDS
 <222> (4)..(1509)

<220>
 <221> mat_peptide
 <222> (46)..(1509)

```

<400> 1
gcg atg tcg ctc gtg ctg cta agc ctg gcc gcg ctg tgc agg agc gcc 48
    Met Ser Leu Val Leu Leu Ser Leu Ala Ala Leu Cys Arg Ser Ala
                    -10                    -5                    -1    1

gta ccc cga gag ccg acc gtt caa tgt ggc tct gaa act ggg cca tct 96
Val Pro Arg Glu Pro Thr Val Gln Cys Gly Ser Glu Thr Gly Pro Ser
                    5                    10                    15

cca gag tgg atg cta caa cat gat cta atc ccg gga gac ttg agg gac 144
Pro Glu Trp Met Leu Gln His Asp Leu Ile Pro Gly Asp Leu Arg Asp
                    20                    25                    30

ctc cga gta gaa cct gtt aca act agt gtt gca aca ggg gac tat tca 192
Leu Arg Val Glu Pro Val Thr Thr Ser Val Ala Thr Gly Asp Tyr Ser
                    35                    40                    45

att ttg atg aat gta agc tgg gta ctc cgg gca gat gcc agc atc cgc 240
Ile Leu Met Asn Val Ser Trp Val Leu Arg Ala Asp Ala Ser Ile Arg
                    50                    55                    60                    65

ttg ttg aag gcc acc aag att tgt gtg acg ggc aaa agc aac ttc cag 288
Leu Leu Lys Ala Thr Lys Ile Cys Val Thr Gly Lys Ser Asn Phe Gln
                    70                    75                    80

tcc tac agc tgt gtg agg tgc aat tac aca gag gcc ttc cag act cag 336
Ser Tyr Ser Cys Val Arg Cys Asn Tyr Thr Glu Ala Phe Gln Thr Gln
                    85                    90                    95

acc aga ccc tct ggt ggt aaa tgg aca ttt tcc tat atc ggc ttc cct 384
Thr Arg Pro Ser Gly Gly Lys Trp Thr Phe Ser Tyr Ile Gly Phe Pro
                    100                    105                    110

gta gag ctg aac aca gtc tat ttc att ggg gcc cat aat att cct aat 432
Val Glu Leu Asn Thr Val Tyr Phe Ile Gly Ala His Asn Ile Pro Asn
                    115                    120                    125

gca aat atg aat gaa gat ggc cct tcc atg tct gtg aat ttc acc tca 480
Ala Asn Met Asn Glu Asp Gly Pro Ser Met Ser Val Asn Phe Thr Ser
                    130                    135                    140                    145

cca ggc tgc cta gac cac ata atg aaa tat aaa aaa aag tgt gtc aag 528
Pro Gly Cys Leu Asp His Ile Met Lys Tyr Lys Lys Lys Cys Val Lys
                    150                    155                    160

gcc gga agc ctg tgg gat ccg aac atc act gct tgt aag aag aat gag 576
Ala Gly Ser Leu Trp Asp Pro Asn Ile Thr Ala Cys Lys Lys Asn Glu
                    165                    170                    175

gag aca gta gaa gtg aac ttc aca acc act ccc ctg gga aac aga tac 624
Glu Thr Val Glu Val Asn Phe Thr Thr Thr Pro Leu Gly Asn Arg Tyr
                    180                    185                    190

```

atg gct ctt atc caa cac agc act atc atc ggg ttt tct cag gtg ttt	672
Met Ala Leu Ile Gln His Ser Thr Ile Ile Gly Phe Ser Gln Val Phe	
195 200 205	
gag cca cac cag aag aaa caa acg cga gct tca gtg gtg att cca gtg	720
Glu Pro His Gln Lys Lys Gln Thr Arg Ala Ser Val Val Ile Pro Val	
210 215 220 225	
act ggg gat agt gaa ggt gct acg gtg cag ctg act cca tat ttt cct	768
Thr Gly Asp Ser Glu Gly Ala Thr Val Gln Leu Thr Pro Tyr Phe Pro	
230 235 240	
act tgt ggc agc gac tgc atc cga cat aaa gga aca gtt gtg ctc tgc	816
Thr Cys Gly Ser Asp Cys Ile Arg His Lys Gly Thr Val Val Leu Cys	
245 250 255	
cca caa aca ggc gtc cct ttc cct ctg gat aac aac aaa agc aag ccg	864
Pro Gln Thr Gly Val Pro Phe Pro Leu Asp Asn Asn Lys Ser Lys Pro	
260 265 270	
gga ggc tgg ctg cct ctc ctc ctg ctg tct ctg ctg gtg gcc aca tgg	912
Gly Gly Trp Leu Pro Leu Leu Leu Leu Ser Leu Leu Val Ala Thr Trp	
275 280 285	
gtg ctg gtg gca ggg atc tat cta atg tgg agg cac gaa agg atc aag	960
Val Leu Val Ala Gly Ile Tyr Leu Met Trp Arg His Glu Arg Ile Lys	
290 295 300 305	
aag act tcc ttt tct acc acc aca cta ctg ccc ccc att aag gtt ctt	1008
Lys Thr Ser Phe Ser Thr Thr Thr Leu Leu Pro Pro Ile Lys Val Leu	
310 315 320	
gtg gtt tac cca tct gaa ata tgt ttc cat cac aca att tgt tac ttc	1056
Val Val Tyr Pro Ser Glu Ile Cys Phe His His Thr Ile Cys Tyr Phe	
325 330 335	
act gaa ttt ctt caa aac cat tgc aga agt gag gtc atc ctt gaa aag	1104
Thr Glu Phe Leu Gln Asn His Cys Arg Ser Glu Val Ile Leu Glu Lys	
340 345 350	
tgg cag aaa aag aaa ata gca gag atg ggt cca gtg cag tgg ctt gcc	1152
Trp Gln Lys Lys Lys Ile Ala Glu Met Gly Pro Val Gln Trp Leu Ala	
355 360 365	
act caa aag aag gca gca gac aaa gtc gtc ttc ctt ctt tcc aat gac	1200
Thr Gln Lys Lys Ala Ala Asp Lys Val Val Phe Leu Leu Ser Asn Asp	
370 375 380 385	
gtc aac agt gtg tgc gat ggt acc tgt ggc aag agc gag ggc agt ccc	1248
Val Asn Ser Val Cys Asp Gly Thr Cys Gly Lys Ser Glu Gly Ser Pro	
390 395 400	
agt gag aac tct caa gac ctc ttc ccc ctt gcc ttt aac ctt ttc tgc	1296
Ser Glu Asn Ser Gln Asp Leu Phe Pro Leu Ala Phe Asn Leu Phe Cys	
405 410 415	
agt gat cta aga agc cag att cat ctg cac aaa tac gtg gtg gtc tac	1344
Ser Asp Leu Arg Ser Gln Ile His Leu His Lys Tyr Val Val Val Tyr	
420 425 430	

```

ttt aga gag att gat aca aaa gac gat tac aat gct ctc agt gtc tgc 1392
Phe Arg Glu Ile Asp Thr Lys Asp Asp Tyr Asn Ala Leu Ser Val Cys
    435                440                445

ccc aag tac cac ctc atg aag gat gcc act gct ttc tgt gca gaa ctt 1440
Pro Lys Tyr His Leu Met Lys Asp Ala Thr Ala Phe Cys Ala Glu Leu
    450                455                460                465

ctc cat gtc aag cag cag gtg tca gca gga aaa aga tca caa gcc tgc 1488
Leu His Val Lys Gln Gln Val Ser Ala Gly Lys Arg Ser Gln Ala Cys
                470                475                480

cac gat ggc tgc tgc tcc ttg tagccacccc atgagaagca agagacctta 1539
His Asp Gly Cys Cys Ser Leu
                485

aaggttcct atcccaccaa ttacagggaa aaaacgtgtg atgatcctga agcttactat 1599

gcagcctaca aacagcctta gtaattaaaa cattttatac caataaaatt ttcaaatt 1659

gctaactaat gtagcattaa ctaacgattg gaaactacat ttacaacttc aaagctgttt 1719

tatacataga aatcaattac agctttaatt gaaaactgta accattttga taatgcaaca 1779

ataaagcatc ttcagcc 1796

<210> 2
<211> 502
<212> PRT
<213> Unknown

<400> 2
Met Ser Leu Val Leu Leu Ser Leu Ala Ala Leu Cys Arg Ser Ala Val
    -10                -5                -1    1

Pro Arg Glu Pro Thr Val Gln Cys Gly Ser Glu Thr Gly Pro Ser Pro
    5                10                15

Glu Trp Met Leu Gln His Asp Leu Ile Pro Gly Asp Leu Arg Asp Leu
    20                25                30

Arg Val Glu Pro Val Thr Thr Ser Val Ala Thr Gly Asp Tyr Ser Ile
    35                40                45                50

Leu Met Asn Val Ser Trp Val Leu Arg Ala Asp Ala Ser Ile Arg Leu
    55                60                65

Leu Lys Ala Thr Lys Ile Cys Val Thr Gly Lys Ser Asn Phe Gln Ser
    70                75                80

Tyr Ser Cys Val Arg Cys Asn Tyr Thr Glu Ala Phe Gln Thr Gln Thr
    85                90                95

Arg Pro Ser Gly Gly Lys Trp Thr Phe Ser Tyr Ile Gly Phe Pro Val
    100                105                110

Glu Leu Asn Thr Val Tyr Phe Ile Gly Ala His Asn Ile Pro Asn Ala
    115                120                125                130

```

Asn Met Asn Glu Asp Gly Pro Ser Met Ser Val Asn Phe Thr Ser Pro
 135 140 145
 Gly Cys Leu Asp His Ile Met Lys Tyr Lys Lys Lys Cys Val Lys Ala
 150 155 160
 Gly Ser Leu Trp Asp Pro Asn Ile Thr Ala Cys Lys Lys Asn Glu Glu
 165 170 175
 Thr Val Glu Val Asn Phe Thr Thr Thr Pro Leu Gly Asn Arg Tyr Met
 180 185 190
 Ala Leu Ile Gln His Ser Thr Ile Ile Gly Phe Ser Gln Val Phe Glu
 195 200 205 210
 Pro His Gln Lys Lys Gln Thr Arg Ala Ser Val Val Ile Pro Val Thr
 215 220 225
 Gly Asp Ser Glu Gly Ala Thr Val Gln Leu Thr Pro Tyr Phe Pro Thr
 230 235 240
 Cys Gly Ser Asp Cys Ile Arg His Lys Gly Thr Val Val Leu Cys Pro
 245 250 255
 Gln Thr Gly Val Pro Phe Pro Leu Asp Asn Asn Lys Ser Lys Pro Gly
 260 265 270
 Gly Trp Leu Pro Leu Leu Leu Leu Ser Leu Leu Val Ala Thr Trp Val
 275 280 285 290
 Leu Val Ala Gly Ile Tyr Leu Met Trp Arg His Glu Arg Ile Lys Lys
 295 300 305
 Thr Ser Phe Ser Thr Thr Thr Leu Leu Pro Pro Ile Lys Val Leu Val
 310 315 320
 Val Tyr Pro Ser Glu Ile Cys Phe His His Thr Ile Cys Tyr Phe Thr
 325 330 335
 Glu Phe Leu Gln Asn His Cys Arg Ser Glu Val Ile Leu Glu Lys Trp
 340 345 350
 Gln Lys Lys Lys Ile Ala Glu Met Gly Pro Val Gln Trp Leu Ala Thr
 355 360 365 370
 Gln Lys Lys Ala Ala Asp Lys Val Val Phe Leu Leu Ser Asn Asp Val
 375 380 385
 Asn Ser Val Cys Asp Gly Thr Cys Gly Lys Ser Glu Gly Ser Pro Ser
 390 395 400
 Glu Asn Ser Gln Asp Leu Phe Pro Leu Ala Phe Asn Leu Phe Cys Ser
 405 410 415
 Asp Leu Arg Ser Gln Ile His Leu His Lys Tyr Val Val Val Tyr Phe
 420 425 430
 Arg Glu Ile Asp Thr Lys Asp Asp Tyr Asn Ala Leu Ser Val Cys Pro
 435 440 445 450

Lys Tyr His Leu Met Lys Asp Ala Thr Ala Phe Cys Ala Glu Leu Leu
 455 460 465

His Val Lys Gln Gln Val Ser Ala Gly Lys Arg Ser Gln Ala Cys His
 470 475 480

Asp Gly Cys Cys Ser Leu
 485

<210> 3
 <211> 1506
 <212> DNA
 <213> reverse translation

<220>
 <221> misc_feature
 <222> (1)..(1506)
 <223> n may be a, c, g, or t

<400> 3
 atgwsnytnng tnytnytnws nytngcngcn ytntgymgnw sngcngtncc nmnggarccn 60
 acngtncart gyggwnwsnga racnggnccn wsnccngart ggatgytnca rcaygayytn 120
 athccnggng ayytnmgnga yytnmgngtn garccngtna cnacnwsngt ngcnacnggn 180
 gaytaywsna thytnatgaa ygtnwsntgg gtnytnmgng cngaygcnws nathmgnytn 240
 ytnaargcna cnaarathtg ygtnacnggn aarwsnaayt tycarwsnta ywsntgygtn 300
 mgntgyaayt ayacngargc nttycaracn caracnmgnc cnwsngggngg naartggacn 360
 ttywsntaya thggnttycc ngtngarytn aayacngtnt ayttyathgg ngcncayaay 420
 athccnaayg cnaayatgaa ygargayggn ccnwsnatgw sngtnaaytt yacnwsnccn 480
 ggntgyytnng aycayathat gaartayaar aaraartgyg tnaargcngg nwsnytnntgg 540
 gayccnaaya thacngcntg yaaraaraay gargaracng tngargtnaa yttyacnacn 600
 acncnytnng gnaaymgnta yatggcnytn athcarcayw snacnathat hggnttywsn 660
 cargtnnttyg arcncayca raaraarcar acnmngncnw sngtnngtnat hccngtnacn 720
 ggngaywsng arggngcnac ngtnrcarytn acncntayt tyccnacntg yggwnwsngay 780
 tgyathmgnc ayaarggnac ngtnngtnytn tgyccncara cnggngtncc nttyccnytn 840
 gayaayaaya arwsnaarcc nggnggntgg ytnccnytny tnytnytnws nytnytnngtn 900
 gcnacntggg tnytnngtngc nggnathtay ytnatgtggm gncaygarmg nathaaraar 960
 acnwsnttyw snacnacnac nytnytnccn ccnathaarg tnytnngtngt ntayccnwsn 1020
 garathtgyt tycaycayac nathtgytay ttyacngart tyytncaraa ycaytgymgn 1080
 wsgargtna thytngaraa rtggcaraar aaraarathg cngaratggg nccngtncar 1140

tggytngcna cncaraaraa rgcngcngay aargtngtnt tyytnytnws naaygaygt 1200
 aaywsngtnt gygayggnac ntgyggnaar wsngarggnw snccnwsnga raaywsncar 1260
 gayytnttyc cnytnngcntt yaayytntty tgywsngayy tnmgnwsnca rathcayytn 1320
 cayaartayg tngtngtnta yttymngar athgayacna argaygayta yaaygcnytn 1380
 wsngtntgyg cnaartayca yytnatgaar gaygcnacng cnttytgygc ngarytnytn 1440
 caygtnaarc arcargtnws ngcnggnaar mgnwsncarg cntgycayga yggntgytgy 1500
 wsnytn 1506

<210> 4
 <211> 637
 <212> DNA
 <213> Unknown

<220>
 <223> Description of Unknown Organism:rodent; surmised
 Mus musculus

<220>
 <221> CDS
 <222> (1)..(210)

<400> 4
 gat ttc agc agc cag acg cat ctg cac aaa tac ctg gag gtc tat ctt 48
 Asp Phe Ser Ser Gln Thr His Leu His Lys Tyr Leu Glu Val Tyr Leu
 1 5 10 15
 ggg gga gca gac ctc aaa ggc gac tat aat gcc ctg agt gtc tgc ccc 96
 Gly Gly Ala Asp Leu Lys Gly Asp Tyr Asn Ala Leu Ser Val Cys Pro
 20 25 30
 caa tat cat ctc atg aag gac gcc aca gct ttc cac aca gaa ctt ctc 144
 Gln Tyr His Leu Met Lys Asp Ala Thr Ala Phe His Thr Glu Leu Leu
 35 40 45
 aag gct acg cag agc atg tca gtg aag aaa cgc tca caa gcc tgc cat 192
 Lys Ala Thr Gln Ser Met Ser Val Lys Lys Arg Ser Gln Ala Cys His
 50 55 60
 gat agc tgt tca ccc ttg tagtccaccc gggggaatag agactctgaa 240
 Asp Ser Cys Ser Pro Leu
 65 70
 gccttcctac tctcccttcc agtgacaaat gctgtgtgac gactctgaaa tgtgtgggag 300
 aggctgtgtg gaggtagtgc tatgtacaaa cttgctttaa aactggagtt tgcaaagtca 360
 acctgagcat acacgcctga ggctagtcat tggctggatt tatgaagaca acacagttac 420
 agacaataat gagtgggacc tacatttggg atatacccaa agctgggtaa tgattatcac 480
 tgagaaccac gcactctggc catgaggtaa tacggcactt ccctgtcagg ctgtctgtca 540
 ggttgggtct gtcttgact gcccatgtct tatgtgtcac gtagaccgtt ttgtaacatt 600

ttaatctgtt aatgaataat ccgtttggga ggctctc

637

<210> 5
 <211> 70
 <212> PRT
 <213> Unknown

<400> 5
 Asp Phe Ser Ser Gln Thr His Leu His Lys Tyr Leu Glu Val Tyr Leu
 1 5 10 15
 Gly Gly Ala Asp Leu Lys Gly Asp Tyr Asn Ala Leu Ser Val Cys Pro
 20 25 30
 Gln Tyr His Leu Met Lys Asp Ala Thr Ala Phe His Thr Glu Leu Leu
 35 40 45
 Lys Ala Thr Gln Ser Met Ser Val Lys Lys Arg Ser Gln Ala Cys His
 50 55 60
 Asp Ser Cys Ser Pro Leu
 65 70

<210> 6
 <211> 210
 <212> DNA
 <213> reverse translation

<220>
 <221> misc_feature
 <222> (1)..(210)
 <223> n may be a, c, g, or t

<400> 6
 gayttywsnw sncaracnca yytncayaar tayytnngarg tntayytnng nggngcngay 60
 ytnaarggng aytayaaygc nytnwsngtn tgyccncart aycayytnat gaargaygcn 120
 acngcnttyc ayacngaryt nytnaargcn acncarwsna tgwsngtnaa raarmgnwsn 180
 cargcntgyc aygaywsntg ywsnccnytn 210

<210> 7
 <211> 2308
 <212> DNA
 <213> Unknown

<220>
 <223> Description of Unknown Organism:primate; surmised
 Homo sapiens

<220>
 <221> CDS
 <222> (181)..(2289)

<220>

<221> mat_peptide
 <222> (241)..(2289)

<220>
 <221> misc_feature
 <222> (664)
 <223> Xaa translation depends on genetic code

<400> 7

```

gagtcaggac tcccaggaca gagagtgcac aaactaccca gcacagcccc ctccgcccc 60
tctggaggct gaagagggat tccagccct gccacccaca gacacgggct gactgggggtg 120
tctgcccccc ttgggggcan ccacagggcc tcaggcctgg gtgccacctg gcactagaag 180

atg cct gtg ccc tgg ttc ttg ctg tcc ttg gca ctg ggc cga agc cag 228
Met Pro Val Pro Trp Phe Leu Leu Ser Leu Ala Leu Gly Arg Ser Gln
-20 -15 -10 -5

tgg atc ctt tct ctg gag agg ctt gtg ggg cct cag gac gct acc cac 276
Trp Ile Leu Ser Leu Glu Arg Leu Val Gly Pro Gln Asp Ala Thr His
-1 1 5 10

tgc tct ccg ggc ctc tcc tgc cgc ctc tgg gac agt gac ata ctc tgc 324
Cys Ser Pro Gly Leu Ser Cys Arg Leu Trp Asp Ser Asp Ile Leu Cys
15 20 25

ctg cct ggg gac atc gtg cct gct ccg ggc ccc gtg ctg gcg cct acg 372
Leu Pro Gly Asp Ile Val Pro Ala Pro Gly Pro Val Leu Ala Pro Thr
30 35 40

cac ctg cag aca gag ctg gtg ctg agg tgc cag aag gag acc gac tgt 420
His Leu Gln Thr Glu Leu Val Leu Arg Cys Gln Lys Glu Thr Asp Cys
45 50 55 60

gac ctc tgt ctg cgt gtg gct gtc cac ttg gcc gtg cat ggg cac tgg 468
Asp Leu Cys Leu Arg Val Ala Val His Leu Ala Val His Gly His Trp
65 70 75

gaa gag cct gaa gat gag gaa aag ttt gga gga gca gct gac tta ggg 516
Glu Glu Pro Glu Asp Glu Glu Lys Phe Gly Gly Ala Ala Asp Leu Gly
80 85 90

gtg gag gag cct agg aat gcc tct ctc cag gcc caa gtc gtg ctc tcc 564
Val Glu Glu Pro Arg Asn Ala Ser Leu Gln Ala Gln Val Val Leu Ser
95 100 105

ttc cag gcc tac cct act gcc cgc tgc gtc ctg ctg gag gtg caa gtg 612
Phe Gln Ala Tyr Pro Thr Ala Arg Cys Val Leu Leu Glu Val Gln Val
110 115 120

cct gct gcc ctt gtg cag ttt ggt cag tct gtg ggc tct gtg gta tat 660
Pro Ala Ala Leu Val Gln Phe Gly Gln Ser Val Gly Ser Val Val Tyr
125 130 135 140

gac tgc ttc gag gct gcc cta ggg agt gag gta cga atc tgg tcc tat 708
Asp Cys Phe Glu Ala Ala Leu Gly Ser Glu Val Arg Ile Trp Ser Tyr
145 150 155

act cag ccc agg tac gag aag gaa ctc aac cac aca cag cag ctg cct 756

```

Thr	Gln	Pro	Arg	Tyr	Glu	Lys	Glu	Leu	Asn	His	Thr	Gln	Gln	Leu	Pro		
			160					165					170				
gac	tgc	agg	ggg	ctc	gaa	gtc	tgg	aac	agc	atc	ccg	agc	tgc	tgg	gcc	804	
Asp	Cys	Arg	Gly	Leu	Glu	Val	Trp	Asn	Ser	Ile	Pro	Ser	Cys	Trp	Ala		
		175					180					185					
ctg	ccc	tgg	ctc	aac	gtg	tca	gca	gat	ggc	gac	aac	gtg	cat	ctg	gtt	852	
Leu	Pro	Trp	Leu	Asn	Val	Ser	Ala	Asp	Gly	Asp	Asn	Val	His	Leu	Val		
	190					195					200						
ctg	aat	gtc	tct	gag	gag	cag	cac	ttc	ggc	ctc	tcc	ctg	tac	tgg	aat	900	
Leu	Asn	Val	Ser	Glu	Glu	Gln	His	Phe	Gly	Leu	Ser	Leu	Tyr	Trp	Asn		
205				210					215						220		
cag	gtc	cag	ggc	ccc	cca	aaa	ccc	cgg	tgg	cac	aaa	aac	ctg	act	gga	948	
Gln	Val	Gln	Gly	Pro	Pro	Lys	Pro	Arg	Trp	His	Lys	Asn	Leu	Thr	Gly		
			225					230						235			
ccg	cag	atc	att	acc	ttg	aac	cac	aca	gac	ctg	gtt	ccc	tgc	ctc	tgt	996	
Pro	Gln	Ile	Ile	Thr	Leu	Asn	His	Thr	Asp	Leu	Val	Pro	Cys	Leu	Cys		
		240						245					250				
att	cag	gtg	tgg	cct	ctg	gaa	cct	gac	tcc	gtt	agg	acg	aac	atc	tgc	1044	
Ile	Gln	Val	Trp	Pro	Leu	Glu	Pro	Asp	Ser	Val	Arg	Thr	Asn	Ile	Cys		
		255				260						265					
ccc	ttc	agg	gag	gac	ccc	cgc	gca	cac	cag	aac	ctc	tgg	caa	gcc	gcc	1092	
Pro	Phe	Arg	Glu	Asp	Pro	Arg	Ala	His	Gln	Asn	Leu	Trp	Gln	Ala	Ala		
	270					275					280						
cga	ctg	cga	ctg	ctg	acc	ctg	cag	agc	tgg	ctg	ctg	gac	gca	ccg	tgc	1140	
Arg	Leu	Arg	Leu	Leu	Thr	Leu	Gln	Ser	Trp	Leu	Leu	Asp	Ala	Pro	Cys		
285					290				295						300		
tcg	ctg	ccc	gca	gaa	gcg	gca	ctg	tgc	tgg	cgg	gct	ccg	ggc	ggg	gac	1188	
Ser	Leu	Pro	Ala	Glu	Ala	Ala	Leu	Cys	Trp	Arg	Ala	Pro	Gly	Gly	Asp		
			305					310						315			
ccc	tgc	cag	cca	ctg	gtc	cca	ccg	ctt	tcc	tgg	gag	aat	gtc	act	gtg	1236	
Pro	Cys	Gln	Pro	Leu	Val	Pro	Pro	Leu	Ser	Trp	Glu	Asn	Val	Thr	Val		
			320					325					330				
gac	gtg	aac	agc	tcg	gag	aag	ctg	cag	ctg	cag	gag	tgc	ttg	tgg	gct	1284	
Asp	Val	Asn	Ser	Ser	Glu	Lys	Leu	Gln	Leu	Gln	Glu	Cys	Leu	Trp	Ala		
		335					340					345					
gac	tcc	ctg	ggg	cct	ctc	aaa	gac	gat	gtg	cta	ctg	ttg	gag	aca	cga	1332	
Asp	Ser	Leu	Gly	Pro	Leu	Lys	Asp	Asp	Val	Leu	Leu	Leu	Glu	Thr	Arg		
	350					355					360						
ggc	ccc	cag	gac	aac	aga	tcc	ctc	tgt	gcc	ttg	gaa	ccc	agt	ggc	tgt	1380	
Gly	Pro	Gln	Asp	Asn	Arg	Ser	Leu	Cys	Ala	Leu	Glu	Pro	Ser	Gly	Cys		
	365				370				375						380		
act	tca	cta	ccc	agc	aaa	gcc	tcc	acg	agg	gca	gct	cgc	ctt	gga	gag	1428	
Thr	Ser	Leu	Pro	Ser	Lys	Ala	Ser	Thr	Arg	Ala	Ala	Arg	Leu	Gly	Glu		
				385				390						395			
tac	tta	cta	caa	gac	ctg	cag	tca	ggc	cag	tgt	ctg	cag	cta	tgg	gac	1476	

Tyr	Leu	Leu	Gln	Asp	Leu	Gln	Ser	Gly	Gln	Cys	Leu	Gln	Leu	Trp	Asp		
			400					405					410				
gat	gac	ttg	gga	gcg	cta	tgg	gcc	tgc	ccc	atg	gac	aaa	tac	atc	cac	1524	
Asp	Asp	Leu	Gly	Ala	Leu	Trp	Ala	Cys	Pro	Met	Asp	Lys	Tyr	Ile	His		
		415					420					425					
aag	cgc	tgg	gcc	ctc	gtg	tgg	ctg	gcc	tgc	cta	ctc	ttt	gcc	gct	gcg	1572	
Lys	Arg	Trp	Ala	Leu	Val	Trp	Leu	Ala	Cys	Leu	Leu	Phe	Ala	Ala	Ala		
	430					435					440						
ctt	tcc	ctc	atc	ctc	ctt	ctc	aaa	aag	gat	cac	gcg	aaa	ggg	tgg	ctg	1620	
Leu	Ser	Leu	Ile	Leu	Leu	Leu	Lys	Lys	Asp	His	Ala	Lys	Gly	Trp	Leu		
445					450					455					460		
agg	ctc	ttg	aaa	cag	gac	gtc	cgc	tgc	ggg	gcg	gcc	gcc	agg	ggc	cgc	1668	
Arg	Leu	Leu	Lys	Gln	Asp	Val	Arg	Ser	Gly	Ala	Ala	Ala	Arg	Gly	Arg		
			465					470					475				
gcg	gct	ctg	ctc	ctc	tac	tca	gcc	gat	gac	tgc	ggg	ttc	gag	cgc	ctg	1716	
Ala	Ala	Leu	Leu	Leu	Tyr	Ser	Ala	Asp	Asp	Ser	Gly	Phe	Glu	Arg	Leu		
		480						485					490				
gtg	ggc	gcc	ctg	gcg	tgc	gcc	ctg	tgc	cag	ctg	ccg	ctg	cgc	gtg	gcc	1764	
Val	Gly	Ala	Leu	Ala	Ser	Ala	Leu	Cys	Gln	Leu	Pro	Leu	Arg	Val	Ala		
	495					500					505						
gta	gac	ctg	tgg	agc	cgt	cgt	gaa	ctg	agc	gcg	cag	ggg	ccc	gtg	gct	1812	
Val	Asp	Leu	Trp	Ser	Arg	Arg	Glu	Leu	Ser	Ala	Gln	Gly	Pro	Val	Ala		
	510					515					520						
tgg	ttt	cac	gcg	cag	cgg	cgc	cag	acc	ctg	cag	gag	ggc	ggc	gtg	gtg	1860	
Trp	Phe	His	Ala	Gln	Arg	Arg	Gln	Thr	Leu	Gln	Glu	Gly	Gly	Val	Val		
525				530					535					540			
gtc	ttg	ctc	ttc	tct	ccc	ggg	gcg	gtg	gcg	ctg	tgc	agc	gag	tgg	cta	1908	
Val	Leu	Leu	Phe	Ser	Pro	Gly	Ala	Val	Ala	Leu	Cys	Ser	Glu	Trp	Leu		
			545					550					555				
cag	gat	ggg	gtg	tcc	ggg	ccc	ggg	gcg	cac	ggc	ccg	cac	gac	gcc	ttc	1956	
Gln	Asp	Gly	Val	Ser	Gly	Pro	Gly	Ala	His	Gly	Pro	His	Asp	Ala	Phe		
		560					565					570					
cgc	gcc	tgc	ctc	agc	tgc	gtg	ctg	ccc	gac	ttc	ttg	cag	ggc	cgg	gcg	2004	
Arg	Ala	Ser	Leu	Ser	Cys	Val	Leu	Pro	Asp	Phe	Leu	Gln	Gly	Arg	Ala		
		575				580						585					
ccc	ggc	agc	tac	gtg	ggg	gcc	tgc	ttc	gac	agg	ctg	ctc	cac	ccg	gac	2052	
Pro	Gly	Ser	Tyr	Val	Gly	Ala	Cys	Phe	Asp	Arg	Leu	Leu	His	Pro	Asp		
	590					595					600						
gcc	gta	ccc	gcc	ctt	ttc	cgc	acc	gtg	ccc	gtc	ttc	aca	ctg	ccc	tcc	2100	
Ala	Val	Pro	Ala	Leu	Phe	Arg	Thr	Val	Pro	Val	Phe	Thr	Leu	Pro	Ser		
	605				610					615					620		
caa	ctg	cca	gac	ttc	ctg	ggg	gcc	ctg	cag	cag	cct	cgc	gcc	ccg	cgt	2148	
Gln	Leu	Pro	Asp	Phe	Leu	Gly	Ala	Leu	Gln	Gln	Pro	Arg	Ala	Pro	Arg		
			625					630					635				
tcc	ggg	cgg	ctc	caa	gag	aga	gcg	gag	caa	gtg	tcc	cgg	gcc	ctt	cag	2196	

Ser Gly Arg Leu Gln Glu Arg Ala Glu Gln Val Ser Arg Ala Leu Gln
 640 645 650

cca gcc ctg gat agc tac ttc cat ccc ccg ggg acn tcc gcg ccg gga 2244
 Pro Ala Leu Asp Ser Tyr Phe His Pro Pro Gly Xaa Ser Ala Pro Gly
 655 660 665

cgc ggg gtg gga cca ggg gcg gga cct ggg gcg ggg gac ggg act 2289
 Arg Gly Val Gly Pro Gly Ala Gly Pro Gly Ala Gly Asp Gly Thr
 670 675 680

taaataaagg cagacgctg 2308

<210> 8
 <211> 703
 <212> PRT
 <213> Unknown

<400> 8
 Met Pro Val Pro Trp Phe Leu Leu Ser Leu Ala Leu Gly Arg Ser Gln
 -20 -15 -10 -5

Trp Ile Leu Ser Leu Glu Arg Leu Val Gly Pro Gln Asp Ala Thr His
 -1 1 5 10

Cys Ser Pro Gly Leu Ser Cys Arg Leu Trp Asp Ser Asp Ile Leu Cys
 15 20 25

Leu Pro Gly Asp Ile Val Pro Ala Pro Gly Pro Val Leu Ala Pro Thr
 30 35 40

His Leu Gln Thr Glu Leu Val Leu Arg Cys Gln Lys Glu Thr Asp Cys
 45 50 55 60

Asp Leu Cys Leu Arg Val Ala Val His Leu Ala Val His Gly His Trp
 65 70 75

Glu Glu Pro Glu Asp Glu Glu Lys Phe Gly Gly Ala Ala Asp Leu Gly
 80 85 90

Val Glu Glu Pro Arg Asn Ala Ser Leu Gln Ala Gln Val Val Leu Ser
 95 100 105

Phe Gln Ala Tyr Pro Thr Ala Arg Cys Val Leu Leu Glu Val Gln Val
 110 115 120

Pro Ala Ala Leu Val Gln Phe Gly Gln Ser Val Gly Ser Val Val Tyr
 125 130 135 140

Asp Cys Phe Glu Ala Ala Leu Gly Ser Glu Val Arg Ile Trp Ser Tyr
 145 150 155

Thr Gln Pro Arg Tyr Glu Lys Glu Leu Asn His Thr Gln Gln Leu Pro
 160 165 170

Asp Cys Arg Gly Leu Glu Val Trp Asn Ser Ile Pro Ser Cys Trp Ala
 175 180 185

Leu Pro Trp Leu Asn Val Ser Ala Asp Gly Asp Asn Val His Leu Val

190	195	200
Leu Asn Val Ser Glu Glu Gln His Phe Gly Leu Ser Leu Tyr Trp Asn 205 210 215 220		
Gln Val Gln Gly Pro Pro Lys Pro Arg Trp His Lys Asn Leu Thr Gly 225 230 235		
Pro Gln Ile Ile Thr Leu Asn His Thr Asp Leu Val Pro Cys Leu Cys 240 245 250		
Ile Gln Val Trp Pro Leu Glu Pro Asp Ser Val Arg Thr Asn Ile Cys 255 260 265		
Pro Phe Arg Glu Asp Pro Arg Ala His Gln Asn Leu Trp Gln Ala Ala 270 275 280		
Arg Leu Arg Leu Leu Thr Leu Gln Ser Trp Leu Leu Asp Ala Pro Cys 285 290 295 300		
Ser Leu Pro Ala Glu Ala Ala Leu Cys Trp Arg Ala Pro Gly Gly Asp 305 310 315		
Pro Cys Gln Pro Leu Val Pro Pro Leu Ser Trp Glu Asn Val Thr Val 320 325 330		
Asp Val Asn Ser Ser Glu Lys Leu Gln Leu Gln Glu Cys Leu Trp Ala 335 340 345		
Asp Ser Leu Gly Pro Leu Lys Asp Asp Val Leu Leu Leu Glu Thr Arg 350 355 360		
Gly Pro Gln Asp Asn Arg Ser Leu Cys Ala Leu Glu Pro Ser Gly Cys 365 370 375 380		
Thr Ser Leu Pro Ser Lys Ala Ser Thr Arg Ala Ala Arg Leu Gly Glu 385 390 395		
Tyr Leu Leu Gln Asp Leu Gln Ser Gly Gln Cys Leu Gln Leu Trp Asp 400 405 410		
Asp Asp Leu Gly Ala Leu Trp Ala Cys Pro Met Asp Lys Tyr Ile His 415 420 425		
Lys Arg Trp Ala Leu Val Trp Leu Ala Cys Leu Leu Phe Ala Ala Ala 430 435 440		
Leu Ser Leu Ile Leu Leu Leu Lys Lys Asp His Ala Lys Gly Trp Leu 445 450 455 460		
Arg Leu Leu Lys Gln Asp Val Arg Ser Gly Ala Ala Ala Arg Gly Arg 465 470 475		
Ala Ala Leu Leu Leu Tyr Ser Ala Asp Asp Ser Gly Phe Glu Arg Leu 480 485 490		
Val Gly Ala Leu Ala Ser Ala Leu Cys Gln Leu Pro Leu Arg Val Ala 495 500 505		
Val Asp Leu Trp Ser Arg Arg Glu Leu Ser Ala Gln Gly Pro Val Ala		

510	515	520
Trp Phe His Ala Gln Arg Arg Gln Thr Leu Gln Glu Gly Gly Val Val		
525	530	535 540
Val Leu Leu Phe Ser Pro Gly Ala Val Ala Leu Cys Ser Glu Trp Leu		
	545	550 555
Gln Asp Gly Val Ser Gly Pro Gly Ala His Gly Pro His Asp Ala Phe		
	560	565 570
Arg Ala Ser Leu Ser Cys Val Leu Pro Asp Phe Leu Gln Gly Arg Ala		
	575	580 585
Pro Gly Ser Tyr Val Gly Ala Cys Phe Asp Arg Leu Leu His Pro Asp		
	590	595 600
Ala Val Pro Ala Leu Phe Arg Thr Val Pro Val Phe Thr Leu Pro Ser		
	605	610 615 620
Gln Leu Pro Asp Phe Leu Gly Ala Leu Gln Gln Pro Arg Ala Pro Arg		
	625	630 635
Ser Gly Arg Leu Gln Glu Arg Ala Glu Gln Val Ser Arg Ala Leu Gln		
	640	645 650
Pro Ala Leu Asp Ser Tyr Phe His Pro Pro Gly Xaa Ser Ala Pro Gly		
	655	660 665
Arg Gly Val Gly Pro Gly Ala Gly Pro Gly Ala Gly Asp Gly Thr		
	670	675 680

<210> 9

<211> 2109

<212> DNA

<213> reverse translation

<220>

<221> misc_feature

<222> (1)..(2109)

<223> n may be a, c, g, or t

<400> 9

atgccngtnc cntggttyt nytnwsnytn gcnytnggnm gnwsncartg gathytnwsn 60

ytngarmgny tngtnggncc ncargaycn acncaytgyw snccnggnytn nwsntgymgn 120

ytntgggayw sngayathyt ntgyytnccn ggngayathg tncngcncc nggnccngtn 180

ytngcncna cncayytnca racngarytn gtnytnmgnt gycaraarga racngaytgy 240

gayytnntgyy tnmngtngc ngtncaaytn gcngtncayg gncaytggga rgarcncgar 300

gaygargara arttyggngg ngcngcngay ytngngtng argarccnmg naaygcwnsn 360

ytncargcnc argtngtnyt nwsnttycar gcntayccna cngcnmgntg ygtnytnytn 420

gargtncarg tncngcngc nytngtncar ttyggncarw sngtnggnws ngtngtntay 480

gaytgyttyg argcngcnyt nggnwsngar gtnmgnaht ggwsntayac ncarccnmgn 540
 taygaraarg arytnaayca yacncarcar ytncngayt gymnggnyt ngargtntgg 600
 aaywsnathc cnwsntgytg ggcnytnccn tggytnaayg tnwsngcnga yggngayaay 660
 gtncayytn gnytnaaygt nwsngargar carcayttyg gnytnwsnyt ntaytggay 720
 cargtncarg gnccnecnaa rccnmgttg cayaaraay tnacnggncc ncarathath 780
 acnytnaayc ayacngayyt ngtnccntgy ytntgyathc argtntggcc nytngarccn 840
 gaywsngtnm gnacnaayat htgyccntty mgngargayc cnmgngcnca ycaraaytn 900
 tggcargcng cnmgnytnmg nytnytnacn ytncarwsnt ggytnytn ga ygcncntgy 960
 wsnytnccng cngargcngc nytnntgytg mgngcncng gngngaycc ntgycarccn 1020
 ytngtnccnc cnytnwsntg ggaraaygt acngtngayg tnaaywsnws ngaraaytn 1080
 carytncarg artgyytn ggcngaywsn ytnggncny tnaargayga ygtnytnytn 1140
 ytngaracnm gnggncnca rgayaaymgn wsnytnntgy cnytngarcc nwsngntgy 1200
 acnwsnytn cnwsnaargc nwsnacnmgn gcngcnmgny tngngarta yytnytn car 1260
 gayytn carw snggncartg yytn carytn tgggaygayg ayytn gngc nytn tgggcn 1320
 tgyccnatgg ayaartayat hcayaarmgn tgggnytn tntggytn ntgyytnytn 1380
 ttygcngcng cnytnwsnyt nathytnytn ytnaaraarg aycaygcnaa rggntgytn 1440
 mgnytnytna arcargaygt nmgnwsngn gcngcngcnm gngnmngnc ngcnytnytn 1500
 ytntaywsng cngaygayws nggnttygar mgnytngtng gngcnytn ntgyytnytn 1560
 tgy carytn cnytnmgnt ngcngtngay ytntggwsnm gnmngaryt nwsngncar 1620
 ggcncngtn cntggtyca ygcncarmgn mgncaracny tncargarg nggngtn 1680
 gtnytnytn tywsncng ngcngtn gnytnytn artggytnca rgayggngtn 1740
 wsnggncng gngcncaygg nccncaygay gcnttymgng cnwsnytnws ntgyytnytn 1800
 ccngaytty tncarggnmg ngcncngn wsntaygtng gngcntgyt ygaymgnytn 1860
 ytncayccng aygngtncc ngcnytnntty mgnacngtn cngnttyac nytnccnwsn 1920
 carytnccng aytyytnng ngcnytn carccnmng cncnmgnws nggnmnytn 1980
 cargarmng cngarcargt nwsnmngcn ytncarccng cnytngayws ntayttyay 2040
 ccncnggna cnwsngcnc nggnmngngn gtnggncng gngcnggnc nggngcngn 2100
 gayggnacn 2109

<210> 10
 <211> 2314
 <212> DNA

<213> Unknown

<220>

<223> Description of Unknown Organism:rodent; surmised
Mus musculus

<220>

<221> CDS

<222> (199) .. (2292)

<220>

<221> mat_peptide

<222> (259) .. (2292)

<400> 10

```

ccaaatcgaa agcacgggag ctgatactgg gcctggagtc caggctcact ggagtgggga 60

agcatggctg gagaggaatt ctagcccttg ctctctccca gggacacggg gctgattgtc 120

agcaggggag aggggtctgc ccccccttgg gggggcagga cggggcctca ggcctgggtg 180

ctgtccggca cctggaag atg cct gtg tcc tgg ttc ctg ctg tcc ttg gca 231
      Met Pro Val Ser Trp Phe Leu Leu Ser Leu Ala
      -20                               -15                -10

ctg ggc cga aac cct gtg gtc gtc tct ctg gag aga ctg atg gag cct 279
Leu Gly Arg Asn Pro Val Val Val Ser Leu Glu Arg Leu Met Glu Pro
      -5                               -1    1                5

cag gac act gca cgc tgc tct cta ggc ctc tcc tgc cac ctc tgg gat 327
Gln Asp Thr Ala Arg Cys Ser Leu Gly Leu Ser Cys His Leu Trp Asp
      10                               15                20

ggg gac gtg ctc tgc ctg cct gga agc ctc cag tct gcc cca ggc cct 375
Gly Asp Val Leu Cys Leu Pro Gly Ser Leu Gln Ser Ala Pro Gly Pro
      25                               30                35

gtg cta gtg cct acc cgc ctg cag acg gag ctg gtg ctg agg tgt cca 423
Val Leu Val Pro Thr Arg Leu Gln Thr Glu Leu Val Leu Arg Cys Pro
      40                               45                50                55

cag aag aca gat tgc gcc ctc tgt gtc cgt gtg gtg gtc cac ttg gcc 471
Gln Lys Thr Asp Cys Ala Leu Cys Val Arg Val Val Val His Leu Ala
      60                               65                70

gtg cat ggg cac tgg gca gag cct gaa gaa gct gga aag tct gat tca 519
Val His Gly His Trp Ala Glu Pro Glu Glu Ala Gly Lys Ser Asp Ser
      75                               80                85

gaa ctc cag gag tct agg aac gcc tct ctc cag gcc cag gtg gtg ctc 567
Glu Leu Gln Glu Ser Arg Asn Ala Ser Leu Gln Ala Gln Val Val Leu
      90                               95                100

tcc ttc cag gcc tac ccc atc gcc cgc tgt gcc ctg ctg gag gtc cag 615
Ser Phe Gln Ala Tyr Pro Ile Ala Arg Cys Ala Leu Leu Glu Val Gln
      105                               110                115

gtg ccc gct gac ctg gtg cag cct ggt cag tcc gtg ggt tct gcg gta 663
Val Pro Ala Asp Leu Val Gln Pro Gly Gln Ser Val Gly Ser Ala Val
      120                               125                130                135

```

ttt gac tgt ttc gag gct agt ctt ggg gct gag gta cag atc tgg tcc	711
Phe Asp Cys Phe Glu Ala Ser Leu Gly Ala Glu Val Gln Ile Trp Ser	
140 145 150	
tac acg aag ccc agg tac cag aaa gag ctc aac ctc aca cag cag ctg	759
Tyr Thr Lys Pro Arg Tyr Gln Lys Glu Leu Asn Leu Thr Gln Gln Leu	
155 160 165	
cct gac tgc agg ggt ctt gaa gtc cgg gac agc atc cag agc tgc tgg	807
Pro Asp Cys Arg Gly Leu Glu Val Arg Asp Ser Ile Gln Ser Cys Trp	
170 175 180	
gtc ctg ccc tgg ctc aat gtg tct aca gat ggt gac aat gtc ctt ctg	855
Val Leu Pro Trp Leu Asn Val Ser Thr Asp Gly Asp Asn Val Leu Leu	
185 190 195	
aca ctg gat gtc tct gag gag cag gac ttt agc ttc tta ctg tac ctg	903
Thr Leu Asp Val Ser Glu Glu Gln Asp Phe Ser Phe Leu Leu Tyr Leu	
200 205 210 215	
cgt cca gtc ccg gat gct ctc aaa tcc ttg tgg tac aaa aac ctg act	951
Arg Pro Val Pro Asp Ala Leu Lys Ser Leu Trp Tyr Lys Asn Leu Thr	
220 225 230	
gga cct cag aac att act tta aac cac aca gac ctg gtt ccc tgc ctc	999
Gly Pro Gln Asn Ile Thr Leu Asn His Thr Asp Leu Val Pro Cys Leu	
235 240 245	
tgc att cag gtg tgg tcg cta gag cca gac tct gag agg gtc gaa ttc	1047
Cys Ile Gln Val Trp Ser Leu Glu Pro Asp Ser Glu Arg Val Glu Phe	
250 255 260	
tgc ccc ttc cgg gaa gat ccc ggt gca cac agg aac ctc tgg cac ata	1095
Cys Pro Phe Arg Glu Asp Pro Gly Ala His Arg Asn Leu Trp His Ile	
265 270 275	
gcc agg ctg cgg gta ctg tcc cca ggg gta tgg cag cta gat gcg cct	1143
Ala Arg Leu Arg Val Leu Ser Pro Gly Val Trp Gln Leu Asp Ala Pro	
280 285 290 295	
tgc tgt ctg ccg ggc aag gta aca ctg tgc tgg cag gca cca gac cag	1191
Cys Cys Leu Pro Gly Lys Val Thr Leu Cys Trp Gln Ala Pro Asp Gln	
300 305 310	
agt ccc tgc cag cca ctt gtg cca cca gtg ccc cag aag aac gcc act	1239
Ser Pro Cys Gln Pro Leu Val Pro Pro Val Pro Gln Lys Asn Ala Thr	
315 320 325	
gtg aat gag cca caa gat ttc cag ttg gtg gca ggc cac ccc aac ctc	1287
Val Asn Glu Pro Gln Asp Phe Gln Leu Val Ala Gly His Pro Asn Leu	
330 335 340	
tgt gtc cag gtg agc acc tgg gag aag gtt cag ctg caa gcg tgc ttg	1335
Cys Val Gln Val Ser Thr Trp Glu Lys Val Gln Leu Gln Ala Cys Leu	
345 350 355	
tgg gct gac tcc ttg ggg ccc ttc aag gat gat atg ctg tta gtg gag	1383
Trp Ala Asp Ser Leu Gly Pro Phe Lys Asp Asp Met Leu Leu Val Glu	
360 365 370 375	

atg aaa acc ggc ctc aac aac aca tca gtc tgt gcc ttg gaa ccc agt	1431
Met Lys Thr Gly Leu Asn Asn Thr Ser Val Cys Ala Leu Glu Pro Ser	
380 385 390	
ggc tgt aca cca ctg ccc agc atg gcc tcc acg aga gct gct cgc ctg	1479
Gly Cys Thr Pro Leu Pro Ser Met Ala Ser Thr Arg Ala Arg Leu	
395 400 405	
gga gag gag ttg ctg caa gac ttc cga tca cac cag tgt atg cag ctg	1527
Gly Glu Glu Leu Leu Gln Asp Phe Arg Ser His Gln Cys Met Gln Leu	
410 415 420	
tgg aac gat gac aac atg gga tgc cta tgg gcc tgc ccc atg gac aag	1575
Trp Asn Asp Asp Asn Met Gly Ser Leu Trp Ala Cys Pro Met Asp Lys	
425 430 435	
tac atc cac agg cgc tgg gtc cta gta tgg ctg gcc tgc cta ctc ttg	1623
Tyr Ile His Arg Arg Trp Val Leu Val Trp Leu Ala Cys Leu Leu Leu	
440 445 450 455	
gct gcg gcg ctt ttc ttc ttc ctc ctt cta aaa aag gac cgc agg aaa	1671
Ala Ala Ala Leu Phe Phe Phe Leu Leu Leu Lys Lys Asp Arg Arg Lys	
460 465 470	
gcg gcc cgt ggc tcc cgc acg gcc ttg ctc ctc cac tcc gcc gac gga	1719
Ala Ala Arg Gly Ser Arg Thr Ala Leu Leu Leu His Ser Ala Asp Gly	
475 480 485	
gcg ggc tac gag cgc ctg gtg gga gca ctg gcg tcc gcg ttg agc cag	1767
Ala Gly Tyr Glu Arg Leu Val Gly Ala Leu Ala Ser Ala Leu Ser Gln	
490 495 500	
atg cca ctg cgc gtg gcc gtg gac ctg tgg agc cgc cgc gag ctg agc	1815
Met Pro Leu Arg Val Ala Val Asp Leu Trp Ser Arg Arg Glu Leu Ser	
505 510 515	
gcg cac gga gcc cta gcc tgg ttc cac cac cag cga cgc cgt atc ctg	1863
Ala His Gly Ala Leu Ala Trp Phe His His Gln Arg Arg Arg Ile Leu	
520 525 530 535	
cag gag ggt ggc gtg gta atc ctt ctc ttc tgc ccc gcg gcc gtg gcg	1911
Gln Glu Gly Gly Val Val Ile Leu Leu Phe Ser Pro Ala Ala Val Ala	
540 545 550	
cag tgt cag cag tgg ctg cag ctc cag aca gtg gag ccc ggg ccg cat	1959
Gln Cys Gln Gln Trp Leu Gln Leu Gln Thr Val Glu Pro Gly Pro His	
555 560 565	
gac gcc ctc gcc gcc tgg ctc agc tgc gtg cta ccc gat ttc ctg caa	2007
Asp Ala Leu Ala Ala Trp Leu Ser Cys Val Leu Pro Asp Phe Leu Gln	
570 575 580	
ggc cgg gcg acc ggc cgc tac gtc ggg gtc tac ttc gac ggg ctg ctg	2055
Gly Arg Ala Thr Gly Arg Tyr Val Gly Val Tyr Phe Asp Gly Leu Leu	
585 590 595	
cac cca gac tct gtg ccc tcc ccg ttc cgc gtc gcc ccg ctc ttc tcc	2103
His Pro Asp Ser Val Pro Ser Pro Phe Arg Val Ala Pro Leu Phe Ser	
600 605 610 615	

ctg ccc tcg cag ctg ccg gct ttc ctg gat gca ctg cag gga ggc tgc 2151
 Leu Pro Ser Gln Leu Pro Ala Phe Leu Asp Ala Leu Gln Gly Gly Cys
 620 625 630
 tcc act tcc gcg ggg cga ccc gcg gac cgg gtg gaa cga gtg acc cag 2199
 Ser Thr Ser Ala Gly Arg Pro Ala Asp Arg Val Glu Arg Val Thr Gln
 635 640 645
 gcg ctg cgg tcc gcc ctg gac agc tgt act tct agc tcg gaa gcc cca 2247
 Ala Leu Arg Ser Ala Leu Asp Ser Cys Thr Ser Ser Ser Glu Ala Pro
 650 655 660
 ggc tgc tgc gag gaa tgg gac ctg gga ccc tgc act aca cta gaa 2292
 Gly Cys Cys Glu Glu Trp Asp Leu Gly Pro Cys Thr Thr Leu Glu
 665 670 675
 taaaagccga tacagtattc ct 2314

<210> 11
 <211> 698
 <212> PRT
 <213> Unknown

<400> 11
 Met Pro Val Ser Trp Phe Leu Leu Ser Leu Ala Leu Gly Arg Asn Pro
 -20 -15 -10 -5
 Val Val Val Ser Leu Glu Arg Leu Met Glu Pro Gln Asp Thr Ala Arg
 -1 1 5 10
 Cys Ser Leu Gly Leu Ser Cys His Leu Trp Asp Gly Asp Val Leu Cys
 15 20 25
 Leu Pro Gly Ser Leu Gln Ser Ala Pro Gly Pro Val Leu Val Pro Thr
 30 35 40
 Arg Leu Gln Thr Glu Leu Val Leu Arg Cys Pro Gln Lys Thr Asp Cys
 45 50 55 60
 Ala Leu Cys Val Arg Val Val Val His Leu Ala Val His Gly His Trp
 65 70 75
 Ala Glu Pro Glu Glu Ala Gly Lys Ser Asp Ser Glu Leu Gln Glu Ser
 80 85 90
 Arg Asn Ala Ser Leu Gln Ala Gln Val Val Leu Ser Phe Gln Ala Tyr
 95 100 105
 Pro Ile Ala Arg Cys Ala Leu Leu Glu Val Gln Val Pro Ala Asp Leu
 110 115 120
 Val Gln Pro Gly Gln Ser Val Gly Ser Ala Val Phe Asp Cys Phe Glu
 125 130 135 140
 Ala Ser Leu Gly Ala Glu Val Gln Ile Trp Ser Tyr Thr Lys Pro Arg
 145 150 155
 Tyr Gln Lys Glu Leu Asn Leu Thr Gln Gln Leu Pro Asp Cys Arg Gly

160										165					170				
Leu	Glu	Val	Arg	Asp	Ser	Ile	Gln	Ser	Cys	Trp	Val	Leu	Pro	Trp	Leu				
		175					180					185							
Asn	Val	Ser	Thr	Asp	Gly	Asp	Asn	Val	Leu	Leu	Thr	Leu	Asp	Val	Ser				
	190					195					200								
Glu	Glu	Gln	Asp	Phe	Ser	Phe	Leu	Leu	Tyr	Leu	Arg	Pro	Val	Pro	Asp				
205					210					215					220				
Ala	Leu	Lys	Ser	Leu	Trp	Tyr	Lys	Asn	Leu	Thr	Gly	Pro	Gln	Asn	Ile				
				225					230					235					
Thr	Leu	Asn	His	Thr	Asp	Leu	Val	Pro	Cys	Leu	Cys	Ile	Gln	Val	Trp				
			240					245					250						
Ser	Leu	Glu	Pro	Asp	Ser	Glu	Arg	Val	Glu	Phe	Cys	Pro	Phe	Arg	Glu				
		255					260					265							
Asp	Pro	Gly	Ala	His	Arg	Asn	Leu	Trp	His	Ile	Ala	Arg	Leu	Arg	Val				
	270					275					280								
Leu	Ser	Pro	Gly	Val	Trp	Gln	Leu	Asp	Ala	Pro	Cys	Cys	Leu	Pro	Gly				
285					290					295					300				
Lys	Val	Thr	Leu	Cys	Trp	Gln	Ala	Pro	Asp	Gln	Ser	Pro	Cys	Gln	Pro				
				305					310					315					
Leu	Val	Pro	Pro	Val	Pro	Gln	Lys	Asn	Ala	Thr	Val	Asn	Glu	Pro	Gln				
			320					325					330						
Asp	Phe	Gln	Leu	Val	Ala	Gly	His	Pro	Asn	Leu	Cys	Val	Gln	Val	Ser				
		335					340					345							
Thr	Trp	Glu	Lys	Val	Gln	Leu	Gln	Ala	Cys	Leu	Trp	Ala	Asp	Ser	Leu				
	350					355					360								
Gly	Pro	Phe	Lys	Asp	Asp	Met	Leu	Leu	Val	Glu	Met	Lys	Thr	Gly	Leu				
365					370					375					380				
Asn	Asn	Thr	Ser	Val	Cys	Ala	Leu	Glu	Pro	Ser	Gly	Cys	Thr	Pro	Leu				
				385					390					395					
Pro	Ser	Met	Ala	Ser	Thr	Arg	Ala	Ala	Arg	Leu	Gly	Glu	Glu	Leu	Leu				
			400					405					410						
Gln	Asp	Phe	Arg	Ser	His	Gln	Cys	Met	Gln	Leu	Trp	Asn	Asp	Asp	Asn				
		415					420					425							
Met	Gly	Ser	Leu	Trp	Ala	Cys	Pro	Met	Asp	Lys	Tyr	Ile	His	Arg	Arg				
	430					435					440								
Trp	Val	Leu	Val	Trp	Leu	Ala	Cys	Leu	Leu	Leu	Ala	Ala	Ala	Leu	Phe				
445					450					455					460				
Phe	Phe	Leu	Leu	Leu	Lys	Lys	Asp	Arg	Arg	Lys	Ala	Ala	Arg	Gly	Ser				
				465					470					475					
Arg	Thr	Ala	Leu	Leu	Leu	His	Ser	Ala	Asp	Gly	Ala	Gly	Tyr	Glu	Arg				

480	485	490
Leu Val Gly Ala Leu Ala Ser Ala Leu Ser Gln Met Pro Leu Arg Val 495 500 505		
Ala Val Asp Leu Trp Ser Arg Arg Glu Leu Ser Ala His Gly Ala Leu 510 515 520		
Ala Trp Phe His His Gln Arg Arg Arg Ile Leu Gln Glu Gly Gly Val 525 530 535 540		
Val Ile Leu Leu Phe Ser Pro Ala Ala Val Ala Gln Cys Gln Gln Trp 545 550 555		
Leu Gln Leu Gln Thr Val Glu Pro Gly Pro His Asp Ala Leu Ala Ala 560 565 570		
Trp Leu Ser Cys Val Leu Pro Asp Phe Leu Gln Gly Arg Ala Thr Gly 575 580 585		
Arg Tyr Val Gly Val Tyr Phe Asp Gly Leu Leu His Pro Asp Ser Val 590 595 600		
Pro Ser Pro Phe Arg Val Ala Pro Leu Phe Ser Leu Pro Ser Gln Leu 605 610 615 620		
Pro Ala Phe Leu Asp Ala Leu Gln Gly Gly Cys Ser Thr Ser Ala Gly 625 630 635		
Arg Pro Ala Asp Arg Val Glu Arg Val Thr Gln Ala Leu Arg Ser Ala 640 645 650		
Leu Asp Ser Cys Thr Ser Ser Ser Glu Ala Pro Gly Cys Cys Glu Glu 655 660 665		
Trp Asp Leu Gly Pro Cys Thr Thr Leu Glu 670 675		

<210> 12

<211> 2094

<212> DNA

<213> reverse translation

<220>

<221> misc_feature

<222> (1)..(2094)

<223> n may be a, c, g, or t

<400> 12

atgccngtnw sntgggttyt nytnwsnytn gcnytnggnm gnaayccngt ngtnngtnwsn 60

ytngarmgny tnatggarcc ncargayacn gcnmgntgyw snytnggnyt nwsntgy cay 120

ytntgggagyg gngaygnyt ntgyytnccn ggnwsnytn arwsngcncc nggncngtn 180

ytngtnccna cnmgnytnca racngarytn gtnytnmgnt gyccncaraa racngaytgy 240

gcnytntgyg tnmngtngt ngtncayytn gcngtncayg gncaytgggc ngarcngar 300

gargcnggna arwsngayws ngarytnear garwsnmgna aygcwnsnyt ncargcnear 360
gtngtnytnw snttycargc ntayccnath gcnmngntgyg cnytnytnga rgtncargtn 420
ccngcngayy tngtncarcc nggncarwsn gtnggnwsng cngtnttyga ytgyttygar 480
gcwnsnytnng gngcngargt ncarathtgg wsntayacna arccnmngta ycaraargar 540
ytnaaaytna cncarcaryt nccngaytgy mgnggnytnng argtnmgnga ywsnathcar 600
wsntgytggg tnytnccntg gytnaaygt nwnacngayg gngayaaygt nytnytnacn 660
ytngaygtnw sngargarca rgayttywsn tyytntytnt ayytnmgnc ngtnccngay 720
gcnytnaarw snytntgga yaaraaytn acnggnccnc araayathac nytnaaycay 780
acngayytnng tnccntgyt ntgyathcar gtntggwsny tngarccnga ywsngarmgn 840
gtngarttyt gyccnttymg ngargayccn ggngcncaym gnaaaytnng gcayathgc 900
mgnytnmgng tnytnwsncc ngngntntgg carytngayg cncntgytg yytnccnggn 960
aargtnacny tntgytggca rgcnccngay carwsnccnt gycarccnyt ngtnccnccn 1020
gtncncara araaygcnc ngtnaaygar ccncargayt tycarytngt ngcnggncay 1080
ccnaaaytn gygtncargt nwnacntgg garaargtnc arytncargc ntgyytnng 1140
gngaywsny tngnccntt yaargaygay atgytnytnng tngaratgaa racnggnytn 1200
aayaayacnw sngtntgygc nytngarccn wsgngntgya cncnytncc nwnatggcn 1260
wsnacnmng cngcnmgnyt ngngargar ytnytnargc ayttymgnws ncaycartgy 1320
atgcarytn ggaaygayga yaayatgggn wsnynntggg cntgyccnat ggayaartay 1380
athcaymgm gntgggntyt ngtnntgytn gcntgyytny tnytnngcnc ngcnytnnty 1440
ttyttytny tnytnaaraa rgaymgmgn aargcngcnm gnggnwsnmg nacngcnytn 1500
ytnytncayw sngcngaygg ngcngntay garmgnytn tngngcnytn ngcnwsngcn 1560
ytnwsncara tgccnytnmg ngtnngcngtn gayytnntggw smgmngnga rytwnsngcn 1620
cayggngcny tngcntggtt ycaycaycar mgmngmngna thytncarga rggngngtn 1680
gtnathytny tnttywsncc ngcngcngtn gcncartgy arcartggyt ncarytnear 1740
acngtnargc cnggncnca ygaygnytn gcngcntggy tnwsntgyt nytnccngay 1800
tgytnargc gnmngcnc nggmngntay gtngngntnt ayttygaygg nytnytnca 1860
ccngaywsng tnccnwsncc nttymgntn gcncnytn tywsnytncc nwnncarytn 1920
ccngcntty tngaygnytn ncargnggn tgywsnca sngcnggmng nccngcngay 1980
mgngtnargm gngtnacna rgcnytnmg wsgcnytn aywsntgyac nwnwnwn 2040
gargcncng gntgytgyga rgartgggay ytnngnccnt gyacnnytn ngar 2094

<210> 13
 <211> 2786
 <212> DNA
 <213> Unknown

<220>
 <223> Description of Unknown Organism: primate; surmised
 Homo sapiens

<220>
 <221> CDS
 <222> (70)..(2283)

<220>
 <221> mat_peptide
 <222> (118)..(2283)

<220>
 <221> misc_feature
 <222> (9)..(134)
 <223> Xaa translation (9, 18, 26, 109, 120, 134) depends
 on genetic code

<400> 13
 cccacgcntc cgggccagca gcgggcggcc ggggcgcaga gaacggcctg gctgggcgag 60

cgcacggcc atg gcc ccg tgg ctg cag ctc tgc tcc gtc ttc ttt acg gtc 111
 Met Ala Pro Trp Leu Gln Leu Cys Ser Val Phe Phe Thr Val
 -15 -10 -5

aac gcc tgc ctc aac ggc tcg cag ctg gct gtn gcc gct ggc ggg tcc 159
 Asn Ala Cys Leu Asn Gly Ser Gln Leu Ala Xaa Ala Ala Gly Gly Ser
 -1 1 5 10

ggc cgc gcg cng ggc gcc gac acc tgt agc tgg ang gga gtg ggg cca 207
 Gly Arg Ala Xaa Gly Ala Asp Thr Cys Ser Trp Xaa Gly Val Gly Pro
 15 20 25 30

gcc agc aga aac agt ggg ctg tac aac atc acc ttc aaa tat gac aat 255
 Ala Ser Arg Asn Ser Gly Leu Tyr Asn Ile Thr Phe Lys Tyr Asp Asn
 35 40 45

tgt acc acc tac ttg aat cca gtg ggg aag cat gtg att gct gac gcc 303
 Cys Thr Thr Tyr Leu Asn Pro Val Gly Lys His Val Ile Ala Asp Ala
 50 55 60

cag aat atc acc atc agc cag tat gct tgc cat gac caa gtg gca gtc 351
 Gln Asn Ile Thr Ile Ser Gln Tyr Ala Cys His Asp Gln Val Ala Val
 65 70 75

acc att ctt tgg tcc cca ggg gcc ctc ggc atc gaa ttc ctg aaa gga 399
 Thr Ile Leu Trp Ser Pro Gly Ala Leu Gly Ile Glu Phe Leu Lys Gly
 80 85 90

ttt cgg gta ata ctg gag gag ctg aag tcg gag gga aga cag ngc caa 447
 Phe Arg Val Ile Leu Glu Leu Lys Ser Glu Gly Arg Gln Xaa Gln
 95 100 105 110

caa ctg att cta aag gat ccg aag cag ntc aac agt agc ttc aaa aga 495

Gln	Leu	Ile	Leu	Lys	Asp	Pro	Lys	Gln	Xaa	Asn	Ser	Ser	Phe	Lys	Arg		
				115					120					125			
act	gga	atg	gaa	tct	caa	cct	ttt	ctg	aat	atg	aaa	ttt	gaa	acg	gat	543	
Thr	Gly	Met	Glu	Ser	Gln	Pro	Xaa	Leu	Asn	Met	Lys	Phe	Glu	Thr	Asp		
			130					135					140				
tat	ttc	gta	agg	ttg	tcc	ttt	tcc	ttc	att	aaa	aac	gaa	agc	aat	tac	591	
Tyr	Phe	Val	Arg	Leu	Ser	Phe	Ser	Phe	Ile	Lys	Asn	Glu	Ser	Asn	Tyr		
		145						150				155					
cac	cct	ttc	ttc	ttt	aga	acc	cga	gcc	tgt	gac	ctg	ttg	tta	cag	ccg	639	
His	Pro	Phe	Phe	Phe	Arg	Thr	Arg	Ala	Cys	Asp	Leu	Leu	Leu	Gln	Pro		
		160					165				170						
gac	aat	cta	gct	tgt	aaa	ccc	ttc	tgg	aag	cct	cgg	aac	ctg	aac	atc	687	
Asp	Asn	Leu	Ala	Cys	Lys	Pro	Phe	Trp	Lys	Pro	Arg	Asn	Leu	Asn	Ile		
175					180					185					190		
agc	cag	cat	ggc	tcg	gac	atg	cag	gtg	tcc	ttc	gac	cac	gca	ccg	cac	735	
Ser	Gln	His	Gly	Ser	Asp	Met	Gln	Val	Ser	Phe	Asp	His	Ala	Pro	His		
				195					200					205			
aac	ttc	ggc	ttc	cgt	ttc	ttc	tat	ctt	cac	tac	aag	ctc	aag	cac	gaa	783	
Asn	Phe	Gly	Phe	Arg	Phe	Phe	Tyr	Leu	His	Tyr	Lys	Leu	Lys	His	Glu		
		210						215					220				
gga	cct	ttc	aag	cga	aag	acc	tgt	aag	cag	gag	caa	act	aca	gag	atg	831	
Gly	Pro	Phe	Lys	Arg	Lys	Thr	Cys	Lys	Gln	Glu	Gln	Thr	Thr	Glu	Met		
		225					230					235					
acc	agc	tgc	ctc	ctt	caa	aat	gtt	tct	cca	ggg	gat	tat	ata	att	gag	879	
Thr	Ser	Cys	Leu	Leu	Gln	Asn	Val	Ser	Pro	Gly	Asp	Tyr	Ile	Ile	Glu		
		240				245					250						
ctg	gtg	gat	gac	act	aac	aca	aca	aga	aaa	gtg	atg	cat	tat	gcc	tta	927	
Leu	Val	Asp	Asp	Thr	Asn	Thr	Thr	Arg	Lys	Val	Met	His	Tyr	Ala	Leu		
255					260					265					270		
aag	cca	gtg	cac	tcc	ccg	tgg	gcc	ggg	ccc	atc	aga	gcc	gtg	gcc	atc	975	
Lys	Pro	Val	His	Ser	Pro	Trp	Ala	Gly	Pro	Ile	Arg	Ala	Val	Ala	Ile		
				275					280					285			
aca	gtg	cca	ctg	gta	gtc	ata	tcg	gca	ttc	gcg	acg	ctc	ttc	act	gtg	1023	
Thr	Val	Pro	Leu	Val	Val	Ile	Ser	Ala	Phe	Ala	Thr	Leu	Phe	Thr	Val		
			290					295					300				
atg	tgc	cgc	aag	aag	caa	caa	gaa	aat	ata	tat	tca	cat	tta	gat	gaa	1071	
Met	Cys	Arg	Lys	Lys	Gln	Gln	Glu	Asn	Ile	Tyr	Ser	His	Leu	Asp	Glu		
		305					310					315					
gag	agc	tct	gag	tct	tcc	aca	tac	act	gca	gca	ctc	cca	aga	gag	agg	1119	
Glu	Ser	Ser	Glu	Ser	Ser	Thr	Tyr	Thr	Ala	Ala	Leu	Pro	Arg	Glu	Arg		
		320				325					330						
ctc	cgg	ccg	cgg	ccg	aag	gtc	ttt	ctc	tgc	tat	tcc	agt	aaa	gat	ggc	1167	
Leu	Arg	Pro	Arg	Pro	Lys	Val	Phe	Leu	Cys	Tyr	Ser	Ser	Lys	Asp	Gly		
335					340				345					350			
cag	aat	cac	atg	aat	gtc	gtc	cag	tgt	ttc	gcc	tac	ttc	ctc	cag	gac	1215	

Gln Asn His Met Asn Val Val Gln Cys Phe Ala Tyr Phe Leu Gln Asp	
355 360 365	
ttc tgt ggc tgt gag gtg gct ctg gac ctg tgg gaa gac ttc agc ctc	1263
Phe Cys Gly Cys Glu Val Ala Leu Asp Leu Trp Glu Asp Phe Ser Leu	
370 375 380	
tgt aga gaa ggg cag aga gaa tgg gtc atc cag aag atc cac gag tcc	1311
Cys Arg Glu Gly Gln Arg Glu Trp Val Ile Gln Lys Ile His Glu Ser	
385 390 395	
cag ttc atc att gtg gtt tgt tcc aaa ggt atg aag tac ttt gtg gac	1359
Gln Phe Ile Ile Val Val Cys Ser Lys Gly Met Lys Tyr Phe Val Asp	
400 405 410	
aag aag aac tac aaa cac aaa gga ggt ggc cga ggc tcg ggg aaa gga	1407
Lys Lys Asn Tyr Lys His Lys Gly Gly Gly Arg Gly Ser Gly Lys Gly	
415 420 425 430	
gag ctc ttc ctg gtg gcg gtg tca gcc att gcc gaa aag ctc cgc cag	1455
Glu Leu Phe Leu Val Ala Val Ser Ala Ile Ala Glu Lys Leu Arg Gln	
435 440 445	
gcc aag cag agt tgc tcc gcg gcg ctc agc aag ttt atc gcc gtc tac	1503
Ala Lys Gln Ser Ser Ser Ala Ala Leu Ser Lys Phe Ile Ala Val Tyr	
450 455 460	
ttt gat tat tcc tgc gag gga gac gtc ccc ggt atc cta gac ctg agt	1551
Phe Asp Tyr Ser Cys Glu Gly Asp Val Pro Gly Ile Leu Asp Leu Ser	
465 470 475	
acc aag tac aga ctc atg gac aat ctt cct cag ctc tgt tcc cac ctg	1599
Thr Lys Tyr Arg Leu Met Asp Asn Leu Pro Gln Leu Cys Ser His Leu	
480 485 490	
cac tcc cga gac cac ggc ctc cag gag ccg ggg cag cac acg cga cag	1647
His Ser Arg Asp His Gly Leu Gln Glu Pro Gly Gln His Thr Arg Gln	
495 500 505 510	
ggc agc aga agg aac tac ttc cgg agc aag tca ggc cgg tcc cta tac	1695
Gly Ser Arg Arg Asn Tyr Phe Arg Ser Lys Ser Gly Arg Ser Leu Tyr	
515 520 525	
gtc gcc att tgc aac atg cac cag ttt att gac gag gag ccc gac tgg	1743
Val Ala Ile Cys Asn Met His Gln Phe Ile Asp Glu Glu Pro Asp Trp	
530 535 540	
ttc gaa aag cag ttc gtt ccc ttc cat cct cct cca ctg cgc tac cgg	1791
Phe Glu Lys Gln Phe Val Pro Phe His Pro Pro Pro Leu Arg Tyr Arg	
545 550 555	
gag cca gtc ttg gag aaa ttt gat tgc ggc ttg gtt tta aat gat gtc	1839
Glu Pro Val Leu Glu Lys Phe Asp Ser Gly Leu Val Leu Asn Asp Val	
560 565 570	
atg tgc aaa cca ggg cct gag agt gac ttc tgc cta aag gta gag gcg	1887
Met Cys Lys Pro Gly Pro Glu Ser Asp Phe Cys Leu Lys Val Glu Ala	
575 580 585 590	
gct gtt ctt ggg gca acc gga cca gcc gac tcc cag cac gag agt cag	1935

Ala Val Leu Gly Ala Thr Gly Pro Ala Asp Ser Gln His Glu Ser Gln
595 600 605

cat ggg ggc ctg gac caa gac ggg gag gcc cgg cct gcc ctt gac ggt 1983
His Gly Gly Leu Asp Gln Asp Gly Glu Ala Arg Pro Ala Leu Asp Gly
610 615 620

agc gcc gcc ctg caa ccc ctg ctg cac acg gtg aaa gcc ggc agc ccc 2031
Ser Ala Ala Leu Gln Pro Leu Leu His Thr Val Lys Ala Gly Ser Pro
625 630 635

tcg gac atg ccg cgg gac tca ggc atc tat gac tcg tct gtg ccc tca 2079
Ser Asp Met Pro Arg Asp Ser Gly Ile Tyr Asp Ser Ser Val Pro Ser
640 645 650

tcc gag ctg tct ctg cca ctg atg gaa gga ctc tcg acg gac cag aca 2127
Ser Glu Leu Ser Leu Pro Leu Met Glu Gly Leu Ser Thr Asp Gln Thr
655 660 665 670

gaa acg tct tcc ctg acg gag agc gtg tcc tcc tct tca ggc ctg ggt 2175
Glu Thr Ser Ser Leu Thr Glu Ser Val Ser Ser Ser Ser Gly Leu Gly
675 680 685

gag gag gaa cct cct gcc ctt cct tcc aag ctc ctc tct tct ggg tca 2223
Glu Glu Glu Pro Pro Ala Leu Pro Ser Lys Leu Leu Ser Ser Gly Ser
690 695 700

tgc aaa gca gat ctt ggt tgc cgc agc tac act gat gaa ctc cac gcg 2271
Cys Lys Ala Asp Leu Gly Cys Arg Ser Tyr Thr Asp Glu Leu His Ala
705 710 715

gtc gcc cct ttg taacaaaacg aaagagtcta agcattgccca ctttagctgc 2323
Val Ala Pro Leu
720

tgcctccctc tgattcccca gctcatctcc ctggttgcac ggcccacttg gagctgaggt 2383

ctcatacaag gatatttgga gtgaaatgct ggccagtact tggtctccct tgccccaacc 2443

ctttaccgga tatcttgaca aactctccaa ttttctaaaa tgatatggag ctctgaaagg 2503

catgtccata aggtctgaca acagcttgcc aaatttggtt agtccttgga tcagagcctg 2563

ttgtgggagg tagggaggaa atatgtaaag aaaaacagga agatacctgc actaatcatt 2623

cagacttcat tgagctctgc aaactttgcc tgtttgctat tggctacctt gatttgaaat 2683

gctttgtgaa aaaaggcact tttaacatca tagccacaga aatcaagtgc cagtctatct 2743

ggaatccatg ttgtattgca gataatgttc tcatttattt ttg 2786

<210> 14
<211> 738
<212> PRT
<213> Unknown

<400> 14
Met Ala Pro Trp Leu Gln Leu Cys Ser Val Phe Phe Thr Val Asn Ala
-15 -10 -5 -1

Cys Leu Asn Gly Ser Gln Leu Ala Xaa Ala Ala Gly Gly Ser Gly Arg
 1 5 10 15
 Ala Xaa Gly Ala Asp Thr Cys Ser Trp Xaa Gly Val Gly Pro Ala Ser
 20 25 30
 Arg Asn Ser Gly Leu Tyr Asn Ile Thr Phe Lys Tyr Asp Asn Cys Thr
 35 40 45
 Thr Tyr Leu Asn Pro Val Gly Lys His Val Ile Ala Asp Ala Gln Asn
 50 55 60
 Ile Thr Ile Ser Gln Tyr Ala Cys His Asp Gln Val Ala Val Thr Ile
 65 70 75 80
 Leu Trp Ser Pro Gly Ala Leu Gly Ile Glu Phe Leu Lys Gly Phe Arg
 85 90 95
 Val Ile Leu Glu Glu Leu Lys Ser Glu Gly Arg Gln Xaa Gln Gln Leu
 100 105 110
 Ile Leu Lys Asp Pro Lys Gln Xaa Asn Ser Ser Phe Lys Arg Thr Gly
 115 120 125
 Met Glu Ser Gln Pro Xaa Leu Asn Met Lys Phe Glu Thr Asp Tyr Phe
 130 135 140
 Val Arg Leu Ser Phe Ser Phe Ile Lys Asn Glu Ser Asn Tyr His Pro
 145 150 155 160
 Phe Phe Phe Arg Thr Arg Ala Cys Asp Leu Leu Leu Gln Pro Asp Asn
 165 170 175
 Leu Ala Cys Lys Pro Phe Trp Lys Pro Arg Asn Leu Asn Ile Ser Gln
 180 185 190
 His Gly Ser Asp Met Gln Val Ser Phe Asp His Ala Pro His Asn Phe
 195 200 205
 Gly Phe Arg Phe Phe Tyr Leu His Tyr Lys Leu Lys His Glu Gly Pro
 210 215 220
 Phe Lys Arg Lys Thr Cys Lys Gln Glu Gln Thr Thr Glu Met Thr Ser
 225 230 235 240
 Cys Leu Leu Gln Asn Val Ser Pro Gly Asp Tyr Ile Ile Glu Leu Val
 245 250 255
 Asp Asp Thr Asn Thr Thr Arg Lys Val Met His Tyr Ala Leu Lys Pro
 260 265 270
 Val His Ser Pro Trp Ala Gly Pro Ile Arg Ala Val Ala Ile Thr Val
 275 280 285
 Pro Leu Val Val Ile Ser Ala Phe Ala Thr Leu Phe Thr Val Met Cys
 290 295 300
 Arg Lys Lys Gln Gln Glu Asn Ile Tyr Ser His Leu Asp Glu Glu Ser
 305 310 315 320

Ser Glu Ser Ser Thr Tyr Thr Ala Ala Leu Pro Arg Glu Arg Leu Arg
 325 330 335
 Pro Arg Pro Lys Val Phe Leu Cys Tyr Ser Ser Lys Asp Gly Gln Asn
 340 345 350
 His Met Asn Val Val Gln Cys Phe Ala Tyr Phe Leu Gln Asp Phe Cys
 355 360 365
 Gly Cys Glu Val Ala Leu Asp Leu Trp Glu Asp Phe Ser Leu Cys Arg
 370 375 380
 Glu Gly Gln Arg Glu Trp Val Ile Gln Lys Ile His Glu Ser Gln Phe
 385 390 395 400
 Ile Ile Val Val Cys Ser Lys Gly Met Lys Tyr Phe Val Asp Lys Lys
 405 410 415
 Asn Tyr Lys His Lys Gly Gly Gly Arg Gly Ser Gly Lys Gly Glu Leu
 420 425 430
 Phe Leu Val Ala Val Ser Ala Ile Ala Glu Lys Leu Arg Gln Ala Lys
 435 440 445
 Gln Ser Ser Ser Ala Ala Leu Ser Lys Phe Ile Ala Val Tyr Phe Asp
 450 455 460
 Tyr Ser Cys Glu Gly Asp Val Pro Gly Ile Leu Asp Leu Ser Thr Lys
 465 470 475 480
 Tyr Arg Leu Met Asp Asn Leu Pro Gln Leu Cys Ser His Leu His Ser
 485 490 495
 Arg Asp His Gly Leu Gln Glu Pro Gly Gln His Thr Arg Gln Gly Ser
 500 505 510
 Arg Arg Asn Tyr Phe Arg Ser Lys Ser Gly Arg Ser Leu Tyr Val Ala
 515 520 525
 Ile Cys Asn Met His Gln Phe Ile Asp Glu Glu Pro Asp Trp Phe Glu
 530 535 540
 Lys Gln Phe Val Pro Phe His Pro Pro Pro Leu Arg Tyr Arg Glu Pro
 545 550 555 560
 Val Leu Glu Lys Phe Asp Ser Gly Leu Val Leu Asn Asp Val Met Cys
 565 570 575
 Lys Pro Gly Pro Glu Ser Asp Phe Cys Leu Lys Val Glu Ala Ala Val
 580 585 590
 Leu Gly Ala Thr Gly Pro Ala Asp Ser Gln His Glu Ser Gln His Gly
 595 600 605
 Gly Leu Asp Gln Asp Gly Glu Ala Arg Pro Ala Leu Asp Gly Ser Ala
 610 615 620
 Ala Leu Gln Pro Leu Leu His Thr Val Lys Ala Gly Ser Pro Ser Asp
 625 630 635 640

Met Pro Arg Asp Ser Gly Ile Tyr Asp Ser Ser Val Pro Ser Ser Glu
645 650 655

Leu Ser Leu Pro Leu Met Glu Gly Leu Ser Thr Asp Gln Thr Glu Thr
660 665 670

Ser Ser Leu Thr Glu Ser Val Ser Ser Ser Ser Gly Leu Gly Glu Glu
675 680 685

Glu Pro Pro Ala Leu Pro Ser Lys Leu Leu Ser Ser Gly Ser Cys Lys
690 695 700

Ala Asp Leu Gly Cys Arg Ser Tyr Thr Asp Glu Leu His Ala Val Ala
705 710 715 720

Pro Leu

<210> 15
<211> 2214
<212> DNA
<213> reverse translation

<220>
<221> misc_feature
<222> (1)..(2214)
<223> n may be a, c, g, or t

<400> 15
atggcncnt ggytncaryt ntgywsngtn ttyttyacng tnaaygcntg yytnaayggn 60
wsncarytng cngtngcngc nggnngnwsn ggnmgngcnn nngngncnga yacntgywsn 120
tggnnnggng tnggncncgc nwsnmgnaay wsnggnytn tayaayathac nttyaartay 180
gayaaytgaa cnacntayyt naayccngtn ggnaarcayg tnathgcnga ygcncaraay 240
athacnathw sncartaygc ntgycaayg cargtngcng tnacnathyt ntggwsncn 300
ggngcnytn gnatgcartt yytnaarggn ttmngntna thytnrga rytnaarwsn 360
garggnmgnc arnnncarca rytnathytn aargayccna arcarnnnaa ywsnwsntty 420
aarmgnacng gnatggarws ncarccnnnn ytnaayatga arttygarac ngaytaytty 480
gtmngnytnw snttywsntt yathaaraay garwsnaayt aycayccntt yttyttymgn 540
acnmngcncnt gygayytnyt nytnarcncn gayaayytn cngtyaarcc nttytggaar 600
ccnmgnaayy tnaayathws ncarcayggn wsngayatgc argtnwsntt ygaycaygcn 660
ccncayaayt tyggnttymg nttytttyay ytncaytaya arytnaarca ygarggnccn 720
ttyaarmgna aracntgyaa rcargarcar acnacngara tgacnwsntg yytnytnear 780
aaygtnwsnc cngngayta yathathgar ytngtngayg ayacnaayac nacnmgnaar 840
gtnatgcayt aygcnytnaa rccngtncay wsncntggg cnggncnat hmgngcngtn 900

gcnathacng tnccnytngt ngtnathwsn gcnttygcna cnytnntyac ngtnatgtgy 960
 mgnaaraarc arcargaraa yathtaywsn cayytngayg argarwsnws ngarwsnwsn 1020
 acntayacng cngcnytncc nmngngarmgn ytnmgncnm gncnaargt nttyytnnty 1080
 taywsnwsna argayggna raaycaytg aaygtngtnc artgyttygc ntayttytn 1140
 cargayttyp gyggntgyga rgtngcnytn gayytnntggg argayttyws nytntgymgn 1200
 garggncarm gngartgggt nathcaraar athcaygarw sncarttyat hathgtngtn 1260
 tgywsnaarg gnatgaarta ytygtngay aaraaraayt ayaarcayaa rggngngngn 1320
 mgnggnwsng gnaarggnga rytnttytn gtngcngtnw sngcnathgc ngaraarytn 1380
 mgncargcna arcarsnws nwsngcngcn ytnwsnaart tyathgcngt ntayttygay 1440
 taywsntgyg arggngaygt nccnggnath ytngayytnw snacnaarta ymgnytnatg 1500
 gayaayytn cncarytntg ywsncayytn caywsnmngn aycayggnyt ncargarccn 1560
 ggncarcaya cnmgncargg nwsnmgnmgn aaytaytym gnwsnaarws nggnmgnwsn 1620
 ytntaygtng cnathtgyaa yatgcaycar ttyathgayg argarccnga ytggttygar 1680
 aarcarttyg tnccnttyca yccnccnccn ytnmgntaym gngarccngt nytngaraar 1740
 ttygaywsng gnytngtnyt naaygaygt atgtgyaarc cnggnccnga rwsngaytty 1800
 tgyytnaarg tngargcngc ngtnytnngn gcnacnggnc cngcngayws ncargaygar 1860
 wsnarcayg gnggnytna yargayggn gargcnmgnc cngcnytna yggnwngcn 1920
 gcnytnarc cnytnytnca yacngtnaar gcnggnwsnc cnwsngayat gccnmngay 1980
 wsggnatht aygaywsnws ngtnccnwsn wsgarytnw snytnccnyt natggarggn 2040
 ytnwsnacng aycaracnga racnwsnwsn ytnacngarw sngtnwsnws nwsnwsngn 2100
 ytnngngarg argarccncc ngcnytnccn wsnaarytny tnwsnwsngg nwsntgyaar 2160
 gcngayytn gntgymgnws ntayacngay garytncayg cngtngcncc nytn 2214

<210> 16

<211> 2012

<212> DNA

<213> Unknown

<220>

<223> Description of Unknown Organism: primate; surmised
Homo sapiens

<220>

<221> CDS

<222> (1)..(1971)

<220>

<221> mat_peptide

<222> (70)..(1971)

<400> 16

atg ggg agc tcc aga ctg gca gcc ctg ctc ctg cct ctc ctc ctc ata	48
Met Gly Ser Ser Arg Leu Ala Ala Leu Leu Leu Pro Leu Leu Leu Ile	
-20 -15 -10	
gtc atc gac ctc tct gac tct gct ggg att ggc ttt cgc cac ctg ccc	96
Val Ile Asp Leu Ser Asp Ser Ala Gly Ile Gly Phe Arg His Leu Pro	
-5 -1 1 5	
cac tgg aac acc cgc tgt cct ctg gcc tcc cac acg gaa gtt ctg cct	144
His Trp Asn Thr Arg Cys Pro Leu Ala Ser His Thr Glu Val Leu Pro	
10 15 20 25	
ata tcc ctt gcc gca cct ggt ggg ccc tct tct cca caa agc ctt ggt	192
Ile Ser Leu Ala Ala Pro Gly Gly Pro Ser Ser Pro Gln Ser Leu Gly	
30 35 40	
gtg tgc gag tct ggc act gtt ccc gct gtt tgt gcc agc atc tgc tgt	240
Val Cys Glu Ser Gly Thr Val Pro Ala Val Cys Ala Ser Ile Cys Cys	
45 50 55	
cag gtg gct cag gtc ttc aac ggg gcc tct tcc acc tcc tgg tgc aga	288
Gln Val Ala Gln Val Phe Asn Gly Ala Ser Ser Thr Ser Trp Cys Arg	
60 65 70	
aat cca aaa agt ctt cca cat tca agt tct ata gga gac aca aga tgc	336
Asn Pro Lys Ser Leu Pro His Ser Ser Ser Ile Gly Asp Thr Arg Cys	
75 80 85	
cag cac ctg ctc aga gga agc tgc tgc ctc gtc gtc acc tgt ctg aga	384
Gln His Leu Leu Arg Gly Ser Cys Cys Leu Val Val Thr Cys Leu Arg	
90 95 100 105	
aga gcc atc aca ttt cca tcc cct ccc cag aca tct ccc aca agg gac	432
Arg Ala Ile Thr Phe Pro Ser Pro Pro Gln Thr Ser Pro Thr Arg Asp	
110 115 120	
ttc gct cta aaa gga ccc aac ctt cgg atc cag aga cat ggg aaa gtc	480
Phe Ala Leu Lys Gly Pro Asn Leu Arg Ile Gln Arg His Gly Lys Val	
125 130 135	
ttc cca gat tgg act cac aaa ggc atg gag gtg ggc act ggg tac aac	528
Phe Pro Asp Trp Thr His Lys Gly Met Glu Val Gly Thr Gly Tyr Asn	
140 145 150	
agg aga tgg gtt cag ctg agt ggt gga ccc gag ttc tcc ttt gat ttg	576
Arg Arg Trp Val Gln Leu Ser Gly Gly Pro Glu Phe Ser Phe Asp Leu	
155 160 165	
ctg cct gag gcc cgg gct att cgg gtg acc ata tct tca ggc cct gag	624
Leu Pro Glu Ala Arg Ala Ile Arg Val Thr Ile Ser Ser Gly Pro Glu	
170 175 180 185	
gtc agc gtg cgt ctt tgt cac cag tgg gca ctg gag tgt gaa gag ctg	672
Val Ser Val Arg Leu Cys His Gln Trp Ala Leu Glu Cys Glu Glu Leu	
190 195 200	

agc agt ccc tat gat gtc cag aaa att gtg tct ggg ggc cac act gta	720
Ser Ser Pro Tyr Asp Val Gln Lys Ile Val Ser Gly Gly His Thr Val	
205 210 215	
gag ctg cct tat gaa ttc ctt ctg ccc tgt ctg tgc ata gag gca tcc	768
Glu Leu Pro Tyr Glu Phe Leu Leu Pro Cys Leu Cys Ile Glu Ala Ser	
220 225 230	
tac ctg caa gag gac act gtg agg cgc aaa aaa tgt ccc ttc cag agc	816
Tyr Leu Gln Glu Asp Thr Val Arg Arg Lys Lys Cys Pro Phe Gln Ser	
235 240 245	
tgg cca gaa gcc tat ggc tcg gac ttc tgg aag tca gtg cac ttc act	864
Trp Pro Glu Ala Tyr Ser Asp Phe Trp Lys Ser Val His Phe Thr	
250 255 260 265	
gac tac agc cag cac act cag atg gtc atg gcc ctg aca ctc cgc tgc	912
Asp Tyr Ser Gln His Thr Gln Met Val Met Ala Leu Thr Leu Arg Cys	
270 275 280	
cca ctg aag ctg gaa gct gcc ctc tgc cag agg cac gac tgg cat acc	960
Pro Leu Lys Leu Glu Ala Ala Leu Cys Gln Arg His Asp Trp His Thr	
285 290 295	
ctt tgc aaa gac ctc ccg aat gcc acg gct cga gag tca gat ggg tgg	1008
Leu Cys Lys Asp Leu Pro Asn Ala Thr Ala Arg Glu Ser Asp Gly Trp	
300 305 310	
tat gtt ttg gag aag gtg gac ctg cac ccc cag ctc tgc ttc aag gta	1056
Tyr Val Leu Glu Lys Val Asp Leu His Pro Gln Leu Cys Phe Lys Val	
315 320 325	
caa cca tgg ttc tct ttt gga aac agc agc cat gtt gaa tgc ccc cac	1104
Gln Pro Trp Phe Ser Phe Gly Asn Ser Ser His Val Glu Cys Pro His	
330 335 340 345	
cag act ggg tct ctc aca tcc tgg aat gta agc atg gat acc caa gcc	1152
Gln Thr Gly Ser Leu Thr Ser Trp Asn Val Ser Met Asp Thr Gln Ala	
350 355 360	
cag cag ctg att ctt cac ttc tcc tca aga atg cat gcc acc ttc agt	1200
Gln Gln Leu Ile Leu His Phe Ser Ser Arg Met His Ala Thr Phe Ser	
365 370 375	
gct gcc tgg agc ctc cca ggc ttg ggg cag gac act ttg gtg ccc ccc	1248
Ala Ala Trp Ser Leu Pro Gly Leu Gly Gln Asp Thr Leu Val Pro Pro	
380 385 390	
gtg tac act gtc agc cag gtg tgg cgg tca gat gtc cag ttt gcc tgg	1296
Val Tyr Thr Val Ser Gln Val Trp Arg Ser Asp Val Gln Phe Ala Trp	
395 400 405	
aag cac ctc ttg tgt cca gat gtc tct tac aga cac ctg ggg ctc ttg	1344
Lys His Leu Leu Cys Pro Asp Val Ser Tyr Arg His Leu Gly Leu Leu	
410 415 420 425	
atc ctg gca ctg ctg gcc ctc ctc acc cta ctg ggt gtt gtt ctg gcc	1392
Ile Leu Ala Leu Leu Ala Leu Leu Thr Leu Leu Gly Val Val Leu Ala	
430 435 440	

ctc acc tgc cgg cgc cca cag tca ggc ccg ggc cca gcg cgg cca gtg 1440
 Leu Thr Cys Arg Arg Pro Gln Ser Gly Pro Gly Pro Ala Arg Pro Val
 445 450 455

ctc ctc ctg cac gcg gcg gac tcg gag gcg cag cgg cgc ctg gtg gga 1488
 Leu Leu Leu His Ala Ala Asp Ser Glu Ala Gln Arg Arg Leu Val Gly
 460 465 470

gcg ctg gct gaa ctg cta cgg gca gcg ctg ggc ggc ggg cgc gac gtg 1536
 Ala Leu Ala Glu Leu Leu Arg Ala Ala Leu Gly Gly Gly Arg Asp Val
 475 480 485

atc gtg gac ctg tgg gag ggg agg cac gtg gcg cgc gtg ggc ccg ctg 1584
 Ile Val Asp Leu Trp Glu Gly Arg His Val Ala Arg Val Gly Pro Leu
 490 495 500 505

ccg tgg ctc tgg gcg gcg cgg acg cgc gta gcg cgg gag cag ggc act 1632
 Pro Trp Leu Trp Ala Ala Arg Thr Arg Val Ala Arg Glu Gln Gly Thr
 510 515 520

gtg ctg ctg ctg tgg agc ggc gcc gac ctt cgc ccg gtc agc ggc ccc 1680
 Val Leu Leu Leu Trp Ser Gly Ala Asp Leu Arg Pro Val Ser Gly Pro
 525 530 535

gac ccc cgc gcc gcg ccc ctg ctc gcc ctg ctc cac gct gcc ccg cgc 1728
 Asp Pro Arg Ala Ala Pro Leu Leu Ala Leu Leu His Ala Ala Pro Arg
 540 545 550

ccg ctg ctg ctg ctc gct tac ttc agt cgc ctc tgc gcc aag ggc gac 1776
 Pro Leu Leu Leu Leu Ala Tyr Phe Ser Arg Leu Cys Ala Lys Gly Asp
 555 560 565

atc ccc ccg ccg ctg cgc gcc ctg ccg cgc tac cgc ctg ctg cgc gac 1824
 Ile Pro Pro Pro Leu Arg Ala Leu Pro Arg Tyr Arg Leu Leu Arg Asp
 570 575 580 585

ctg ccg cgt ctg ctg cgg gcg ctg gac gcg cgg cct ttc gca gag gcc 1872
 Leu Pro Arg Leu Leu Arg Ala Leu Asp Ala Arg Pro Phe Ala Glu Ala
 590 595 600

acc agc tgg ggc cgc ctt ggg gcg cgg cag cgc agg cag agc cgc cta 1920
 Thr Ser Trp Gly Arg Leu Gly Ala Arg Gln Arg Arg Gln Ser Arg Leu
 605 610 615

gag ctg tgc agc cgg ctc gaa cga gag gcc gcc cga ctt gca gac cta 1968
 Glu Leu Cys Ser Arg Leu Glu Arg Glu Ala Ala Arg Leu Ala Asp Leu
 620 625 630

ggt tgagcagagc tccaccgcag tcccgggtgt ctgcggccgc t 2012
 Gly

<210> 17

<211> 657

<212> PRT

<213> Unknown

<400> 17

Met Gly Ser Ser Arg Leu Ala Ala Leu Leu Leu Pro Leu Leu Leu Ile
 -20 -15 -10

Tyr Val Leu Glu Lys Val Asp Leu His Pro Gln Leu Cys Phe Lys Val
 315 320 325
 Gln Pro Trp Phe Ser Phe Gly Asn Ser Ser His Val Glu Cys Pro His
 330 335 340 345
 Gln Thr Gly Ser Leu Thr Ser Trp Asn Val Ser Met Asp Thr Gln Ala
 350 355 360
 Gln Gln Leu Ile Leu His Phe Ser Ser Arg Met His Ala Thr Phe Ser
 365 370 375
 Ala Ala Trp Ser Leu Pro Gly Leu Gly Gln Asp Thr Leu Val Pro Pro
 380 385 390
 Val Tyr Thr Val Ser Gln Val Trp Arg Ser Asp Val Gln Phe Ala Trp
 395 400 405
 Lys His Leu Leu Cys Pro Asp Val Ser Tyr Arg His Leu Gly Leu Leu
 410 415 420 425
 Ile Leu Ala Leu Leu Ala Leu Leu Thr Leu Leu Gly Val Val Leu Ala
 430 435 440
 Leu Thr Cys Arg Arg Pro Gln Ser Gly Pro Gly Pro Ala Arg Pro Val
 445 450 455
 Leu Leu Leu His Ala Ala Asp Ser Glu Ala Gln Arg Arg Leu Val Gly
 460 465 470
 Ala Leu Ala Glu Leu Leu Arg Ala Ala Leu Gly Gly Gly Arg Asp Val
 475 480 485
 Ile Val Asp Leu Trp Glu Gly Arg His Val Ala Arg Val Gly Pro Leu
 490 495 500 505
 Pro Trp Leu Trp Ala Ala Arg Thr Arg Val Ala Arg Glu Gln Gly Thr
 510 515 520
 Val Leu Leu Leu Trp Ser Gly Ala Asp Leu Arg Pro Val Ser Gly Pro
 525 530 535
 Asp Pro Arg Ala Ala Pro Leu Leu Ala Leu Leu His Ala Ala Pro Arg
 540 545 550
 Pro Leu Leu Leu Leu Ala Tyr Phe Ser Arg Leu Cys Ala Lys Gly Asp
 555 560 565
 Ile Pro Pro Pro Leu Arg Ala Leu Pro Arg Tyr Arg Leu Leu Arg Asp
 570 575 580 585
 Leu Pro Arg Leu Leu Arg Ala Leu Asp Ala Arg Pro Phe Ala Glu Ala
 590 595 600
 Thr Ser Trp Gly Arg Leu Gly Ala Arg Gln Arg Arg Gln Ser Arg Leu
 605 610 615
 Glu Leu Cys Ser Arg Leu Glu Arg Glu Ala Ala Arg Leu Ala Asp Leu
 620 625 630

Gly

<210> 18

<211> 1971

<212> DNA

<213> reverse translation

<220>

<221> misc_feature

<222> (1)..(1971)

<223> n may be a, c, g, or t

<400> 18

atgggnwsnw snmgnytn gc ngcnytnytn ytnccnytny tnytnathgt nathgayytn 60
wsngaywsng cnggnathgg nttymgncay ytnccncayt ggaayacnmg ntgyccnytn 120
gcwnsncaaya cngargtnyt nccnathwsn ytngcngcnc cnggnggncc nwsnwsnccn 180
carwsnytn gngtntgyga rwsnggnacn gtncncngcng tntgygcnws nathtgytgy 240
cargtngcnc argtnnttyaa yggngcnwsn wsnacnwsnt ggtgymgnaa yccnaarwsn 300
ytncncncayw snwsnwsnat hggngayacn mgntgyarc ayytnytnmg nggnwsntgy 360
tgyytngtng tnaentgyt nmgnmgngcn athacnttyc cnwsnccncc ncaracnwsn 420
ccnacnmng aytytgcn yt naarggnccn aaytnmgna thcarmgna yggnaargtn 480
ttyccngayt ggacncayaa rggnatggar gtnggnacng gntayaaymg nmngtgggt 540
carytnwsng gnggnccnga rtywsntty gayytnytn cngargcnmg ngcnathmgn 600
gtnacnathw snwsnggncc ngargtnwsn gtnmgnytn gycaycartg ggcnytn gar 660
tgygargary tnwsnwsncc ntaygaytn caraarathg tnwsngngng ncayacngtn 720
garytnccnt aygarttyt nytnccntgy ytntgyathg argcnwsnta yytnccargar 780
gayacngtnm gnmgnaaraa rtgyccntty carwsntggc cngargcnta yggnwsngay 840
ttytggaarw sngtncaytt yacngaytay wsnarcaya cncaratggt natggcn ytn 900
acnytnmgnt gyccnytnaa rytngargcn gcnytntgyc armgncayga ytggcayacn 960
ytntgyaarg ayytnccnaa ygcncngcn mgngarwsng ayggntggta ygtnytn gar 1020
aargtngayy tncayccna rytntgyt aargtnarc cntggtyws nttyggnaay 1080
wsnwsncayg tngartgycc ncaycaracn ggnwsnytna cnwsntggaa ygtnwsnatg 1140
gayacncarg cncarcaryt nathytnca tywsnwsnm gnatgcaygc nacnttywsn 1200
gcngcntggw snytnccngg nytnngncar gayacnytn tncncncngt ntayacngtn 1260
wsncargtn ggmgnwsnga ygtncartty gcntggaarc ayytnytntg yccngaygtn 1320

wsttaymgnc ayytnggny nytnathytn gcnynytng cnytnytnac nytnytnngn 1380
 gtngtnytng cnytnacntg ymgngmgnccn carwsnggnc cnggncngc nmgnccngtn 1440
 ytnytnytn cnyngcngc ywsngargcn carmgmgnny tngtnggngc nytnngcngar 1500
 ytnytnmgng cngcnytnng nggnggmgn gaygtathg tngayytnng ggarggmgn 1560
 caygtngcnm gngtnggncc nytnccntgg ytnngggcng cmgnacnmg ngtnngcnmgn 1620
 garcarggna cngtnytnyt nytnnggwsn ggngcngayy tnmgnccngt nwsnggnccn 1680
 gayccnmng cngcncnny nytnngcnytn ytnaygng cncnccnny nmngcnytn 1740
 ytnngcntayt tywsnmgny ntgygcnaar ggngayathc cncncnny nmngcnytn 1800
 ccnmngtaym gnytnytnmg ngayytnccn mgnynytnm gngcnytna ygnmgnccn 1860
 ttygcngarg cnacwstng gggngmnytn ggngcnmgnc armgmgnca rwsnmgnytn 1920
 garytnngyw snmngnytna rmngargcn gcnmngnytn cngayytnng n 1971

<210> 19

<211> 808

<212> DNA

<213> Unknown

<220>

<223> Description of Unknown Organism: rodent; surmised
Mus musculus

<220>

<221> CDS

<222> (78)..(806)

<220>

<221> mat_peptide

<222> (147)..(806)

<400> 19

cagctccggg ccagccctg ctgccctctt gcagacagga aagacatggt ctctgcgccc 60
 tgatcctaca gaagctc atg ggg agc ccc aga ctg gca gcc ttg ctc ctg 110
 Met Gly Ser Pro Arg Leu Ala Ala Leu Leu Leu
 -20 -15
 tct ctc ccg cta ctg ctc atc ggc ctc gct gtg tct gct cgg gtt gcc 158
 Ser Leu Pro Leu Leu Leu Ile Gly Leu Ala Val Ser Ala Arg Val Ala
 -10 -5 -1 1
 tgc ccc tgc ctg cgg agt tgg acc agc cac tgt ctc ctg gcc tac cgt 206
 Cys Pro Cys Leu Arg Ser Trp Thr Ser His Cys Leu Leu Ala Tyr Arg
 5 10 15 20
 gtg gat aaa cgt ttt gct ggc ctt cag tgg ggc tgg ttc cct ctc ttg 254
 Val Asp Lys Arg Phe Ala Gly Leu Gln Trp Gly Trp Phe Pro Leu Leu
 25 30 35
 gtg agg aaa tct aaa agt cct cct aaa ttt gaa gac tat tgg agg cac 302

Val	Arg	Lys	Ser	Lys	Ser	Pro	Pro	Lys	Phe	Glu	Asp	Tyr	Trp	Arg	His		
			40					45					50				
agg	aca	cca	gca	tcc	ttc	cag	agg	aag	ctg	cta	ggc	agc	cct	tcc	ctg	350	
Arg	Thr	Pro	Ala	Ser	Phe	Gln	Arg	Lys	Leu	Leu	Gly	Ser	Pro	Ser	Leu		
		55					60					65					
tct	gag	gaa	agc	cat	cga	att	tcc	atc	ccc	tcc	tca	gcc	atc	tcc	cac	398	
Ser	Glu	Glu	Ser	His	Arg	Ile	Ser	Ile	Pro	Ser	Ser	Ala	Ile	Ser	His		
	70					75					80						
aga	ggc	caa	cgc	acc	aaa	agg	gcc	cag	cct	tca	gct	gca	gaa	gga	aga	446	
Arg	Gly	Gln	Arg	Thr	Lys	Arg	Ala	Gln	Pro	Ser	Ala	Ala	Glu	Gly	Arg		
	85				90					95					100		
gaa	cat	ctc	cct	gaa	gca	ggg	tca	caa	aag	tgt	gga	gga	cct	gaa	ttc	494	
Glu	His	Leu	Pro	Glu	Ala	Gly	Ser	Gln	Lys	Cys	Gly	Gly	Pro	Glu	Phe		
				105				110						115			
tcc	ttt	gat	ttg	ctg	ccc	gag	gtg	cag	gct	gtt	cgg	gtg	act	att	cct	542	
Ser	Phe	Asp	Leu	Leu	Pro	Glu	Val	Gln	Ala	Val	Arg	Val	Thr	Ile	Pro		
			120					125					130				
gca	ggc	ccc	aag	gca	cgt	gtg	cgc	ctt	tgt	tat	cag	tgg	gca	ctg	gaa	590	
Ala	Gly	Pro	Lys	Ala	Arg	Val	Arg	Leu	Cys	Tyr	Gln	Trp	Ala	Leu	Glu		
		135				140						145					
tgt	gaa	gac	ttg	agt	agc	cct	ttt	gat	acc	cag	aaa	att	gtg	tct	gga	638	
Cys	Glu	Asp	Leu	Ser	Ser	Pro	Phe	Asp	Thr	Gln	Lys	Ile	Val	Ser	Gly		
	150					155					160						
ggg	cac	act	gta	gac	ctg	cct	tat	gaa	ttc	ctt	ctg	ccc	tgc	atg	tgc	686	
Gly	His	Thr	Val	Asp	Leu	Pro	Tyr	Glu	Phe	Leu	Leu	Pro	Cys	Met	Cys		
	165				170					175					180		
ata	gag	gcc	tcc	tac	ctg	caa	gag	gac	act	gtg	agg	cgc	aaa	agt	gtc	734	
Ile	Glu	Ala	Ser	Tyr	Leu	Gln	Glu	Asp	Thr	Val	Arg	Arg	Lys	Ser	Val		
				185					190					195			
cct	tcc	aga	gct	ggc	ctg	aag	ctt	atg	gct	cag	act	tct	ggc	agt	caa	782	
Pro	Ser	Arg	Ala	Gly	Leu	Lys	Leu	Met	Ala	Gln	Thr	Ser	Gly	Ser	Gln		
			200					205					210				
tac	gct	tca	ctg	act	aca	gcc	agc	ac								808	
Tyr	Ala	Ser	Leu	Thr	Thr	Ala	Ser										
		215				220											
<210> 20																	
<211> 243																	
<212> PRT																	
<213> Unknown																	
<400> 20																	
Met	Gly	Ser	Pro	Arg	Leu	Ala	Ala	Leu	Leu	Leu	Ser	Leu	Pro	Leu	Leu		
			-20					-15					-10				
Leu	Ile	Gly	Leu	Ala	Val	Ser	Ala	Arg	Val	Ala	Cys	Pro	Cys	Leu	Arg		
	-5					-1	1				5						

Ser Trp Thr Ser His Cys Leu Leu Ala Tyr Arg Val Asp Lys Arg Phe
 10 15 20 25
 Ala Gly Leu Gln Trp Gly Trp Phe Pro Leu Leu Val Arg Lys Ser Lys
 30 35 40
 Ser Pro Pro Lys Phe Glu Asp Tyr Trp Arg His Arg Thr Pro Ala Ser
 45 50 55
 Phe Gln Arg Lys Leu Leu Gly Ser Pro Ser Leu Ser Glu Glu Ser His
 60 65 70
 Arg Ile Ser Ile Pro Ser Ser Ala Ile Ser His Arg Gly Gln Arg Thr
 75 80 85
 Lys Arg Ala Gln Pro Ser Ala Ala Glu Gly Arg Glu His Leu Pro Glu
 90 95 100 105
 Ala Gly Ser Gln Lys Cys Gly Gly Pro Glu Phe Ser Phe Asp Leu Leu
 110 115 120
 Pro Glu Val Gln Ala Val Arg Val Thr Ile Pro Ala Gly Pro Lys Ala
 125 130 135
 Arg Val Arg Leu Cys Tyr Gln Trp Ala Leu Glu Cys Glu Asp Leu Ser
 140 145 150
 Ser Pro Phe Asp Thr Gln Lys Ile Val Ser Gly Gly His Thr Val Asp
 155 160 165
 Leu Pro Tyr Glu Phe Leu Leu Pro Cys Met Cys Ile Glu Ala Ser Tyr
 170 175 180 185
 Leu Gln Glu Asp Thr Val Arg Arg Lys Ser Val Pro Ser Arg Ala Gly
 190 195 200
 Leu Lys Leu Met Ala Gln Thr Ser Gly Ser Gln Tyr Ala Ser Leu Thr
 205 210 215
 Thr Ala Ser
 220

<210> 21

<211> 729

<212> DNA

<213> reverse translation

<220>

<221> misc_feature

<222> (1)..(729)

<223> n may be a, c, g, or t

<400> 21

atgggnwsnc cnmgnytnge ngcnytnytn ytawsnytn cnytnytnyt nathggnytn 60

gcngtnwsng cnmgngtnge ntgyccntgy ytnmgnwsnt ggacnwsnca ytggytnytn 120

gcntaymgng tngayaarmg nttygcnggn ytncartggg gntgggttycc nytnytnngtn 180

mgnaarwsna arwsnccncc naarttygar gaytaytggm gncaymgnac nccngcnwsn 240
 ttycarmgna arytnytngg nwsnccnwsn ytnwsngarg arwsncaymg nathwsnath 300
 ccnwsnwsng cnathwsnca ymgnggncar mgnacnaarm gngcncarcc nwsngcngcn 360
 garggnmgng arcayytnc ngargcnggn wncaraart gyggnggncc ngarttywsn 420
 ttygayytny tncngargt ncargcngtn mgngtnacna thcngcngg nccnaargcn 480
 mgngtnmgny tntgytayca rtgggcnyn gartgygarg ayytnwsnws nccnttygay 540
 acncaraara thgtnwsngg nggncayacn gtngayytnc cntaygartt yytnytncn 600
 tgyatgtgya thgargcnws ntayytncar gargayacng tnmgnmgnaa rwsngtnccn 660
 wsnmgngcng gnytnaaryt natggcncar acnwsnggnw sncartaygc nwsnytnacn 720
 acngcnwsn 729

<210> 22

<211> 2377

<212> DNA

<213> Unknown

<220>

<223> Description of Unknown Organism: primate; surmised
Homo sapiens

<220>

<221> CDS

<222> (180)..(1874)

<400> 22

ttttgagcag aggccttccta ggctccgtag aaatttgcac acagcttcca cttcctgctt 60
 cagagcctgt tcttctactt acctgggccc ggagaagggtg gagggagacg agaagccgcc 120
 gagagccgac taccctccgg gccagctctg tctgtccgtg gtggatctaa gaaactaga 179
 atg aac cga agc att cct gtg gag gtt gat gaa tca gaa cca tac cca 227
 Met Asn Arg Ser Ile Pro Val Glu Val Asp Glu Ser Glu Pro Tyr Pro
 1 5 10 15
 agt cag ttg ctg aaa cca atc cca gaa tat tcc ccg gaa gag gaa tca 275
 Ser Gln Leu Leu Lys Pro Ile Pro Glu Tyr Ser Pro Glu Glu Glu Ser
 20 25 30
 gaa cca cct gct cca aat ata agg aac atg gca ccc aac agc ttg tct 323
 Glu Pro Pro Ala Pro Asn Ile Arg Asn Met Ala Pro Asn Ser Leu Ser
 35 40 45
 gca ccc aca atg ctt cac aat tcc tcc gga gac ttt tct caa gct cac 371
 Ala Pro Thr Met Leu His Asn Ser Ser Gly Asp Phe Ser Gln Ala His
 50 55 60
 tca acc ctg aaa ctt gca aat cac cag cgg cct gta tcc cgg cag gtc 419
 Ser Thr Leu Lys Leu Ala Asn His Gln Arg Pro Val Ser Arg Gln Val
 65 70 75 80

acc tgc ctg cgc act caa gtt ctg gag gac agt gaa gac agt ttc tgc	467
Thr Cys Leu Arg Thr Gln Val Leu Glu Asp Ser Glu Asp Ser Phe Cys	
85 90 95	
agg aga cac cca ggc ctg ggc aaa gct ttc cct tct ggg tgc tct gca	515
Arg Arg His Pro Gly Leu Gly Lys Ala Phe Pro Ser Gly Cys Ser Ala	
100 105 110	
gtc agc gag cct gcg tct gag tct gtg gtt gga gcc ctc cct gca gag	563
Val Ser Glu Pro Ala Ser Glu Ser Val Val Gly Ala Leu Pro Ala Glu	
115 120 125	
cat cag ttt tca ttt atg gaa aaa cgt aat caa tgg ctg gta tct cag	611
His Gln Phe Ser Phe Met Glu Lys Arg Asn Gln Trp Leu Val Ser Gln	
130 135 140	
ctt tca gcg gct tct cct gac act ggc cat gac tca gac aaa tca gac	659
Leu Ser Ala Ala Ser Pro Asp Thr Gly His Asp Ser Asp Lys Ser Asp	
145 150 155 160	
caa agt tta cct aat gcc tca gca gac tcc ttg ggc ggt agc cag gag	707
Gln Ser Leu Pro Asn Ala Ser Ala Asp Ser Leu Gly Gly Ser Gln Glu	
165 170 175	
atg gtg caa cgg ccc cag cct cac agg aac cga gca ggc ctg gat ctg	755
Met Val Gln Arg Pro Gln Pro His Arg Asn Arg Ala Gly Leu Asp Leu	
180 185 190	
cca acc ata gac acg gga tat gat tcc cag ccc cag gat gtc ctg ggc	803
Pro Thr Ile Asp Thr Gly Tyr Asp Ser Gln Pro Gln Asp Val Leu Gly	
195 200 205	
atc agg cag ctg gaa agg ccc ctg ccc ctc acc tcc gtg tgt tac ccc	851
Ile Arg Gln Leu Glu Arg Pro Leu Pro Leu Thr Ser Val Cys Tyr Pro	
210 215 220	
cag gac ctc ccc aga cct ctc agg tcc agg gag ttc cct cag ttt gaa	899
Gln Asp Leu Pro Arg Pro Leu Arg Ser Arg Glu Phe Pro Gln Phe Glu	
225 230 235 240	
cct cag agg tat cca gca tgt gca cag atg ctg cct ccc aat ctt tcc	947
Pro Gln Arg Tyr Pro Ala Cys Ala Gln Met Leu Pro Pro Asn Leu Ser	
245 250 255	
cca cat gct cca tgg aac tat cat tac cat tgt cct gga agt ccc gat	995
Pro His Ala Pro Trp Asn Tyr His Tyr His Cys Pro Gly Ser Pro Asp	
260 265 270	
cac cag gtg cca tat ggc cat gac tac cct cga gca gcc tac cag caa	1043
His Gln Val Pro Tyr Gly His Asp Tyr Pro Arg Ala Ala Tyr Gln Gln	
275 280 285	
gtg atc cag ccg gct ctg cct ggg cag ccc ctg cct gga gcc agt gtg	1091
Val Ile Gln Pro Ala Leu Pro Gly Gln Pro Leu Pro Gly Ala Ser Val	
290 295 300	
aga ggc ctg cac cct gtg cag aag gtt atc ctg aat tat ccc agc ccc	1139
Arg Gly Leu His Pro Val Gln Lys Val Ile Leu Asn Tyr Pro Ser Pro	
305 310 315 320	

tgg gac caa gaa gag agg ccc gca cag aga gac tgc tcc ttt ccg ggg	1187
Trp Asp Gln Glu Glu Arg Pro Ala Gln Arg Asp Cys Ser Phe Pro Gly	
325 330 335	
ctt cca agg cac cag gac cag cca cat cac cag cca cct aat aga gct	1235
Leu Pro Arg His Gln Asp Gln Pro His His Gln Pro Pro Asn Arg Ala	
340 345 350	
ggg gct cct ggg gag tcc ttg gag tgc cct gca gag ctg aga cca cag	1283
Gly Ala Pro Gly Glu Ser Leu Glu Cys Pro Ala Glu Leu Arg Pro Gln	
355 360 365	
gtt ccc cag cct ccg tcc cca gct gct gtg cct aga ccc cct agc aac	1331
Val Pro Gln Pro Pro Ser Pro Ala Ala Val Pro Arg Pro Pro Ser Asn	
370 375 380	
cct cca gcc aga gga act cta aaa aca agc aat ttg cca gaa gaa ttg	1379
Pro Pro Ala Arg Gly Thr Leu Lys Thr Ser Asn Leu Pro Glu Glu Leu	
385 390 395 400	
cgg aaa gtc ttt atc act tat tcg atg gac aca gct atg gag gtg gtg	1427
Arg Lys Val Phe Ile Thr Tyr Ser Met Asp Thr Ala Met Glu Val Val	
405 410 415	
aaa ttc gtg aac ttt ttg ttg gta aat ggc ttc caa act gca att gac	1475
Lys Phe Val Asn Phe Leu Leu Val Asn Gly Phe Gln Thr Ala Ile Asp	
420 425 430	
ata ttt gag gat aga atc cga ggc att gat atc att aaa ttg atg gag	1523
Ile Phe Glu Asp Arg Ile Arg Gly Ile Asp Ile Ile Lys Trp Met Glu	
435 440 445	
cgc tac ctt agg gat aag acc gtg atg ata atc gta gca atc agc ccc	1571
Arg Tyr Leu Arg Asp Lys Thr Val Met Ile Ile Val Ala Ile Ser Pro	
450 455 460	
aaa tac aaa cag gac gtg gaa ggc gct gag tcg cag ctg gac gag gat	1619
Lys Tyr Lys Gln Asp Val Glu Gly Ala Glu Ser Gln Leu Asp Glu Asp	
465 470 475 480	
gag cat ggc tta cat act aag tac att cat cga atg atg cag att gag	1667
Glu His Gly Leu His Thr Lys Tyr Ile His Arg Met Met Gln Ile Glu	
485 490 495	
ttc ata aaa caa gga agc atg aat ttc aga ttc atc cct gtg ctg ttc	1715
Phe Ile Lys Gln Gly Ser Met Asn Phe Arg Phe Ile Pro Val Leu Phe	
500 505 510	
cca aat gct aag aag gag cat gtg ccc acc tgg ctt cag aac act cat	1763
Pro Asn Ala Lys Lys Glu His Val Pro Thr Trp Leu Gln Asn Thr His	
515 520 525	
gtc tac agc tgg ccc aag aat aaa aaa aac atc ctg ctg cgg ctg ctg	1811
Val Tyr Ser Trp Pro Lys Asn Lys Lys Asn Ile Leu Leu Arg Leu Leu	
530 535 540	
aga gag gaa gag tat gtg gct cct cca cgg ggg cct ctg ccc acc ctt	1859
Arg Glu Glu Glu Tyr Val Ala Pro Pro Arg Gly Pro Leu Pro Thr Leu	
545 550 555 560	

cag gtg gtt ccc ttg tgacaccgtt catccccaga tcaactgaggc caggccatgt 1914
 Gln Val Val Pro Leu
 565

ttggggcctt gttctgacag cattctggct gaggctggtc ggtagcactc ctggctgggt 1974
 tttttctgtt cctccccgag aggccctctg gccccagga aacctgttgt gcagagctct 2034
 tccccggaga cctccacaca ccctggcttt gaagtggagt ctgtgactgc tctgcattct 2094
 ctgcttttaa aaaaaccatt gcaggtgcc a gtgtcccata tgttcctcct gacagtttga 2154
 tgtgtccatt ctgggcctct cagtgccttag caagtagata atgtaaggga tgtggcagca 2214
 aatggaaatg actacaaaca ctctcctatc aatcacttca ggctactttt atgagttagc 2274
 cagatgcttg tgtatcctca gaccaaactg attcatgtac aaataataaa atgtttactc 2334
 ttttgtaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaa 2377

<210> 23
 <211> 565
 <212> PRT
 <213> Unknown

<400> 23
 Met Asn Arg Ser Ile Pro Val Glu Val Asp Glu Ser Glu Pro Tyr Pro
 1 5 10 15
 Ser Gln Leu Leu Lys Pro Ile Pro Glu Tyr Ser Pro Glu Glu Glu Ser
 20 25 30
 Glu Pro Pro Ala Pro Asn Ile Arg Asn Met Ala Pro Asn Ser Leu Ser
 35 40 45
 Ala Pro Thr Met Leu His Asn Ser Ser Gly Asp Phe Ser Gln Ala His
 50 55 60
 Ser Thr Leu Lys Leu Ala Asn His Gln Arg Pro Val Ser Arg Gln Val
 65 70 75 80
 Thr Cys Leu Arg Thr Gln Val Leu Glu Asp Ser Glu Asp Ser Phe Cys
 85 90 95
 Arg Arg His Pro Gly Leu Gly Lys Ala Phe Pro Ser Gly Cys Ser Ala
 100 105 110
 Val Ser Glu Pro Ala Ser Glu Ser Val Val Gly Ala Leu Pro Ala Glu
 115 120 125
 His Gln Phe Ser Phe Met Glu Lys Arg Asn Gln Trp Leu Val Ser Gln
 130 135 140
 Leu Ser Ala Ala Ser Pro Asp Thr Gly His Asp Ser Asp Lys Ser Asp
 145 150 155 160
 Gln Ser Leu Pro Asn Ala Ser Ala Asp Ser Leu Gly Gly Ser Gln Glu
 165 170 175

Met Val Gln Arg Pro Gln Pro His Arg Asn Arg Ala Gly Leu Asp Leu
 180 185 190
 Pro Thr Ile Asp Thr Gly Tyr Asp Ser Gln Pro Gln Asp Val Leu Gly
 195 200 205
 Ile Arg Gln Leu Glu Arg Pro Leu Pro Leu Thr Ser Val Cys Tyr Pro
 210 215 220
 Gln Asp Leu Pro Arg Pro Leu Arg Ser Arg Glu Phe Pro Gln Phe Glu
 225 230 235 240
 Pro Gln Arg Tyr Pro Ala Cys Ala Gln Met Leu Pro Pro Asn Leu Ser
 245 250 255
 Pro His Ala Pro Trp Asn Tyr His Tyr His Cys Pro Gly Ser Pro Asp
 260 265 270
 His Gln Val Pro Tyr Gly His Asp Tyr Pro Arg Ala Ala Tyr Gln Gln
 275 280 285
 Val Ile Gln Pro Ala Leu Pro Gly Gln Pro Leu Pro Gly Ala Ser Val
 290 295 300
 Arg Gly Leu His Pro Val Gln Lys Val Ile Leu Asn Tyr Pro Ser Pro
 305 310 315 320
 Trp Asp Gln Glu Glu Arg Pro Ala Gln Arg Asp Cys Ser Phe Pro Gly
 325 330 335
 Leu Pro Arg His Gln Asp Gln Pro His His Gln Pro Pro Asn Arg Ala
 340 345 350
 Gly Ala Pro Gly Glu Ser Leu Glu Cys Pro Ala Glu Leu Arg Pro Gln
 355 360 365
 Val Pro Gln Pro Pro Ser Pro Ala Ala Val Pro Arg Pro Pro Ser Asn
 370 375 380
 Pro Pro Ala Arg Gly Thr Leu Lys Thr Ser Asn Leu Pro Glu Glu Leu
 385 390 395 400
 Arg Lys Val Phe Ile Thr Tyr Ser Met Asp Thr Ala Met Glu Val Val
 405 410 415
 Lys Phe Val Asn Phe Leu Leu Val Asn Gly Phe Gln Thr Ala Ile Asp
 420 425 430
 Ile Phe Glu Asp Arg Ile Arg Gly Ile Asp Ile Ile Lys Trp Met Glu
 435 440 445
 Arg Tyr Leu Arg Asp Lys Thr Val Met Ile Ile Val Ala Ile Ser Pro
 450 455 460
 Lys Tyr Lys Gln Asp Val Glu Gly Ala Glu Ser Gln Leu Asp Glu Asp
 465 470 475 480
 Glu His Gly Leu His Thr Lys Tyr Ile His Arg Met Met Gln Ile Glu
 485 490 495

Phe Ile Lys Gln Gly Ser Met Asn Phe Arg Phe Ile Pro Val Leu Phe
 500 505 510
 Pro Asn Ala Lys Lys Glu His Val Pro Thr Trp Leu Gln Asn Thr His
 515 520 525
 Val Tyr Ser Trp Pro Lys Asn Lys Lys Asn Ile Leu Leu Arg Leu Leu
 530 535 540
 Arg Glu Glu Glu Tyr Val Ala Pro Pro Arg Gly Pro Leu Pro Thr Leu
 545 550 555 560
 Gln Val Val Pro Leu
 565

<210> 24
 <211> 1695
 <212> DNA
 <213> reverse translation

<220>
 <221> misc_feature
 <222> (1)..(1695)
 <223> n may be a, c, g, or t

<400> 24
 atgaaymgmw snathccngt ngargtngay garwsngarc cntayccnws ncarytnytn 60
 aarccnathc cngartayws nccngargar garwsngarc cncngcncnc naayathmgn 120
 aayatggcnc cnaaywsnyt nwsngcncnc acnatgytnc ayaaywsnws nggngaytty 180
 wsnarcgcnc aywsnacnyt naarytngcn aaycaycarm gncngtnws nmgnarcgtn 240
 acntgyytnm gnacncargt nytngargay wsngargayw snttytgymg nmgnacayccn 300
 ggnytnnggna argcnttycc nwsnggntgy wsngcngtnw sngarccngc nwsngarwsn 360
 gtngtnnggng cnytnccngc ngarcaycar ttywsnttya tggaraarmg naaycartgg 420
 ytngtnwsnc arytnwsngc ngcnwsnccn gayacnggnc aygaywsnga yaarwsngay 480
 carwsnytn cnaaygcnws ngcngaywsn ytnggnggnw sncargarat ggtncarmgn 540
 ccncarccnc aymgnaaymg ngcnggnytn gayytnccna cnathgayac nggntaygay 600
 wsnarcccnc argaygtnyt nggnathmgn carytngarm gncnytncc nytnacnwsn 660
 gtntgytayc cncargayt nccnmgnccn ytnmgnwsnm gngarttycc ncarttygar 720
 ccncarmgnt ayccngcntg ygncaratg ytnccncna ayytnwsncc ncaygcncn 780
 tggaaytayc aytaaycayt gccnggnwsn ccngaycayc argtnccnta yggncaygay 840
 tayccnmgng cngcntayca rcargtnath carccngcny tncnggnca rccnytnccn 900
 ggngcnwsng tnmnggnytn ncayccngtn caraargtna thytnaayta yccnwsnccn 960

tgggaycarg argarmgncc ngcncarmgn gaytgywsnt tyccnggnynt nccnmgn cay 1020
 cargaycarc cncaycayca rccnccnaay mgngcnggng cncnggnga rwsnytn gar 1080
 tgyccngcng arytnmgnc ncartnccn carccnccnw snccngcngc ngtnccnmgn 1140
 ccnccnwsna ayccnccngc nmngngnacn ytnaaracnw snaaytncc ngargarytn 1200
 mgnaargtnt tyathacnta ywsnatggay acngcnatgg argtngtnaa rttygtnaay 1260
 ttyytnytn g tnaayggntt ycaracngcn athgayatht tygargaymg nathmgnggn 1320
 athgayatha thaartggat ggarmgntay ytnmgngaya aracngtnat gathathgtn 1380
 gcnathwsnc cnaartayaa rcargaygtn garggngcng arwsncaryt ngaygargay 1440
 garcayggny tncayacnaa rtayathcay mgntatgatgc arathgartt yathaarcar 1500
 ggnwsnatga ayttymgntt yathccngtn ytnttyccna aygcnaaraa rgarcaygtn 1560
 ccnacntggy tncaraayac ncaygtntay wsntggccna araayaaraa raayathytn 1620
 ytnmgnytny tnmngngarga rgartaygtn gcncnccnm gnggncnynt nccnacnytn 1680
 cargtngtnc cnytn 1695

<210> 25

<211> 1323

<212> DNA

<213> Unknown

<220>

<223> Description of Unknown Organism: rodent; surmised
Mus musculus

<220>

<221> CDS

<222> (1)..(1026)

<400> 25

cag gac ctc cct ggg cct ctg agg tcc agg gaa ttg cca cct cag ttt	48
Gln Asp Leu Pro Gly Pro Leu Arg Ser Arg Glu Leu Pro Pro Gln Phe	
1 5 10 15	
gaa ctt gag agg tat cca atg aac gcc cag ctg ctg ccg ccc cat cct	96
Glu Leu Glu Arg Tyr Pro Met Asn Ala Gln Leu Leu Pro Pro His Pro	
20 25 30	
tcc cca cag gcc cca tgg aac tgt cag tac tac tgc ccc gga ggg ccc	144
Ser Pro Gln Ala Pro Trp Asn Cys Gln Tyr Tyr Cys Pro Gly Gly Pro	
35 40 45	
tac cac cac cag gtg cca cac ggc cat ggc tac cct cca gca gca gcc	192
Tyr His His Gln Val Pro His Gly His Gly Tyr Pro Pro Ala Ala Ala	
50 55 60	
tac cag caa gta ctc cag cct gct ctg cct ggg cag gtc ctt cct ggg	240
Tyr Gln Gln Val Leu Gln Pro Ala Leu Pro Gly Gln Val Leu Pro Gly	
65 70 75 80	

gca agg gca aga ggc cca cgc cct gtg cag aag gtc atc ctg aat gac	288
Ala Arg Ala Arg Gly Pro Arg Pro Val Gln Lys Val Ile Leu Asn Asp	
85 90 95	
tcc agc ccc caa gac caa gaa gag aga cct gca cag aga gac ttc tct	336
Ser Ser Pro Gln Asp Gln Glu Glu Arg Pro Ala Gln Arg Asp Phe Ser	
100 105 110	
ttc ccg agg ctc ccg agg gac cag ctc tac cgc cca cca tct aat gga	384
Phe Pro Arg Leu Pro Arg Asp Gln Leu Tyr Arg Pro Pro Ser Asn Gly	
115 120 125	
gtg gaa gcc cct gag gag tcc ttg gac ctt cct gca gag ctg aga cca	432
Val Glu Ala Pro Glu Glu Ser Leu Asp Leu Pro Ala Glu Leu Arg Pro	
130 135 140	
cat ggt ccc cag gct cca tcc cta gct gcc gtg cct aga ccc cct agc	480
His Gly Pro Gln Ala Pro Ser Leu Ala Ala Val Pro Arg Pro Pro Ser	
145 150 155 160	
aac ccc tta gcc cga gga act cta aga acc agc aat ttg cca gaa gaa	528
Asn Pro Leu Ala Arg Gly Thr Leu Arg Thr Ser Asn Leu Pro Glu Glu	
165 170 175	
tta cgg aaa gtc ttt atc act tat tct atg gac aca gcc atg gag gtg	576
Leu Arg Lys Val Phe Ile Thr Tyr Ser Met Asp Thr Ala Met Glu Val	
180 185 190	
gtg aaa ttt gtg aac ttt ctg ttg gtg aac ggc ttc caa act gcg att	624
Val Lys Phe Val Asn Phe Leu Leu Val Asn Gly Phe Gln Thr Ala Ile	
195 200 205	
gac ata ttt gag gat aga atc cgg ggt att gat atc att aaa tgg atg	672
Asp Ile Phe Glu Asp Arg Ile Arg Gly Ile Asp Ile Ile Lys Trp Met	
210 215 220	
gag cgc tat ctt cga gat aag aca gtg atg ata atc gta gca atc agc	720
Glu Arg Tyr Leu Arg Asp Lys Thr Val Met Ile Ile Val Ala Ile Ser	
225 230 235 240	
ccc aaa tac aaa cag gat gtg gaa ggc gct gag tcg cag ctg gac gag	768
Pro Lys Tyr Lys Gln Asp Val Glu Gly Ala Glu Ser Gln Leu Asp Glu	
245 250 255	
gac gag cat ggc tta cat act aag tac att cat cgg atg atg cag att	816
Asp Glu His Gly Leu His Thr Lys Tyr Ile His Arg Met Met Gln Ile	
260 265 270	
gag ttc ata agt cag gga agc atg aac ttc aga ttc atc cct gtg ctc	864
Glu Phe Ile Ser Gln Gly Ser Met Asn Phe Arg Phe Ile Pro Val Leu	
275 280 285	
ttc cca aat gcc aag aag gag cat gtg ccg acc tgg ctt cag aac act	912
Phe Pro Asn Ala Lys Lys Glu His Val Pro Thr Trp Leu Gln Asn Thr	
290 295 300	
cat gtt tac agc tgg ccc aag aat aag aaa aac atc ctg ctg cgg ctg	960
His Val Tyr Ser Trp Pro Lys Asn Lys Lys Asn Ile Leu Leu Arg Leu	
305 310 315 320	

ctc agg gag gaa gag tat gtg gct cct ccc cga ggc cct ctg ccc acc 1008
 Leu Arg Glu Glu Glu Tyr Val Ala Pro Pro Arg Gly Pro Leu Pro Thr
 325 330 335

ctt cag gtg gta ccc ttg tgacgatggc cactccagct cagtgccagc 1056
 Leu Gln Val Val Pro Leu
 340

ctgtttctcac agcattcttc tagcggagct ggctggtggc acccaggccc tggaacacct 1116

cttctacaga gtcctctgtc tcctgagtct gagttgtcct cgctgggctt ccagagcttc 1176

agtgccctgga tgctgcaggt gacagaaaca aacatctatg accacaaaaa ctctcatcac 1236

ttcagctact tttatgagtc ggtcagatgc tctgtgtcct tagaccagtc taaatcatgc 1296

tcaaataata aaatgattat tctttgt 1323

<210> 26

<211> 342

<212> PRT

<213> Unknown

<400> 26

Gln Asp Leu Pro Gly Pro Leu Arg Ser Arg Glu Leu Pro Pro Gln Phe
 1 5 10 15

Glu Leu Glu Arg Tyr Pro Met Asn Ala Gln Leu Leu Pro Pro His Pro
 20 25 30

Ser Pro Gln Ala Pro Trp Asn Cys Gln Tyr Tyr Cys Pro Gly Gly Pro
 35 40 45

Tyr His His Gln Val Pro His Gly His Gly Tyr Pro Pro Ala Ala Ala
 50 55 60

Tyr Gln Gln Val Leu Gln Pro Ala Leu Pro Gly Gln Val Leu Pro Gly
 65 70 75 80

Ala Arg Ala Arg Gly Pro Arg Pro Val Gln Lys Val Ile Leu Asn Asp
 85 90 95

Ser Ser Pro Gln Asp Gln Glu Glu Arg Pro Ala Gln Arg Asp Phe Ser
 100 105 110

Phe Pro Arg Leu Pro Arg Asp Gln Leu Tyr Arg Pro Pro Ser Asn Gly
 115 120 125

Val Glu Ala Pro Glu Glu Ser Leu Asp Leu Pro Ala Glu Leu Arg Pro
 130 135 140

His Gly Pro Gln Ala Pro Ser Leu Ala Ala Val Pro Arg Pro Pro Ser
 145 150 155 160

Asn Pro Leu Ala Arg Gly Thr Leu Arg Thr Ser Asn Leu Pro Glu Glu
 165 170 175

Leu Arg Lys Val Phe Ile Thr Tyr Ser Met Asp Thr Ala Met Glu Val

180	185	190
Val Lys Phe Val Asn Phe Leu Leu Val Asn Gly Phe Gln Thr Ala Ile		
195	200	205
Asp Ile Phe Glu Asp Arg Ile Arg Gly Ile Asp Ile Ile Lys Trp Met		
210	215	220
Glu Arg Tyr Leu Arg Asp Lys Thr Val Met Ile Ile Val Ala Ile Ser		
225	230	235
Pro Lys Tyr Lys Gln Asp Val Glu Gly Ala Glu Ser Gln Leu Asp Glu		
245	250	255
Asp Glu His Gly Leu His Thr Lys Tyr Ile His Arg Met Met Gln Ile		
260	265	270
Glu Phe Ile Ser Gln Gly Ser Met Asn Phe Arg Phe Ile Pro Val Leu		
275	280	285
Phe Pro Asn Ala Lys Lys Glu His Val Pro Thr Trp Leu Gln Asn Thr		
290	295	300
His Val Tyr Ser Trp Pro Lys Asn Lys Lys Asn Ile Leu Leu Arg Leu		
305	310	315
Leu Arg Glu Glu Glu Tyr Val Ala Pro Pro Arg Gly Pro Leu Pro Thr		
325	330	335
Leu Gln Val Val Pro Leu		
340		

<210> 27
 <211> 1026
 <212> DNA
 <213> reverse translation

<220>
 <221> misc_feature
 <222> (1)..(1026)
 <223> n any be a, c, g, or t

<400> 27
 cargayytnc cnggnccnyt nmgnwsnmgn garytncnc cncarttyga rytngarmgn 60
 tayccnatga aygcncaryt nytcccncn cayccnwsnc ncargcncc ntggaaytgy 120
 cartaytayt gyccngngg nccntaycay caycargtnc ncayggncay yggntayccn 180
 ccngcngcng cntaycarca rgtnytncar ccngcnytn cnggncargt nytcccnggn 240
 gcnmgngcnm gnggnccnmg nccngtn car aargtnathy tnaaygayws nwsnccncar 300
 gaycargarg armgncngc ncarmngay ttywsnttyc cnmgnytncc nmngaycar 360
 ytnatymgnc cncnwsnaa yggngtn gar gcnccngarg arwsnytn ga yytnccngcn 420
 garytnmgnc ncayggnc ncargcnccn wsnytn gcn gngtnccnmg nccnccnwsn 480

aayccnytn g cnmgnggnac nytnmgnacn wsnaayytnc cngargaryt nmgnaargtn 540
 ttyathacnt aywsnatgga yacngcnatg gargtngtna arttygtnaa ytttytnytn 600
 gtnaayggnt tyacaracngc nathgayath ttygargaym gnathmgngg nathgayath 660
 athaartgga tggarmgnta yytnmgngay aaracngtna tgathathgt ngcnathwsn 720
 ccnaartaya arcargaygt ngarggngcn garwsncary tngaygarga ygar cayggn 780
 ytncaayacna artayathca ymgnatgatg carathgart tyathwsnca rggwnsnatg 840
 aayttymgnt tyathccngt nytnnttyccn aaygcnaara argarcaygt nccnacntgg 900
 ytncaraaya cncaygtnta ywsntggccn aaraayaara araayathyt nytnmgnytn 960
 ytnmgngarg argartaygt ngcnccnccn mgnggngccny tncnacny ncartngtn 1020
 ccnytn 1026

<210> 28

<211> 207

<212> PRT

<213> Unknown

<220>

<223> Description of Unknown Organism: primate; surmised
Homo sapiens

<400> 28

Arg	Lys	Val	Trp	Ile	Ile	Tyr	Ser	Ala	Asp	His	Pro	Leu	Tyr	Val	Asp
1				5					10					15	

Val	Val	Leu	Lys	Phe	Ala	Gln	Phe	Leu	Leu	Thr	Ala	Cys	Gly	Thr	Glu
			20					25					30		

Val	Ala	Leu	Asp	Leu	Leu	Glu	Glu	Gln	Ala	Ile	Ser	Glu	Ala	Gly	Val
		35				40						45			

Met	Thr	Trp	Val	Gly	Arg	Gln	Lys	Gln	Glu	Met	Val	Glu	Ser	Asn	Ser
	50				55						60				

Lys	Ile	Ile	Val	Leu	Cys	Ser	Arg	Gly	Thr	Arg	Ala	Lys	Trp	Gln	Ala
65					70					75				80	

Leu	Leu	Gly	Arg	Gly	Ala	Pro	Val	Arg	Leu	Arg	Cys	Asp	His	Gly	Lys
			85						90					95	

Pro	Val	Gly	Asp	Leu	Phe	Thr	Ala	Ala	Met	Asn	Met	Ile	Leu	Pro	Asp
		100					105						110		

Phe	Lys	Arg	Pro	Ala	Cys	Phe	Gly	Thr	Tyr	Val	Val	Cys	Tyr	Phe	Ser
		115					120					125			

Glu	Val	Ser	Cys	Asp	Gly	Asp	Val	Pro	Asp	Leu	Phe	Gly	Ala	Ala	Pro
	130				135						140				

Arg	Tyr	Pro	Leu	Met	Asp	Arg	Phe	Glu	Glu	Val	Tyr	Phe	Arg	Ile	Gln
145					150					155					160

```
<220>
<223> Description of Unknown Organism:rodent; surmised
      Mus musculus
```

<400> 29																
Arg	Lys	Val	Trp	Ile	Val	Tyr	Ser	Ala	Asp	His	Pro	Leu	Tyr	Val	Glu	
1				5					10					15		
Val	Val	Leu	Lys	Phe	Ala	Gln	Phe	Leu	Ile	Thr	Ala	Cys	Gly	Thr	Glu	
			20					25					30			
Val	Ala	Leu	Asp	Leu	Leu	Glu	Glu	Gln	Val	Ile	Ser	Glu	Val	Gly	Val	
		35					40					45				
Met	Thr	Trp	Val	Ser	Arg	Gln	Lys	Gln	Glu	Met	Val	Glu	Ser	Asn	Ser	
	50					55					60					
Lys	Ile	Ile	Ile	Leu	Cys	Ser	Arg	Gly	Thr	Gln	Ala	Lys	Trp	Lys	Ala	
65					70					75					80	
Ile	Leu	Gly	Trp	Ala	Glu	Pro	Ala	Val	Gln	Leu	Arg	Cys	Asp	His	Trp	
				85					90					95		
Lys	Pro	Ala	Gly	Asp	Leu	Phe	Thr	Ala	Ala	Met	Asn	Met	Ile	Leu	Pro	
			100					105					110			
Asp	Phe	Lys	Arg	Pro	Ala	Cys	Phe	Gly	Thr	Tyr	Val	Val	Cys	Tyr	Phe	
		115					120					125				
Ser	Gly	Ile	Cys	Ser	Glu	Arg	Asp	Val	Pro	Asp	Leu	Phe	Asn	Ile	Thr	
	130					135					140					
Ser	Arg	Tyr	Pro	Leu	Met	Asp	Arg	Phe	Glu	Glu	Val	Tyr	Phe	Arg	Ile	
145					150					155					160	
Gln	Asp	Leu	Glu	Met	Phe	Glu	Pro	Gly	Arg	Met	His	His	Val	Arg	Glu	
				165					170					175		
Leu	Thr	Gly	Asp	Asn	Tyr	Leu	Gln	Ser	Pro	Ser	Gly	Arg	Gln	Leu	Lys	
			180					185					190			
Glu	Ala	Val	Leu	Arg	Phe	Gln	Glu	Trp	Gln	Thr	Gln	Cys	Pro	Asp	Trp	
		195					200					205				

<210> 30
 <211> 190
 <212> PRT
 <213> Unknown

<220>

<223> Description of Unknown Organism:worm; surmised
 Caenorabditis elegans

<400> 30

Val	Lys	Val	Met	Ile	Val	Tyr	Ala	Asp	Asp	Asn	Asp	Leu	His	Thr	Asp	1	5	10	15
Cys	Val	Lys	Lys	Leu	Val	Glu	Asn	Leu	Arg	Asn	Cys	Ala	Ser	Cys	Asp	20	25	30	
Pro	Val	Phe	Asp	Leu	Glu	Lys	Leu	Ile	Thr	Ala	Glu	Ile	Val	Pro	Ser	35	40	45	
Arg	Trp	Leu	Val	Asp	Gln	Ile	Ser	Ser	Leu	Lys	Lys	Phe	Ile	Ile	Val	50	55	60	
Val	Ser	Asp	Cys	Ala	Glu	Lys	Ile	Leu	Asp	Thr	Glu	Ala	Ser	Glu	Thr	65	70	75	80
His	Gln	Leu	Val	Gln	Ala	Arg	Pro	Phe	Ala	Asp	Leu	Phe	Gly	Pro	Ala	85	90	95	
Met	Glu	Met	Ile	Ile	Arg	Asp	Ala	Thr	His	Asn	Phe	Pro	Glu	Ala	Arg	100	105	110	
Lys	Lys	Tyr	Ala	Val	Val	Arg	Phe	Asn	Tyr	Ser	Pro	His	Val	Pro	Pro	115	120	125	
Asn	Leu	Ala	Ile	Leu	Asn	Leu	Pro	Thr	Phe	Ile	Pro	Glu	Gln	Phe	Ala	130	135	140	
Gln	Leu	Thr	Ala	Phe	Leu	His	Asn	Val	Glu	His	Thr	Glu	Arg	Ala	Asn	145	150	155	160
Val	Thr	Gln	Asn	Ile	Ser	Glu	Ala	Gln	Ile	His	Glu	Trp	Asn	Leu	Cys	165	170	175	
Ala	Ser	Arg	Met	Met	Ser	Phe	Phe	Val	Arg	Asn	Pro	Asn	Trp	180	185	190			

<210> 31
 <211> 178
 <212> PRT
 <213> Unknown

<220>

<223> Description of Unknown Organism:worm; surmised
 Caenorabditis elegans

<400> 31

```

Phe Lys Val Met Leu Val Cys Pro Glu Val Ser Gly Arg Asp Glu Asp
 1             5             10             15

Phe Met Met Arg Ile Ala Asp Ala Leu Lys Lys Ser Asn Asn Lys Val
      20             25             30

Val Cys Asp Arg Trp Phe Glu Asp Ser Lys Asn Ala Glu Glu Asn Met
      35             40             45

Leu His Trp Val Tyr Glu Gln Thr Lys Ile Ala Glu Lys Ile Ile Val
 50             55             60

Phe His Ser Ala Tyr Tyr His Pro Arg Cys Gly Ile Tyr Asp Val Ile
 65             70             75             80

Asn Asn Phe Phe Pro Cys Thr Asp Pro Arg Leu Ala His Ile Ala Leu
      85             90             95

Thr Pro Glu Ala Gln Arg Ser Val Pro Lys Glu Val Glu Tyr Val Leu
      100             105             110

Pro Arg Asp Gln Lys Leu Leu Glu Asp Ala Phe Asp Ile Thr Ile Ala
      115             120             125

Asp Pro Leu Val Ile Asp Ile Pro Ile Glu Asp Val Ala Ile Pro Glu
      130             135             140

Asn Val Pro Ile His His Glu Ser Cys Asp Ser Ile Asp Ser Arg Asn
      145             150             155             160

Asn Ser Lys Thr His Ser Thr Asp Ser Gly Val Ser Ser Leu Ser Ser
      165             170             175

Asn Ser

```


(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 November 2001 (29.11.2001)

PCT

(10) International Publication Number
WO 01/090358 A3

(51) International Patent Classification⁷: **C12N 15/12**,
C07K 14/715, 16/18, G01N 33/53, C12N 5/10

(21) International Application Number: PCT/US01/16767

(22) International Filing Date: 23 May 2001 (23.05.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/206,862 24 May 2000 (24.05.2000) US

(71) Applicant: **SCHERING CORPORATION** [US/US];
2000 Galloping Hill Road, Kenilworth, NJ 07033-0530
(US).

(72) Inventor: **GORMAN, Daniel, M.**; 6371 Central Avenue,
Newark, CA 94560 (US).

(74) Agent: **ZARADIC, Sandy**; Schering-Plough Corpora-
tion, Patent Department, K-6-1, 1990, 2000 Galloping Hill
Road, Kenilworth, NJ 07033-0530 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CZ,

DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, HR, HU,
ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LU, LV,
MA, MD, MG, MK, MN, MX, MZ, NO, NZ, PL, PT, RO,
RU, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UZ, VN,
YU, ZA.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

*as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii)) for all designations*

Published:

— with international search report

(88) Date of publication of the international search report:

23 January 2003

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

WO 01/090358 A3

(54) Title: MAMMALIAN RECEPTOR PROTEINS; RELATED REAGENTS AND METHODS

(57) Abstract: Nucleic acids encoding mammalian, e.g., primate, receptors, purified receptor proteins and fragments thereof. Anti-
bodies, both polyclonal and monoclonal, are also provided. Methods of using the compositions for both diagnostic and therapeutic
utilities are described.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/16767

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C07K14/715 C07K16/18 G01N33/53 C12N5/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

SEQUENCE SEARCH, EMBL, EPO-Internal, MEDLINE, BIOSIS, WPI Data, PAJ, CHEM ABS Data, SCISEARCH, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 29408 A (IMMUNEX CORP) 26 September 1996 (1996-09-26) page 2, line 35 -page 15, line 4 ---	1-18
X	YAO Z ET AL: "MOLECULAR CHARACTERIZATION OF THE HUMAN INTERLEUKIN (IL)-17 RECEPTOR" CYTOKINE, ACADEMIC PRESS LTD, PHILADELPHIA, PA, US, vol. 9, no. 11, November 1997 (1997-11), pages 794-800, XP000867704 ISSN: 1043-4666 page 795; figure 2 --- -/--	1-4, 6, 12-15

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "I" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the International search

12 August 2002

Date of mailing of the International search report

29.08.02

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3018

Authorized officer

Steffen, P

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/16767

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL 'Online! EBI; 18 February 2000 (2000-02-18) BLOECKER ET AL.: "Homo sapiens mRNA; cDNA DKFZp434N1928" Database accession no. AL133097 XP002183453 the whole document</p> <p>---</p>	1-4, 6, 12-15
A	<p>WO 99 14240 A (HUMAN GENOME SCIENCES INC ; RUBEN STEVEN M (US); SHI YANGGU (US)) 25 March 1999 (1999-03-25) the whole document</p> <p>---</p>	
A	<p>TIAN E ET AL: "EVI27 ENCODES A NOVEL MEMBRANE PROTEIN WITH HOMOLOGY TO THE IL17 RECEPOR" ONCOGENE, BASINGSTOKE, HANTS, GB, vol. 19, no. 17, 20 April 2000 (2000-04-20), pages 2098-2109, XP008000240 ISSN: 0950-9232 the whole document</p> <p>---</p>	
A	<p>SHI YANGGU ET AL: "A novel cytokine receptor-ligand pair: Identification, molecular characterization, and in vivo immunomodulatory activity." JOURNAL OF BIOLOGICAL CHEMISTRY (JBC PAPERS IN PRESS, DOI 10.1074/JBC.M910228199), vol. 275, no. 25, 3 April 2000 (2000-04-03), pages 19167-19176, XP002197927 ISSN: 0021-9258 the whole document</p> <p>---</p>	
A	<p>FOSSIEZ F ET AL: "INTERLEUKIN-17" INTERNATIONAL REVIEWS OF IMMUNOLOGY, HARWOOD ACADEMIC PUBLISHERS, LONDON, GB, vol. 16, no. 5/6, 1998, pages 541-551, XP000867763 ISSN: 0883-0185 the whole document</p> <p>---</p>	
E	<p>WO 01 68859 A (AMGEN INC ; JING SHUQIAN (US)) 20 September 2001 (2001-09-20) page 2, line 19 -page 10, line 27; examples 1-4</p> <p>---</p>	1-18
E	<p>WO 01 46420 A (GENENTECH INC) 28 June 2001 (2001-06-28) page 5, line 1 -page 16, line 17; figures 17, 18</p> <p>---</p>	1-18

-/-

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/16767

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 55865 A (GENESIS RES & DEV CORP LTD) 4 November 1999 (1999-11-04) SEQ ID NO's 125 and 303 and corresponding cDNA's page 3 -page 17 ----	1-18
X	DATABASE EMBL 'Online! EBI; 22 July 1999 (1999-07-22) NCI-CGAP: "ty30c03.x1 NCI_CGAP_UT2 Homo sapiens cDNA clone IMAGE:2280580 3' mRNA sequence" Database accession no. AI861981 XP002209553 the whole document ----	12-18
X	DATABASE EMBL 'Online! EBI; 21 October 1999 (1999-10-21) MARRA ET AL.: "u191g04.y1 Sugano mouse kidney mkia Mus musculus cDNA clone IMAGE:2159478 5', mRNA sequence" Database accession no. AW107583 XP002209554 the whole document -----	12-18

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 01/16767

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 19, 20
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-18 (all partly)

Compositions comprising primate DCRS8 polypeptides and nucleic acid sequences (SEQ ID NO's 14 and 13, respectively) as well as further embodiments relating to the said polypeptides and nucleic acid sequences.

2. Claims: 1-18 (all partly)

Compositions comprising primate or rodent DCRS9 polypeptides and nucleic acid sequences (SEQ ID NO's 16, 19 and 17, 20, respectively) as well as further embodiments relating to the said polypeptides and nucleic acid sequences.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 19, 20

Present claims 19 and 20 relate to a method defined by reference to a desirable characteristic or property, namely contacting a cell with an unspecified agonist or antagonist of a mammalian protein of the application (e.g. DCRS8 or DCRS9).

The claims cover all methods having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such methods. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the method by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, no search has been carried out for claims 19 and 20.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/16767

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9629408	A	26-09-1996	AU 696775 B2	17-09-1998
			AU 5526396 A	08-10-1996
			CA 2215394 A1	26-09-1996
			EP 0817847 A1	14-01-1998
			JP 11503309 T	26-03-1999
			NO 974258 A	24-11-1997
			NZ 306653 A	29-03-1999
			US 6096305 A	01-08-2000
			US 6072033 A	06-06-2000
			US 6197525 B1	06-03-2001
			US 6191104 B1	20-02-2001
			US 6100235 A	08-08-2000
			US 6072037 A	06-06-2000
			WO 9629408 A1	26-09-1996
			US 5869286 A	09-02-1999
WO 9914240	A	25-03-1999	AU 9482498 A	05-04-1999
			EP 1015488 A1	05-07-2000
			WO 9914240 A1	25-03-1999
WO 0168859	A	20-09-2001	AU 4754501 A	24-09-2001
			WO 0168859 A2	20-09-2001
			US 2002045213 A1	18-04-2002
WO 0146420	A	28-06-2001	AU 2590901 A	03-07-2001
			WO 0146420 A2	28-06-2001
			AU 2055401 A	12-06-2001
			AU 2224800 A	28-09-2000
			AU 2596700 A	28-09-2000
			AU 2877900 A	28-09-2000
			AU 3514400 A	28-09-2000
			EP 1220905 A2	10-07-2002
			WO 0053754 A1	14-09-2000
			WO 0053756 A2	14-09-2000
			WO 0053758 A2	14-09-2000
			WO 0140466 A2	07-06-2001
			US 2002094416 A1	18-07-2002
			AU 3434601 A	12-06-2001
			AU 7573000 A	26-03-2001
			EP 1208202 A2	29-05-2002
			WO 0116318 A2	08-03-2001
			WO 0140465 A2	07-06-2001
			AU 2883600 A	28-09-2000
			AU 3381600 A	28-09-2000
			AU 3774300 A	18-12-2000
			AU 5152700 A	02-01-2001
			AU 5459900 A	18-12-2000
			AU 6391000 A	19-02-2001
			EP 1159419 A1	05-12-2001
			EP 1210418 A1	05-06-2002
			EP 1208195 A2	29-05-2002
			EP 1212417 A2	12-06-2002
			EP 1185648 A2	13-03-2002
			WO 0053757 A2	14-09-2000
			WO 0073454 A1	07-12-2000
			WO 0073445 A2	07-12-2000
			WO 0077037 A2	21-12-2000
			WO 0073348 A2	07-12-2000

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/16767

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0146420	A	WO 0073452 A2	07-12-2000
		WO 0109327 A2	08-02-2001
		US 2002072496 A1	13-06-2002
		US 2002072067 A1	13-06-2002
		US 2002072092 A1	13-06-2002
		US 2002072497 A1	13-06-2002
		AU 1916701 A	16-07-2001
		AU 6802801 A	24-09-2001
		WO 0149715 A2	12-07-2001
		WO 0168848 A2	20-09-2001
		US 2002098505 A1	25-07-2002
		US 2002098506 A1	25-07-2002
		US 2002098507 A1	25-07-2002
		US 2002090681 A1	11-07-2002
		US 2002010137 A1	24-01-2002
		US 2002058309 A1	16-05-2002
WO 9955865	A	04-11-1999	
		US 6150502 A	21-11-2000
		AU 3961499 A	16-11-1999
		CA 2327317 A1	04-11-1999
		CN 1307637 T	08-08-2001
		EP 1073740 A1	07-02-2001
		JP 2002512798 T	08-05-2002
		WO 9955865 A1	04-11-1999
		NZ 507728 A	28-06-2002

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 January 2002 (03.01.2002)

PCT

(10) International Publication Number
WO 02/00690 A2

- (51) International Patent Classification⁷: **C07K 14/00**
- (21) International Application Number: PCT/US01/19692
- (22) International Filing Date: 20 June 2001 (20.06.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
- | | | |
|----------------|--------------------------------|----|
| 60/213,637 | 23 June 2000 (23.06.2000) | US |
| 60/219,556 | 20 July 2000 (20.07.2000) | US |
| 60/220,624 | 25 July 2000 (25.07.2000) | US |
| 60/220,664 | 25 July 2000 (25.07.2000) | US |
| PCT/US00/20710 | 28 July 2000 (28.07.2000) | US |
| 60/222,695 | 2 August 2000 (02.08.2000) | US |
| 09/643,657 | 17 August 2000 (17.08.2000) | US |
| PCT/US00/23522 | 23 August 2000 (23.08.2000) | US |
| PCT/US00/23328 | 24 August 2000 (24.08.2000) | US |
| 60/230,978 | 7 September 2000 (07.09.2000) | US |
| 60/000,000 | 15 September 2000 (15.09.2000) | US |
| 09/664,610 | 18 September 2000 (18.09.2000) | US |
| 09/665,350 | 18 September 2000 (18.09.2000) | US |
| 60/242,922 | 24 October 2000 (24.10.2000) | US |
| 09/709,238 | 8 November 2000 (08.11.2000) | US |
| PCT/US00/30952 | 8 November 2000 (08.11.2000) | US |
| PCT/US00/30873 | 10 November 2000 (10.11.2000) | US |
| PCT/US00/32678 | 1 December 2000 (01.12.2000) | US |
| 09/747,259 | 20 December 2000 (20.12.2000) | US |
| PCT/US00/34956 | 20 December 2000 (20.12.2000) | US |
| 09/767,609 | 22 January 2001 (22.01.2001) | US |
| 09/796,498 | 28 February 2001 (28.02.2001) | US |
| PCT/US01/06520 | 28 February 2001 (28.02.2001) | US |
| PCT/US01/06666 | 1 March 2001 (01.03.2001) | US |
| 09/802,706 | 9 March 2001 (09.03.2001) | US |
| 09/808,689 | 14 March 2001 (14.03.2001) | US |
| 09/816,744 | 22 March 2001 (22.03.2001) | US |
| 09/828,366 | 5 April 2001 (05.04.2001) | US |
- (71) Applicant (for all designated States except US): **GENENTECH, INC.** [US/US]; 1 DNA Way, South San Francisco, CA 94080-4990 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **BAKER, Kevin, P.** [GB/US]; 14006 Indian Run Drive, Darnestown, MD 20878 (US). **FERRARA, Napoleone** [US/US]; #704, 2090 Pacific Avenue, San Francisco, CA 94109 (US). **GERBER, Hanspeter** [CH/US]; #5, 1121 Tennessee Street, San Francisco, CA 94107 (US). **GERRITSEN, Mary, E.** [CA/US]; 541 Parrott Drive, San Mateo, CA 94402 (US). **GODDARD, Audrey** [CA/US]; 110 Congo Street, San Francisco, CA 94131 (US). **GODOWSKI, Paul, J.** [US/US]; 2627 Easton Drive, Burlingame, CA 94010 (US). **GURNEY, Austin, L.** [US/US]; 1 Debbie Lane, Belmont, CA 94002 (US). **HILLAN, Kenneth, J.** [GB/US]; 64 Seward Street, San Francisco, CA 94114 (US). **MARSTERS, Scot, A.** [US/US]; 990 Cherry Street, San Carlos, CA 94070 (US). **PAN, James** [CA/US]; 2705 Coronet Boulevard, Belmont, CA 94002 (US). **PAONI, Nicholas, F.** [US/US]; 1756 Terrace Drive, Belmont, CA 94002 (US). **STEPHAN, Jean-Philippe, F.** [FR/US]; 320 C Lansdale Avenue, Millbrae, CA 94030 (US). **WATANABE, Colin, K.** [US/US]; 128 Corliss Drive, Moraga, CA 94556 (US). **WILLIAMS, P., Mickey** [US/US]; 509 Alto Avenue, Half Moon Bay, CA 94019 (US). **WOOD, William, I.** [US/US]; 35 Southdown Court, Hillsborough, CA 94010 (US). **YE, Weilan** [CN/US]; 119 Barkentine Street, Foster City, CA 94404 (US).
- (74) Agents: **AGARWAL, Atulya, R.** et al.; c/o GENENTECH, INC., MS49, 1 DNA Way, South San Francisco, CA 94080-4990 (US).

[Continued on next page]

WO 02/00690 A2

(54) Title: COMPOSITIONS AND METHODS FOR THE DIAGNOSIS AND TREATMENT OF DISORDERS INVOLVING ANGIOGENESIS

(57) Abstract: Compositions and methods are disclosed for stimulating or inhibiting angiogenesis and/or cardiovascularization in mammals, including humans. Pharmaceutical compositions are based on polypeptides or antagonists thereto that have been identified for one or more of these uses. Disorders that can be diagnosed, prevented, or treated by the compositions herein include trauma such as wounds, various cancers, and disorders of the vessels including atherosclerosis and cardiac hypertrophy. In addition, the present invention is directed to novel polypeptides and to nucleic acid molecules encoding those polypeptides. Also provided herein are vectors and host cells comprising those nucleic acid sequences, chimeric polypeptide molecules comprising the polypeptides of the present invention fused to heterologous polypeptide sequences, antibodies which bind to the polypeptides of the present invention and to methods for producing the polypeptides of the present invention.



(81) **Designated States (national):** AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

**COMPOSITIONS AND METHODS FOR THE DIAGNOSIS AND TREATMENT OF DISORDERS
INVOLVING ANGIOGENESIS**

5 1. Field of the Invention

The present invention relates to compositions and methods useful for the modulation (e.g., promotion or inhibition) of angiogenesis and/or cardiovascularization in mammals in need of such biological effect. The present invention further relates to the diagnosis and treatment of disorders involving angiogenesis (e.g., cardiovascular as well as oncological disorders).

10

2. Background of the Invention

2.1. Angiogenesis

Angiogenesis, defined as the growth or sprouting of new blood vessels from existing vessels, is a complex process that primarily occurs during embryonic development. Under normal physiological conditions in adults, angiogenesis takes place only in very restricted situations such as hair growth and wounding healing (Auerbach, W. and Auerbach, R., 1994, *Pharmacol Ther* 63(3):265-3 11; Ribatti et al., 1991, *Haematologica* 76(4):3 11-20; Risau, 1997, *Nature* 386(6626):67 1-4). Unregulated angiogenesis has gradually been recognized to be responsible for a wide range of disorders, including, but not limited to cardiovascular disease, cancer, rheumatoid arthritis, psoriasis and diabetic retinopathy (Folkman, 1995, *Nat Med* 1(1):27-31; Isner, 1999, *Circulation* 99(13): 1653-5; Koch, 1998, *Arthritis Rheum* 41(6):951-62; Walsh, 1999, *Rheumatology (Oxford)* 38(2):103-12; Ware and Simons, 1997, *Nat Med* 3(2): 158-64).

20

2.2. Cardiac Disorders and Factors

Heart failure affects approximately five million Americans, and new cases of heart failure number about 400,000 each year. It is the single most frequent cause of hospitalization for people age 65 and older in the United States. Recent advances in the management of acute cardiac diseases, including acute myocardial infarction, are resulting in an expanding patient population that will eventually develop chronic heart failure. From 1979 to 1995, hospitalizations for congestive heart failure (CHF) rose from 377,000 to 872,000 (a 130 percent increase) and CHF deaths increased 116 percent.

25

CHF is a syndrome characterized by left ventricular dysfunction, reduced exercise tolerance, impaired quality of life, and markedly shortened life expectancy. The sine qua non of heart failure is an inability of the heart to pump blood at a rate sufficient to meet the metabolic needs of the body's tissues (in other words, there is insufficient cardiac output).

30

At least four major compensatory mechanisms are activated in the setting of heart failure to boost cardiac output, including peripheral vasoconstriction, increased heart rate, increased cardiac contractility, and increased plasma volume. These effects are mediated primarily by the sympathetic nervous system and the renin-angiotensin system. See, Eichhorn, American Journal of Medicine, 104: 163-169 (1998). Increased output from the

35

sympathetic nervous system increases vascular tone, heart rate, and contractility. Angiotensin II elevates blood pressure by 1) directly stimulating vascular smooth muscle contraction, 2) promoting plasma volume expansion by stimulating aldosterone and antidiuretic hormone secretion, 3) stimulating sympathetic-mediated vascular tone, and 4) catalyzing the degradation of bradykinin, which has vasodilatory and natriuretic activity. *See*, review by Brown and Vaughan, Circulation, 97: 1411-1420 (1998). As noted below, angiotensin II may also have directly deleterious effects on the heart by promoting myocyte necrosis (impairing systolic function) and intracardiac fibrosis (impairing diastolic and in some cases systolic function). *See*, Weber, Circulation, 96: 4065-4082 (1998).

A consistent feature of congestive heart failure (CHF) is cardiac hypertrophy, an enlargement of the heart that is activated by both mechanical and hormonal stimuli and enables the heart to adapt to demands for increased cardiac output. Morgan and Baker, Circulation, 83: 13-25 (1991). This hypertrophic response is frequently associated with a variety of distinct pathological conditions such as hypertension, aortic stenosis, myocardial infarction, cardiomyopathy, valvular regurgitation, and intracardiac shunt, all of which result in chronic hemodynamic overload.

Hypertrophy is generally defined as an increase in size of an organ or structure independent of natural growth that does not involve tumor formation. Hypertrophy of the heart is due either to an increase in the mass of the individual cells (myocytes), or to an increase in the number of cells making up the tissue (hyperplasia), or both. While the enlargement of an embryonic heart is largely dependent on an increase in myocyte number (which continues until shortly after birth), post-natal cardiac myocytes lose their proliferative capacity. Further growth occurs through hypertrophy of the individual cells.

Adult myocyte hypertrophy is initially beneficial as a short term response to impaired cardiac function by permitting a decrease in the load on individual muscle fibers. With severe, long-standing overload, however, the hypertrophied cells begin to deteriorate and die. Katz, "Heart Failure", in: Katz A.M. ed., Physiology of the Heart (New York: Raven Press, 1992) pp. 638-668. Cardiac hypertrophy is a significant risk factor for both mortality and morbidity in the clinical course of heart failure. Katz, Trends Cardiovasc. Med., 5: 37-44 (1995). For further details of the causes and pathology of cardiac hypertrophy *see, e.g.*, Heart Disease. A Textbook of Cardiovascular Medicine, Braunwald, E. ed. (W.B. Saunders Co., 1988), Chapter 14, "Pathophysiology of Heart Failure."

On a cellular level, the heart is composed of myocytes and surrounding support cells, generically called non-myocytes. While non-myocytes are primarily fibroblast/mesenchymal cells, they also include endothelial and smooth muscle cells. Indeed, although myocytes make up most of the adult myocardial mass, they represent only about 30% of the total cell numbers present in heart. In response to hormonal, physiological, hemodynamic, and pathological stimuli, adult ventricular muscle cells can adapt to increased workloads through the activation of a hypertrophic process. This response is characterized by an increase in myocyte cell size and contractile protein content of individual cardiac muscle cells, without concomitant cell division and activation of embryonic genes, including the gene for atrial natriuretic peptide (ANP). Chien *et al.*, FASEB J., 5: 3037-3046 (1991); Chien *et al.*, Annu. Rev. Physiol., 55: 77-95 (1993). An increment in myocardial mass as a result of an increase in myocyte size that is associated with an accumulation of interstitial collagen within the extracellular matrix and around intramyocardial coronary arteries has been described in left ventricular hypertrophy secondary to pressure overload

in humans. Caspari *et al.*, Cardiovasc. Res., **11**: 554-558 (1977); Schwarz *et al.*, Am. J. Cardiol., **42**: 895-903 (1978); Hess *et al.*, Circulation, **63**: 360-371 (1981); Pearlman *et al.*, Lab. Invest., **46**: 158-164 (1982).

It has also been suggested that paracrine factors produced by non-myocyte supporting cells may additionally be involved in the development of cardiac hypertrophy, and various non-myocyte derived hypertrophic factors, such as, leukocyte inhibitory factor (LIF) and endothelin, have been identified. Metcalf, Growth Factors, **7**: 169-173 (1992); Kurzrock *et al.*, Endocrine Reviews, **12**: 208-217 (1991); Inoue *et al.*, Proc. Natl. Acad. Sci. USA, **86**: 2863-2867 (1989); Yanagisawa and Masaki, Trends Pharm. Sci., **10**: 374-378 (1989); U.S. Patent No. 5,573,762 (issued November 12, 1996). Further exemplary factors that have been identified as potential mediators of cardiac hypertrophy include cardiotrophin-1 (CT-1) (Pennica *et al.*, Proc. Nat. Acad. Sci. USA, **92**: 1142-1146 (1995)), catecholamines, adrenocorticosteroids, angiotensin, and prostaglandins.

At present, the treatment of cardiac hypertrophy varies depending on the underlying cardiac disease. Catecholamines, adrenocorticosteroids, angiotensin, prostaglandins, LIF, endothelin (including endothelin-1, -2, and -3 and big endothelin), and CT-1 are among the factors identified as potential mediators of hypertrophy. For example, beta-adrenergic receptor blocking drugs (beta-blockers, *e.g.*, propranolol, timolol, tertalolol, carteolol, nadolol, betaxolol, penbutolol, acetobutolol, atenolol, metoprolol, carvedilol, etc.) and verapamil have been used extensively in the treatment of hypertrophic cardiomyopathy. The beneficial effects of beta-blockers on symptoms (*e.g.*, chest pain) and exercise tolerance are largely due to a decrease in the heart rate with a consequent prolongation of diastole and increased passive ventricular filling. Thompson *et al.*, Br. Heart J., **44**: 488-98 (1980); Harrison *et al.*, Circulation, **29**: 84-98 (1964). Verapamil has been described to improve ventricular filling and probably reducing myocardial ischemia. Bonow *et al.*, Circulation, **72**: 853-64 (1985).

Nifedipine and diltiazem have also been used occasionally in the treatment of hypertrophic cardiomyopathy. Lorell *et al.*, Circulation, **65**: 499-507 (1982); Betocchi *et al.*, Am. J. Cardiol., **78**: 451-457 (1996). However, because of its potent vasodilating properties, nifedipine may be harmful, especially in patients with outflow obstruction. Disopyramide has been used to relieve symptoms by virtue of its negative inotropic properties. Pollick, N. Engl. J. Med., **307**: 997-999 (1982). In many patients, however, the initial benefits decrease with time. Wigle *et al.*, Circulation, **92**: 1680-1692 (1995). Antihypertensive drug therapy has been reported to have beneficial effects on cardiac hypertrophy associated with elevated blood pressure. Examples of drugs used in antihypertensive therapy, alone or in combination, are calcium antagonists, *e.g.*, nitrendipine; adrenergic receptor blocking agents, *e.g.*, those listed above; angiotensin converting enzyme (ACE) inhibitors such as quinapril, captopril, enalapril, ramipril, benazepril, fosinopril, and lisinopril; diuretics, *e.g.*, chlorothiazide, hydrochlorothiazide, hydroflumethazide, methylchlorothiazide, benzthiazide, dichlorphenamide, acetazolamide, and indapamide; and calcium channel blockers, *e.g.*, diltiazem, nifedipine, verapamil, and nicardipine.

For example, treatment of hypertension with diltiazem and captopril showed a decrease in left ventricular muscle mass, but the Doppler indices of diastolic function did not normalize. Szlachet *et al.*, Am. J. Cardiol., **63**: 198-201 (1989); Shahi *et al.*, Lancet, **336**: 458-461 (1990). These findings were interpreted to indicate that excessive amounts of interstitial collagen may remain after regression of left ventricular hypertrophy. Rossi *et al.*, Am. Heart J., **124**: 700-709 (1992). Rossi *et al.*, *supra*, investigated the effect of captopril on the prevention and

regression of myocardial cell hypertrophy and interstitial fibrosis in pressure overload cardiac hypertrophy, in experimental rats.

Agents that increase cardiac contractility directly (ionotropic agents) were initially thought to benefit patients with heart failure because they improved cardiac output in the short term. However, all positive inotropic agents except digoxigenin have been found to result in increased long-term mortality, in spite of short-term improvements in cardiac performance. Massie, Curr. Op. in Cardiology, 12: 209-217 (1997); Reddy *et al.*, Curr. Opin. Cardiol., 12: 233-241 (1997). Beta-adrenergic receptor blockers have recently been advocated for use in heart failure. Evidence from clinical trials suggests that improvements in cardiac function can be achieved without increased mortality, though documented improvements of patient survival have not yet been demonstrated. See also, U.S. Pat. Nos. 5,935,924, 5,624,806; 5,661,122; and 5,610,134 and WO 95/28173 regarding the use of cardiotropin-1 or antagonists thereof, or growth hormone and/or insulin-like growth factor-I in the treatment of CHF. Another treatment modality is heart transplantation, but this is limited by the availability of donor hearts.

Endothelin is a vasoconstricting peptide comprising 21 amino acids, isolated from swine arterial endothelial culture supernatant and structurally determined. Yanagisawa *et al.*, Nature, 332: 411-415 (1988). Endothelin was later found to exhibit various actions, and endothelin antibodies as endothelin antagonists have proven effective in the treatment of myocardial infarction, renal failure, and other diseases. Since endothelin is present in live bodies and exhibits vasoconstricting action, it is expected to be an endogenous factor involved in the regulation of the circulatory system, and may be associated with hypertension, cardiovascular diseases such as myocardial infarction, and renal diseases such as acute renal failure. Endothelin antagonists are described, for example, in U.S. Pat. No. 5,773,414; JP Pat. Publ. 3130299/1991, EP 457,195; EP 460,679; and EP 552,489. A new endothelin B receptor for identifying endothelin receptor antagonists is described in U.S. Pat. No. 5,773,223.

Current therapy for heart failure is primarily directed to using angiotensin-converting enzyme (ACE) inhibitors, such as captopril, and diuretics. These drugs improve hemodynamic profile and exercise tolerance and reduce the incidence of morbidity and mortality in patients with CHF. Kramer *et al.*, Circulation, 67(4): 807-816 (1983); Captopril Multicenter Research Group, J.A.C.C., 2(4): 755-763 (1983); The CONSENSUS Trial Study Group, N. Engl. J. Med., 316(23): 1429-1435 (1987); The SOLVD Investigators, N. Engl. J. Med., 325(5): 293-302 (1991). Further, they are useful in treating hypertension, left ventricular dysfunction, atherosclerotic vascular disease, and diabetic nephropathy. Brown and Vaughan, *supra*. However, despite proven efficacy, response to ACE inhibitors has been limited. For example, while prolonging survival in the setting of heart failure, ACE inhibitors appear to slow the progression towards end-stage heart failure, and substantial numbers of patients on ACE inhibitors have functional class III heart failure.

Moreover, improvement of functional capacity and exercise time is only small and mortality, although reduced, continues to be high. The CONSENSUS Trial Study Group, N. Engl. J. Med., 316(23): 1429-1435 (1987); The SOLVD Investigators, N. Engl. J. Med., 325(5): 293-302 (1991); Cohn *et al.*, N. Engl. J. Med., 325(5): 303-310 (1991); The Captopril-Digoxin Multicenter Research Group, JAMA, 259(4): 539-544 (1988). Hence, ACE inhibitors consistently appear unable to relieve symptoms in more than 60% of heart failure patients and reduce mortality of heart failure only by approximately 15-20%. For further adverse effects, see Brown and Vaughan,

supra.

An alternative to ACE inhibitors is represented by specific AT1 receptor antagonists. Clinical studies are planned to compare the efficacy of these two modalities in the treatment of cardiovascular and renal disease. However, animal model data suggests that the ACE/Ang II pathway, while clearly involved in cardiac hypertrophy, is not the only, or even the primary pathway active in this role. Mouse genetic "knockout" models have been made to test individual components of the pathway. In one such model, the primary cardiac receptor for Ang II, AT sub 1A, has been genetically deleted; these mice do not develop hypertrophy when Ang II is given experimentally (confirming the basic success of the model in eliminating hypertrophy secondary to Ang II). However, when the aorta is constricted in these animals (a model of hypertensive cardiac stress), the hearts still become hypertrophic. This suggests that alternative signaling pathways, not depending on this receptor (AT sub 1A), are activated in hypertension. ACE inhibitors would presumably not be able to inhibit these pathways. See, Harada *et al.*, Circulation, 97: 1952-1959 (1998). See also, Homcy, Circulation, 97: 1890-1892 (1998) regarding the enigma associated with the process and mechanism of cardiac hypertrophy.

About 750,000 patients suffer from acute myocardial infarction (AMI) annually, and approximately one-fourth of all deaths in the United States are due to AMI. In recent years, thrombolytic agents, *e.g.*, streptokinase, urokinase, and in particular tissue plasminogen activator (t-PA) have significantly increased the survival of patients who suffered myocardial infarction. When administered as a continuous intravenous infusion over 1.5 to 4 hours, t-PA produces coronary patency at 90 minutes in 69% to 90% of the treated patients. Topol *et al.*, Am. J. Cardiol., 61: 723-728 (1988); Neuhaus *et al.*, J. Am. Coll. Cardiol., 12: 581-587 (1988); Neuhaus *et al.*, J. Am. Coll. Cardiol., 14: 1566-1569 (1989). The highest patency rates have been reported with high dose or accelerated dosing regimens. Topol, J. Am. Coll. Cardiol., 15: 922-924 (1990). t-PA may also be administered as a single bolus, although due to its relatively short half-life, it is better suited for infusion therapy. Tebbe *et al.*, Am. J. Cardiol., 64: 448-453 (1989). A t-PA variant, specifically designed to have longer half-life and very high fibrin specificity, TNK t-PA (a T103N, N117Q, KHR(296-299)AAAA t-PA variant, Keyt *et al.*, Proc. Natl. Acad. Sci. USA, 91: 3670-3674 (1994)) is particularly suitable for bolus administration. However, despite all these advances, the long-term prognosis of patient survival depends greatly on the post-infarction monitoring and treatment of the patients, which should include monitoring and treatment of cardiac hypertrophy.

2.3. Growth Factors

Various naturally occurring polypeptides reportedly induce the proliferation of endothelial cells. Among those polypeptides are the basic and acidic fibroblast growth factors (FGF) (Burgess and Maciag, Annual Rev. Biochem., 58: 575 (1989)), platelet-derived endothelial cell growth factor (PD-ECGF) (Ishikawa *et al.*, Nature, 338: 557 (1989)), and vascular endothelial growth factor (VEGF). Leung *et al.*, Science, 246: 1306 (1989); Ferrara and Henzel, Biochem. Biophys. Res. Commun., 161: 851 (1989); Tischer *et al.*, Biochem. Biophys. Res. Commun., 165: 1198 (1989); EP 471,754B granted July 31, 1996.

Media conditioned by cells transfected with the human VEGF (hVEGF) cDNA promoted the proliferation of capillary endothelial cells, whereas control cells did not. Leung *et al.*, Science, 246: 1306 (1989). Several

additional cDNAs were identified in human cDNA libraries that encode 121-, 189-, and 206-amino acid isoforms of hVEGF (also collectively referred to as hVEGF-related proteins). The 121-amino acid protein differs from hVEGF by virtue of the deletion of the 44 amino acids between residues 116 and 159 in hVEGF. The 189-amino acid protein differs from hVEGF by virtue of the insertion of 24 amino acids at residue 116 in hVEGF, and apparently is identical to human vascular permeability factor (hVPF). The 206-amino acid protein differs from hVEGF by virtue of an insertion of 41 amino acids at residue 116 in hVEGF. Houck *et al.*, Mol. Endocrin., **5**: 1806 (1991); Ferrara *et al.*, J. Cell. Biochem., **47**: 211 (1991); Ferrara *et al.*, Endocrine Reviews, **13**: 18 (1992); Keck *et al.*, Science, **246**: 1309 (1989); Connolly *et al.*, J. Biol. Chem., **264**: 20017 (1989); EP 370,989 published May 30, 1990.

It is now well established that angiogenesis, which involves the formation of new blood vessels from preexisting endothelium, is implicated in the pathogenesis of a variety of disorders. These include solid tumors and metastasis, atherosclerosis, retrolental fibroplasia, hemangiomas, chronic inflammation, intraocular neovascular syndromes such as proliferative retinopathies, *e.g.*, diabetic retinopathy, age-related macular degeneration (AMD), neovascular glaucoma, immune rejection of transplanted corneal tissue and other tissues, rheumatoid arthritis, and psoriasis. Folkman *et al.*, J. Biol. Chem., **267**: 10931-10934 (1992); Klagsbrun *et al.*, Annu. Rev. Physiol., **53**: 217-239 (1991); and Garner A., "Vascular diseases", In: Pathobiology of Ocular Disease. A Dynamic Approach, Garner A., Klintworth GK, eds., 2nd Edition (Marcel Dekker, NY, 1994), pp 1625-1710.

In the case of tumor growth, angiogenesis appears to be crucial for the transition from hyperplasia to neoplasia, and for providing nourishment for the growth and metastasis of the tumor. Folkman *et al.*, Nature, **339**: 58 (1989). The neovascularization allows the tumor cells to acquire a growth advantage and proliferative autonomy compared to the normal cells. A tumor usually begins as a single aberrant cell which can proliferate only to a size of a few cubic millimeters due to the distance from available capillary beds, and it can stay 'dormant' without further growth and dissemination for a long period of time. Some tumor cells then switch to the angiogenic phenotype to activate endothelial cells, which proliferate and mature into new capillary blood vessels. These newly formed blood vessels not only allow for continued growth of the primary tumor, but also for the dissemination and recolonization of metastatic tumor cells. Accordingly, a correlation has been observed between density of microvessels in tumor sections and patient survival in breast cancer as well as in several other tumors. Weidner *et al.*, N. Engl. J. Med., **324**: 1-6 (1991); Horak *et al.*, Lancet, **340**: 1120-1124 (1992); Macchiarini *et al.*, Lancet, **340**: 145-146 (1992). The precise mechanisms that control the angiogenic switch is not well understood, but it is believed that neovascularization of tumor mass results from the net balance of a multitude of angiogenesis stimulators and inhibitors (Folkman, 1995, Nat Med **1**(1):27-31).

The search for positive regulators of angiogenesis has yielded many candidates, including aFGF, bFGF, TGF- α , TGF- β , HGF, TNF- α , angiogenin, IL-8, etc. Folkman *et al.*, J.B.C., *supra*, and Klagsbrun *et al.*, *supra*. The negative regulators so far identified include thrombospondin (Good *et al.*, Proc. Natl. Acad. Sci. USA., **87**: 6624-6628 (1990)), the 16-kilodalton N-terminal fragment of prolactin (Clapp *et al.*, Endocrinology, **133**: 1292-1299 (1993)), angiostatin (O'Reilly *et al.*, Cell, **79**: 315-328 (1994)), and endostatin. O'Reilly *et al.*, Cell, **88**: 277-285 (1996).

Work done over the last several years has established the key role of VEGF, not only in stimulating vascular endothelial cell proliferation, but also in inducing vascular permeability and angiogenesis. Ferrara *et al.*, Endocr. Rev., **18**: 4-25 (1997). The finding that the loss of even a single VEGF allele results in embryonic lethality points to an irreplaceable role played by this factor in the development and differentiation of the vascular system.

Furthermore, VEGF has been shown to be a key mediator of neovascularization associated with tumors and intraocular disorders. Ferrara *et al.*, Endocr. Rev., *supra*. The VEGF mRNA is overexpressed by the majority of human tumors examined. Berkman *et al.*, J. Clin. Invest., **91**: 153-159 (1993); Brown *et al.*, Human Pathol., **26**: 86-91 (1995); Brown *et al.*, Cancer Res., **53**: 4727-4735 (1993); Mattern *et al.*, Brit. J. Cancer, **73**: 931-934 (1996); Dvorak *et al.*, Am. J. Pathol., **146**: 1029-1039 (1995).

Also, the concentration levels of VEGF in eye fluids are highly correlated to the presence of active proliferation of blood vessels in patients with diabetic and other ischemia-related retinopathies. Aiello *et al.*, N. Engl. J. Med., **331**: 1480-1487 (1994). Furthermore, recent studies have demonstrated the localization of VEGF in choroidal neovascular membranes in patients affected by AMD. Lopez *et al.*, Invest. Ophthalmol. Vis. Sci., **37**: 855-868 (1996).

Anti-VEGF neutralizing antibodies suppress the growth of a variety of human tumor cell lines in nude mice (Kim *et al.*, Nature, **362**: 841-844 (1993); Warren *et al.*, J. Clin. Invest., **95**: 1789-1797 (1995); Borgström *et al.*, Cancer Res., **56**: 4032-4039 (1996); Melnyk *et al.*, Cancer Res., **56**: 921-924 (1996)) and also inhibit intraocular angiogenesis in models of ischemic retinal disorders. Adamis *et al.*, Arch. Ophthalmol., **114**: 66-71 (1996). Therefore, anti-VEGF monoclonal antibodies or other inhibitors of VEGF action are promising candidates for the treatment of solid tumors and various intraocular neovascular disorders. Such antibodies are described, for example, in EP 817,648 published January 14, 1998 and in WO98/45331 and WO98/45332 both published October 15, 1998.

There exist several other growth factors and mitogens, including transforming oncogenes, that are capable of rapidly inducing a complex set of genes to be expressed by certain cells. Lau and Nathans, Molecular Aspects of Cellular Regulation, **6**: 165-202 (1991). These genes, which have been named immediate-early- or early-response genes, are transcriptionally activated within minutes after contact with a growth factor or mitogen, independent of *de novo* protein synthesis. A group of these intermediate-early genes encodes secreted, extracellular proteins that are needed for coordination of complex biological processes such as differentiation and proliferation, regeneration, and wound healing. Ryseck *et al.*, Cell Growth Differ., **2**: 235-233 (1991).

Highly-related proteins that belong to this group include *cef10* (Simmons *et al.*, Proc. Natl. Acad. Sci. USA, **86**: 1178-1182 (1989)), *cyr61*, which is rapidly activated by serum- or platelet-derived growth factor (PDGF) (O'Brien *et al.*, Mol. Cell Biol., **10**: 3569-3577 (1990)), human connective tissue growth factor (CTGF) (Bradham *et al.*, J. Cell. Biol., **114**: 1285-1294 (1991)), which is secreted by human vascular endothelial cells in high levels after activation with transforming growth factor beta (TGF- β), exhibits PDGF-like biological and immunological activities, and competes with PDGF for a particular cell surface receptor, *fisp-12* (Ryseck *et al.*, Cell Growth Differ., **2**: 235-233 (1991)), human vascular IBP-like growth factor (VIGF) (WO 96/17931), and *nov*, normally arrested in adult kidney cells, which was found to be overexpressed in myeloblastosis-associated-virus-type-1-induced nephroblastomas. Joliet *et al.*, Mol. Cell Biol., **12**: 10-21 (1992).

The expression of these immediate-early genes acts as "third messengers" in the cascade of events triggered by growth factors. It is also thought that they are needed to integrate and coordinate complex biological processes, such as differentiation and wound healing in which cell proliferation is a common event.

As additional mitogens, insulin-like growth factor binding proteins (IGFBPs) have been shown, in complex with insulin-like growth factor (IGF), to stimulate increased binding of IGF to fibroblast and smooth muscle cell surface receptors. Clemmons *et al.*, *J. Clin. Invest.*, **77**: 1548 (1986). Inhibitory effects of IGFBP on various IGF actions *in vitro* include stimulation of glucose transport by adipocytes, sulfate incorporation by chondrocytes, and thymidine incorporation in fibroblast. Zapf *et al.*, *J. Clin. Invest.*, **63**: 1077 (1979). In addition, inhibitory effects of IGFBPs on growth factor-mediated mitogen activity in normal cells have been shown.

2.4. Need for Further Treatments

In view of the role of vascular endothelial cell growth and angiogenesis in many diseases and disorders, it is desirable to have a means of reducing or inhibiting one or more of the biological effects causing these processes. It is also desirable to have a means of assaying for the presence of pathogenic polypeptides in normal and diseased conditions, and especially cancer. Further, in a specific aspect, as there is no generally applicable therapy for the treatment of cardiac hypertrophy, the identification of factors that can prevent or reduce cardiac myocyte hypertrophy is of primary importance in the development of new therapeutic strategies to inhibit pathophysiological cardiac growth. While there are several treatment modalities for various cardiovascular and oncologic disorders, there is still a need for additional therapeutic approaches.

3. Summary of the Invention

The present invention provides compositions and methods for modulating (*e.g.*, promoting or inhibiting) angiogenesis and/or cardiovascularization in mammals. The present invention is based on the identification of compounds (*i.e.*, proteins) that test positive in various cardiovascular assays that test modulation (*e.g.*, promotion or inhibition) of certain biological activities. Accordingly, the compounds are believed to be useful drugs and/or drug components for the diagnosis and/or treatment (including prevention and amelioration) of disorders where such effects are desired, such as the promotion or inhibition of angiogenesis, inhibition or stimulation of vascular endothelial cell growth, stimulation of growth or proliferation of vascular endothelial cells, inhibition of tumor growth, inhibition of angiogenesis-dependent tissue growth, stimulation of angiogenesis-dependent tissue growth, inhibition of cardiac hypertrophy and stimulation of cardiac hypertrophy, *e.g.*, for the treatment of congestive heart failure. In addition, the compositions and methods of the invention provide for the diagnostic monitoring of patients undergoing clinical evaluation for the treatment of angiogenesis-related disorders, for monitoring the efficacy of compounds in clinical trials and for identifying subjects who may be predisposed to such angiogenic-related disorders.

In one embodiment, the present invention provides a composition comprising a PRO polypeptide, an agonist or antagonist thereof, or an anti-PRO antibody in admixture with a pharmaceutically acceptable carrier. In one aspect, the composition comprises a therapeutically effective amount of the polypeptide, agonist, antagonist

or antibody. In another aspect, the composition comprises a further active ingredient, namely, a cardiovascular, endothelial or angiogenic agent or an angiostatic agent, preferably an angiogenic or angiostatic agent. Preferably, the composition is sterile. The PRO polypeptide, agonist, antagonist or antibody may be administered in the form of a liquid pharmaceutical formulation, which may be preserved to achieve extended storage stability. Preserved liquid pharmaceutical formulations might contain multiple doses of PRO polypeptide, agonist, antagonist or antibody, and might, therefore, be suitable for repeated use. In a preferred embodiment, where the composition comprises an antibody, the antibody is a monoclonal antibody, an antibody fragment, a humanized antibody, or a single-chain antibody.

In a further embodiment, the present invention provides a method for preparing such a composition useful for the treatment of a cardiovascular, endothelial or angiogenic disorder comprising admixing a therapeutically effective amount of a PRO polypeptide, agonist, antagonist or antibody with a pharmaceutically acceptable carrier.

In a still further aspect, the present invention provides an article of manufacture comprising:

- (a) a composition of matter comprising a PRO polypeptide or agonist or antagonist thereof;
- (b) a container containing said composition; and

(c) a label affixed to said container, or a package insert included in said container referring to the use of said PRO polypeptide or agonist or antagonist thereof in the treatment of a cardiovascular, endothelial or angiogenic disorder, wherein the agonist or antagonist may be an antibody which binds to the PRO polypeptide. The composition may comprise a therapeutically effective amount of the PRO polypeptide or the agonist or antagonist thereof.

In another embodiment, the present invention provides a method for identifying an agonist of a PRO polypeptide comprising:

- (a) contacting cells and a test compound to be screened under conditions suitable for the induction of a cellular response normally induced by a PRO polypeptide; and
- (b) determining the induction of said cellular response to determine if the test compound is an effective agonist, wherein the induction of said cellular response is indicative of said test compound being an effective agonist.

In another embodiment, the present invention provides a method for identifying an agonist of a PRO polypeptide comprising:

- (a) contacting cells and a test compound to be screened under conditions suitable for the stimulation of cell proliferation by a PRO polypeptide; and
- (b) measuring the proliferation of said cells to determine if the test compound is an effective agonist, wherein the stimulation of cell proliferation is indicative of said test compound being an effective agonist.

In another embodiment, the invention provides a method for identifying a compound that inhibits the activity of a PRO polypeptide comprising contacting a test compound with a PRO polypeptide under conditions and for a time sufficient to allow the test compound and polypeptide to interact and determining whether the activity of the PRO polypeptide is inhibited. In a specific preferred aspect, either the test compound or the PRO polypeptide is immobilized on a solid support. In another preferred aspect, the non-immobilized component carries a detectable

label. In a preferred aspect, this method comprises the steps of:

(a) contacting cells and a test compound to be screened in the presence of a PRO polypeptide under conditions suitable for the induction of a cellular response normally induced by a PRO polypeptide; and

5 (b) determining the induction of said cellular response to determine if the test compound is an effective antagonist.

In another preferred aspect, this process comprises the steps of:

(a) contacting cells and a test compound to be screened in the presence of a PRO polypeptide under conditions suitable for the stimulation of cell proliferation by a PRO polypeptide; and

(b) measuring the proliferation of the cells to determine if the test compound is an effective antagonist.

10 In another embodiment, the invention provides a method for identifying a compound that inhibits the expression of a PRO polypeptide in cells that normally expresses the polypeptide, wherein the method comprises contacting the cells with a test compound and determining whether the expression of the PRO polypeptide is inhibited. In a preferred aspect, this method comprises the steps of:

15 (a) contacting cells and a test compound to be screened under conditions suitable for allowing expression of the PRO polypeptide; and

(b) determining the inhibition of expression of said polypeptide.

In a still further embodiment, the invention provides a compound that inhibits the expression of a PRO polypeptide, such as a compound that is identified by the methods set forth above.

20 Another aspect of the present invention is directed to an agonist or an antagonist of a PRO polypeptide which may optionally be identified by the methods described above.

One type of antagonist of a PRO polypeptide that inhibits one or more of the functions or activities of the PRO polypeptide is an antibody. Hence, in another aspect, the invention provides an isolated antibody that binds a PRO polypeptide. In a preferred aspect, the antibody is a monoclonal antibody, which preferably has non-human complementarity-determining-region (CDR) residues and human framework-region (FR) residues. The antibody
25 may be labeled and may be immobilized on a solid support. In a further aspect, the antibody is an antibody fragment, a single-chain antibody, or a humanized antibody. Preferably, the antibody specifically binds to the polypeptide.

30 In a still further aspect, the present invention provides a method for diagnosing a disease or susceptibility to a disease which is related to a mutation in a PRO polypeptide-encoding nucleic acid sequence comprising determining the presence or absence of said mutation in the PRO polypeptide nucleic acid sequence, wherein the presence or absence of said mutation is indicative of the presence of said disease or susceptibility to said disease.

35 In a still further aspect, the invention provides a method of diagnosing a cardiovascular, endothelial or angiogenic disorder in a mammal which comprises analyzing the level of expression of a gene encoding a PRO polypeptide (a) in a test sample of tissue cells obtained from said mammal, and (b) in a control sample of known normal tissue cells of the same cell type, wherein a higher or lower expression level in the test sample as compared to the control sample is indicative of the presence of a cardiovascular, endothelial or angiogenic disorder in said mammal. The expression of a gene encoding a PRO polypeptide may optionally be accomplished by measuring

the level of mRNA or the polypeptide in the test sample as compared to the control sample.

5 In a still further aspect, the present invention provides a method of diagnosing a cardiovascular, endothelial or angiogenic disorder in a mammal which comprises detecting the presence or absence of a PRO polypeptide in a test sample of tissue cells obtained from said mammal, wherein the presence or absence of said PRO polypeptide in said test sample is indicative of the presence of a cardiovascular, endothelial or angiogenic disorder in said mammal.

10 In a still further embodiment, the invention provides a method of diagnosing a cardiovascular, endothelial or angiogenic disorder in a mammal comprising (a) contacting an anti-PRO antibody with a test sample of tissue cells obtained from the mammal, and (b) detecting the formation of a complex between the antibody and the PRO polypeptide in the test sample, wherein the formation of said complex is indicative of the presence of a cardiovascular, endothelial or angiogenic disorder in the mammal. The detection may be qualitative or quantitative, and may be performed in comparison with monitoring the complex formation in a control sample of known normal tissue cells of the same cell type. A larger or smaller quantity of complexes formed in the test sample indicates the presence of a cardiovascular, endothelial or angiogenic dysfunction in the mammal from which the test tissue cells were obtained. The antibody preferably carries a detectable label. Complex formation can be monitored, for example, by light microscopy, flow cytometry, fluorimetry, or other techniques known in the art. The test sample is usually obtained from an individual suspected to have a cardiovascular, endothelial or angiogenic disorder.

15 In another embodiment, the invention provides a method for determining the presence of a PRO polypeptide in a sample comprising exposing a sample suspected of containing the PRO polypeptide to an anti-PRO antibody and determining binding of said antibody to a component of said sample. In a specific aspect, the sample comprises a cell suspected of containing the PRO polypeptide and the antibody binds to the cell. The antibody is preferably detectably labeled and/or bound to a solid support.

20 In further aspects, the invention provides a cardiovascular, endothelial or angiogenic disorder diagnostic kit comprising an anti-PRO antibody and a carrier in suitable packaging. Preferably, such kit further comprises instructions for using said antibody to detect the presence of the PRO polypeptide. Preferably, the carrier is a buffer, for example. Preferably, the cardiovascular, endothelial or angiogenic disorder is cancer.

25 In yet another embodiment, the present invention provides a method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal comprising administering to the mammal an effective amount of a PRO polypeptide. Preferably, the disorder is cardiac hypertrophy, trauma such as wounds or burns, or a type of cancer. In a further aspect, the mammal is further exposed to angioplasty or a drug that treats cardiovascular, endothelial or angiogenic disorders such as ACE inhibitors or chemotherapeutic agents if the cardiovascular, endothelial or angiogenic disorder is a type of cancer. Preferably, the mammal is human, preferably one who is at risk of developing cardiac hypertrophy and more preferably has suffered myocardial infarction.

30 In another preferred aspect, the cardiac hypertrophy is characterized by the presence of an elevated level of $\text{PGF}_{2\alpha}$. Alternatively, the cardiac hypertrophy may be induced by myocardial infarction, wherein preferably the administration of the PRO polypeptide is initiated within 48 hours, more preferably within 24 hours, following myocardial infarction.

In another preferred embodiment, the cardiovascular, endothelial or angiogenic disorder is cardiac hypertrophy and said PRO polypeptide is administered together with a cardiovascular, endothelial or angiogenic agent. The preferred cardiovascular, endothelial or angiogenic agent for this purpose is selected from the group consisting of an antihypertensive drug, an ACE inhibitor, an endothelin receptor antagonist and a thrombolytic agent. If a thrombolytic agent is administered, preferably the PRO polypeptide is administered following administration of such agent. More preferably, the thrombolytic agent is recombinant human tissue plasminogen activator.

In another preferred aspect, the cardiovascular, endothelial or angiogenic disorder is cardiac hypertrophy and the PRO polypeptide is administered following primary angioplasty for the treatment of acute myocardial infarction, preferably wherein the mammal is further exposed to angioplasty or a cardiovascular, endothelial, or angiogenic agent.

In another preferred embodiment, the cardiovascular, endothelial or angiogenic disorder is a cancer and the PRO polypeptide is administered in combination with a chemotherapeutic agent, a growth inhibitory agent or a cytotoxic agent.

In a further embodiment, the invention provides a method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal comprising administering to the mammal an effective amount of a PRO polypeptide agonist, antagonist or anti-PRO antibody. Preferably, the cardiovascular, endothelial or angiogenic disorder is cardiac hypertrophy, trauma, a cancer, or age-related macular degeneration. Also preferred is where the mammal is human, and where an effective amount of an angiogenic or angiostatic agent is administered in conjunction with the agonist, antagonist or anti-PRO antibody.

In still further embodiments, the invention provides a method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal that suffers therefrom comprising administering to the mammal a nucleic acid molecule that codes for either (a) a PRO polypeptide, (b) an agonist of a PRO polypeptide or (c) an antagonist of a PRO polypeptide, wherein said agonist or antagonist may be an anti-PRO antibody. In a preferred embodiment, the mammal is human. In another preferred embodiment, the gene is administered via *ex vivo* gene therapy. In a further preferred embodiment, the gene is comprised within a vector, more preferably an adenoviral, adeno-associated viral, lentiviral, or retroviral vector.

In yet another aspect, the invention provides a recombinant retroviral particle comprising a retroviral vector consisting essentially of a promoter, nucleic acid encoding (a) a PRO polypeptide, (b) an agonist polypeptide of a PRO polypeptide, or (c) an antagonist polypeptide of a PRO polypeptide, and a signal sequence for cellular secretion of the polypeptide, wherein the retroviral vector is in association with retroviral structural proteins. Preferably, the signal sequence is from a mammal, such as from a native PRO polypeptide.

In a still further embodiment, the invention supplies an *ex vivo* producer cell comprising a nucleic acid construct that expresses retroviral structural proteins and also comprises a retroviral vector consisting essentially of a promoter, nucleic acid encoding (a) a PRO polypeptide, (b) an agonist polypeptide of a PRO polypeptide or (c) an antagonist polypeptide of a PRO polypeptide, and a signal sequence for cellular secretion of the polypeptide, wherein said producer cell packages the retroviral vector in association with the structural proteins to produce

recombinant retroviral particles.

In yet another embodiment, the invention provides a method for inhibiting endothelial cell growth in a mammal comprising administering to the mammal (a) a PRO polypeptide, (b) an agonist of a PRO polypeptide, or (c) an antagonist of a PRO polypeptide, wherein endothelial cell growth in said mammal is inhibited, and wherein said agonist or antagonist may be an anti-PRO antibody. Preferably, the mammal is human and the endothelial cell growth is associated with a tumor or a retinal disorder.

In yet another embodiment, the invention provides a method for stimulating endothelial cell growth in a mammal comprising administering to the mammal (a) a PRO polypeptide, (b) an agonist of a PRO polypeptide, or (c) an antagonist of a PRO polypeptide, wherein endothelial cell growth in said mammal is stimulated, and wherein said agonist or antagonist may be an anti-PRO antibody. Preferably, the mammal is human.

In yet another embodiment, the invention provides a method for inhibiting cardiac hypertrophy in a mammal comprising administering to the mammal (a) a PRO polypeptide, (b) an agonist of a PRO polypeptide, or (c) an antagonist of a PRO polypeptide, wherein cardiac hypertrophy in said mammal is inhibited, and wherein said agonist or antagonist may be an anti-PRO antibody. Preferably, the mammal is human and the cardiac hypertrophy has been induced by myocardial infarction.

In yet another embodiment, the invention provides a method for stimulating cardiac hypertrophy in a mammal comprising administering to the mammal (a) a PRO polypeptide, (b) an agonist of a PRO polypeptide, or (c) an antagonist of a PRO polypeptide, wherein cardiac hypertrophy in said mammal is stimulated, and wherein said agonist or antagonist may be an anti-PRO antibody. Preferably, the mammal is human who suffers from congestive heart failure.

In yet another embodiment, the invention provides a method for inhibiting angiogenesis induced by a PRO polypeptide in a mammal comprising administering a therapeutically effective amount of an anti-PRO antibody to the mammal. Preferably, the mammal is a human, and more preferably the mammal has a tumor or a retinal disorder.

In yet another embodiment, the invention provides a method for stimulating angiogenesis induced by a PRO polypeptide in a mammal comprising administering a therapeutically effective amount of a PRO polypeptide to the mammal. Preferably, the mammal is a human, and more preferably angiogenesis would promote tissue regeneration or wound healing.

In yet another embodiment, the invention provides a method for modulating (e.g., inhibiting or stimulating) endothelial cell growth in a mammal comprising administering to the mammal a PRO21, PRO181, PRO205, PRO214, PRO221, PRO229, PRO231, PRO238, PRO241, PRO247, PRO256, PRO258, PRO263, PRO265, PRO295, PRO321, PRO322, PRO337, PRO363, PRO365, PRO444, PRO533, PRO697, PRO720, PRO725, PRO771, PRO788, PRO791, PRO819, PRO827, PRO828, PRO836, PRO846, PRO865, PRO1005, PRO1006, PRO1007, PRO1025, PRO1029, PRO1054, PRO1071, PRO1075, PRO1079, PRO1080, PRO1114, PRO1131, PRO1155, PRO1160, PRO1184, PRO1186, PRO1190, PRO1192, PRO1195, PRO1244, PRO1272, PRO1273, PRO1274, PRO1279, PRO1283, PRO1286, PRO1306, PRO1309, PRO1325, PRO1329, PRO1347, PRO1356, PRO1376, PRO1382, PRO1411, PRO1412, PRO1419, PRO1474, PRO1477, PRO1488, PRO1508, PRO1550,

PRO1556, PRO1760, PRO1782, PRO1787, PRO1801, PRO1868, PRO1887, PRO1890, PRO3438, PRO3444, PRO4302, PRO4324, PRO4333, PRO4341, PRO4342, PRO4353, PRO4354, PRO4356, PRO4371, PRO4405, PRO4408, PRO4422, PRO4425, PRO4499, PRO5723, PRO5725, PRO5737, PRO5776, PRO6006, PRO6029, PRO6071, PRO7436, PRO9771, PRO9821, PRO9873, PRO10008, PRO10096, PRO19670, PRO20040, PRO20044, PRO21055, PRO21384 or PRO28631 polypeptide, agonist or antagonist thereof, wherein endothelial cell growth in said mammal is modulated.

In yet another embodiment, the invention provides a method for modulating (*e.g.*, inhibiting or stimulating) smooth muscle cell growth in a mammal comprising administering to the mammal a PRO162, PRO181, PRO182, PRO195, PRO204, PRO221, PRO230, PRO256, PRO258, PRO533, PRO697, PRO725, PRO738, PRO826, PRO836, PRO840, PRO846, PRO865, PRO982, PRO1025, PRO1029, PRO1071, PRO1080, PRO1083, PRO1134, PRO1160, PRO1182, PRO1184, PRO1186, PRO1192, PRO1265, PRO1274, PRO1279, PRO1283, PRO1306, PRO1308, PRO1309, PRO1325, PRO1337, PRO1338, PRO1343, PRO1376, PRO1387, PRO1411, PRO1412, PRO1415, PRO1434, PRO1474, PRO1488, PRO1550, PRO1556, PRO1567, PRO1600, PRO1754, PRO1758, PRO1760, PRO1787, PRO1865, PRO1868, PRO1917, PRO1928, PRO3438, PRO3562, PRO4302, PRO4333, PRO4345, PRO4353, PRO4354, PRO4405, PRO4408, PRO4430, PRO4503, PRO5725, PRO6714, PRO9771, PRO9820, PRO9940, PRO10096, PRO21055, PRO21184 or PRO21366 polypeptide, agonist or antagonist thereof, wherein endothelial cell growth in said mammal is modulated.

In yet another embodiment, the invention provides a method for modulating (*e.g.*, inducing or reducing) cardiac hypertrophy in a mammal comprising administering to the mammal a PRO21 polypeptide, agonist or antagonist thereof, wherein cardiac hypertrophy in said mammal is modulated.

In yet another embodiment, the invention provides a method for modulating (*e.g.*, inducing or reducing) endothelial cell apoptosis in a mammal comprising administering to the mammal a PRO4302 polypeptide, agonist or antagonist thereof, wherein cardiac hypertrophy in said mammal is modulated.

In yet another embodiment, the invention provides a method for modulating (*e.g.*, stimulating or inhibiting) angiogenesis in a mammal comprising administering a therapeutically effective amount of a PRO1376 or PRO1449 polypeptide, agonist or antagonist thereof to the mammal, wherein said angiogenesis is modulated.

In yet another embodiment, the invention provides a method for modulating (*e.g.*, inducing or reducing) angiogenesis by modulating (*e.g.*, inducing or reducing) endothelial cell tube formation in a mammal comprising administering to the mammal a PRO178, PRO195, PRO228, PRO301, PRO302, PRO532, PRO724, PRO730, PRO734, PRO793, PRO871, PRO938, PRO1012, PRO1120, PRO1139, PRO1198, PRO1287, PRO1361, PRO1864, PRO1873, PRO2010, PRO3579, PRO4313, PRO4527, PRO4538, PRO4553, PRO4995, PRO5730, PRO6008, PRO7223, PRO7248 or PRO7261 polypeptide, agonist or antagonist thereof, wherein endothelial cell tube formation in said mammal is modulated.

In other embodiments of the present invention, the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence that encodes a PRO polypeptide.

In one aspect, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% or 98%

nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule encoding a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein, an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of the full-length amino acid sequence as disclosed herein, or (b) the complement of the DNA molecule of (a).

In other aspects, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% or 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule comprising the coding sequence of a full-length PRO polypeptide cDNA as disclosed herein, the coding sequence of a PRO polypeptide lacking the signal peptide as disclosed herein, the coding sequence of an extracellular domain of a transmembrane PRO polypeptide, with or without the signal peptide, as disclosed herein or the coding sequence of any other specifically defined fragment of the full-length amino acid sequence as disclosed herein, or (b) the complement of the DNA molecule of (a).

In a further aspect, the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence having at least about 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% or 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule that encodes the same mature polypeptide encoded by any of the human protein cDNAs deposited with the ATCC as disclosed herein, or (b) the complement of the DNA molecule of (a).

Another aspect of the present invention provides an isolated nucleic acid molecule comprising a nucleotide sequence encoding a PRO polypeptide which is either transmembrane domain-deleted or transmembrane domain-inactivated, or is complementary to such encoding nucleotide sequence, wherein the transmembrane domain(s) of such polypeptide are disclosed herein. Therefore, soluble extracellular domains of the herein described PRO polypeptides are contemplated.

Another embodiment is directed to fragments of a PRO polypeptide coding sequence, or the complement thereof, that may find use as, for example, hybridization probes, for encoding fragments of a PRO polypeptide that may optionally encode a polypeptide comprising a binding site for an anti-PRO antibody or as antisense oligonucleotide probes. Such nucleic acid fragments are usually at least about 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 250, 300, 350, 400, 450, 500, 600, 700 or 800 nucleotides in length and alternatively at least about 1000 nucleotides in length, wherein in this context the term "about" means the referenced nucleotide sequence length plus or minus 10% of that referenced length. It is noted that novel fragments of a PRO polypeptide-encoding nucleotide sequence may be determined in a routine manner by aligning the PRO polypeptide-encoding nucleotide sequence with other known nucleotide sequences using any of a number of well known sequence alignment programs and determining which PRO polypeptide-encoding nucleotide sequence fragment(s) are novel. All of such PRO polypeptide-encoding nucleotide sequences are contemplated herein. Also contemplated are the PRO polypeptide fragments encoded by these nucleotide molecule fragments, preferably those PRO polypeptide fragments that comprise a binding site for an anti-PRO antibody.

In another embodiment, the invention provides an isolated PRO polypeptide encoded by any of the isolated

nucleic acid sequences hereinabove identified.

5 In a certain aspect, the invention provides an isolated PRO polypeptide comprising an amino acid sequence having at least about 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% or 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein, an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of the full-length amino acid sequence as disclosed herein.

10 In a further aspect, the invention provides an isolated PRO polypeptide comprising an amino acid sequence having at least about 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% or 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to an amino acid sequence encoded by any of the human protein cDNAs deposited with the ATCC as disclosed herein.

15 In a specific aspect, the invention provides an isolated PRO polypeptide without the N-terminal signal sequence and/or the initiating methionine and that is encoded by a nucleotide sequence that encodes such an amino acid sequence as hereinbefore described. Processes for producing the same are also herein described, wherein those processes comprise culturing a host cell comprising a vector which comprises the appropriate encoding nucleic acid molecule under conditions suitable for expression of the PRO polypeptide and recovering the PRO polypeptide from the cell culture.

20 Another aspect of the invention provides an isolated PRO polypeptide which is either transmembrane domain-deleted or transmembrane domain-inactivated. Processes for producing the same are also herein described, wherein those processes comprise culturing a host cell comprising a vector which comprises the appropriate encoding nucleic acid molecule under conditions suitable for expression of the PRO polypeptide and recovering the PRO polypeptide from the cell culture.

25 In yet another embodiment, the invention provides agonists and antagonists of a native PRO polypeptide as defined herein. In a particular embodiment, the agonist or antagonist is an anti-PRO antibody or a small molecule.

30 In a further embodiment, the invention provides a method of identifying agonists or antagonists to a PRO polypeptide which comprise contacting the PRO polypeptide with a candidate molecule and monitoring a biological activity mediated by said PRO polypeptide. Preferably, the PRO polypeptide is a native PRO polypeptide.

In a still further embodiment, the invention provides a composition of matter comprising a PRO polypeptide, or an agonist or antagonist of a PRO polypeptide as herein described, or an anti-PRO antibody, in combination with a carrier. Optionally, the carrier is a pharmaceutically acceptable carrier.

35 Another embodiment of the present invention is directed to the use of a PRO polypeptide, or an agonist or antagonist thereof as hereinbefore described, or an anti-PRO antibody, for the preparation of a medicament useful in the treatment of a condition which is responsive to the PRO polypeptide, an agonist or antagonist thereof or an anti-PRO antibody.

In additional embodiments of the present invention, the invention provides vectors comprising DNA encoding any of the herein described polypeptides. Host cells comprising any such vector are also provided. By way of example, the host cells may be CHO cells, *E. coli*, yeast, or Baculovirus-infected insect cells. A process for producing any of the herein described polypeptides is further provided and comprises culturing host cells under conditions suitable for expression of the desired polypeptide and recovering the desired polypeptide from the cell culture.

In other embodiments, the invention provides chimeric molecules comprising any of the herein described polypeptides fused to a heterologous polypeptide or amino acid sequence. Example of such chimeric molecules comprise any of the herein described polypeptides fused to an epitope tag sequence or a Fc region of an immunoglobulin.

In yet another embodiment, the invention provides an antibody which specifically binds to any of the above or below described polypeptides. Optionally, the antibody is a monoclonal antibody, humanized antibody, antibody fragment or single-chain antibody.

In yet other embodiments, the invention provides oligonucleotide probes useful for isolating genomic and cDNA nucleotide sequences or as antisense probes, wherein those probes may be derived from any of the above or below described nucleotide sequences.

4. Brief Description of the Drawings

Figure 1 shows a nucleotide sequence (SEQ ID NO:1) of a native sequence PRO181 cDNA, wherein SEQ ID NO:1 is a clone designated herein as "DNA23330-1390".

Figure 2 shows the amino acid sequence (SEQ ID NO:2) derived from the coding sequence of SEQ ID NO:1 shown in Figure 1.

Figure 3 shows a nucleotide sequence (SEQ ID NO:3) of a native sequence PRO178 cDNA, wherein SEQ ID NO:3 is a clone designated herein as "DNA23339-1130".

Figure 4 shows the amino acid sequence (SEQ ID NO:4) derived from the coding sequence of SEQ ID NO:3 shown in Figure 3.

Figure 5 shows a nucleotide sequence (SEQ ID NO:5) of a native sequence PRO444 cDNA, wherein SEQ ID NO:5 is a clone designated herein as "DNA26846-1397".

Figure 6 shows the amino acid sequence (SEQ ID NO:6) derived from the coding sequence of SEQ ID NO:5 shown in Figure 5.

Figure 7 shows a nucleotide sequence (SEQ ID NO:7) of a native sequence PRO195 cDNA, wherein SEQ ID NO:7 is a clone designated herein as "DNA26847-1395".

Figure 8 shows the amino acid sequence (SEQ ID NO:8) derived from the coding sequence of SEQ ID NO:7 shown in Figure 7.

Figure 9 shows a nucleotide sequence (SEQ ID NO:9) of a native sequence PRO182 cDNA, wherein SEQ ID NO:9 is a clone designated herein as "DNA27865-1091".

Figure 10 shows the amino acid sequence (SEQ ID NO:10) derived from the coding sequence of SEQ ID

NO:9 shown in Figure 9.

Figure 11 shows a nucleotide sequence (SEQ ID NO:11) of a native sequence PRO205 cDNA, wherein SEQ ID NO:11 is a clone designated herein as "DNA30868-1156".

5 Figure 12 shows the amino acid sequence (SEQ ID NO:12) derived from the coding sequence of SEQ ID NO:11 shown in Figure 11.

Figure 13 shows a nucleotide sequence (SEQ ID NO:13) of a native sequence PRO204 cDNA, wherein SEQ ID NO:13 is a clone designated herein as "DNA30871-1157".

Figure 14 shows the amino acid sequence (SEQ ID NO:14) derived from the coding sequence of SEQ ID NO:13 shown in Figure 13.

10 Figure 15 shows a nucleotide sequence (SEQ ID NO:15) of a native sequence PRO1873 cDNA, wherein SEQ ID NO:15 is a clone designated herein as "DNA30880".

Figure 16 shows the amino acid sequence (SEQ ID NO:16) derived from the coding sequence of SEQ ID NO:15 shown in Figure 15.

15 Figure 17 shows a nucleotide sequence (SEQ ID NO:17) of a native sequence PRO214 cDNA, wherein SEQ ID NO:17 is a clone designated herein as "DNA32286-1191".

Figure 18 shows the amino acid sequence (SEQ ID NO:18) derived from the coding sequence of SEQ ID NO:17 shown in Figure 17.

Figure 19 shows a nucleotide sequence (SEQ ID NO:19) of a native sequence PRO221 cDNA, wherein SEQ ID NO:19 is a clone designated herein as "DNA33089-1132".

20 Figure 20 shows the amino acid sequence (SEQ ID NO:20) derived from the coding sequence of SEQ ID NO:19 shown in Figure 19.

Figure 21 shows a nucleotide sequence (SEQ ID NO:21) of a native sequence PRO228 cDNA, wherein SEQ ID NO:21 is a clone designated herein as "DNA33092-1202".

25 Figure 22 shows the amino acid sequence (SEQ ID NO:22) derived from the coding sequence of SEQ ID NO:21 shown in Figure 21.

Figure 23 shows a nucleotide sequence (SEQ ID NO:23) of a native sequence PRO229 cDNA, wherein SEQ ID NO:23 is a clone designated herein as "DNA33100-1159".

Figure 24 shows the amino acid sequence (SEQ ID NO:24) derived from the coding sequence of SEQ ID NO:23 shown in Figure 23.

30 Figure 25 shows a nucleotide sequence (SEQ ID NO:25) of a native sequence PRO230 cDNA, wherein SEQ ID NO:25 is a clone designated herein as "DNA33223-1136".

Figure 26 shows the amino acid sequence (SEQ ID NO:26) derived from the coding sequence of SEQ ID NO:25 shown in Figure 25.

35 Figure 27 shows a nucleotide sequence (SEQ ID NO:27) of a native sequence PRO7223 cDNA, wherein SEQ ID NO:27 is a clone designated herein as "DNA34385".

Figure 28 shows the amino acid sequence (SEQ ID NO:28) derived from the coding sequence of SEQ ID NO:27 shown in Figure 27.

Figure 29 shows a nucleotide sequence (SEQ ID NO:29) of a native sequence PRO241 cDNA, wherein SEQ ID NO:29 is a clone designated herein as "DNA34392-1170".

Figure 30 shows the amino acid sequence (SEQ ID NO:30) derived from the coding sequence of SEQ ID NO:29 shown in Figure 29.

5 Figure 31 shows a nucleotide sequence (SEQ ID NO:31) of a native sequence PRO263 cDNA, wherein SEQ ID NO:31 is a clone designated herein as "DNA34431-1177".

Figure 32 shows the amino acid sequence (SEQ ID NO:32) derived from the coding sequence of SEQ ID NO:31 shown in Figure 31.

10 Figure 33 shows a nucleotide sequence (SEQ ID NO:33) of a native sequence PRO321 cDNA, wherein SEQ ID NO:33 is a clone designated herein as "DNA34433-1308".

Figure 34 shows the amino acid sequence (SEQ ID NO:34) derived from the coding sequence of SEQ ID NO:33 shown in Figure 33.

Figure 35 shows a nucleotide sequence (SEQ ID NO:35) of a native sequence PRO231 cDNA, wherein SEQ ID NO:35 is a clone designated herein as "DNA34434-1139".

15 Figure 36 shows the amino acid sequence (SEQ ID NO:36) derived from the coding sequence of SEQ ID NO:35 shown in Figure 35.

Figure 37 shows a nucleotide sequence (SEQ ID NO:37) of a native sequence PRO238 cDNA, wherein SEQ ID NO:37 is a clone designated herein as "DNA35600-1162".

20 Figure 38 shows the amino acid sequence (SEQ ID NO:38) derived from the coding sequence of SEQ ID NO:37 shown in Figure 37.

Figure 39 shows a nucleotide sequence (SEQ ID NO:39) of a native sequence PRO247 cDNA, wherein SEQ ID NO:39 is a clone designated herein as "DNA35673-1201".

Figure 40 shows the amino acid sequence (SEQ ID NO:40) derived from the coding sequence of SEQ ID NO:39 shown in Figure 39.

25 Figure 41 shows a nucleotide sequence (SEQ ID NO:41) of a native sequence PRO256 cDNA, wherein SEQ ID NO:41 is a clone designated herein as "DNA35880-1160".

Figure 42 shows the amino acid sequence (SEQ ID NO:42) derived from the coding sequence of SEQ ID NO:41 shown in Figure 41.

30 Figure 43 shows a nucleotide sequence (SEQ ID NO:43) of a native sequence PRO258 cDNA, wherein SEQ ID NO:43 is a clone designated herein as "DNA35918-1174".

Figure 44 shows the amino acid sequence (SEQ ID NO:44) derived from the coding sequence of SEQ ID NO:43 shown in Figure 43.

Figure 45 shows a nucleotide sequence (SEQ ID NO:45) of a native sequence PRO265 cDNA, wherein SEQ ID NO:45 is a clone designated herein as "DNA36350-1158".

35 Figure 46 shows the amino acid sequence (SEQ ID NO:46) derived from the coding sequence of SEQ ID NO:45 shown in Figure 45.

Figure 47 shows a nucleotide sequence (SEQ ID NO:47) of a native sequence PRO21 cDNA, wherein SEQ

ID NO:47 is a clone designated herein as "DNA36638-1056".

Figure 48 shows the amino acid sequence (SEQ ID NO:48) derived from the coding sequence of SEQ ID NO:47 shown in Figure 47.

5 Figure 49 shows a nucleotide sequence (SEQ ID NO:49) of a native sequence PRO295 cDNA, wherein SEQ ID NO:49 is a clone designated herein as "DNA38268-1188".

Figure 50 shows the amino acid sequence (SEQ ID NO:50) derived from the coding sequence of SEQ ID NO:49 shown in Figure 49.

Figure 51 shows a nucleotide sequence (SEQ ID NO:51) of a native sequence PRO302 cDNA, wherein SEQ ID NO:51 is a clone designated herein as "DNA40370-1217".

10 Figure 52 shows the amino acid sequence (SEQ ID NO:52) derived from the coding sequence of SEQ ID NO:51 shown in Figure 51.

Figure 53 shows a nucleotide sequence (SEQ ID NO:53) of a native sequence PRO301 cDNA, wherein SEQ ID NO:53 is a clone designated herein as "DNA40628-1216".

15 Figure 54 shows the amino acid sequence (SEQ ID NO:54) derived from the coding sequence of SEQ ID NO:53 shown in Figure 53.

Figure 55 shows a nucleotide sequence (SEQ ID NO:55) of a native sequence PRO337 cDNA, wherein SEQ ID NO:55 is a clone designated herein as "DNA43316-1237".

Figure 56 shows the amino acid sequence (SEQ ID NO:56) derived from the coding sequence of SEQ ID NO:55 shown in Figure 55.

20 Figure 57 shows a nucleotide sequence (SEQ ID NO:57) of a native sequence PRO7248 cDNA, wherein SEQ ID NO:57 is a clone designated herein as "DNA44195".

Figure 58 shows the amino acid sequence (SEQ ID NO:58) derived from the coding sequence of SEQ ID NO:57 shown in Figure 57.

25 Figure 59 shows a nucleotide sequence (SEQ ID NO:59) of a native sequence PRO846 cDNA, wherein SEQ ID NO:59 is a clone designated herein as "DNA44196-1353".

Figure 60 shows the amino acid sequence (SEQ ID NO:60) derived from the coding sequence of SEQ ID NO:59 shown in Figure 59.

Figure 61 shows a nucleotide sequence (SEQ ID NO:61) of a native sequence PRO1864 cDNA, wherein SEQ ID NO:61 is a clone designated herein as "DNA45409-2511".

30 Figure 62 shows the amino acid sequence (SEQ ID NO:62) derived from the coding sequence of SEQ ID NO:61 shown in Figure 61.

Figure 63 shows a nucleotide sequence (SEQ ID NO:63) of a native sequence PRO363 cDNA, wherein SEQ ID NO:63 is a clone designated herein as "DNA45419-1252".

35 Figure 64 shows the amino acid sequence (SEQ ID NO:64) derived from the coding sequence of SEQ ID NO:63 shown in Figure 63.

Figure 65 shows a nucleotide sequence (SEQ ID NO:65) of a native sequence PRO730 cDNA, wherein SEQ ID NO:65 is a clone designated herein as "DNA45624-1400".

Figure 66 shows the amino acid sequence (SEQ ID NO:66) derived from the coding sequence of SEQ ID NO:65 shown in Figure 65.

Figure 67 shows a nucleotide sequence (SEQ ID NO:67) of a native sequence PRO365 cDNA, wherein SEQ ID NO:67 is a clone designated herein as "DNA46777-1253".

5 Figure 68 shows the amino acid sequence (SEQ ID NO:68) derived from the coding sequence of SEQ ID NO:67 shown in Figure 67.

Figure 69 shows a nucleotide sequence (SEQ ID NO:69) of a native sequence PRO532 cDNA, wherein SEQ ID NO:69 is a clone designated herein as "DNA48335".

10 Figure 70 shows the amino acid sequence (SEQ ID NO:70) derived from the coding sequence of SEQ ID NO:69 shown in Figure 69.

Figure 71 shows a nucleotide sequence (SEQ ID NO:71) of a native sequence PRO322 cDNA, wherein SEQ ID NO:71 is a clone designated herein as "DNA48336-1309".

Figure 72 shows the amino acid sequence (SEQ ID NO:72) derived from the coding sequence of SEQ ID NO:71 shown in Figure 71.

15 Figure 73 shows a nucleotide sequence (SEQ ID NO:73) of a native sequence PRO1120 cDNA, wherein SEQ ID NO:73 is a clone designated herein as "DNA48606-1479".

Figure 74 shows the amino acid sequence (SEQ ID NO:74) derived from the coding sequence of SEQ ID NO:73 shown in Figure 73.

20 Figure 75 shows a nucleotide sequence (SEQ ID NO:75) of a native sequence PRO7261 cDNA, wherein SEQ ID NO:75 is a clone designated herein as "DNA49149".

Figure 76 shows the amino acid sequence (SEQ ID NO:76) derived from the coding sequence of SEQ ID NO:75 shown in Figure 75.

Figure 77 shows a nucleotide sequence (SEQ ID NO:77) of a native sequence PRO533 cDNA, wherein SEQ ID NO:77 is a clone designated herein as "DNA49435-1219".

25 Figure 78 shows the amino acid sequence (SEQ ID NO:78) derived from the coding sequence of SEQ ID NO:77 shown in Figure 77.

Figure 79 shows a nucleotide sequence (SEQ ID NO:79) of a native sequence PRO724 cDNA, wherein SEQ ID NO:79 is a clone designated herein as "DNA49631-1328".

30 Figure 80 shows the amino acid sequence (SEQ ID NO:80) derived from the coding sequence of SEQ ID NO:79 shown in Figure 79.

Figure 81 shows a nucleotide sequence (SEQ ID NO:81) of a native sequence PRO734 cDNA, wherein SEQ ID NO:81 is a clone designated herein as "DNA49817".

Figure 82 shows the amino acid sequence (SEQ ID NO:82) derived from the coding sequence of SEQ ID NO:81 shown in Figure 81.

35 Figure 83 shows a nucleotide sequence (SEQ ID NO:83) of a native sequence PRO771 cDNA, wherein SEQ ID NO:83 is a clone designated herein as "DNA49829-1346".

Figure 84 shows the amino acid sequence (SEQ ID NO:84) derived from the coding sequence of SEQ ID

NO:83 shown in Figure 83.

Figure 85 shows a nucleotide sequence (SEQ ID NO:85) of a native sequence PRO2010 cDNA, wherein SEQ ID NO:85 is a clone designated herein as "DNA50792".

5 Figure 86 shows the amino acid sequence (SEQ ID NO:86) derived from the coding sequence of SEQ ID NO:85 shown in Figure 85.

Figure 87 shows a nucleotide sequence (SEQ ID NO:87) of a native sequence PRO871 cDNA, wherein SEQ ID NO:87 is a clone designated herein as "DNA50919-1361".

Figure 88 shows the amino acid sequence (SEQ ID NO:88) derived from the coding sequence of SEQ ID NO:87 shown in Figure 87.

10 Figure 89 shows a nucleotide sequence (SEQ ID NO:89) of a native sequence PRO697 cDNA, wherein SEQ ID NO:89 is a clone designated herein as "DNA50920-1325".

Figure 90 shows the amino acid sequence (SEQ ID NO:90) derived from the coding sequence of SEQ ID NO:89 shown in Figure 89.

15 Figure 91 shows a nucleotide sequence (SEQ ID NO:91) of a native sequence PRO1083 cDNA, wherein SEQ ID NO:91 is a clone designated herein as "DNA50921-1458".

Figure 92 shows the amino acid sequence (SEQ ID NO:22) derived from the coding sequence of SEQ ID NO:91 shown in Figure 91.

Figure 93 shows a nucleotide sequence (SEQ ID NO:93) of a native sequence PRO725 cDNA, wherein SEQ ID NO:93 is a clone designated herein as "DNA52758-1399".

20 Figure 94 shows the amino acid sequence (SEQ ID NO:94) derived from the coding sequence of SEQ ID NO:93 shown in Figure 93.

Figure 95 shows a nucleotide sequence (SEQ ID NO:95) of a native sequence PRO720 cDNA, wherein SEQ ID NO:95 is a clone designated herein as "DNA53517-1366-1".

25 Figure 96 shows the amino acid sequence (SEQ ID NO:96) derived from the coding sequence of SEQ ID NO:95 shown in Figure 95.

Figure 97 shows a nucleotide sequence (SEQ ID NO:97) of a native sequence PRO738 cDNA, wherein SEQ ID NO:97 is a clone designated herein as "DNA53915-1258".

Figure 98 shows the amino acid sequence (SEQ ID NO:98) derived from the coding sequence of SEQ ID NO:97 shown in Figure 97.

30 Figure 99 shows a nucleotide sequence (SEQ ID NO:99) of a native sequence PRO865 cDNA, wherein SEQ ID NO:99 is a clone designated herein as "DNA53974-1401".

Figure 100 shows the amino acid sequence (SEQ ID NO:100) derived from the coding sequence of SEQ ID NO:99 shown in Figure 99.

35 Figure 101 shows a nucleotide sequence (SEQ ID NO:101) of a native sequence PRO840 cDNA, wherein SEQ ID NO:101 is a clone designated herein as "DNA53987-1438".

Figure 102 shows the amino acid sequence (SEQ ID NO:102) derived from the coding sequence of SEQ ID NO:101 shown in Figure 101.

Figure 103 shows a nucleotide sequence (SEQ ID NO:103) of a native sequence PRO1080 cDNA, wherein SEQ ID NO:103 is a clone designated herein as "DNA56047-1456".

Figure 104 shows the amino acid sequence (SEQ ID NO:104) derived from the coding sequence of SEQ ID NO:103 shown in Figure 103.

5 Figure 105 shows a nucleotide sequence (SEQ ID NO:105) of a native sequence PRO1079 cDNA, wherein SEQ ID NO:105 is a clone designated herein as "DNA56050-1455".

Figure 106 shows the amino acid sequence (SEQ ID NO:106) derived from the coding sequence of SEQ ID NO:105 shown in Figure 105.

10 Figure 107 shows a nucleotide sequence (SEQ ID NO:107) of a native sequence PRO793 cDNA, wherein SEQ ID NO:107 is a clone designated herein as "DNA56110-1437".

Figure 108 shows the amino acid sequence (SEQ ID NO:108) derived from the coding sequence of SEQ ID NO:107 shown in Figure 107.

Figure 109 shows a nucleotide sequence (SEQ ID NO:109) of a native sequence PRO788 cDNA, wherein SEQ ID NO:109 is a clone designated herein as "DNA56405-1357".

15 Figure 110 shows the amino acid sequence (SEQ ID NO:110) derived from the coding sequence of SEQ ID NO:109 shown in Figure 109.

Figure 111 shows a nucleotide sequence (SEQ ID NO:111) of a native sequence PRO938 cDNA, wherein SEQ ID NO:111 is a clone designated herein as "DNA56433-1406".

20 Figure 112 shows the amino acid sequence (SEQ ID NO:112) derived from the coding sequence of SEQ ID NO:111 shown in Figure 111.

Figure 113 shows a nucleotide sequence (SEQ ID NO:113) of a native sequence PRO1012 cDNA, wherein SEQ ID NO:113 is a clone designated herein as "DNA56439-1376".

Figure 114 shows the amino acid sequence (SEQ ID NO:114) derived from the coding sequence of SEQ ID NO:113 shown in Figure 113.

25 Figure 115 shows a nucleotide sequence (SEQ ID NO:115) of a native sequence PRO1477 cDNA, wherein SEQ ID NO:115 is a clone designated herein as "DNA56529-1647".

Figure 116 shows the amino acid sequence (SEQ ID NO:116) derived from the coding sequence of SEQ ID NO:115 shown in Figure 115.

30 Figure 117 shows a nucleotide sequence (SEQ ID NO:117) of a native sequence PRO1134 cDNA, wherein SEQ ID NO:117 is a clone designated herein as "DNA56865-1491".

Figure 118 shows the amino acid sequence (SEQ ID NO:118) derived from the coding sequence of SEQ ID NO:117 shown in Figure 117.

Figure 119 shows a nucleotide sequence (SEQ ID NO:119) of a native sequence PRO162 cDNA, wherein SEQ ID NO:119 is a clone designated herein as "DNA56965-1356".

35 Figure 120 shows the amino acid sequence (SEQ ID NO:120) derived from the coding sequence of SEQ ID NO:119 shown in Figure 119.

Figure 121 shows a nucleotide sequence (SEQ ID NO:121) of a native sequence PRO1114 cDNA, wherein

SEQ ID NO:121 is a clone designated herein as "DNA57033-1403-1".

Figure 122 shows the amino acid sequence (SEQ ID NO:122) derived from the coding sequence of SEQ ID NO:121 shown in Figure 121.

5 Figure 123 shows a nucleotide sequence (SEQ ID NO:123) of a native sequence PRO828 cDNA, wherein SEQ ID NO:123 is a clone designated herein as "DNA57037-1444".

Figure 124 shows the amino acid sequence (SEQ ID NO:124) derived from the coding sequence of SEQ ID NO:123 shown in Figure 123.

Figure 125 shows a nucleotide sequence (SEQ ID NO:125) of a native sequence PRO827 cDNA, wherein SEQ ID NO:125 is a clone designated herein as "DNA57039-1402".

10 Figure 126 shows the amino acid sequence (SEQ ID NO:126) derived from the coding sequence of SEQ ID NO:125 shown in Figure 125.

Figure 127 shows a nucleotide sequence (SEQ ID NO:127) of a native sequence PRO1075 cDNA, wherein SEQ ID NO:127 is a clone designated herein as "DNA57689-1385".

15 Figure 128 shows the amino acid sequence (SEQ ID NO:128) derived from the coding sequence of SEQ ID NO:127 shown in Figure 127.

Figure 129 shows a nucleotide sequence (SEQ ID NO:129) of a native sequence PRO1007 cDNA, wherein SEQ ID NO:129 is a clone designated herein as "DNA57690-1374".

Figure 130 shows the amino acid sequence (SEQ ID NO:130) derived from the coding sequence of SEQ ID NO:129 shown in Figure 129.

20 Figure 131 shows a nucleotide sequence (SEQ ID NO:131) of a native sequence PRO826 cDNA, wherein SEQ ID NO:131 is a clone designated herein as "DNA57694-1341".

Figure 132 shows the amino acid sequence (SEQ ID NO:132) derived from the coding sequence of SEQ ID NO:131 shown in Figure 131.

25 Figure 133 shows a nucleotide sequence (SEQ ID NO:133) of a native sequence PRO819 cDNA, wherein SEQ ID NO:132 is a clone designated herein as "DNA57695-1340".

Figure 134 shows the amino acid sequence (SEQ ID NO:134) derived from the coding sequence of SEQ ID NO:133 shown in Figure 133.

Figure 135 shows a nucleotide sequence (SEQ ID NO:135) of a native sequence PRO1006 cDNA, wherein SEQ ID NO:135 is a clone designated herein as "DNA57699-1412".

30 Figure 136 shows the amino acid sequence (SEQ ID NO:136) derived from the coding sequence of SEQ ID NO:135 shown in Figure 135.

Figure 137 shows a nucleotide sequence (SEQ ID NO:137) of a native sequence PRO982 cDNA, wherein SEQ ID NO:137 is a clone designated herein as "DNA57700-1408".

35 Figure 138 shows the amino acid sequence (SEQ ID NO:138) derived from the coding sequence of SEQ ID NO:137 shown in Figure 137.

Figure 139 shows a nucleotide sequence (SEQ ID NO:139) of a native sequence PRO1005 cDNA, wherein SEQ ID NO:139 is a clone designated herein as "DNA57708-1411".

Figure 140 shows the amino acid sequence (SEQ ID NO:140) derived from the coding sequence of SEQ ID NO:139 shown in Figure 139.

Figure 141 shows a nucleotide sequence (SEQ ID NO:141) of a native sequence PRO791 cDNA, wherein SEQ ID NO:141 is a clone designated herein as "DNA57838-1337".

5 Figure 142 shows the amino acid sequence (SEQ ID NO:142) derived from the coding sequence of SEQ ID NO:141 shown in Figure 141.

Figure 143 shows a nucleotide sequence (SEQ ID NO:143) of a native sequence PRO1071 cDNA, wherein SEQ ID NO:143 is a clone designated herein as "DNA58847-1383".

10 Figure 144 shows the amino acid sequence (SEQ ID NO:144) derived from the coding sequence of SEQ ID NO:143 shown in Figure 43.

Figure 145 shows a nucleotide sequence (SEQ ID NO:145) of a native sequence PRO1415 cDNA, wherein SEQ ID NO:145 is a clone designated herein as "DNA58852-1637".

Figure 146 shows the amino acid sequence (SEQ ID NO:146) derived from the coding sequence of SEQ ID NO:145 shown in Figure 145.

15 Figure 147 shows a nucleotide sequence (SEQ ID NO:147) of a native sequence PRO1054 cDNA, wherein SEQ ID NO:147 is a clone designated herein as "DNA58853-1423".

Figure 148 shows the amino acid sequence (SEQ ID NO:148) derived from the coding sequence of SEQ ID NO:147 shown in Figure 147.

20 Figure 149 shows a nucleotide sequence (SEQ ID NO:149) of a native sequence PRO1411 cDNA, wherein SEQ ID NO:149 is a clone designated herein as "DNA59212-1627".

Figure 150 shows the amino acid sequence (SEQ ID NO:150) derived from the coding sequence of SEQ ID NO:149 shown in Figure 149.

Figure 151 shows a nucleotide sequence (SEQ ID NO:151) of a native sequence PRO1184 cDNA, wherein SEQ ID NO:151 is a clone designated herein as "DNA59220-1514".

25 Figure 152 shows the amino acid sequence (SEQ ID NO:152) derived from the coding sequence of SEQ ID NO:151 shown in Figure 151.

Figure 153 shows a nucleotide sequence (SEQ ID NO:153) of a native sequence PRO1029 cDNA, wherein SEQ ID NO:153 is a clone designated herein as "DNA59493-1420".

30 Figure 154 shows the amino acid sequence (SEQ ID NO:154) derived from the coding sequence of SEQ ID NO:153 shown in Figure 153.

Figure 155 shows a nucleotide sequence (SEQ ID NO:155) of a native sequence PRO1139 cDNA, wherein SEQ ID NO:155 is a clone designated herein as "DNA59497-1496".

Figure 156 shows the amino acid sequence (SEQ ID NO:156) derived from the coding sequence of SEQ ID NO:155 shown in Figure 155.

35 Figure 157 shows a nucleotide sequence (SEQ ID NO:157) of a native sequence PRO1190 cDNA, wherein SEQ ID NO:157 is a clone designated herein as "DNA59586-1520".

Figure 158 shows the amino acid sequence (SEQ ID NO:158) derived from the coding sequence of SEQ

ID NO:157 shown in Figure 157.

Figure 159 shows a nucleotide sequence (SEQ ID NO:159) of a native sequence PRO1309 cDNA, wherein SEQ ID NO:159 is a clone designated herein as "DNA59588-1571".

5 Figure 160 shows the amino acid sequence (SEQ ID NO:160) derived from the coding sequence of SEQ ID NO:159 shown in Figure 159.

Figure 161 shows a nucleotide sequence (SEQ ID NO:161) of a native sequence PRO836 cDNA, wherein SEQ ID NO:161 is a clone designated herein as "DNA59620-1463".

Figure 162 shows the amino acid sequence (SEQ ID NO:162) derived from the coding sequence of SEQ ID NO:161 shown in Figure 161.

10 Figure 163 shows a nucleotide sequence (SEQ ID NO:163) of a native sequence PRO1025 cDNA, wherein SEQ ID NO:163 is a clone designated herein as "DNA59622-1334".

Figure 164 shows the amino acid sequence (SEQ ID NO:164) derived from the coding sequence of SEQ ID NO:163 shown in Figure 163.

15 Figure 165 shows a nucleotide sequence (SEQ ID NO:165) of a native sequence PRO1131 cDNA, wherein SEQ ID NO:165 is a clone designated herein as "DNA59777-1480".

Figure 166 shows the amino acid sequence (SEQ ID NO:166) derived from the coding sequence of SEQ ID NO:165 shown in Figure 165.

Figure 167 shows a nucleotide sequence (SEQ ID NO:167) of a native sequence PRO1182 cDNA, wherein SEQ ID NO:167 is a clone designated herein as "DNA59848-1512".

20 Figure 168 shows the amino acid sequence (SEQ ID NO:168) derived from the coding sequence of SEQ ID NO:167 shown in Figure 167.

Figure 169 shows a nucleotide sequence (SEQ ID NO:169) of a native sequence PRO1155 cDNA, wherein SEQ ID NO:169 is a clone designated herein as "DNA59849-1504".

25 Figure 170 shows the amino acid sequence (SEQ ID NO:170) derived from the coding sequence of SEQ ID NO:169 shown in Figure 169.

Figure 171 shows a nucleotide sequence (SEQ ID NO:171) of a native sequence PRO1186 cDNA, wherein SEQ ID NO:171 is a clone designated herein as "DNA60621-1516".

Figure 172 shows the amino acid sequence (SEQ ID NO:172) derived from the coding sequence of SEQ ID NO:171 shown in Figure 171.

30 Figure 173 shows a nucleotide sequence (SEQ ID NO:173) of a native sequence PRO1198 cDNA, wherein SEQ ID NO:173 is a clone designated herein as "DNA60622-1525".

Figure 174 shows the amino acid sequence (SEQ ID NO:174) derived from the coding sequence of SEQ ID NO:173 shown in Figure 173.

35 Figure 175 shows a nucleotide sequence (SEQ ID NO:175) of a native sequence PRO1265 cDNA, wherein SEQ ID NO:175 is a clone designated herein as "DNA60764-1533".

Figure 176 shows the amino acid sequence (SEQ ID NO:176) derived from the coding sequence of SEQ ID NO:175 shown in Figure 175.

Figure 177 shows a nucleotide sequence (SEQ ID NO:177) of a native sequence PRO1361 cDNA, wherein SEQ ID NO:177 is a clone designated herein as "DNA60783-1611".

Figure 178 shows the amino acid sequence (SEQ ID NO:178) derived from the coding sequence of SEQ ID NO:177 shown in Figure 177.

5 Figure 179 shows a nucleotide sequence (SEQ ID NO:179) of a native sequence PRO1287 cDNA, wherein SEQ ID NO:179 is a clone designated herein as "DNA61755-1554".

Figure 180 shows the amino acid sequence (SEQ ID NO:180) derived from the coding sequence of SEQ ID NO:179 shown in Figure 179.

10 Figure 181 shows a nucleotide sequence (SEQ ID NO:181) of a native sequence PRO1308 cDNA, wherein SEQ ID NO:181 is a clone designated herein as "DNA62306-1570".

Figure 182 shows the amino acid sequence (SEQ ID NO:182) derived from the coding sequence of SEQ ID NO:181 shown in Figure 181.

Figure 183 shows a nucleotide sequence (SEQ ID NO:183) of a native sequence PRO4313 cDNA, wherein SEQ ID NO:183 is a clone designated herein as "DNA62312-2558".

15 Figure 184 shows the amino acid sequence (SEQ ID NO:184) derived from the coding sequence of SEQ ID NO:183 shown in Figure 183.

Figure 185 shows a nucleotide sequence (SEQ ID NO:185) of a native sequence PRO1192 cDNA, wherein SEQ ID NO:185 is a clone designated herein as "DNA62814-1521".

20 Figure 186 shows the amino acid sequence (SEQ ID NO:186) derived from the coding sequence of SEQ ID NO:185 shown in Figure 185.

Figure 187 shows a nucleotide sequence (SEQ ID NO:187) of a native sequence PRO1160 cDNA, wherein SEQ ID NO:187 is a clone designated herein as "DNA62872-1509".

Figure 188 shows the amino acid sequence (SEQ ID NO:188) derived from the coding sequence of SEQ ID NO:187 shown in Figure 187.

25 Figure 189 shows a nucleotide sequence (SEQ ID NO:189) of a native sequence PRO1244 cDNA, wherein SEQ ID NO:189 is a clone designated herein as "DNA64883-1526".

Figure 190 shows the amino acid sequence (SEQ ID NO:190) derived from the coding sequence of SEQ ID NO:189 shown in Figure 189.

30 Figure 191 shows a nucleotide sequence (SEQ ID NO:191) of a native sequence PRO1356 cDNA, wherein SEQ ID NO:191 is a clone designated herein as "DNA64886-1601".

Figure 192 shows the amino acid sequence (SEQ ID NO:192) derived from the coding sequence of SEQ ID NO:191 shown in Figure 191.

Figure 193 shows a nucleotide sequence (SEQ ID NO:193) of a native sequence PRO1274 cDNA, wherein SEQ ID NO:193 is a clone designated herein as "DNA64889-1541".

35 Figure 194 shows the amino acid sequence (SEQ ID NO:194) derived from the coding sequence of SEQ ID NO:193 shown in Figure 193.

Figure 195 shows a nucleotide sequence (SEQ ID NO:195) of a native sequence PRO1272 cDNA, wherein

SEQ ID NO:195 is a clone designated herein as "DNA64896-1539".

Figure 196 shows the amino acid sequence (SEQ ID NO:196) derived from the coding sequence of SEQ ID NO:195 shown in Figure 195.

5 Figure 197 shows a nucleotide sequence (SEQ ID NO:197) of a native sequence PRO1412 cDNA, wherein SEQ ID NO:197 is a clone designated herein as "DNA64897-1628".

Figure 198 shows the amino acid sequence (SEQ ID NO:198) derived from the coding sequence of SEQ ID NO:197 shown in Figure 197.

Figure 199 shows a nucleotide sequence (SEQ ID NO:199) of a native sequence PRO1286 cDNA, wherein SEQ ID NO:199 is a clone designated herein as "DNA64903-1553".

10 Figure 200 shows the amino acid sequence (SEQ ID NO:200) derived from the coding sequence of SEQ ID NO:199 shown in Figure 199.

Figure 201 shows a nucleotide sequence (SEQ ID NO:201) of a native sequence PRO1347 cDNA, wherein SEQ ID NO:201 is a clone designated herein as "DNA64950-1590".

15 Figure 202 shows the amino acid sequence (SEQ ID NO:202) derived from the coding sequence of SEQ ID NO:201 shown in Figure 201.

Figure 203 shows a nucleotide sequence (SEQ ID NO:203) of a native sequence PRO1273 cDNA, wherein SEQ ID NO:203 is a clone designated herein as "DNA65402-1540".

Figure 204 shows the amino acid sequence (SEQ ID NO:204) derived from the coding sequence of SEQ ID NO:203 shown in Figure 203.

20 Figure 205 shows a nucleotide sequence (SEQ ID NO:205) of a native sequence PRO1283 cDNA, wherein SEQ ID NO:205 is a clone designated herein as "DNA65404-1551".

Figure 206 shows the amino acid sequence (SEQ ID NO:206) derived from the coding sequence of SEQ ID NO:205 shown in Figure 205.

25 Figure 207 shows a nucleotide sequence (SEQ ID NO:207) of a native sequence PRO1279 cDNA, wherein SEQ ID NO:207 is a clone designated herein as "DNA65405-1547".

Figure 208 shows the amino acid sequence (SEQ ID NO:208) derived from the coding sequence of SEQ ID NO:207 shown in Figure 207.

Figure 209 shows a nucleotide sequence (SEQ ID NO:209) of a native sequence PRO1306 cDNA, wherein SEQ ID NO:209 is a clone designated herein as "DNA65410-1569".

30 Figure 210 shows the amino acid sequence (SEQ ID NO:210) derived from the coding sequence of SEQ ID NO:209 shown in Figure 209.

Figure 211 shows a nucleotide sequence (SEQ ID NO:211) of a native sequence PRO1195 cDNA, wherein SEQ ID NO:211 is a clone designated herein as "DNA65412-1523".

35 Figure 212 shows the amino acid sequence (SEQ ID NO:212) derived from the coding sequence of SEQ ID NO:211 shown in Figure 211.

Figure 213 shows a nucleotide sequence (SEQ ID NO:213) of a native sequence PRO4995 cDNA, wherein SEQ ID NO:213 is a clone designated herein as "DNA66307-2661".

Figure 214 shows the amino acid sequence (SEQ ID NO:214) derived from the coding sequence of SEQ ID NO:213 shown in Figure 213.

Figure 215 shows a nucleotide sequence (SEQ ID NO:215) of a native sequence PRO1382 cDNA, wherein SEQ ID NO:215 is a clone designated herein as "DNA66526-1616".

5 Figure 216 shows the amino acid sequence (SEQ ID NO:216) derived from the coding sequence of SEQ ID NO:215 shown in Figure 215.

Figure 217 shows a nucleotide sequence (SEQ ID NO:217) of a native sequence PRO1325 cDNA, wherein SEQ ID NO:217 is a clone designated herein as "DNA66659-1593".

10 Figure 218 shows the amino acid sequence (SEQ ID NO:218) derived from the coding sequence of SEQ ID NO:217 shown in Figure 217.

Figure 219 shows a nucleotide sequence (SEQ ID NO:219) of a native sequence PRO1329 cDNA, wherein SEQ ID NO:219 is a clone designated herein as "DNA66660-1585".

Figure 220 shows the amino acid sequence (SEQ ID NO:220) derived from the coding sequence of SEQ ID NO:219 shown in Figure 219.

15 Figure 221 shows a nucleotide sequence (SEQ ID NO:221) of a native sequence PRO1338 cDNA, wherein SEQ ID NO:221 is a clone designated herein as "DNA66667-1596".

Figure 222 shows the amino acid sequence (SEQ ID NO:222) derived from the coding sequence of SEQ ID NO:221 shown in Figure 221.

20 Figure 223 shows a nucleotide sequence (SEQ ID NO:223) of a native sequence PRO1337 cDNA, wherein SEQ ID NO:223 is a clone designated herein as "DNA66672-1586".

Figure 224 shows the amino acid sequence (SEQ ID NO:224) derived from the coding sequence of SEQ ID NO:223 shown in Figure 223.

Figure 225 shows a nucleotide sequence (SEQ ID NO:225) of a native sequence PRO1343 cDNA, wherein SEQ ID NO:225 is a clone designated herein as "DNA66675-1587".

25 Figure 226 shows the amino acid sequence (SEQ ID NO:226) derived from the coding sequence of SEQ ID NO:225 shown in Figure 225.

Figure 227 shows a nucleotide sequence (SEQ ID NO:227) of a native sequence PRO1376 cDNA, wherein SEQ ID NO:227 is a clone designated herein as "DNA67300-1605".

30 Figure 228 shows the amino acid sequence (SEQ ID NO:228) derived from the coding sequence of SEQ ID NO:227 shown in Figure 227.

Figure 229 shows a nucleotide sequence (SEQ ID NO:229) of a native sequence PRO1434 cDNA, wherein SEQ ID NO:229 is a clone designated herein as "DNA68818-2536".

Figure 230 shows the amino acid sequence (SEQ ID NO:230) derived from the coding sequence of SEQ ID NO:229 shown in Figure 229.

35 Figure 231 shows a nucleotide sequence (SEQ ID NO:231) of a native sequence PRO3579 cDNA, wherein SEQ ID NO:231 is a clone designated herein as "DNA68862-2546".

Figure 232 shows the amino acid sequence (SEQ ID NO:232) derived from the coding sequence of SEQ

ID NO:231 shown in Figure 231.

Figure 233 shows a nucleotide sequence (SEQ ID NO:233) of a native sequence PRO1387 cDNA, wherein SEQ ID NO:233 is a clone designated herein as "DNA68872-1620".

5 Figure 234 shows the amino acid sequence (SEQ ID NO:234) derived from the coding sequence of SEQ ID NO:233 shown in Figure 233.

Figure 235 shows a nucleotide sequence (SEQ ID NO:235) of a native sequence PRO1419 cDNA, wherein SEQ ID NO:235 is a clone designated herein as "DNA71290-1630".

Figure 236 shows the amino acid sequence (SEQ ID NO:236) derived from the coding sequence of SEQ ID NO:235 shown in Figure 235.

10 Figure 237 shows a nucleotide sequence (SEQ ID NO:237) of a native sequence PRO1488 cDNA, wherein SEQ ID NO:237 is a clone designated herein as "DNA73736-1657".

Figure 238 shows the amino acid sequence (SEQ ID NO:238) derived from the coding sequence of SEQ ID NO:237 shown in Figure 237.

15 Figure 239 shows a nucleotide sequence (SEQ ID NO:239) of a native sequence PRO1474 cDNA, wherein SEQ ID NO:239 is a clone designated herein as "DNA73739-1645".

Figure 240 shows the amino acid sequence (SEQ ID NO:240) derived from the coding sequence of SEQ ID NO:239 shown in Figure 239.

Figure 241 shows a nucleotide sequence (SEQ ID NO:241) of a native sequence PRO1508 cDNA, wherein SEQ ID NO:241 is a clone designated herein as "DNA73742-1662".

20 Figure 242 shows the amino acid sequence (SEQ ID NO:242) derived from the coding sequence of SEQ ID NO:241 shown in Figure 241.

Figure 243 shows a nucleotide sequence (SEQ ID NO:243) of a native sequence PRO1754 cDNA, wherein SEQ ID NO:243 is a clone designated herein as "DNA76385-1692".

25 Figure 244 shows the amino acid sequence (SEQ ID NO:244) derived from the coding sequence of SEQ ID NO:243 shown in Figure 243.

Figure 245 shows a nucleotide sequence (SEQ ID NO:245) of a native sequence PRO1550 cDNA, wherein SEQ ID NO:245 is a clone designated herein as "DNA76393-1664".

Figure 246 shows the amino acid sequence (SEQ ID NO:246) derived from the coding sequence of SEQ ID NO:245 shown in Figure 245.

30 Figure 247 shows a nucleotide sequence (SEQ ID NO:247) of a native sequence PRO1758 cDNA, wherein SEQ ID NO:247 is a clone designated herein as "DNA76399-1700".

Figure 248 shows the amino acid sequence (SEQ ID NO:248) derived from the coding sequence of SEQ ID NO:247 shown in Figure 247.

35 Figure 249 shows a nucleotide sequence (SEQ ID NO:249) of a native sequence PRO1917 cDNA, wherein SEQ ID NO:249 is a clone designated herein as "DNA76400-2528".

Figure 250 shows the amino acid sequence (SEQ ID NO:250) derived from the coding sequence of SEQ ID NO:249 shown in Figure 249.

Figure 251 shows a nucleotide sequence (SEQ ID NO:251) of a native sequence PRO1787 cDNA, wherein SEQ ID NO:251 is a clone designated herein as "DNA76510-2504".

Figure 252 shows the amino acid sequence (SEQ ID NO:252) derived from the coding sequence of SEQ ID NO:251 shown in Figure 251.

5 Figure 253 shows a nucleotide sequence (SEQ ID NO:253) of a native sequence PRO1556 cDNA, wherein SEQ ID NO:253 is a clone designated herein as "DNA76529-1666".

Figure 254 shows the amino acid sequence (SEQ ID NO:254) derived from the coding sequence of SEQ ID NO:253 shown in Figure 253.

10 Figure 255 shows a nucleotide sequence (SEQ ID NO:255) of a native sequence PRO1760 cDNA, wherein SEQ ID NO:255 is a clone designated herein as "DNA76532-1702".

Figure 256 shows the amino acid sequence (SEQ ID NO:256) derived from the coding sequence of SEQ ID NO:255 shown in Figure 255.

Figure 257 shows a nucleotide sequence (SEQ ID NO:257) of a native sequence PRO1567 cDNA, wherein SEQ ID NO:257 is a clone designated herein as "DNA76541-1675".

15 Figure 258 shows the amino acid sequence (SEQ ID NO:258) derived from the coding sequence of SEQ ID NO:257 shown in Figure 257.

Figure 259 shows a nucleotide sequence (SEQ ID NO:259) of a native sequence PRO1600 cDNA, wherein SEQ ID NO:259 is a clone designated herein as "DNA77503-1686".

20 Figure 260 shows the amino acid sequence (SEQ ID NO:260) derived from the coding sequence of SEQ ID NO:259 shown in Figure 259.

Figure 261 shows a nucleotide sequence (SEQ ID NO:261) of a native sequence PRO1868 cDNA, wherein SEQ ID NO:261 is a clone designated herein as "DNA77624-2515".

Figure 262 shows the amino acid sequence (SEQ ID NO:262) derived from the coding sequence of SEQ ID NO:261 shown in Figure 261.

25 Figure 263 shows a nucleotide sequence (SEQ ID NO:263) of a native sequence PRO1890 cDNA, wherein SEQ ID NO:263 is a clone designated herein as "DNA79230-2525".

Figure 264 shows the amino acid sequence (SEQ ID NO:264) derived from the coding sequence of SEQ ID NO:263 shown in Figure 263.

30 Figure 265 shows a nucleotide sequence (SEQ ID NO:265) of a native sequence PRO1887 cDNA, wherein SEQ ID NO:265 is a clone designated herein as "DNA79862-2522".

Figure 266 shows the amino acid sequence (SEQ ID NO:265) derived from the coding sequence of SEQ ID NO:265 shown in Figure 265.

Figure 267 shows a nucleotide sequence (SEQ ID NO:267) of a native sequence PRO4353 cDNA, wherein SEQ ID NO:267 is a clone designated herein as "DNA80145-2594".

35 Figure 268 shows the amino acid sequence (SEQ ID NO:268) derived from the coding sequence of SEQ ID NO:267 shown in Figure 267.

Figure 269 shows a nucleotide sequence (SEQ ID NO:269) of a native sequence PRO1782 cDNA, wherein

SEQ ID NO:269 is a clone designated herein as "DNA80899-2501".

Figure 270 shows the amino acid sequence (SEQ ID NO:270) derived from the coding sequence of SEQ ID NO:269 shown in Figure 269.

5 Figure 271 shows a nucleotide sequence (SEQ ID NO:271) of a native sequence PRO1928 cDNA, wherein SEQ ID NO:271 is a clone designated herein as "DNA81754-2532".

Figure 272 shows the amino acid sequence (SEQ ID NO:272) derived from the coding sequence of SEQ ID NO:271 shown in Figure 271.

Figure 273 shows a nucleotide sequence (SEQ ID NO:273) of a native sequence PRO1865 cDNA, wherein SEQ ID NO:273 is a clone designated herein as "DNA81757-2512".

10 Figure 274 shows the amino acid sequence (SEQ ID NO:274) derived from the coding sequence of SEQ ID NO:273 shown in Figure 273.

Figure 275 shows a nucleotide sequence (SEQ ID NO:275) of a native sequence PRO4341 cDNA, wherein SEQ ID NO:275 is a clone designated herein as "DNA81761-2583".

15 Figure 276 shows the amino acid sequence (SEQ ID NO:276) derived from the coding sequence of SEQ ID NO:275 shown in Figure 275.

Figure 277 shows a nucleotide sequence (SEQ ID NO:277) of a native sequence PRO6714 cDNA, wherein SEQ ID NO:277 is a clone designated herein as "DNA82358-2738".

Figure 278 shows the amino acid sequence (SEQ ID NO:278) derived from the coding sequence of SEQ ID NO:277 shown in Figure 277.

20 Figure 279 shows a nucleotide sequence (SEQ ID NO:279) of a native sequence PRO5723 cDNA, wherein SEQ ID NO:279 is a clone designated herein as "DNA82361".

Figure 280 shows the amino acid sequence (SEQ ID NO:280) derived from the coding sequence of SEQ ID NO:279 shown in Figure 279.

25 Figure 281 shows a nucleotide sequence (SEQ ID NO:281) of a native sequence PRO3438 cDNA, wherein SEQ ID NO:281 is a clone designated herein as "DNA82364-2538".

Figure 282 shows the amino acid sequence (SEQ ID NO:282) derived from the coding sequence of SEQ ID NO:281 shown in Figure 281.

Figure 283 shows a nucleotide sequence (SEQ ID NO:283) of a native sequence PRO6071 cDNA, wherein SEQ ID NO:283 is a clone designated herein as "DNA82403-2959".

30 Figure 284 shows the amino acid sequence (SEQ ID NO:284) derived from the coding sequence of SEQ ID NO:283 shown in Figure 283.

Figure 285 shows a nucleotide sequence (SEQ ID NO:285) of a native sequence PRO1801 cDNA, wherein SEQ ID NO:285 is a clone designated herein as "DNA83500-2506".

35 Figure 286 shows the amino acid sequence (SEQ ID NO:286) derived from the coding sequence of SEQ ID NO:285 shown in Figure 285.

Figure 287 shows a nucleotide sequence (SEQ ID NO:287) of a native sequence PRO4324 cDNA, wherein SEQ ID NO:287 is a clone designated herein as "DNA83560-2569".

Figure 288 shows the amino acid sequence (SEQ ID NO:288) derived from the coding sequence of SEQ ID NO:287 shown in Figure 287.

Figure 289 shows a nucleotide sequence (SEQ ID NO:289) of a native sequence PRO4333 cDNA, wherein SEQ ID NO:289 is a clone designated herein as "DNA84210-2576".

5 Figure 290 shows the amino acid sequence (SEQ ID NO:290) derived from the coding sequence of SEQ ID NO:289 shown in Figure 289.

Figure 291 shows a nucleotide sequence (SEQ ID NO:291) of a native sequence PRO4405 cDNA, wherein SEQ ID NO:291 is a clone designated herein as "DNA84920-2614".

10 Figure 292 shows the amino acid sequence (SEQ ID NO:292) derived from the coding sequence of SEQ ID NO:291 shown in Figure 291.

Figure 293 shows a nucleotide sequence (SEQ ID NO:293) of a native sequence PRO4356 cDNA, wherein SEQ ID NO:293 is a clone designated herein as "DNA86576-2595".

Figure 294 shows the amino acid sequence (SEQ ID NO:294) derived from the coding sequence of SEQ ID NO:293 shown in Figure 293.

15 Figure 295 shows a nucleotide sequence (SEQ ID NO:295) of a native sequence PRO3444 cDNA, wherein SEQ ID NO:295 is a clone designated herein as "DNA87997".

Figure 296 shows the amino acid sequence (SEQ ID NO:296) derived from the coding sequence of SEQ ID NO:295 shown in Figure 295.

20 Figure 297 shows a nucleotide sequence (SEQ ID NO:297) of a native sequence PRO4302 cDNA, wherein SEQ ID NO:297 is a clone designated herein as "DNA92218-2554".

Figure 298 shows the amino acid sequence (SEQ ID NO:298) derived from the coding sequence of SEQ ID NO:297 shown in Figure 297.

Figure 299 shows a nucleotide sequence (SEQ ID NO:299) of a native sequence PRO4371 cDNA, wherein SEQ ID NO:299 is a clone designated herein as "DNA92233-2599".

25 Figure 300 shows the amino acid sequence (SEQ ID NO:300) derived from the coding sequence of SEQ ID NO:299 shown in Figure 299.

Figure 301 shows a nucleotide sequence (SEQ ID NO:301) of a native sequence PRO4354 cDNA, wherein SEQ ID NO:301 is a clone designated herein as "DNA92256-2596".

30 Figure 302 shows the amino acid sequence (SEQ ID NO:302) derived from the coding sequence of SEQ ID NO:301 shown in Figure 301.

Figure 303 shows a nucleotide sequence (SEQ ID NO:303) of a native sequence PRO5725 cDNA, wherein SEQ ID NO:303 is a clone designated herein as "DNA92265-2669".

Figure 304 shows the amino acid sequence (SEQ ID NO:304) derived from the coding sequence of SEQ ID NO:303 shown in Figure 303.

35 Figure 305 shows a nucleotide sequence (SEQ ID NO:305) of a native sequence PRO4408 cDNA, wherein SEQ ID NO:305 is a clone designated herein as "DNA92274-2617".

Figure 306 shows the amino acid sequence (SEQ ID NO:306) derived from the coding sequence of SEQ

ID NO:305 shown in Figure 305.

Figure 307 shows a nucleotide sequence (SEQ ID NO:307) of a native sequence PRO9940 cDNA, wherein SEQ ID NO:307 is a clone designated herein as "DNA92282".

5 Figure 308 shows the amino acid sequence (SEQ ID NO:308) derived from the coding sequence of SEQ ID NO:307 shown in Figure 307.

Figure 309 shows a nucleotide sequence (SEQ ID NO:309) of a native sequence PRO5737 cDNA, wherein SEQ ID NO:309 is a clone designated herein as "DNA92929-2534-1".

Figure 310 shows the amino acid sequence (SEQ ID NO:310) derived from the coding sequence of SEQ ID NO:309 shown in Figure 309.

10 Figure 311 shows a nucleotide sequence (SEQ ID NO:311) of a native sequence PRO4425 cDNA, wherein SEQ ID NO:311 is a clone designated herein as "DNA93011-2637".

Figure 312 shows the amino acid sequence (SEQ ID NO:312) derived from the coding sequence of SEQ ID NO:311 shown in Figure 311.

15 Figure 313 shows a nucleotide sequence (SEQ ID NO:313) of a native sequence PRO4345 cDNA, wherein SEQ ID NO:313 is a clone designated herein as "DNA94854-2586".

Figure 314 shows the amino acid sequence (SEQ ID NO:314) derived from the coding sequence of SEQ ID NO:313 shown in Figure 313.

Figure 315 shows a nucleotide sequence (SEQ ID NO:315) of a native sequence PRO4342 cDNA, wherein SEQ ID NO:315 is a clone designated herein as "DNA96787-2534-1".

20 Figure 316 shows the amino acid sequence (SEQ ID NO:316) derived from the coding sequence of SEQ ID NO:315 shown in Figure 315.

Figure 317 shows a nucleotide sequence (SEQ ID NO:317) of a native sequence PRO3562 cDNA, wherein SEQ ID NO:317 is a clone designated herein as "DNA96791".

25 Figure 318 shows the amino acid sequence (SEQ ID NO:318) derived from the coding sequence of SEQ ID NO:317 shown in Figure 317.

Figure 319 shows a nucleotide sequence (SEQ ID NO:319) of a native sequence PRO4422 cDNA, wherein SEQ ID NO:319 is a clone designated herein as "DNA96867-2620".

Figure 320 shows the amino acid sequence (SEQ ID NO:320) derived from the coding sequence of SEQ ID NO:319 shown in Figure 319.

30 Figure 321 shows a nucleotide sequence (SEQ ID NO:321) of a native sequence PRO5776 cDNA, wherein SEQ ID NO:321 is a clone designated herein as "DNA96872-2674".

Figure 322 shows the amino acid sequence (SEQ ID NO:322) derived from the coding sequence of SEQ ID NO:321 shown in Figure 321.

35 Figure 323 shows a nucleotide sequence (SEQ ID NO:323) of a native sequence PRO4430 cDNA, wherein SEQ ID NO:323 is a clone designated herein as "DNA96878-2626".

Figure 324 shows the amino acid sequence (SEQ ID NO:324) derived from the coding sequence of SEQ ID NO:323 shown in Figure 323.

Figure 325 shows a nucleotide sequence (SEQ ID NO:325) of a native sequence PRO4499 cDNA, wherein SEQ ID NO:325 is a clone designated herein as "DNA96889-2641".

Figure 326 shows the amino acid sequence (SEQ ID NO:326) derived from the coding sequence of SEQ ID NO:325 shown in Figure 325.

5 Figure 327 shows a nucleotide sequence (SEQ ID NO:327) of a native sequence PRO4503 cDNA, wherein SEQ ID NO:327 is a clone designated herein as "DNA100312-2645".

Figure 328 shows the amino acid sequence (SEQ ID NO:328) derived from the coding sequence of SEQ ID NO:327 shown in Figure 327.

10 Figure 329 shows a nucleotide sequence (SEQ ID NO:329) of a native sequence PRO10008 cDNA, wherein SEQ ID NO:329 is a clone designated herein as "DNA101921".

Figure 330 shows the amino acid sequence (SEQ ID NO:330) derived from the coding sequence of SEQ ID NO:329 shown in Figure 329.

Figure 331 shows a nucleotide sequence (SEQ ID NO:331) of a native sequence PRO5730 cDNA, wherein SEQ ID NO:331 is a clone designated herein as "DNA101926".

15 Figure 332 shows the amino acid sequence (SEQ ID NO:332) derived from the coding sequence of SEQ ID NO:331 shown in Figure 331.

Figure 333 shows a nucleotide sequence (SEQ ID NO:333) of a native sequence PRO6008 cDNA, wherein SEQ ID NO:333 is a clone designated herein as "DNA102844".

20 Figure 334 shows the amino acid sequence (SEQ ID NO:334) derived from the coding sequence of SEQ ID NO:333 shown in Figure 333.

Figure 335 shows a nucleotide sequence (SEQ ID NO:335) of a native sequence PRO4527 cDNA, wherein SEQ ID NO:335 is a clone designated herein as "DNA103197".

Figure 336 shows the amino acid sequence (SEQ ID NO:336) derived from the coding sequence of SEQ ID NO:335 shown in Figure 335.

25 Figure 337 shows a nucleotide sequence (SEQ ID NO:337) of a native sequence PRO4538 cDNA, wherein SEQ ID NO:337 is a clone designated herein as "DNA103208".

Figure 338 shows the amino acid sequence (SEQ ID NO:338) derived from the coding sequence of SEQ ID NO:337 shown in Figure 337.

30 Figure 339 shows a nucleotide sequence (SEQ ID NO:339) of a native sequence PRO4553 cDNA, wherein SEQ ID NO:339 is a clone designated herein as "DNA103223".

Figure 340 shows the amino acid sequence (SEQ ID NO:340) derived from the coding sequence of SEQ ID NO:339 shown in Figure 339.

Figure 341 shows a nucleotide sequence (SEQ ID NO:341) of a native sequence PRO6006 cDNA, wherein SEQ ID NO:341 is a clone designated herein as "DNA105782-2693".

35 Figure 342 shows the amino acid sequence (SEQ ID NO:342) derived from the coding sequence of SEQ ID NO:341 shown in Figure 341.

Figure 343 shows a nucleotide sequence (SEQ ID NO:343) of a native sequence PRO6029 cDNA, wherein

SEQ ID NO:343 is a clone designated herein as "DNA105849-2704".

Figure 344 shows the amino acid sequence (SEQ ID NO:344) derived from the coding sequence of SEQ ID NO:343 shown in Figure 343.

5 Figure 345 shows a nucleotide sequence (SEQ ID NO:345) of a native sequence PRO9821 cDNA, wherein SEQ ID NO:345 is a clone designated herein as "DNA108725-2766".

Figure 346 shows the amino acid sequence (SEQ ID NO:346) derived from the coding sequence of SEQ ID NO:345 shown in Figure 345.

Figure 347 shows a nucleotide sequence (SEQ ID NO:347) of a native sequence PRO9820 cDNA, wherein SEQ ID NO:347 is a clone designated herein as "DNA108769-2765".

10 Figure 348 shows the amino acid sequence (SEQ ID NO:348) derived from the coding sequence of SEQ ID NO:347 shown in Figure 347.

Figure 349 shows a nucleotide sequence (SEQ ID NO:349) of a native sequence PRO9771 cDNA, wherein SEQ ID NO:349 is a clone designated herein as "DNA119498-2965".

15 Figure 350 shows the amino acid sequence (SEQ ID NO:350) derived from the coding sequence of SEQ ID NO:349 shown in Figure 349.

Figure 351 shows a nucleotide sequence (SEQ ID NO:351) of a native sequence PRO7436 cDNA, wherein SEQ ID NO:351 is a clone designated herein as "DNA119535-2756".

Figure 352 shows the amino acid sequence (SEQ ID NO:352) derived from the coding sequence of SEQ ID NO:351 shown in Figure 351.

20 Figure 353 shows a nucleotide sequence (SEQ ID NO:353) of a native sequence PRO10096 cDNA, wherein SEQ ID NO:353 is a clone designated herein as "DNA125185-2806".

Figure 354 shows the amino acid sequence (SEQ ID NO:354) derived from the coding sequence of SEQ ID NO:353 shown in Figure 353.

25 Figure 355 shows a nucleotide sequence (SEQ ID NO:355) of a native sequence PRO19670 cDNA, wherein SEQ ID NO:355 is a clone designated herein as "DNA131639-2874".

Figure 356 shows the amino acid sequence (SEQ ID NO:356) derived from the coding sequence of SEQ ID NO:355 shown in Figure 355.

Figure 357 shows a nucleotide sequence (SEQ ID NO:357) of a native sequence PRO20044 cDNA, wherein SEQ ID NO:357 is a clone designated herein as "DNA139623-2893".

30 Figure 358 shows the amino acid sequence (SEQ ID NO:358) derived from the coding sequence of SEQ ID NO:357 shown in Figure 357.

Figure 359 shows a nucleotide sequence (SEQ ID NO:359) of a native sequence PRO9873 cDNA, wherein SEQ ID NO:359 is a clone designated herein as "DNA143076-2787".

35 Figure 360 shows the amino acid sequence (SEQ ID NO:360) derived from the coding sequence of SEQ ID NO:359 shown in Figure 359.

Figure 361 shows a nucleotide sequence (SEQ ID NO:361) of a native sequence PRO21366 cDNA, wherein SEQ ID NO:361 is a clone designated herein as "DNA143276-2975".

Figure 362 shows the amino acid sequence (SEQ ID NO:362) derived from the coding sequence of SEQ ID NO:361 shown in Figure 361.

Figure 363 shows a nucleotide sequence (SEQ ID NO:363) of a native sequence PRO20040 cDNA, wherein SEQ ID NO:363 is a clone designated herein as "DNA164625-2890".

5 Figure 364 shows the amino acid sequence (SEQ ID NO:364) derived from the coding sequence of SEQ ID NO:363 shown in Figure 363.

Figure 365 shows a nucleotide sequence (SEQ ID NO:365) of a native sequence PRO21184 cDNA, wherein SEQ ID NO:365 is a clone designated herein as "DNA167678-2963".

10 Figure 366 shows the amino acid sequence (SEQ ID NO:366) derived from the coding sequence of SEQ ID NO:365 shown in Figure 365.

Figure 367 shows a nucleotide sequence (SEQ ID NO:367) of a native sequence PRO21055 cDNA, wherein SEQ ID NO:367 is a clone designated herein as "DNA170021-2923".

Figure 368 shows the amino acid sequence (SEQ ID NO:368) derived from the coding sequence of SEQ ID NO:367 shown in Figure 367.

15 Figure 369 shows a nucleotide sequence (SEQ ID NO:369) of a native sequence PRO28631 cDNA, wherein SEQ ID NO:369 is a clone designated herein as "DNA170212-3000".

Figure 370 shows the amino acid sequence (SEQ ID NO:370) derived from the coding sequence of SEQ ID NO:369 shown in Figure 369.

20 Figure 371 shows a nucleotide sequence (SEQ ID NO:371) of a native sequence PRO21384 cDNA, wherein SEQ ID NO:371 is a clone designated herein as "DNA177313-2982".

Figure 372 shows the amino acid sequence (SEQ ID NO:372) derived from the coding sequence of SEQ ID NO:371 shown in Figure 371.

Figure 373 shows a nucleotide sequence (SEQ ID NO:373) of a native sequence PRO1449 cDNA, wherein SEQ ID NO:373 is a clone designated herein as "DNA64908-1163-1".

25 Figure 374 shows the amino acid sequence (SEQ ID NO:374) derived from the coding sequence of SEQ ID NO:373 shown in Figure 373.

30 Figure 375 shows wholemount in situ hybridization results on mouse embryos using a mouse orthologue of PRO1449 which has about 78% amino acid identity with PRO1449. The results show that PRO1449 orthologue is expressed in the developing vasculature. The cross-section further shows expression in endothelial cells and progenitors of endothelial cells.

Figure 376 shows that a PRO1449 orthologue having about 78% amino acid identity with PRO1449 is expressed in vasculature of many inflamed and diseased tissues, but is very low, or lacking, in normal adult vessels.

Figure 377 shows that a PRO1449 orthologue having about 78% amino acid identity with PRO1449 induces ectopic vessels in the eyes of chicken embryos.

35

5. Detailed Description of the Invention

5.1. Definitions

The phrases "cardiovascular, endothelial and angiogenic disorder", "cardiovascular, endothelial and angiogenic dysfunction", "cardiovascular, endothelial or angiogenic disorder" and "cardiovascular, endothelial or angiogenic dysfunction" are used interchangeably and refer in part to systemic disorders that affect vessels, such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins, and/or lymphatics. This would include indications that stimulate angiogenesis and/or cardiovascularization, and those that inhibit angiogenesis and/or cardiovascularization. Such disorders include, for example, arterial disease, such as atherosclerosis, hypertension, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, and arterial restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, cancer such as vascular tumors, *e.g.*, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, Kaposi's sarcoma, lymphangioma, and lymphangiosarcoma, tumor angiogenesis, trauma such as wounds, burns, and other injured tissue, implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. This would also include angina, myocardial infarctions such as acute myocardial infarctions, cardiac hypertrophy, and heart failure such as CHF.

"Hypertrophy", as used herein, is defined as an increase in mass of an organ or structure independent of natural growth that does not involve tumor formation. Hypertrophy of an organ or tissue is due either to an increase in the mass of the individual cells (true hypertrophy), or to an increase in the number of cells making up the tissue (hyperplasia), or both. Certain organs, such as the heart, lose the ability to divide shortly after birth. Accordingly, "cardiac hypertrophy" is defined as an increase in mass of the heart, which, in adults, is characterized by an increase in myocyte cell size and contractile protein content without concomitant cell division. The character of the stress responsible for inciting the hypertrophy, (*e.g.*, increased preload, increased afterload, loss of myocytes, as in myocardial infarction, or primary depression of contractility), appears to play a critical role in determining the nature of the response. The early stage of cardiac hypertrophy is usually characterized morphologically by increases in the size of myofibrils and mitochondria, as well as by enlargement of mitochondria and nuclei. At this stage, while muscle cells are larger than normal, cellular organization is largely preserved. At a more advanced stage of cardiac hypertrophy, there are preferential increases in the size or number of specific organelles, such as mitochondria, and new contractile elements are added in localized areas of the cells, in an irregular manner. Cells subjected to long-standing hypertrophy show more obvious disruptions in cellular organization, including markedly enlarged nuclei with highly lobulated membranes, which displace adjacent myofibrils and cause breakdown of normal Z-band registration. The phrase "cardiac hypertrophy" is used to include all stages of the progression of this condition, characterized by various degrees of structural damage of the heart muscle, regardless of the underlying cardiac disorder. Hence, the term also includes physiological conditions instrumental in the development of cardiac hypertrophy, such as elevated blood pressure, aortic stenosis, or myocardial infarction.

"Heart failure" refers to an abnormality of cardiac function where the heart does not pump blood at the rate

needed for the requirements of metabolizing tissues. The heart failure can be caused by a number of factors, including ischemic, congenital, rheumatic, or idiopathic forms.

“Congestive heart failure” (CHF) is a progressive pathologic state where the heart is increasingly unable to supply adequate cardiac output (the volume of blood pumped by the heart over time) to deliver the oxygenated blood to peripheral tissues. As CHF progresses, structural and hemodynamic damages occur. While these damages have a variety of manifestations, one characteristic symptom is ventricular hypertrophy. CHF is a common end result of a number of various cardiac disorders.

“Myocardial infarction” generally results from atherosclerosis of the coronary arteries, often with superimposed coronary thrombosis. It may be divided into two major types: transmural infarcts, in which myocardial necrosis involves the full thickness of the ventricular wall, and subendocardial (nontransmural) infarcts, in which the necrosis involves the subendocardium, the intramural myocardium, or both, without extending all the way through the ventricular wall to the epicardium. Myocardial infarction is known to cause both a change in hemodynamic effects and an alteration in structure in the damaged and healthy zones of the heart. Thus, for example, myocardial infarction reduces the maximum cardiac output and the stroke volume of the heart. Also associated with myocardial infarction is a stimulation of the DNA synthesis occurring in the interstice as well as an increase in the formation of collagen in the areas of the heart not affected.

As a result of the increased stress or strain placed on the heart in prolonged hypertension due, for example, to the increased total peripheral resistance, cardiac hypertrophy has long been associated with “hypertension”. A characteristic of the ventricle that becomes hypertrophic as a result of chronic pressure overload is an impaired diastolic performance. Fouad *et al.*, J. Am. Coll. Cardiol., 4: 1500-1506 (1984); Smith *et al.*, J. Am. Coll. Cardiol., 5: 869-874 (1985). A prolonged left ventricular relaxation has been detected in early essential hypertension, in spite of normal or supranormal systolic function. Hartford *et al.*, Hypertension, 6: 329-338 (1984). However, there is no close parallelism between blood pressure levels and cardiac hypertrophy. Although improvement in left ventricular function in response to antihypertensive therapy has been reported in humans, patients variously treated with a diuretic (hydrochlorothiazide), a β -blocker (propranolol), or a calcium channel blocker (diltiazem), have shown reversal of left ventricular hypertrophy, without improvement in diastolic function. Inouye *et al.*, Am. J. Cardiol., 53: 1583-7 (1984).

Another complex cardiac disease associated with cardiac hypertrophy is “hypertrophic cardiomyopathy”. This condition is characterized by a great diversity of morphologic, functional, and clinical features (Maron *et al.*, N. Engl. J. Med., 316: 780-789 (1987); Spirito *et al.*, N. Engl. J. Med., 320: 749-755 (1989); Louie and Edwards, Prog. Cardiovasc. Dis., 36: 275-308 (1994); Wigle *et al.*, Circulation, 92: 1680-1692 (1995)), the heterogeneity of which is accentuated by the fact that it afflicts patients of all ages. Spirito *et al.*, N. Engl. J. Med., 336: 775-785 (1997). The causative factors of hypertrophic cardiomyopathy are also diverse and little understood. In general, mutations in genes encoding sarcomeric proteins are associated with hypertrophic cardiomyopathy. Recent data suggest that β -myosin heavy chain mutations may account for approximately 30 to 40 percent of cases of familial hypertrophic cardiomyopathy. Watkins *et al.*, N. Engl. J. Med., 326: 1108-1114 (1992); Schwartz *et al.*, Circulation, 91: 532-540 (1995); Marian and Roberts, Circulation, 92: 1336-1347 (1995); Thierfelder *et al.*, Cell, 77: 701-712

(1994); Watkins *et al.*, Nat. Gen., 11: 434-437 (1995). Besides β -myosin heavy chain, other locations of genetic mutations include cardiac troponin T, alpha topomyosin, cardiac myosin binding protein C, essential myosin light chain, and regulatory myosin light chain. See, Malik and Watkins, Curr. Opin. Cardiol., 12: 295-302 (1997).

5 Supravalvular "aortic stenosis" is an inherited vascular disorder characterized by narrowing of the ascending aorta, but other arteries, including the pulmonary arteries, may also be affected. Untreated aortic stenosis may lead to increased intracardiac pressure resulting in myocardial hypertrophy and eventually heart failure and death. The pathogenesis of this disorder is not fully understood, but hypertrophy and possibly hyperplasia of medial smooth muscle are prominent features of this disorder. It has been reported that molecular variants of the elastin gene are involved in the development and pathogenesis of aortic stenosis. U.S. Patent No. 5,650,282 issued July 10 22, 1997.

 "Valvular regurgitation" occurs as a result of heart diseases resulting in disorders of the cardiac valves. Various diseases, like rheumatic fever, can cause the shrinking or pulling apart of the valve orifice, while other diseases may result in endocarditis, an inflammation of the endocardium or lining membrane of the atrioventricular orifices and operation of the heart. Defects such as the narrowing of the valve stenosis or the defective closing of 15 the valve result in an accumulation of blood in the heart cavity or regurgitation of blood past the valve. If uncorrected, prolonged valvular stenosis or insufficiency may result in cardiac hypertrophy and associated damage to the heart muscle, which may eventually necessitate valve replacement.

 The treatment of all these, and other cardiovascular, endothelial and angiogenic disorders, which may or may not be accompanied by cardiac hypertrophy, is encompassed by the present invention.

20 The terms "cancer", "cancerous", and "malignant" refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth. Examples of cancer include but are not limited to, carcinoma including adenocarcinoma, lymphoma, blastoma, melanoma, sarcoma, and leukemia. More particular examples of such cancers include squamous cell cancer, small-cell lung cancer, non-small cell lung cancer, gastrointestinal cancer, Hodgkin's and non-Hodgkin's lymphoma, pancreatic cancer, glioblastoma, cervical cancer, 25 ovarian cancer, liver cancer such as hepatic carcinoma and hepatoma, bladder cancer, breast cancer, colon cancer, colorectal cancer, endometrial carcinoma, salivary gland carcinoma, kidney cancer such as renal cell carcinoma and Wilms' tumors, basal cell carcinoma, melanoma, prostate cancer, vulval cancer, thyroid cancer, testicular cancer, esophageal cancer, and various types of head and neck cancer. The preferred cancers for treatment herein are breast, colon, lung, melanoma, ovarian, and others involving vascular tumors as noted above.

30 The term "cytotoxic agent" as used herein refers to a substance that inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes (*e.g.*, ^{131}I , ^{125}I , ^{90}Y , and ^{186}Re), chemotherapeutic agents, and toxins such as enzymatically active toxins of bacterial, fungal, plant, or animal origin, or fragments thereof.

35 A "chemotherapeutic agent" is a chemical compound useful in the treatment of cancer. Examples of chemotherapeutic agents include alkylating agents, folic acid antagonists, anti-metabolites of nucleic acid metabolism, antibiotics, pyrimidine analogs, 5-fluorouracil, cisplatin, purine nucleosides, amines, amino acids, triazol nucleosides, or corticosteroids. Specific examples include Adriamycin, Doxorubicin, 5-Fluorouracil,

Cytosine arabinoside ("Ara-C"), Cyclophosphamide, Thiotepa, Busulfan, Cytosin, Taxol, Toxotere, Methotrexate, Cisplatin, Melphalan, Vinblastine, Bleomycin, Etoposide, Ifosfamide, Mitomycin C, Mitoxantrone, Vincristine, Vinorelbine, Carboplatin, Teniposide, Daunomycin, Carminomycin, Aminopterin, Dactinomycin, Mitomycins, Esperamicins (see U.S. Pat. No. 4,675,187), Melphalan, and other related nitrogen mustards. Also included in this definition are hormonal agents that act to regulate or inhibit hormone action on tumors, such as tamoxifen and onapristone.

A "growth-inhibitory agent" when used herein refers to a compound or composition that inhibits growth of a cell, such as an Wnt-overexpressing cancer cell, either *in vitro* or *in vivo*. Thus, the growth-inhibitory agent is one which significantly reduces the percentage of malignant cells in S phase. Examples of growth-inhibitory agents include agents that block cell cycle progression (at a place other than S phase), such as agents that induce G1 arrest and M-phase arrest. Classical M-phase blockers include the vincas (vincristine and vinblastine), taxol, and topo II inhibitors such as doxorubicin, daunorubicin, etoposide, and bleomycin. Those agents that arrest G1 also spill over into S-phase arrest, for example, DNA alkylating agents such as tamoxifen, prednisone, dacarbazine, mechlorethamine, cisplatin, methotrexate, 5-fluorouracil, and ara-C. Further information can be found in The Molecular Basis of Cancer, Mendelsohn and Israel, eds., Chapter 1, entitled "Cell cycle regulation, oncogenes, and antineoplastic drugs" by Murakami *et al.* (WB Saunders: Philadelphia, 1995), especially p. 13. Additional examples include tumor necrosis factor (TNF), an antibody capable of inhibiting or neutralizing the angiogenic activity of acidic or basic FGF or hepatocyte growth factor (HGF), an antibody capable of inhibiting or neutralizing the coagulant activities of tissue factor, protein C, or protein S (see, WO 91/01753, published 21 February 1991), or an antibody capable of binding to HER2 receptor (WO 89/06692), such as the 4D5 antibody (and functional equivalents thereof) (e.g., WO 92/22653).

"Treatment" is an intervention performed with the intention of preventing the development or altering the pathology of a cardiovascular, endothelial, and angiogenic disorder. The concept of treatment is used in the broadest sense, and specifically includes the prevention (prophylaxis), moderation, reduction, and curing of cardiovascular, endothelial, and angiogenic disorders of any stage. Accordingly, "treatment" refers to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) or ameliorate a cardiovascular, endothelial, and angiogenic disorder such as hypertrophy. Those in need of treatment include those already with the disorder as well as those prone to have the disorder or those in whom the disorder is to be prevented. The disorder may result from any cause, including idiopathic, cardiopathic, or myotrophic causes, or ischemia or ischemic insults, such as myocardial infarction.

"Chronic" administration refers to administration of the agent(s) in a continuous mode as opposed to an acute mode, so as to maintain the initial effect, such as an anti-hypertrophic effect, for an extended period of time.

"Mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, horses, cats, cows, sheep, pigs, etc. Preferably, the mammal is human.

Administration "in combination with" one or more further therapeutic agents includes simultaneous (concurrent) and consecutive administration in any order.

The phrase "cardiovascular, endothelial or angiogenic agents" refers generically to any drug that acts in treating cardiovascular, endothelial, and angiogenic disorders. Examples of cardiovascular agents are those that promote vascular homeostasis by modulating blood pressure, heart rate, heart contractility, and endothelial and smooth muscle biology, all of which factors have a role in cardiovascular disease. Specific examples of these include angiotensin-II receptor antagonists; endothelin receptor antagonists such as, for example, BOSENTANTM and MOXONODINTM; interferon-gamma (IFN- γ); des-aspartate-angiotensin I; thrombolytic agents, *e.g.*, streptokinase, urokinase, t-PA, and a t-PA variant specifically designed to have longer half-life and very high fibrin specificity, TNK t-PA (a T103N, N117Q, KHRR(296-299)AAAA t-PA variant, Keyt *et al.*, Proc. Natl. Acad. Sci. USA, 91: 3670-3674 (1994)); inotropic or hypertensive agents such as digoxigenin and β -adrenergic receptor blocking agents, *e.g.*, propranolol, timolol, tertalolol, carteolol, nadolol, betaxolol, penbutolol, acetobutolol, atenolol, metoprolol, and carvedilol; angiotensin converting enzyme (ACE) inhibitors, *e.g.*, quinapril, captopril, enalapril, ramipril, benazepril, fosinopril, and lisinopril; diuretics, *e.g.*, chlorothiazide, hydrochlorothiazide, hydroflumethazide, methylchlorothiazide, benzthiazide, dichlorphenamide, acetazolamide, and indapamide; and calcium channel blockers, *e.g.*, diltiazem, nifedipine, verapamil, nicardipine. One preferred category of this type is a therapeutic agent used for the treatment of cardiac hypertrophy or of a physiological condition instrumental in the development of cardiac hypertrophy, such as elevated blood pressure, aortic stenosis, or myocardial infarction.

"Angiogenic agents" and "endothelial agents" are active agents that promote angiogenesis and/or endothelial cell growth, or, if applicable, vasculogenesis. This would include factors that accelerate wound healing, such as growth hormone, insulin-like growth factor-I (IGF-I), VEGF, VIGF, PDGF, epidermal growth factor (EGF), CTGF and members of its family, FGF, and TGF- α and TGF- β .

"Angiostatic agents" are active agents that inhibit angiogenesis or vasculogenesis or otherwise inhibit or prevent growth of cancer cells. Examples include antibodies or other antagonists to angiogenic agents as defined above, such as antibodies to VEGF. They additionally include cytotherapeutic agents such as cytotoxic agents, chemotherapeutic agents, growth-inhibitory agents, apoptotic agents, and other agents to treat cancer, such as anti-HER-2, anti-CD20, and other bioactive and organic chemical agents.

In a pharmacological sense, in the context of the present invention, a "therapeutically effective amount" of an active agent such as a PRO polypeptide or agonist or antagonist thereto or an anti-PRO antibody, refers to an amount effective in the treatment of a cardiovascular, endothelial or angiogenic disorder in a mammal and can be determined empirically.

As used herein, an "effective amount" of an active agent such as a PRO polypeptide or agonist or antagonist thereto or an anti-PRO antibody, refers to an amount effective for carrying out a stated purpose, wherein such amounts may be determined empirically for the desired effect.

The terms "PRO polypeptide" and "PRO" as used herein and when immediately followed by a numerical designation refer to various polypeptides, wherein the complete designation (*i.e.*, PRO/number) refers to specific polypeptide sequences as described herein. The terms "PRO/number polypeptide" and "PRO/number" wherein the term "number" is provided as an actual numerical designation as used herein encompass native sequence polypeptides and polypeptide variants (which are further defined herein). The PRO polypeptides described herein

may be isolated from a variety of sources, such as from human tissue types or from another source, or prepared by recombinant or synthetic methods.

5 A "native sequence PRO polypeptide" comprises a polypeptide having the same amino acid sequence as the corresponding PRO polypeptide derived from nature. Such native sequence PRO polypeptides can be isolated from nature or can be produced by recombinant or synthetic means. The term "native sequence PRO polypeptide" specifically encompasses naturally-occurring truncated or secreted forms of the specific PRO polypeptide (e.g., an extracellular domain sequence), naturally-occurring variant forms (e.g., alternatively spliced forms) and naturally-occurring allelic variants of the polypeptide. In various embodiments of the invention, the native sequence PRO polypeptides disclosed herein are mature or full-length native sequence polypeptides comprising the full-length amino acids sequences shown in the accompanying figures. Start and stop codons are shown in bold font and underlined in the figures. However, while the PRO polypeptide disclosed in the accompanying figures are shown to begin with methionine residues designated herein as amino acid position 1 in the figures, it is conceivable and possible that other methionine residues located either upstream or downstream from the amino acid position 1 in the figures may be employed as the starting amino acid residue for the PRO polypeptides.

15 The PRO polypeptide "extracellular domain" or "ECD" refers to a form of the PRO polypeptide which is essentially free of the transmembrane and cytoplasmic domains. Ordinarily, a PRO polypeptide ECD will have less than 1% of such transmembrane and/or cytoplasmic domains and preferably, will have less than 0.5% of such domains. It will be understood that any transmembrane domains identified for the PRO polypeptides of the present invention are identified pursuant to criteria routinely employed in the art for identifying that type of hydrophobic domain. The exact boundaries of a transmembrane domain may vary but most likely by no more than about 5 amino acids at either end of the domain as initially identified herein. Optionally, therefore, an extracellular domain of a PRO polypeptide may contain from about 5 or fewer amino acids on either side of the transmembrane domain/extracellular domain boundary as identified in the Examples or specification and such polypeptides, with or without the associated signal peptide, and nucleic acid encoding them, are contemplated by the present invention.

25 The approximate location of the "signal peptides" of the various PRO polypeptides disclosed herein are shown in the present specification and/or the accompanying figures. It is noted, however, that the C-terminal boundary of a signal peptide may vary, but most likely by no more than about 5 amino acids on either side of the signal peptide C-terminal boundary as initially identified herein, wherein the C-terminal boundary of the signal peptide may be identified pursuant to criteria routinely employed in the art for identifying that type of amino acid sequence element (e.g., Nielsen *et al.*, Prot. Eng., 10:1-6 (1997) and von Heinje *et al.*, Nucl. Acids Res., 14:4683-4690 (1986)). Moreover, it is also recognized that, in some cases, cleavage of a signal sequence from a secreted polypeptide is not entirely uniform, resulting in more than one secreted species. These mature polypeptides, where the signal peptide is cleaved within no more than about 5 amino acids on either side of the C-terminal boundary of the signal peptide as identified herein, and the polynucleotides encoding them, are contemplated by the present invention.

35 "PRO polypeptide variant" means an active PRO polypeptide as defined above or below having at least about 80% amino acid sequence identity with a full-length native sequence PRO polypeptide sequence as disclosed

herein, a PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Such PRO polypeptide variants include, for instance, PRO polypeptides wherein one or more amino acid residues are added, or deleted, at the N- or C-terminus of the full-length native amino acid sequence. Ordinarily, a PRO polypeptide variant will have at least about 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% or 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to a full-length native sequence PRO polypeptide sequence as disclosed herein, a PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of a full-length PRO polypeptide sequence as disclosed herein. Ordinarily, PRO variant polypeptides are at least about 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 150 or 200 amino acids in length and alternatively at least about 300 amino acids in length, or more.

"Percent (%) amino acid sequence identity" with respect to the PRO polypeptide sequences identified herein is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in a PRO sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN, ALIGN-2 or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full-length of the sequences being compared. For purposes herein, however, % amino acid sequence identity values are obtained as described below by using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc., and the source code shown in Table 1 has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, California or may be compiled from the source code provided in Table 1. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

For purposes herein, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

$$100 \text{ times the fraction } X/Y$$

where X is the number of amino acid residues scored as identical matches by the sequence alignment program

ALIGN-2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A. As examples of % amino acid sequence identity calculations, Tables 2-3 demonstrate how to calculate the % amino acid sequence identity of the amino acid sequence designated "Comparison Protein" to the amino acid sequence designated "PRO".

Unless specifically stated otherwise, all % amino acid sequence identity values used herein are obtained as described above using the ALIGN-2 sequence comparison computer program. However, % amino acid sequence identity may also be determined using the sequence comparison program NCBI-BLAST2 (Altschul *et al.*, Nucleic Acids Res., 25:3389-3402 (1997)). The NCBI-BLAST2 sequence comparison program may be downloaded from <http://www.ncbi.nlm.nih.gov> or otherwise obtained from the National Institute of Health, Bethesda, MD. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask = yes, strand = all, expected occurrences = 10, minimum low complexity length = 15/5, multi-pass e-value = 0.01, constant for multi-pass = 25, dropoff for final gapped alignment = 25 and scoring matrix = BLOSUM62.

In situations where NCBI-BLAST2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

$$100 \text{ times the fraction } X/Y$$

where X is the number of amino acid residues scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A.

In addition, % amino acid sequence identity may also be determined using the WU-BLAST-2 computer program (Altschul *et al.*, Methods in Enzymology, 266:460-480 (1996)). Most of the WU-BLAST-2 search parameters are set to the default values. Those not set to default values, *i.e.*, the adjustable parameters, are set with the following values: overlap span = 1, overlap fraction = 0.125, word threshold (T) = 11, and scoring matrix = BLOSUM62. For purposes herein, a % amino acid sequence identity value is determined by dividing (a) the number of matching identical amino acids residues between the amino acid sequence of the PRO polypeptide of interest having a sequence derived from the native PRO polypeptide and the comparison amino acid sequence of interest (*i.e.*, the sequence against which the PRO polypeptide of interest is being compared which may be a PRO variant polypeptide) as determined by WU-BLAST-2 by (b) the total number of amino acid residues of the PRO polypeptide of interest. For example, in the statement "a polypeptide comprising an amino acid sequence A which

has or having at least 80% amino acid sequence identity to the amino acid sequence B", the amino acid sequence A is the comparison amino acid sequence of interest and the amino acid sequence B is the amino acid sequence of the PRO polypeptide of interest.

5 "PRO variant polynucleotide" or "PRO variant nucleic acid sequence" means a nucleic acid molecule which encodes an active PRO polypeptide as defined below and which has at least about 80% nucleic acid sequence identity with a nucleotide acid sequence encoding a full-length native sequence PRO polypeptide sequence as disclosed herein, a full-length native sequence PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Ordinarily, a PRO variant
10 polynucleotide will have at least about 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% or 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity with a nucleic acid sequence encoding a full-length native sequence PRO polypeptide sequence as disclosed herein, a full-length native sequence PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal sequence, as disclosed herein or
15 any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Variants do not encompass the native nucleotide sequence.

Ordinarily, PRO variant polynucleotides are at least about 30, 60, 90, 120, 150, 180, 210, 240, 270, 300, 450, or 600 nucleotides in length and alternatively at least about 900 nucleotides in length, or more.

"Percent (%) nucleic acid sequence identity" with respect to the PRO polypeptide-encoding nucleic acid
20 sequences identified herein is defined as the percentage of nucleotides in a candidate sequence that are identical with the nucleotides in a PRO polypeptide-encoding nucleic acid sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity. Alignment for purposes of determining percent nucleic acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN, ALIGN-2 or Megalign
25 (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full-length of the sequences being compared. For purposes herein, however, % nucleic acid sequence identity values are obtained as described below by using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1. The ALIGN-2 sequence comparison computer program was authored by
30 Genentech, Inc., and the source code shown in Table 1 has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, California or may be compiled from the source code provided in Table 1. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2
35 program and do not vary.

For purposes herein, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that

has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

100 times the fraction W/Z

5

where W is the number of nucleotides scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C. As examples of %
10 nucleic acid sequence identity calculations, Tables 4-5 demonstrate how to calculate the % nucleic acid sequence identity of the nucleic acid sequence designated "Comparison DNA" to the nucleic acid sequence designated "PRO-DNA".

Unless specifically stated otherwise, all % nucleic acid sequence identity values used herein are obtained as described above using the ALIGN-2 sequence comparison computer program. However, % nucleic acid
15 sequence identity may also be determined using the sequence comparison program NCBI-BLAST2 (Altschul *et al.*, Nucleic Acids Res., 25:3389-3402 (1997)). The NCBI-BLAST2 sequence comparison program may be downloaded from <http://www.ncbi.nlm.nih.gov> or otherwise obtained from the National Institute of Health, Bethesda, MD. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask = yes, strand = all, expected occurrences = 10, minimum low
20 complexity length = 15/5, multi-pass e-value = 0.01, constant for multi-pass = 25, dropoff for final gapped alignment = 25 and scoring matrix = BLOSUM62.

In situations where NCBI-BLAST2 is employed for sequence comparisons, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence
25 identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

100 times the fraction W/Z

where W is the number of nucleotides scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C.
30

In addition, % nucleic acid sequence identity values may also be generated using the WU-BLAST-2 computer program (Altschul *et al.*, Methods in Enzymology, 266:460-480 (1996)). Most of the WU-BLAST-2 search parameters are set to the default values. Those not set to default values, *i.e.*, the adjustable parameters, are set with the following values: overlap span = 1, overlap fraction = 0.125, word threshold (T) = 11, and scoring
35 matrix = BLOSUM62. For purposes herein, a % nucleic acid sequence identity value is determined by dividing (a)

the number of matching identical nucleotides between the nucleic acid sequence of the PRO polypeptide-encoding nucleic acid molecule of interest having a sequence derived from the native sequence PRO polypeptide-encoding nucleic acid and the comparison nucleic acid molecule of interest (i.e., the sequence against which the PRO polypeptide-encoding nucleic acid molecule of interest is being compared which may be a variant PRO polynucleotide) as determined by WU-BLAST-2 by (b) the total number of nucleotides of the PRO polypeptide-encoding nucleic acid molecule of interest. For example, in the statement "an isolated nucleic acid molecule comprising a nucleic acid sequence A which has or having at least 80% nucleic acid sequence identity to the nucleic acid sequence B", the nucleic acid sequence A is the comparison nucleic acid molecule of interest and the nucleic acid sequence B is the nucleic acid sequence of the PRO polypeptide-encoding nucleic acid molecule of interest.

In other embodiments, PRO variant polynucleotides are nucleic acid molecules that encode an active PRO polypeptide and which are capable of hybridizing, preferably under stringent hybridization and wash conditions, to nucleotide sequences encoding the full-length PRO polypeptide as shown in the specification herein and accompanying figures. PRO variant polypeptides may be those that are encoded by a PRO variant polynucleotide.

"Isolated", when used to describe the various polypeptides disclosed herein, means a polypeptide that has been identified and separated and/or recovered from a component of its natural environment. Preferably, the isolated polypeptide is free of association with all components with which it is naturally associated. Contaminant components of its natural environment are materials that would typically interfere with diagnostic or therapeutic uses for the polypeptide, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the polypeptide will be purified (1) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (2) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Isolated polypeptide includes polypeptide *in situ* within recombinant cells, since at least one component of the PRO natural environment will not be present. Ordinarily, however, isolated polypeptide will be prepared by at least one purification step.

An "isolated" nucleic acid molecule encoding a PRO polypeptide or an "isolated" nucleic acid molecule encoding an anti-PRO antibody is a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the natural source of the PRO-encoding nucleic acid or the natural source of the anti-PRO-encoding nucleic acid. Preferably, the isolated nucleic acid is free of association with all components with which it is naturally associated. An isolated PRO-encoding nucleic acid molecule or an isolated anti-PRO-encoding nucleic acid molecule is other than in the form or setting in which it is found in nature. Isolated nucleic acid molecules therefore are distinguished from the PRO-encoding nucleic acid molecule or from the anti-PRO-encoding nucleic acid molecule as it exists in natural cells. However, an isolated nucleic acid molecule encoding a PRO polypeptide or an isolated nucleic acid molecule encoding an anti-PRO antibody includes PRO-nucleic acid molecules or anti-PRO-nucleic acid molecules contained in cells that ordinarily express PRO polypeptides or anti-PRO antibodies where, for example, the nucleic acid molecule is in a chromosomal location different from that of natural cells.

The term "control sequences" refers to DNA sequences necessary for the expression of an operably linked

coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize, for example, promoters, polyadenylation signals, and enhancers.

5 Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a PRO polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, 10 contiguous and in the same reading frame. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

"Stringency" of hybridization reactions is readily determinable by one of ordinary skill in the art, and generally is an empirical calculation dependent upon probe length, washing temperature, and salt concentration. 15 In general, longer probes require higher temperatures for proper annealing, while shorter probes need lower temperatures. Hybridization generally depends on the ability of denatured DNA to reanneal when complementary strands are present in an environment below their melting temperature. The higher the degree of desired homology between the probe and hybridizable sequence, the higher the relative temperature that can be used. As a result, it follows that higher relative temperatures would tend to make the reaction conditions more stringent, while lower 20 temperatures less so. For additional details and explanation of stringency of hybridization reactions, *see*, Ausubel *et al.*, Current Protocols in Molecular Biology (Wiley Interscience Publishers, 1995).

"Stringent conditions" or "high-stringency conditions", as defined herein, may be identified by those that: (1) employ low ionic strength and high temperature for washing, for example, 0.015 M sodium chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate at 50°C; (2) employ during hybridization a denaturing agent, such as 25 formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50mM sodium phosphate buffer at pH 6.5 with 750 mM sodium chloride, 75 mM sodium citrate at 42°C; or (3) employ 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2 x SSC (sodium chloride/sodium citrate) and 50% formamide at 55°C, followed by a high-stringency wash consisting of 0.1 x SSC containing EDTA 30 at 55°C.

"Moderately-stringent conditions" may be identified as described by Sambrook *et al.*, Molecular Cloning: A Laboratory Manual (New York: Cold Spring Harbor Press, 1989), and include the use of washing solution and hybridization conditions (*e.g.*, temperature, ionic strength, and % SDS) less stringent than those described above. 35 An example of moderately stringent conditions is overnight incubation at 37°C in a solution comprising: 20% formamide, 5 x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5 x Denhardt's solution, 10% dextran sulfate, and 20 mg/ml denatured sheared salmon sperm DNA, followed by washing the filters

in 1 x SSC at about 37-50°C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc. as necessary to accommodate factors such as probe length and the like.

The modifier "epitope-tagged" when used herein refers to a chimeric polypeptide comprising a PRO polypeptide fused to a "tag polypeptide". The tag polypeptide has enough residues to provide an epitope against which an antibody can be made, yet is short enough such that it does not interfere with activity of the polypeptide to which it is fused. The tag polypeptide preferably also is fairly unique so that the antibody does not substantially cross-react with other epitopes. Suitable tag polypeptides generally have at least six amino acid residues and usually between about 8 and 50 amino acid residues (preferably, between about 10 and 20 amino acid residues).

"Active" or "activity" in the context of PRO variants refers to form(s) of PRO proteins that retain the biologic and/or immunologic activities of a native or naturally-occurring PRO polypeptide.

"Biological activity" in the context of a molecule that antagonizes a PRO polypeptide that can be identified by the screening assays disclosed herein (*e.g.*, an organic or inorganic small molecule, peptide, etc.) is used to refer to the ability of such molecules to bind or complex with the PRO polypeptide identified herein, or otherwise interfere with the interaction of the PRO polypeptide with other cellular proteins or otherwise inhibits the transcription or translation of the PRO polypeptide. Particularly preferred biological activity includes cardiac hypertrophy, activity that acts on systemic disorders that affect vessels, such as diabetes mellitus, as well as diseases of the arteries, capillaries, veins, and/or lymphatics, and cancer.

The term "antagonist" is used in the broadest sense, and includes any molecule that partially or fully blocks, inhibits, or neutralizes one or more of the biological activities of a native PRO polypeptide disclosed herein, for example, if applicable, its mitogenic or angiogenic activity. Antagonists of a PRO polypeptide may act by interfering with the binding of a PRO polypeptide to a cellular receptor, by incapacitating or killing cells that have been activated by a PRO polypeptide, or by interfering with vascular endothelial cell activation after binding of a PRO polypeptide to a cellular receptor. All such points of intervention by a PRO polypeptide antagonist shall be considered equivalent for purposes of this invention. The antagonists inhibit the mitogenic, angiogenic, or other biological activity of PRO polypeptides, and thus are useful for the treatment of diseases or disorders characterized by undesirable excessive neovascularization, including by way of example tumors, and especially solid malignant tumors, rheumatoid arthritis, psoriasis, atherosclerosis, diabetic and other retinopathies, retrolental fibroplasia, age-related macular degeneration, neovascular glaucoma, hemangiomas, thyroid hyperplasias (including Grave's disease), corneal and other tissue transplantation, and chronic inflammation. The antagonists also are useful for the treatment of diseases or disorders characterized by undesirable excessive vascular permeability, such as edema associated with brain tumors, ascites associated with malignancies, Meigs' syndrome, lung inflammation, nephrotic syndrome, pericardial effusion (such as that associated with pericarditis), and pleural effusion. In a similar manner, the term "agonist" is used in the broadest sense and includes any molecule that mimics a biological activity of a native PRO polypeptide disclosed herein. Suitable agonist or antagonist molecules specifically include agonist or antagonist antibodies or antibody fragments, fragments, or amino acid sequence variants of native PRO polypeptides, peptides, small organic molecules, etc.

A "small molecule" is defined herein to have a molecular weight below about 500 daltons.

The term "PRO polypeptide receptor" as used herein refers to a cellular receptor for a PRO polypeptide, ordinarily a cell-surface receptor found on vascular endothelial cells, as well as variants thereof that retain the ability to bind a PRO polypeptide.

"Antibodies" (Abs) and "immunoglobulins" (Igs) are glycoproteins having the same structural characteristics. While antibodies exhibit binding specificity to a specific antigen, immunoglobulins include both antibodies and other antibody-like molecules that lack antigen specificity. Polypeptides of the latter kind are, for example, produced at low levels by the lymph system and at increased levels by myelomas. The term "antibody" is used in the broadest sense and specifically covers, without limitation, intact monoclonal antibodies, polyclonal antibodies, multispecific antibodies (*e.g.*, bispecific antibodies) formed from at least two intact antibodies, and antibody fragments, so long as they exhibit the desired biological activity.

"Native antibodies" and "native immunoglobulins" are usually heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies among the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at one end a variable domain (V_H) followed by a number of constant domains. Each light chain has a variable domain at one end (V_L) and a constant domain at its other end; the constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light-chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface between the light- and heavy-chain variable domains.

The term "variable" refers to the fact that certain portions of the variable domains differ extensively in sequence among antibodies and are used in the binding and specificity of each particular antibody to and for its particular antigen. However, the variability is not evenly distributed throughout the variable domains of antibodies. It is concentrated in three segments called complementarity-determining regions (CDRs) or hypervariable regions both in the light-chain and the heavy-chain variable domains. The more highly conserved portions of variable domains are called the framework regions (FR). The variable domains of native heavy and light chains each comprise four FR regions, largely adopting a β -sheet configuration, connected by three CDRs, which form loops connecting, and in some cases forming part of, the β -sheet structure. The CDRs in each chain are held together in close proximity by the FR regions and, with the CDRs from the other chain, contribute to the formation of the antigen-binding site of antibodies. See, Kabat *et al.*, NIH Publ. No.91-3242, Vol. I, pages 647-669 (1991). The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody-dependent cellular toxicity.

"Antibody fragments" comprise a portion of an intact antibody, preferably the antigen-binding or variable region of the intact antibody. Examples of antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments; diabodies; linear antibodies (Zapata *et al.*, Protein Eng., **8**(10): 1057-1062 (1995)); single-chain antibody molecules; and multispecific antibodies formed from antibody fragments.

Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, each with a single antigen-binding site, and a residual "Fc" fragment, whose name reflects its ability to crystallize

readily. Pepsin treatment yields an $F(ab')_2$ fragment that has two antigen-combining sites and is still capable of cross-linking antigen.

"Fv" is the minimum antibody fragment that contains a complete antigen-recognition and -binding site. This region consists of a dimer of one heavy- and one light-chain variable domain in tight, non-covalent association. It is in this configuration that the three CDRs of each variable domain interact to define an antigen-binding site on the surface of the V_H - V_L dimer. Collectively, the six CDRs confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab' fragments differ from Fab fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. $F(ab')_2$ antibody fragments originally were produced as pairs of Fab' fragments that have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

The "light chains" of antibodies (immunoglobulins) from any vertebrate species can be assigned to one of two clearly distinct types, called kappa (κ) and lambda (λ), based on the amino acid sequences of their constant domains.

Depending on the amino acid sequence of the constant domain of their heavy chains, immunoglobulins can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM; and several of these may be further divided into subclasses (isotypes), e.g., IgG1, IgG2, IgG3, IgG4, IgA, and IgA2. The heavy-chain constant domains that correspond to the different classes of immunoglobulins are called α , δ , ϵ , γ , and μ , respectively. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known.

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally-occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to conventional (polyclonal) antibody preparations that typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they are synthesized by the hybridoma culture, uncontaminated by other immunoglobulins. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler *et al.*, Nature, 256: 495 (1975), or may be made by recombinant DNA methods (see, e.g., U.S. Patent No. 4,816,567). The "monoclonal antibodies" may also be isolated from phage antibody libraries using the techniques described in Clackson *et al.*, Nature, 352: 624-628 (1991) and Marks *et al.*, J. Mol. Biol., 222: 581-597 (1991), for example.

The monoclonal antibodies herein specifically include "chimeric" antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity. U.S. Patent No. 4,816,567; Morrison *et al.*, Proc. Natl. Acad. Sci. USA, **81**: 6851-6855 (1984).

"Humanized" forms of non-human (*e.g.*, murine) antibodies are chimeric immunoglobulins, immunoglobulin chains, or fragments thereof (such as Fv, Fab, Fab', F(ab')₂, or other antigen-binding subsequences of antibodies) that contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a CDR of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity, and capacity. In some instances, Fv FR residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues that are found neither in the recipient antibody nor in the imported CDR or framework sequences. These modifications are made to further refine and maximize antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin sequence. The humanized antibody preferably also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones *et al.*, Nature, **321**: 522-525 (1986); Reichmann *et al.*, Nature, **332**: 323-329 (1988); and Presta, Curr. Op. Struct. Biol., **2**: 593-596 (1992). The humanized antibody includes a PRIMATIZED™ antibody wherein the antigen-binding region of the antibody is derived from an antibody produced by immunizing macaque monkeys with the antigen of interest.

"Single-chain Fv" or "sFv" antibody fragments comprise the V_H and V_L domains of an antibody, wherein these domains are present in a single polypeptide chain. Preferably, the Fv polypeptide further comprises a polypeptide linker between the V_H and V_L domains that enables the sFv to form the desired structure for antigen binding. For a review of sFv see, Pluckthun in The Pharmacology of Monoclonal Antibodies, Vol. 113, Rosenberg and Moore, eds. (Springer-Verlag: New York, 1994), pp. 269-315.

The term "diabodies" refers to small antibody fragments with two antigen-binding sites, which fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) in the same polypeptide chain (V_H - V_L). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger *et al.*, Proc. Natl. Acad. Sci. USA, **90**: 6444-6448 (1993).

An "isolated" antibody is one that has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would interfere

with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, the antibody will be purified (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the antibody *in situ* within recombinant cells, since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

An antibody that "specifically binds to" or is "specific for" a particular polypeptide or an epitope on a particular polypeptide is one that binds to that particular polypeptide or epitope on a particular polypeptide without substantially binding to any other polypeptide or polypeptide epitope.

The word "label" when used herein refers to a detectable compound or other composition that is conjugated directly or indirectly to the antibody so as to generate a "labeled" antibody. The label may be detectable by itself (e.g., radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition that is detectable. Radionuclides that can serve as detectable labels include, for example, I-131, I-123, I-125, Y-90, Re-188, At-211, Cu-67, Bi-212, and Pd-109. The label may also be a non-detectable entity such as a toxin.

By "solid phase" is meant a non-aqueous matrix to which an antibody of the present invention can adhere. Examples of solid phases encompassed herein include those formed partially or entirely of glass (e.g., controlled pore glass), polysaccharides (e.g., agarose), polyacrylamides, polystyrene, polyvinyl alcohol and silicones. In certain embodiments, depending on the context, the solid phase can comprise the well of an assay plate; in others it is a purification column (e.g., an affinity chromatography column). This term also includes a discontinuous solid phase of discrete particles, such as those described in U.S. Patent No. 4,275,149.

A "liposome" is a small vesicle composed of various types of lipids, phospholipids and/or surfactant that is useful for delivery of a drug (such as the PRO polypeptide or antibodies thereto disclosed herein) to a mammal. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes.

As used herein, the term "immunoadhesin" designates antibody-like molecules that combine the binding specificity of a heterologous protein (an "adhesin") with the effector functions of immunoglobulin constant domains. Structurally, the immunoadhesins comprise a fusion of an amino acid sequence with the desired binding specificity that is other than the antigen recognition and binding site of an antibody (i.e., is "heterologous"), and an immunoglobulin constant domain sequence. The adhesin part of an immunoadhesin molecule typically is a contiguous amino acid sequence comprising at least the binding site of a receptor or a ligand. The immunoglobulin constant domain sequence in the immunoadhesin may be obtained from any immunoglobulin, such as IgG-1, IgG-2, IgG-3, or IgG-4 subtypes, IgA (including IgA-1 and IgA-2), IgE, IgD, or IgM.

As shown below, Table 1 provides the complete source code for the ALIGN-2 sequence comparison computer program. This source code may be routinely compiled for use on a UNIX operating system to provide

the ALIGN-2 sequence comparison computer program.

5 In addition, Tables 2-5 show hypothetical exemplifications for using the below described method to determine % amino acid sequence identity (Tables 2-3) and % nucleic acid sequence identity (Tables 4-5) using the ALIGN-2 sequence comparison computer program, wherein "PRO" represents the amino acid sequence of a
10 hypothetical PRO polypeptide of interest, "Comparison Protein" represents the amino acid sequence of a polypeptide against which the "PRO" polypeptide of interest is being compared, "PRO-DNA" represents a hypothetical PRO-encoding nucleic acid sequence of interest, "Comparison DNA" represents the nucleotide sequence of a nucleic acid molecule against which the "PRO-DNA" nucleic acid molecule of interest is being compared, "X", "Y", and "Z" each represent different hypothetical amino acid residues and "N", "L" and "V" each represent different hypothetical nucleotides.

Table 1

```

/*
 *
 * C-C increased from 12 to 15
 * Z is average of EQ
 * B is average of ND
 * match with stop is _M; stop-stop = 0; J (joker) match = 0
 */
#define _M      -8      /* value of a match with a stop */

int _day[26][26] = {
/* A B C D E F G H I J K L M N O P Q R S T U V W X Y Z */
/* A */ { 2, 0, -2, 0, 0, 4, 1, -1, -1, 0, -1, -2, -1, 0, _M, 1, 0, -2, 1, 1, 0, 0, -6, 0, -3, 0},
/* B */ { 0, 3, -4, 3, 2, -5, 0, 1, -2, 0, 0, -3, -2, 2, _M, -1, 1, 0, 0, 0, 0, -2, -5, 0, -3, 1},
/* C */ { -2, -4, 15, -5, -5, -4, -3, -3, -2, 0, -5, -6, -5, -4, _M, -3, -5, -4, 0, -2, 0, -2, -8, 0, 0, -5},
/* D */ { 0, 3, -5, 4, 3, -6, 1, 1, -2, 0, 0, -4, -3, 2, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 2},
/* E */ { 0, 2, -5, 3, 4, -5, 0, 1, -2, 0, 0, -3, -2, 1, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 3},
/* F */ { -4, -5, -4, -6, -5, 9, -5, -2, 1, 0, -5, 2, 0, -4, _M, -5, -5, -4, -3, -3, 0, -1, 0, 0, 7, -5},
/* G */ { 1, 0, -3, 1, 0, -5, 5, -2, -3, 0, -2, -4, -3, 0, _M, -1, -1, -3, 1, 0, 0, -1, -7, 0, -5, 0},
/* H */ { -1, 1, -3, 1, 1, -2, -2, 6, -2, 0, 0, -2, -2, 2, _M, 0, 3, 2, -1, -1, 0, -2, -3, 0, 0, 2},
/* I */ { -1, -2, -2, -2, -2, 1, -3, -2, 5, 0, -2, 2, 2, -2, _M, -2, -2, -2, -1, 0, 0, 4, -5, 0, -1, -2},
/* J */ { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* K */ { -1, 0, -5, 0, 0, -5, -2, 0, -2, 0, 5, -3, 0, 1, _M, -1, 1, 3, 0, 0, 0, -2, -3, 0, -4, 0},
/* L */ { -2, -3, -6, -4, -3, 2, -4, -2, 2, 0, -3, 6, 4, -3, _M, -3, -2, -3, -3, -1, 0, 2, -2, 0, -1, -2},
/* M */ { -1, -2, -5, -3, -2, 0, -3, -2, 2, 0, 0, 4, 6, -2, _M, -2, -1, 0, -2, -1, 0, 2, -4, 0, -2, -1},
/* N */ { 0, 2, -4, 2, 1, -4, 0, 2, -2, 0, 1, -3, -2, 2, _M, -1, 1, 0, 1, 0, 0, -2, -4, 0, -2, 1},
/* O */ { _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, 0, _M, _M, _M, _M, _M, _M, _M, _M, _M},
/* P */ { 1, -1, -3, -1, -1, -5, -1, 0, -2, 0, -1, -3, -2, -1, _M, 6, 0, 0, 1, 0, 0, -1, -6, 0, -5, 0},
/* Q */ { 0, 1, -5, 2, 2, -5, -1, 3, -2, 0, 1, -2, -1, 1, _M, 0, 4, 1, -1, -1, 0, -2, -5, 0, -4, 3},
/* R */ { -2, 0, -4, -1, -1, -4, -3, 2, -2, 0, 3, -3, 0, 0, _M, 0, 1, 6, 0, -1, 0, -2, 2, 0, -4, 0},
/* S */ { 1, 0, 0, 0, 0, -3, 1, -1, -1, 0, 0, -3, -2, 1, _M, 1, -1, 0, 2, 1, 0, -1, -2, 0, -3, 0},
/* T */ { 1, 0, -2, 0, 0, -3, 0, -1, 0, 0, 0, -1, -1, 0, _M, 0, -1, -1, 1, 3, 0, 0, -5, 0, -3, 0},
/* U */ { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* V */ { 0, -2, -2, -2, -1, -1, -2, 4, 0, -2, 2, 2, -2, _M, -1, -2, -2, -1, 0, 0, 4, -6, 0, -2, -2},
/* W */ { -6, -5, -8, -7, -7, 0, -7, -3, -5, 0, -3, -2, -4, -4, _M, -6, -5, 2, -2, -5, 0, -6, 17, 0, 0, -6},
/* X */ { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* Y */ { -3, -3, 0, -4, -4, 7, -5, 0, -1, 0, -4, -1, -2, -2, _M, -5, -4, -4, -3, -3, 0, -2, 0, 0, 10, -4},
/* Z */ { 0, 1, -5, 2, 3, -5, 0, 2, -2, 0, 0, -2, -1, 1, _M, 0, 3, 0, 0, 0, 0, -2, -6, 0, -4, 4}
};

```

Table 1 (cont')

```

/*
*/
#include <stdio.h>
#include <ctype.h>

#define MAXJMP      16      /* max jumps in a diag */
#define MAXGAP      24      /* don't continue to penalize gaps larger than this */
#define JMPS        1024    /* max jmps in an path */
#define MX          4       /* save if there's at least MX-1 bases since last jmp */

#define DMAT         3       /* value of matching bases */
#define DMIS         0       /* penalty for mismatched bases */
#define DINS0        8       /* penalty for a gap */
#define DINS1         1       /* penalty per base */
#define PINS0         8       /* penalty for a gap */
#define PINS1         4       /* penalty per residue */

struct jmp {
    short      n[MAXJMP];    /* size of jmp (neg for dely) */
    unsigned short x[MAXJMP]; /* base no. of jmp in seq x */
};
/* limits seq to 216-1 */

struct diag {
    int      score;          /* score at last jmp */
    long     offset;         /* offset of prev block */
    short    ijmp;          /* current jmp index */
    struct jmp jp;           /* list of jmps */
};

struct path {
    int      spc;            /* number of leading spaces */
    short    n[JMPS];        /* size of jmp (gap) */
    int      x[JMPS];        /* loc of jmp (last elem before gap) */
};

char      *ofile;           /* output file name */
char      *namex[2];        /* seq names: getseqs( ) */
char      *prog;            /* prog name for err msgs */
char      *seqx[2];         /* seqs: getseqs( ) */
int      dmax;              /* best diag: nw( ) */
int      dmax0;             /* final diag */
int      dna;               /* set if dna: main( ) */
int      endgaps;           /* set if penalizing end gaps */
int      gapx, gapy;        /* total gaps in seqs */
int      len0, len1;        /* seq lens */
int      ngapx, ngapy;      /* total size of gaps */
int      smax;              /* max score: nw( ) */
int      *xbm;              /* bitmap for matching */
long     offset;            /* current offset in jmp file */
struct    diag              /* holds diagonals */
struct    path              /* holds path for seqs */
    *dx;
    pp[2];

char      *calloc( ), *malloc( ), *index( ), *strcpy( );
char      *getseq( ), *g_calloc( );

```

Table 1 (cont')

```

/* Needleman-Wunsch alignment program
*
* usage: progs file1 file2
* where file1 and file2 are two dna or two protein sequences.
* The sequences can be in upper- or lower-case and may contain ambiguity
* Any lines beginning with ';', '>' or '<' are ignored
* Max file length is 65535 (limited by unsigned short x in the jmp struct)
* A sequence with 1/3 or more of its elements ACGTU is assumed to be DNA
* Output is in the file "align.out"
*
* The program may create a tmp file in /tmp to hold info about traceback.
* Original version developed under BSD 4.3 on a vax 8650
*/
#include "nw.h"
#include "day.h"

static  _dbval[26] = {
1,14,2,13,0,0,4,11,0,0,12,0,3,15,0,0,0,5,6,8,8,7,9,0,10,0
};

static  _pbval[26] = {
1, 2|(1<<('D'-'A'))|(1<<('N'-'A')), 4, 8, 16, 32, 64,
128, 256, 0xFFFFFFFF, 1<<10, 1<<11, 1<<12, 1<<13, 1<<14,
1<<15, 1<<16, 1<<17, 1<<18, 1<<19, 1<<20, 1<<21, 1<<22,
1<<23, 1<<24, 1<<25|(1<<('E'-'A'))|(1<<('Q'-'A'))
};

main(ac, av)                                main
{
    int      ac;
    char     *av[];

    prog = av[0];
    if (ac != 3) {
        fprintf(stderr, "usage: %s file1 file2\n", prog);
        fprintf(stderr, "where file1 and file2 are two dna or two protein sequences.\n");
        fprintf(stderr, "The sequences can be in upper- or lower-case\n");
        fprintf(stderr, "Any lines beginning with ';', '>' or '<' are ignored\n");
        fprintf(stderr, "Output is in the file \"align.out\"\n");
        exit(1);
    }
    namex[0] = av[1];
    namex[1] = av[2];
    seqx[0] = getseq(namex[0], &len0);
    seqx[1] = getseq(namex[1], &len1);
    xbm = (dna)? _dbval : _pbval;

    endgaps = 0;                                /* 1 to penalize endgaps */
    ofile = "align.out";                        /* output file */

    nw( );                                /* fill in the matrix, get the possible jumps */
    readjumps( );                            /* get the actual jumps */
    print( );                                /* print stats, alignment */

    cleanup(0);                                /* unlink any tmp files */
}

```

Table 1 (cont')

```

/* do the alignment, return best score: main( )
* dna: values in Fitch and Smith, PNAS, 80, 1382-1386, 1983
* pro: PAM 250 values
* When scores are equal, we prefer mismatches to any gap, prefer
* a new gap to extending an ongoing gap, and prefer a gap in seqx
* to a gap in seq y.
*/
nw( )
{
    char      *px, *py;           /* seqs and ptrs */
    int       *ndely, *dely;      /* keep track of dely */
    int       ndelx, delx;        /* keep track of delx */
    int       *tmp;              /* for swapping row0, row1 */
    int       mis;               /* score for each type */
    int       ins0, ins1;        /* insertion penalties */
    register  id;                /* diagonal index */
    register  ij;                /* jmp index */
    register  *col0, *col1;       /* score for curr, last row */
    register  xx, yy;            /* index into seqs */

    dx = (struct diag *)g_calloc("to get diags", len0+len1+1, sizeof(struct diag));

    ndely = (int *)g_calloc("to get ndely", len1+1, sizeof(int));
    dely = (int *)g_calloc("to get dely", len1+1, sizeof(int));
    col0 = (int *)g_calloc("to get col0", len1+1, sizeof(int));
    col1 = (int *)g_calloc("to get col1", len1+1, sizeof(int));
    ins0 = (dna)? DINS0 : PINS0;
    ins1 = (dna)? DINS1 : PINS1;

    smax = -10000;
    if (endgaps) {
        for (col0[0] = dely[0] = -ins0, yy = 1; yy <= len1; yy++) {
            col0[yy] = dely[yy] = col0[yy-1] - ins1;
            ndely[yy] = yy;
        }
        col0[0] = 0;          /* Waterman Bull Math Biol 84 */
    }
    else
        for (yy = 1; yy <= len1; yy++)
            dely[yy] = -ins0;

    /* fill in match matrix
    */
    for (px = seqx[0], xx = 1; xx <= len0; px++, xx++) {
        /* initialize first entry in col
        */
        if (endgaps) {
            if (xx == 1)
                col1[0] = delx = -(ins0+ins1);
            else
                col1[0] = delx = col0[0] - ins1;
            ndelx = xx;
        }
        else {
            col1[0] = 0;
            delx = -ins0;
            ndelx = 0;
        }
    }
}

```

nw

Table 1 (cont')

...nw

```

for (py = seqx[1], yy = 1; yy <= len1; py++, yy++) {
    mis = col0[yy-1];
    if (dna)
        mis += (xbm[*px-'A']&xbm[*py-'A'])? DMAT : DMIS;
    else
        mis += _day[*px-'A'][*py-'A'];

    /* update penalty for del in x seq;
     * favor new del over ongong del
     * ignore MAXGAP if weighting endgaps
     */
    if (endgaps || ndely[yy] < MAXGAP) {
        if (col0[yy] - ins0 >= dely[yy]) {
            dely[yy] = col0[yy] - (ins0+ins1);
            ndely[yy] = 1;
        } else {
            dely[yy] -= ins1;
            ndely[yy]++;
        }
    } else {
        if (col0[yy] - (ins0+ins1) >= dely[yy]) {
            dely[yy] = col0[yy] - (ins0+ins1);
            ndely[yy] = 1;
        } else
            ndely[yy]++;
    }

    /* update penalty for del in y seq;
     * favor new del over ongong del
     */
    if (endgaps || ndelx < MAXGAP) {
        if (col1[yy-1] - ins0 >= delx) {
            delx = col1[yy-1] - (ins0+ins1);
            ndelx = 1;
        } else {
            delx -= ins1;
            ndelx++;
        }
    } else {
        if (col1[yy-1] - (ins0+ins1) >= delx) {
            delx = col1[yy-1] - (ins0+ins1);
            ndelx = 1;
        } else
            ndelx++;
    }

    /* pick the maximum score; we're favoring
     * mis over any del and delx over dely
     */

```

Table 1 (cont')

...nw

```

id = xx - yy + len1 - 1;
if (mis >= delx && mis >= dely[yy])
    coll[yy] = mis;
else if (delx >= dely[yy]) {
    coll[yy] = delx;
    ij = dx[id].ijmp;
    if (dx[id].jp.n[0] && (!dna || (ndelx >= MAXJMP
    && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
        dx[id].ijmp++;
        if (++ij >= MAXJMP) {
            writeimps(id);
            ij = dx[id].ijmp = 0;
            dx[id].offset = offset;
            offset += sizeof(struct jmp) + sizeof(offset);
        }
        dx[id].jp.n[ij] = ndelx;
        dx[id].jp.x[ij] = xx;
        dx[id].score = delx;
    }
} else {
    coll[yy] = dely[yy];
    ij = dx[id].ijmp;
    if (dx[id].jp.n[0] && (!dna || (ndely[yy] >= MAXJMP
    && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
        dx[id].ijmp++;
        if (++ij >= MAXJMP) {
            writeimps(id);
            ij = dx[id].ijmp = 0;
            dx[id].offset = offset;
            offset += sizeof(struct jmp) + sizeof(offset);
        }
        dx[id].jp.n[ij] = -ndely[yy];
        dx[id].jp.x[ij] = xx;
        dx[id].score = dely[yy];
    }
}
if (xx == len0 && yy < len1) {
    /* last col
    */
    if (endgaps)
        coll[yy] -= ins0+ins1*(len1-yy);
    if (coll[yy] > smax) {
        smax = coll[yy];
        dmax = id;
    }
}
if (endgaps && xx < len0)
    coll[yy-1] -= ins0+ins1*(len0-xx);
if (coll[yy-1] > smax) {
    smax = coll[yy-1];
    dmax = id;
}
tmp = col0; col0 = coll; coll = tmp;
}
(void) free((char *)ndely);
(void) free((char *)dely);
(void) free((char *)col0);
(void) free((char *)coll);
}

```


Table 1 (cont')

```

/*
 *
 * print( ) -- only routine visible outside this module
 *
 * static:
 * getmat( ) -- trace back best path, count matches: print( )
 * pr_align( ) -- print alignment of described in array p[]: print( )
 * dumpblock( ) -- dump a block of lines with numbers, stars: pr_align( )
 * nums( ) -- put out a number line: dumpblock( )
 * putline( ) -- put out a line (name, [num], seq, [num]): dumpblock( )
 * stars( ) -- put a line of stars: dumpblock( )
 * stripname( ) -- strip any path and prefix from a seqname
 */

#include "nw.h"

#define SPC      3
#define P_LINE  256    /* maximum output line */
#define P_SPC    3      /* space between name or num and seq */

extern _day[26][26];
int      olen;          /* set output line length */
FILE     *fx;           /* output file */

print( )
{
    int      lx, ly, firstgap, lastgap;    /* overlap */

    if ((fx = fopen(ofile, "w")) == 0) {
        fprintf(stderr, "%s: can't write %s\n", prog, ofile);
        cleanup(1);
    }
    fprintf(fx, "< first sequence: %s (length = %d)\n", namex[0], len0);
    fprintf(fx, "< second sequence: %s (length = %d)\n", namex[1], len1);
    olen = 60;
    lx = len0;
    ly = len1;
    firstgap = lastgap = 0;
    if (dmax < len1 - 1) { /* leading gap in x */
        pp[0].spc = firstgap = len1 - dmax - 1;
        ly -= pp[0].spc;
    }
    else if (dmax > len1 - 1) { /* leading gap in y */
        pp[1].spc = firstgap = dmax - (len1 - 1);
        lx -= pp[1].spc;
    }
    if (dmax0 < len0 - 1) { /* trailing gap in x */
        lastgap = len0 - dmax0 - 1;
        lx -= lastgap;
    }
    else if (dmax0 > len0 - 1) { /* trailing gap in y */
        lastgap = dmax0 - (len0 - 1);
        ly -= lastgap;
    }
    getmat(lx, ly, firstgap, lastgap);
    pr_align( );
}

```

print

Table 1 (cont')

```

/*
 * trace back the best path, count matches
 */
static
getmat(lx, ly, firstgap, lastgap)                                getmat
{
    int      lx, ly;                                           /* "core" (minus endgaps) */
    int      firstgap, lastgap;                                /* leading trailing overlap */

    int      nm, i0, i1, siz0, siz1;
    char     outx[32];
    double   pct;
    register n0, n1;
    register char *p0, *p1;

    /* get total matches, score
     */
    i0 = i1 = siz0 = siz1 = 0;
    p0 = seqx[0] + pp[1].spc;
    p1 = seqx[1] + pp[0].spc;
    n0 = pp[1].spc + 1;
    n1 = pp[0].spc + 1;

    nm = 0;
    while ( *p0 && *p1 ) {
        if (siz0) {
            p1++;
            n1++;
            siz0--;
        }
        else if (siz1) {
            p0++;
            n0++;
            siz1--;
        }
        else {
            if (xbm[*p0-'A'] & xbm[*p1-'A'])
                nm++;
            if (n0++ == pp[0].x[i0])
                siz0 = pp[0].n[i0++];
            if (n1++ == pp[1].x[i1])
                siz1 = pp[1].n[i1++];
            p0++;
            p1++;
        }
    }

    /* pct homology:
     * if penalizing endgaps, base is the shorter seq
     * else, knock off overhangs and take shorter core
     */
    if (endgaps)
        lx = (len0 < len1)? len0 : len1;
    else
        lx = (lx < ly)? lx : ly;
    pct = 100.*((double)nm/(double)lx;
    fprintf(fx, "\n");
    fprintf(fx, "< %d match%s in an overlap of %d: %.2f percent similarity\n",
        nm, (nm == 1)? "" : "es", lx, pct);

```

Table 1 (cont')

```

fprintf(fx, "< gaps in first sequence: %d", gapx);
if (gapx) {
    (void) sprintf(outx, " (%d %s%s)",
        ngapx, (dna)? "base": "residue", (ngapx == 1)? "" : "s");
    fprintf(fx, "%s", outx);

    fprintf(fx, ", gaps in second sequence: %d", gapy);
    if (gapy) {
        (void) sprintf(outx, " (%d %s%s)",
            ngapy, (dna)? "base": "residue", (ngapy == 1)? "" : "s");
        fprintf(fx, "%s", outx);
    }
    if (dna)
        fprintf(fx,
            "\n< score: %d (match = %d, mismatch = %d, gap penalty = %d + %d per base)\n",
            smax, DMAT, DMIS, DINSO, DINS1);
    else
        fprintf(fx,
            "\n< score: %d (Dayhoff PAM 250 matrix, gap penalty = %d + %d per residue)\n",
            smax, PINSO, PINS1);
    if (endgaps)
        fprintf(fx,
            "< endgaps penalized. left endgap: %d %s%s, right endgap: %d %s%s\n",
            firstgap, (dna)? "base" : "residue", (firstgap == 1)? "" : "s",
            lastgap, (dna)? "base" : "residue", (lastgap == 1)? "" : "s");
    else
        fprintf(fx, "< endgaps not penalized\n");
}

static      nm;          /* matches in core -- for checking */
static      lmax;        /* lengths of stripped file names */
static      ij[2];       /* jmp index for a path */
static      nc[2];       /* number at start of current line */
static      ni[2];       /* current elem number -- for gapping */
static      siz[2];
static char *ps[2];      /* ptr to current element */
static char *po[2];      /* ptr to next output char slot */
static char out[2][P_LINE]; /* output line */
static char star[P_LINE]; /* set by stars() */

/*
 * print alignment of described in struct path pp[]
 */
static
pr_align()
{
    int      nn;          /* char count */
    int      more;
    register i;

    for (i = 0, lmax = 0; i < 2; i++) {
        nn = stripname(namex[i]);
        if (nn > lmax)
            lmax = nn;

        nc[i] = 1;
        ni[i] = 1;
        siz[i] = ij[i] = 0;
        ps[i] = seqx[i];
        po[i] = out[i];
    }
}

```

...getmat

pr_align

Table 1 (cont')

```

for (nn = nm = 0, more = 1; more; ) {
    for (i = more = 0; i < 2; i++) {
        /*
         * do we have more of this sequence?
         */
        if (!*ps[i])
            continue;

        more++;

        if (pp[i].spc) { /* leading space */
            *po[i]++ = ' ';
            pp[i].spc--;
        }
        else if (siz[i]) { /* in a gap */
            *po[i]++ = '-';
            siz[i]--;
        }
        else { /* we're putting a seq element
            */
            *po[i] = *ps[i];
            if (islower(*ps[i]))
                *ps[i] = toupper(*ps[i]);
            po[i]++;
            ps[i]++;

            /*
             * are we at next gap for this seq?
             */
            if (ni[i] == pp[i].x[ij[i]]) {
                /*
                 * we need to merge all gaps
                 * at this location
                 */
                siz[i] = pp[i].n[ij[i]] + +;
                while (ni[i] == pp[i].x[ij[i]])
                    siz[i] += pp[i].n[ij[i]] + +;
            }
            ni[i]++;
        }
    }
    if (++nn == olen || !more && nn) {
        dumpblock();
        for (i = 0; i < 2; i++)
            po[i] = out[i];
        nn = 0;
    }
}

/*
 * dump a block of lines, including numbers, stars: pr_align( )
 */
static
dumpblock( )
{
    register i;

    for (i = 0; i < 2; i++)
        *po[i] = '\0';
}

```

...pr_align

dumpblock

Table 1 (cont')

...dumpblock

```

(void) putc('\n', fx);
for (i = 0; i < 2; i++) {
    if (*out[i] && (*out[i] != ' ' || *(po[i]) != ' ')) {
        if (i == 0)
            nums(i);
        if (i == 0 && *out[1])
            stars( );
        putline(i);
        if (i == 0 && *out[1])
            fprintf(fx, star);
        if (i == 1)
            nums(i);
    }
}

/*
 * put out a number line: dumpblock( )
 */
static
nums(ix)
int    ix;    /* index in out[] holding seq line */
{
    char    nline[P_LINE];
    register i, j;
    register char    *pn, *px, *py;

    for (pn = nline, i = 0; i < lmax+P_SPC; i++, pn++)
        *pn = ' ';
    for (i = nc[ix], py = out[ix]; *py; py++, pn++) {
        if (*py == ' ' || *py == '-')
            *pn = ' ';
        else {
            if (i%10 == 0 || (i == 1 && nc[ix] != 1)) {
                j = (i < 0)? -i : i;
                for (px = pn; j /= 10, px--)
                    *px = j%10 + '0';
                if (i < 0)
                    *px = '-';
            }
            else
                *pn = ' ';
            i++;
        }
    }
    *pn = '\0';
    nc[ix] = i;
    for (pn = nline; *pn; pn++)
        (void) putc(*pn, fx);
    (void) putc('\n', fx);
}

/*
 * put out a line (name, [num], seq, [num]): dumpblock( )
 */
static
putline(ix)
int    ix;
{
    putline

```

Table 1 (cont')

```

int          i;
register char *px;

for (px = namex[ix], i = 0; *px && *px != ':'; px++, i++)
    (void) putc(*px, fx);
for (; i < lmax+P_SPC; i++)
    (void) putc(' ', fx);

/* these count from 1:
 * ni[] is current element (from 1)
 * nc[] is number at start of current line
 */
for (px = out[ix]; *px; px++)
    (void) putc(*px&0x7F, fx);
(void) putc('\n', fx);
}

/*
 * put a line of stars (seqs always in out[0], out[1]): dumpblock( )
 */
static
stars( )
{
    int          i;
    register char *p0, *p1, cx, *px;

    if (!(*out[0] || (*out[0] == ' ' && *(po[0]) == ' ') ||
        !(*out[1] || (*out[1] == ' ' && *(po[1]) == ' ')))
        return;
    px = star;
    for (i = lmax+P_SPC; i; i--)
        *px++ = ' ';

    for (p0 = out[0], p1 = out[1]; *p0 && *p1; p0++, p1++) {
        if (isalpha(*p0) && isalpha(*p1)) {
            if (xbm[*p0-'A']&xbm[*p1-'A']) {
                cx = '*';
                nm++;
            }
            else if (!dna && _day[*p0-'A'][*p1-'A'] > 0)
                cx = '.';
            else
                cx = ' ';
        }
        else
            cx = ' ';
        *px++ = cx;
    }
    *px++ = '\n';
    *px = '\0';
}

```

...putline

stars

Table 1 (cont')

```

/*
 * strip path or prefix from pn, return len: pr_align( )
 */
static
stripname(pn)
char *pn; /* file name (may be path) */
{
    register char *px, *py;

    py = 0;
    for (px = pn; *px; px++)
        if (*px == '/')
            py = px + 1;
    if (py)
        (void) strcpy(pn, py);
    return(strlen(pn));
}

```

stripname

Table 1 (cont')

```

/*
 * cleanup( ) -- cleanup any tmp file
 * getseq( ) -- read in seq, set dna, len, maxlen
 * g_calloc( ) -- calloc( ) with error checkin
 * readjumps( ) -- get the good jumps, from tmp file if necessary
 * writejumps( ) -- write a filled array of jumps to a tmp file: nw( )
 */
#include "nw.h"
#include <sys/file.h>

char    *jname = "/tmp/homgXXXXXX";          /* tmp file for jumps */
FILE    *fj;

int      cleanup( );                          /* cleanup tmp file */
long     lseek( );

/*
 * remove any tmp file if we blow
 */
cleanup(i)
    int    i;
{
    if (fj)
        (void) unlink(jname);
    exit(i);
}

/*
 * read, return ptr to seq, set dna, len, maxlen
 * skip lines starting with ';', '<', or '>'
 * seq in upper or lower case
 */
char    *
getseq(file, len)
    char    *file;    /* file name */
    int     *len;     /* seq len */
{
    char    line[1024], *pseq;
    register char    *px, *py;
    int     natgc, tlen;
    FILE    *fp;

    if ((fp = fopen(file, "r")) == 0) {
        fprintf(stderr, "%s: can't read %s\n", prog, file);
        exit(1);
    }
    tlen = natgc = 0;
    while (fgets(line, 1024, fp)) {
        if (*line == ';' || *line == '<' || *line == '>')
            continue;
        for (px = line; *px != '\n'; px++)
            if (isupper(*px) || islower(*px))
                tlen++;
    }
    if ((pseq = malloc((unsigned)(tlen+6))) == 0) {
        fprintf(stderr, "%s: malloc( ) failed to get %d bytes for %s\n", prog, tlen+6, file);
        exit(1);
    }
    pseq[0] = pseq[1] = pseq[2] = pseq[3] = '\0';
}

```

cleanup

getseq

Table 1 (cont')

...getseq

```

py = pseq + 4;
*len = tlen;
rewind(fp);

while (fgets(line, 1024, fp)) {
    if (*line == ';' || *line == '<' || *line == '>')
        continue;
    for (px = line; *px != '\n'; px++) {
        if (isupper(*px))
            *py++ = *px;
        else if (islower(*px))
            *py++ = toupper(*px);
        if (index("ATGCU", *(py-1)))
            natgc++;
    }
    *py++ = '\0';
    *py = '\0';
    (void) fclose(fp);
    dna = natgc > (tlen/3);
    return(pseq+4);
}

```

```

char *
g_calloc(msg, nx, sz)
char *msg;          /* program, calling routine */
int nx, sz;          /* number and size of elements */
{
    char *px, *calloc();

    if ((px = calloc((unsigned)nx, (unsigned)sz)) == 0) {
        if (*msg) {
            fprintf(stderr, "%s: g_calloc() failed %s (n=%d, sz=%d)\n", prog, msg, nx, sz);
            exit(1);
        }
    }
    return(px);
}

```

g_calloc

```

/*
 * get final jmps from dx[] or tmp file, set pp[], reset dmax: main( )
 */

```

```

readjumps()
{
    int fd = -1;
    int siz, i0, i1;
    register i, j, xx;

    if (fj) {
        (void) fclose(fj);
        if ((fd = open(jname, O_RDONLY, 0)) < 0) {
            fprintf(stderr, "%s: can't open() %s\n", prog, jname);
            cleanup(1);
        }
    }
    for (i = i0 = i1 = 0, dmax0 = dmax, xx = len0; i++) {
        while (1) {
            for (j = dx[dmax].ijmp; j >= 0 && dx[dmax].jp.x[j] >= xx; j--)
                ;
        }
    }
}

```

readjumps

Table 1 (cont')

...readjumps

```

    if (j < 0 && dx[dmax].offset && fj) {
        (void) lseek(fd, dx[dmax].offset, 0);
        (void) read(fd, (char *)&dx[dmax].jp, sizeof(struct jmp));
        (void) read(fd, (char *)&dx[dmax].offset, sizeof(dx[dmax].offset));
        dx[dmax].ijmp = MAXJMP-1;
    }
    else
        break;
}
if (i >= JMPS) {
    fprintf(stderr, "%s: too many gaps in alignment\n", prog);
    cleanup(1);
}
if (j >= 0) {
    siz = dx[dmax].jp.n[j];
    xx = dx[dmax].jp.x[j];
    dmax += siz;
    if (siz < 0) { /* gap in second seq */
        pp[1].n[i1] = -siz;
        xx += siz;
        /* id = xx - yy + len1 - 1 */
        /*
        pp[1].x[i1] = xx - dmax + len1 - 1;
        gapy++;
        ngapy -= siz;
        */
        /* ignore MAXGAP when doing endgaps */
        siz = (-siz < MAXGAP || endgaps)? -siz : MAXGAP;
        i1++;
    }
    else if (siz > 0) { /* gap in first seq */
        pp[0].n[i0] = siz;
        pp[0].x[i0] = xx;
        gapx++;
        ngapx += siz;
        /* ignore MAXGAP when doing endgaps */
        siz = (siz < MAXGAP || endgaps)? siz : MAXGAP;
        i0++;
    }
}
else
    break;
}

/* reverse the order of jumps */
*/
for (j = 0, i0--; j < i0; j++, i0--) {
    i = pp[0].n[j]; pp[0].n[j] = pp[0].n[i0]; pp[0].n[i0] = i;
    i = pp[0].x[j]; pp[0].x[j] = pp[0].x[i0]; pp[0].x[i0] = i;
}
for (j = 0, i1--; j < i1; j++, i1--) {
    i = pp[1].n[j]; pp[1].n[j] = pp[1].n[i1]; pp[1].n[i1] = i;
    i = pp[1].x[j]; pp[1].x[j] = pp[1].x[i1]; pp[1].x[i1] = i;
}
if (fd >= 0)
    (void) close(fd);
if (fj) {
    (void) unlink(jname);
    fj = 0;
    offset = 0;
}
}

```

Table 1 (cont')

```

/*
 * write a filled jmp struct offset of the prev one (if any): nw( )
 */
writejumps(ix)
{
    int ix;
    char *mktemp( );

    if (!fj) {
        if (mktemp(jname) < 0) {
            fprintf(stderr, "%s: can't mktemp( ) %s\n", prog, jname);
            cleanup(1);
        }
        if ((fj = fopen(jname, "w")) == 0) {
            fprintf(stderr, "%s: can't write %s\n", prog, jname);
            exit(1);
        }
    }
    (void) fwrite((char *)&dx[ix].jp, sizeof(struct jmp), 1, fj);
    (void) fwrite((char *)&dx[ix].offset, sizeof(dx[ix].offset), 1, fj);
}

```

writejumps

Table 2

PRO	XXXXXXXXXXXXXXXXXX	(Length = 15 amino acids)
Comparison Protein	XXXXXXYYYYYYY	(Length = 12 amino acids)

5 % amino acid sequence identity =
 (the number of identically matching amino acid residues between the two polypeptide sequences as determined by
 ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) =
 5 divided by 15 = 33.3%

10 Table 3

PRO	XXXXXXXXXXXX	(Length = 10 amino acids)
Comparison Protein	XXXXXXYYYYYYZZYZ	(Length = 15 amino acids)

15 % amino acid sequence identity =
 (the number of identically matching amino acid residues between the two polypeptide sequences as determined by
 ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) =
 5 divided by 10 = 50%

20 Table 4

PRO-DNA	NNNNNNNNNNNNNN	(Length = 14 nucleotides)
Comparison DNA	NNNNNNLLLLLLLLLL	(Length = 16 nucleotides)

25 % nucleic acid sequence identity =
 (the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-
 2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) =
 6 divided by 14 = 42.9%

30 Table 5

PRO-DNA	NNNNNNNNNNNN	(Length = 12 nucleotides)
Comparison DNA	NNNNLLL VV	(Length = 9 nucleotides)

35 % nucleic acid sequence identity =
 (the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-
 2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) =
 4 divided by 12 = 33.3%

5.2. Compositions and Methods of the Invention

5.2.1. PRO Variants

5 In addition to the full-length native sequence PRO polypeptides described herein, it is contemplated that PRO variants can be prepared. PRO variants can be prepared by introducing appropriate nucleotide changes into the PRO DNA, and/or by synthesis of the desired PRO polypeptide. Those skilled in the art will appreciate that amino acid changes may alter post-translational processes of the PRO polypeptide such as changing the number or position of glycosylation sites or altering the membrane anchoring characteristics.

10 Variations in the native full-length sequence PRO polypeptide or in various domains of the PRO polypeptide described herein, can be made, for example, using any of the techniques and guidelines for conservative and non-conservative mutations set forth, for instance, in U.S. Patent No. 5,364,934. Variations may be a substitution, deletion or insertion of one or more codons encoding the PRO polypeptide that results in a change in the amino acid sequence of the PRO polypeptide as compared with the native sequence PRO polypeptide. Optionally the variation is by substitution of at least one amino acid with any other amino acid in one or more of the domains of the PRO polypeptide. Guidance in determining which amino acid residue may be inserted, substituted or deleted without adversely affecting the desired activity may be found by comparing the sequence of the PRO polypeptide with that of homologous known protein molecules and minimizing the number of amino acid sequence changes made in regions of high homology. Amino acid substitutions can be the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, such as the replacement of a leucine with a serine, *i.e.*, conservative amino acid replacements. Insertions or deletions may optionally be in the range of about 15 1 to 5 amino acids. The variation allowed may be determined by systematically making insertions, deletions or substitutions of amino acids in the sequence and testing the resulting variants for activity exhibited by the full-length or mature native sequence.

20 In particular embodiments, conservative substitutions of interest are shown in Table 6 under the heading of preferred substitutions. If such substitutions result in a change in biological activity, then more substantial changes, denominated exemplary substitutions in Table 6, or as further described below in reference to amino acid classes, are introduced and the products screened.

25

Table 6

	Original <u>Residue</u>	Exemplary <u>Substitutions</u>	Preferred <u>Substitutions</u>
	Ala (A)	val; leu; ile	val
5	Arg (R)	lys; gln; asn	lys
	Asn (N)	gln; his; lys; arg	gln
	Asp (D)	glu	glu
	Cys (C)	ser	ser
	Gln (Q)	asn	asn
10	Glu (E)	asp	asp
	Gly (G)	pro; ala	ala
	His (H)	asn; gln; lys; arg	arg
	Ile (I)	leu; val; met; ala; phe; norleucine	leu
15	Leu (L)	norleucine; ile; val; met; ala; phe	ile
	Lys (K)	arg; gln; asn	arg
	Met (M)	leu; phe; ile	leu
	Phe (F)	leu; val; ile; ala; tyr	leu
20	Pro (P)	ala	ala
	Ser (S)	thr	thr
	Thr (T)	ser	ser
	Trp (W)	tyr; phe	tyr
	Tyr (Y)	trp; phe; thr; ser	phe
25	Val (V)	ile; leu; met; phe; ala; norleucine	leu

Substantial modifications in function or immunological identity of the PRO polypeptide are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties:

- (1) hydrophobic: norleucine, met, ala, val, leu, ile;
- (2) neutral hydrophilic: cys, ser, thr;
- (3) acidic: asp, glu;
- 35 (4) basic: asn, gln, his, lys, arg;
- (5) residues that influence chain orientation: gly, pro; and
- (6) aromatic: trp, tyr, phe.

Non-conservative substitutions will entail exchanging a member of one of these classes for another class. Such substituted residues also may be introduced into the conservative substitution sites or, more preferably, into the remaining (non-conserved) sites.

The variations can be made using methods known in the art such as oligonucleotide-mediated (site-directed) mutagenesis, alanine scanning, and PCR mutagenesis. Site-directed mutagenesis [Carter *et al.*, Nucl. Acids Res., 13:4331 (1986); Zoller *et al.*, Nucl. Acids Res., 10:6487 (1987)], cassette mutagenesis [Wells *et al.*, Gene, 34:315 (1985)], restriction selection mutagenesis [Wells *et al.*, Philos. Trans. R. Soc. London SerA, 317:415

(1986)] or other known techniques can be performed on the cloned DNA to produce the PRO variant DNA.

Scanning amino acid analysis can also be employed to identify one or more amino acids along a contiguous sequence. Among the preferred scanning amino acids are relatively small, neutral amino acids. Such amino acids include alanine, glycine, serine, and cysteine. Alanine is typically a preferred scanning amino acid among this group because it eliminates the side-chain beyond the beta-carbon and is less likely to alter the main-chain conformation of the variant [Cunningham and Wells, Science, 244: 1081-1085 (1989)]. Alanine is also typically preferred because it is the most common amino acid. Further, it is frequently found in both buried and exposed positions [Creighton, The Proteins, (W.H. Freeman & Co., N.Y.); Chothia, J. Mol. Biol., 150:1 (1976)]. If alanine substitution does not yield adequate amounts of variant, an isoteric amino acid can be used.

5.2.2. Modifications of PRO Polypeptides

Covalent modifications of PRO polypeptides are included within the scope of this invention. One type of covalent modification includes reacting targeted amino acid residues of a PRO polypeptide with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C- terminal residues of the PRO polypeptide. Derivatization with bifunctional agents is useful, for instance, for crosslinking the PRO polypeptide to a water-insoluble support matrix or surface for use in the method for purifying anti-PRO antibodies, and vice-versa. Commonly used crosslinking agents include, *e.g.*, 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidylpropionate), bifunctional maleimides such as bis-N-maleimido-1,8-octane and agents such as methyl-3-[(p-azidophenyl)dithio]propioimide.

Other modifications include deamidation of glutamyl and asparaginy residues to the corresponding glutamyl and aspartyl residues, respectively, hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the α -amino groups of lysine, arginine, and histidine side chains [T.E. Creighton, Proteins: Structure and Molecular Properties, W.H. Freeman & Co., San Francisco, pp. 79-86 (1983)], acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

Another type of covalent modification of the PRO polypeptide included within the scope of this invention comprises altering the native glycosylation pattern of the polypeptide. "Altering the native glycosylation pattern" is intended for purposes herein to mean deleting one or more carbohydrate moieties found in the native sequence PRO polypeptide (either by removing the underlying glycosylation site or by deleting the glycosylation by chemical and/or enzymatic means), and/or adding one or more glycosylation sites that are not present in the native sequence PRO polypeptide. In addition, the phrase includes qualitative changes in the glycosylation of the native proteins, involving a change in the nature and proportions of the various carbohydrate moieties present.

Addition of glycosylation sites to the PRO polypeptide may be accomplished by altering the amino acid sequence. The alteration may be made, for example, by the addition of, or substitution by, one or more serine or threonine residues to the native sequence PRO polypeptide (for O-linked glycosylation sites). The PRO amino acid sequence may optionally be altered through changes at the DNA level, particularly by mutating the DNA encoding the PRO polypeptide at preselected bases such that codons are generated that will translate into the desired amino

acids.

Another means of increasing the number of carbohydrate moieties on the PRO polypeptide is by chemical or enzymatic coupling of glycosides to the polypeptide. Such methods are described in the art, e.g., in WO 87/05330 published 11 September 1987, and in Aplin and Wriston, CRC Crit. Rev. Biochem., pp. 259-306 (1981).

Removal of carbohydrate moieties present on the PRO polypeptide may be accomplished chemically or enzymatically or by mutational substitution of codons encoding for amino acid residues that serve as targets for glycosylation. Chemical deglycosylation techniques are known in the art and described, for instance, by Hakimuddin, *et al.*, Arch. Biochem. Biophys., 259:52 (1987) and by Edge *et al.*, Anal. Biochem., 118:131 (1981). Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura *et al.*, Meth. Enzymol., 138:350 (1987).

Another type of covalent modification of the PRO polypeptide comprises linking the PRO polypeptide to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol (PEG), polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Patent Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

The PRO polypeptide of the present invention may also be modified in a way to form a chimeric molecule comprising the PRO polypeptide fused to another, heterologous polypeptide or amino acid sequence.

In one embodiment, such a chimeric molecule comprises a fusion of the PRO polypeptide with a protein transduction domain which targets the PRO polypeptide for delivery to various tissues and more particularly across the brain blood barrier, using, for example, the protein transduction domain of human immunodeficiency virus TAT protein (Schwarze *et al.*, 1999, *Science* 285: 1569-72).

In another embodiment, such a chimeric molecule comprises a fusion of the PRO polypeptide with a tag polypeptide which provides an epitope to which an anti-tag antibody can selectively bind. The epitope tag is generally placed at the amino- or carboxyl- terminus of the PRO polypeptide. The presence of such epitope-tagged forms of the PRO polypeptide can be detected using an antibody against the tag polypeptide. Also, provision of the epitope tag enables the PRO polypeptide to be readily purified by affinity purification using an anti-tag antibody or another type of affinity matrix that binds to the epitope tag. Various tag polypeptides and their respective antibodies are well known in the art. Examples include poly-histidine (poly-His) or poly-histidine-glycine (poly-His-gly) tags; the flu HA tag polypeptide and its antibody 12CA5 [Field *et al.*, Mol. Cell. Biol., 8:2159-2165 (1988)]; the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto [Evan *et al.*, Molecular and Cellular Biology, 5:3610-3616 (1985)]; and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody [Paborsky *et al.*, Protein Engineering, 3(6):547-553 (1990)]. Other tag polypeptides include the Flag-peptide [Hopp *et al.*, BioTechnology, 6:1204-1210 (1988)]; the KT3 epitope peptide [Martin *et al.*, Science, 255:192-194 (1992)]; an α -tubulin epitope peptide [Skinner *et al.*, J. Biol. Chem., 266:15163-15166 (1991)]; and the T7 gene 10 protein peptide tag [Lutz-Freyermuth *et al.*, Proc. Natl. Acad. Sci. USA, 87:6393-6397 (1990)].

In an alternative embodiment, the chimeric molecule may comprise a fusion of the PRO polypeptide with an immunoglobulin or a particular region of an immunoglobulin. For a bivalent form of the chimeric molecule (also referred to as an "immunoadhesin"), such a fusion could be to the Fc region of an IgG molecule. The Ig fusions

preferably include the substitution of a soluble (transmembrane domain deleted or inactivated) form of a PRO polypeptide in place of at least one variable region within an Ig molecule. In a particularly preferred embodiment, the immunoglobulin fusion includes the hinge, CH2 and CH3, or the hinge, CH1, CH2 and CH3 regions of an IgG1 molecule. For the production of immunoglobulin fusions *see also*, U.S. Patent No. 5,428,130 issued June 27, 1995.

5 5.2.3. Preparation of the PRO Polypeptide

10 The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO polypeptides. In particular, cDNAs encoding PRO polypeptides have been identified and isolated, as disclosed in further detail in the Examples below. It is noted that proteins produced in separate expression rounds may be given different PRO numbers but the UNQ number is unique for any given DNA and the encoded protein, and will not be changed. However, for sake of simplicity, in the present specification the protein encoded by the PRO DNA as well as all further native homologues and variants included in the foregoing definition of PRO polypeptides, will be referred to as "PRO" regardless of their origin or mode of preparation.

15 The description below relates primarily to production of PRO polypeptides by culturing cells transformed or transfected with a vector containing nucleic acid encoding PRO polypeptides. It is, of course, contemplated that alternative methods that are well known in the art may be employed to prepare the PRO polypeptide. For instance, the PRO polypeptide sequence, or portions thereof, may be produced by direct peptide synthesis using solid-phase techniques. *See, e.g., Stewart et al., Solid-Phase Peptide Synthesis* (W.H. Freeman Co.: San Francisco, CA, 1969); Merrifield, *J. Am. Chem. Soc.*, **85**: 2149-2154 (1963). *In vitro* protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be accomplished, for instance, with an Applied Biosystems Peptide Synthesizer (Foster City, CA) using manufacturer's instructions. Various portions of the PRO polypeptide may be chemically synthesized separately and combined using chemical or enzymatic methods to produce the full-length PRO polypeptide.

20 5.2.3.1. Isolation of DNA Encoding PRO Polypeptides

25 DNA encoding the PRO polypeptide may be obtained from a cDNA library prepared from tissue believed to possess the mRNA encoding the PRO polypeptide and to express it at a detectable level. Accordingly, DNAs encoding the human PRO polypeptide can be conveniently obtained from cDNA libraries prepared from human tissues, such as described in the Examples. The gene encoding the PRO polypeptide may also be obtained from a genomic library or by oligonucleotide synthesis.

30 Libraries can be screened with probes (such as antibodies to the PRO polypeptide or oligonucleotides of at least about 20-80 bases) designed to identify the gene of interest or the protein encoded by it. Screening the cDNA or genomic library with the selected probe may be conducted using standard procedures, such as described in Sambrook *et al., supra*. An alternative means to isolate the gene encoding the PRO polypeptide is to use PCR methodology. Sambrook *et al., supra*; Dieffenbach *et al., PCR Primer: A Laboratory Manual* (New York: Cold

Spring Harbor Laboratory Press, 1995).

The Examples below describe techniques for screening a cDNA library. The oligonucleotide sequences selected as probes should be of sufficient length and sufficiently unambiguous that false positives are minimized. The oligonucleotide is preferably labeled such that it can be detected upon hybridization to DNA in the library being
5 screened. Methods of labeling are well known in the art, and include the use of radiolabels like ^{32}P -labeled ATP, biotinylation, or enzyme labeling. Hybridization conditions, including moderate stringency and high stringency, are provided in Sambrook *et al.*, *supra*.

Sequences identified in such library screening methods can be compared and aligned to other known sequences deposited and available in public databases such as GenBank or other private sequence databases.
10 Sequence identity (at either the amino acid or nucleotide level) within defined regions of the molecule or across the full-length sequence can be determined through sequence alignment using computer software programs such as ALIGN, DNASTar, and INHERIT, which employ various algorithms to measure homology.

Nucleic acid having protein coding sequence may be obtained by screening selected cDNA or genomic libraries using the deduced amino acid sequence disclosed herein for the first time, and, if necessary, using
15 conventional primer extension procedures as described in Sambrook *et al.*, *supra*, to detect precursors and processing intermediates of mRNA that may not have been reverse-transcribed into cDNA.

5.2.3.2. Selection and Transformation of Host Cells

Host cells are transfected or transformed with expression or cloning vectors described herein for PRO polypeptide production and cultured in conventional nutrient media modified as appropriate for inducing promoters,
20 selecting transformants, or amplifying the genes encoding the desired sequences. The culture conditions, such as media, temperature, pH, and the like, can be selected by the skilled artisan without undue experimentation. In general, principles, protocols, and practical techniques for maximizing the productivity of cell cultures can be found in Mammalian Cell Biotechnology: A Practical Approach, M. Butler, ed. (IRL Press, 1991) and Sambrook *et al.*, *supra*.

Methods of transfection are known to the ordinarily skilled artisan, for example, CaPO_4 treatment and electroporation. Depending on the host cell used, transformation is performed using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in Sambrook *et al.*, *supra*, or electroporation is generally used for prokaryotes or other cells that contain substantial cell-wall barriers. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw *et al.*, Gene,
25 23: 315 (1983) and WO 89/05859 published 29 June 1989. For mammalian cells without such cell walls, the calcium phosphate precipitation method of Graham and van der Eb, Virology, 52:456-457 (1978) can be employed. General aspects of mammalian cell host system transformations have been described in U.S. Patent No. 4,399,216. Transformations into yeast are typically carried out according to the method of Van Solingen *et al.*, J. Bact., 130:
30 946 (1977) and Hsiao *et al.*, Proc. Natl. Acad. Sci. (USA), 76: 3829 (1979). However, other methods for introducing DNA into cells, such as by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, *e.g.*, polybrene or polyornithine, may also be used. For various techniques for
35

transforming mammalian cells, *see*, Keown *et al.*, Methods in Enzymology, 185: 527-537 (1990) and Mansour *et al.*, Nature, 336: 348-352 (1988).

Suitable host cells for cloning or expressing the DNA in the vectors herein include prokaryote, yeast, or higher eukaryote cells. Suitable prokaryotes include, but are not limited to, eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as *E. coli*. Various *E. coli* strains are publicly available, such as *E. coli* K12 strain MM294 (ATCC 31,446); *E. coli* X1776 (ATCC 31,537); *E. coli* strain W3110 (ATCC 27,325); and K5 772 (ATCC 53,635). Other suitable prokaryotic host cells include Enterobacteriaceae such as *Escherichia*, *e.g.*, *E. coli*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*, *e.g.*, *Salmonella typhimurium*, *Serratia*, *e.g.*, *Serratia marcescans*, and *Shigella*, as well as *Bacilli* such as *B. subtilis* and *B. licheniformis* (*e.g.*, *B. licheniformis* 41P disclosed in DD 266,710 published 12 April 1989), *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*. These examples are illustrative rather than limiting. Strain W3110 is one particularly preferred host or parent host because it is a common host strain for recombinant DNA product fermentations. Preferably, the host cell secretes minimal amounts of proteolytic enzymes. For example, strain W3110 may be modified to effect a genetic mutation in the genes encoding proteins endogenous to the host, with examples of such hosts including *E. coli* W3110 strain 1A2, which has the complete genotype *tonA*; *E. coli* W3110 strain 9E4, which has the complete genotype *tonA ptr3*; *E. coli* W3110 strain 27C7 (ATCC 55,244), which has the complete genotype *tonA ptr3 phoA E15 (argF-lac)169 degP ompT kan'*; *E. coli* W3110 strain 37D6, which has the complete genotype *tonA ptr3 phoA E15 (argF-lac)169 degP ompT rbs7 ilvG kan'*; *E. coli* W3110 strain 40B4, which is strain 37D6 with a non-kanamycin resistant *degP* deletion mutation; and an *E. coli* strain having mutant periplasmic protease disclosed in U.S. Patent No. 4,946,783 issued 7 August 1990. Alternatively, *in vitro* methods of cloning, *e.g.*, PCR or other nucleic acid polymerase reactions, are suitable.

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for vectors encoding the PRO polypeptide. *Saccharomyces cerevisiae* is a commonly used lower eukaryotic host microorganism. Others include *Schizosaccharomyces pombe* (Beach and Nurse, Nature, 290: 140 [1981]; EP 139,383 published 2 May 1985); *Kluyveromyces* hosts (U.S. Patent No. 4,943,529; Fleer *et al.*, Bio/Technology, 9: 968-975 (1991)) such as, *e.g.*, *K. lactis* (MW98-8C, CBS683, CBS4574; Louvencourt *et al.*, J. Bacteriol., 737 [1983]), *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilum* (ATCC 36,906; Van den Berg *et al.*, Bio/Technology, 8: 135 (1990)), *K. thermotolerans*, and *K. marxianus*; *yarrowia* (EP 402,226); *Pichia pastoris* (EP 183,070; Sreekrishna *et al.*, J. Basic Microbiol., 28: 265-278 [1988]); *Candida*; *Trichoderma reesia* (EP 244,234); *Neurospora crassa* (Case *et al.*, Proc. Natl. Acad. Sci. USA, 76: 5259-5263 [1979]); *Schwanniomyces* such as *Schwanniomyces occidentalis* (EP 394,538 published 31 October 1990); and filamentous fungi such as, *e.g.*, *Neurospora*, *Penicillium*, *Tolyposcladium* (WO 91/00357 published 10 January 1991), and *Aspergillus* hosts such as *A. nidulans* (Ballance *et al.*, Biochem. Biophys. Res. Commun., 112: 284-289 [1983]; Tilburn *et al.*, Gene, 26: 205-221 [1983]; Yelton *et al.*, Proc. Natl. Acad. Sci. USA, 81: 1470-1474 [1984]) and *A. niger* (Kelly and Hynes, EMBO J., 4: 475-479 [1985]). Methylophilic yeasts are suitable herein and include, but are not limited to, yeast capable of growth on methanol selected from the genera consisting of *Hansenula*, *Candida*, *Kloeckera*, *Pichia*, *Saccharomyces*,

Torulopsis, and *Rhodotorula*. A list of specific species that are exemplary of this class of yeasts may be found in C. Anthony, The Biochemistry of Methylophilic, 269 (1982).

Suitable host cells for the expression of nucleic acid encoding glycosylated PRO polypeptides are derived from multicellular organisms. Examples of invertebrate cells include insect cells such as *Drosophila* S2 and *Spodoptera* Sf9, as well as plant cells. Examples of useful mammalian host cell lines include Chinese hamster ovary (CHO) and COS cells. More specific examples include monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham *et al.*, J. Gen. Virol., 36: 59 (1977)); Chinese hamster ovary cells/-DHFR (CHO, Urlaub and Chasin, Proc. Natl. Acad. Sci. USA, 77:4216 (1980)); mouse sertoli cells (TM4, Mather, Biol. Reprod., 23:243-251 (1980)); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); and mouse mammary tumor (MMT 060562, ATCC CCL51). The selection of the appropriate host cell is deemed to be within the skill in the art.

5.2.3.3. Selection and Use of a Replicable Vector

The nucleic acid (*e.g.*, cDNA or genomic DNA) encoding the PRO polypeptide may be inserted into a replicable vector for cloning (amplification of the DNA) or for expression. Various vectors are publicly available. The vector may, for example, be in the form of a plasmid, cosmid, viral particle, or phage. The appropriate nucleic acid sequence may be inserted into the vector by a variety of procedures. In general, DNA is inserted into an appropriate restriction endonuclease site(s) using techniques known in the art. Vector components generally include, but are not limited to, one or more of a signal sequence if the sequence is to be secreted, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence. Construction of suitable vectors containing one or more of these components employs standard ligation techniques that are known to the skilled artisan.

The PRO polypeptide may be produced recombinantly not only directly, but also as a fusion polypeptide with a heterologous polypeptide, which may be a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. In general, the signal sequence may be a component of the vector, or it may be a part of the DNA encoding the PRO polypeptide that is inserted into the vector. The signal sequence may be a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. For yeast secretion the signal sequence may be, *e.g.*, the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces* α -factor leaders, the latter described in U.S. Patent No. 5,010,182), or acid phosphatase leader, the *C. albicans* glucoamylase leader (EP 362,179 published 4 April 1990), or the signal described in WO 90/13646 published 15 November 1990. In mammalian cell expression, mammalian signal sequences may be used to direct secretion of the protein, such as signal sequences from secreted polypeptides of the same or related species, as well as viral secretory leaders.

Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2 μ plasmid origin

is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV, or BPV) are useful for cloning vectors in mammalian cells.

Expression and cloning vectors will typically contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, *e.g.*, ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, *e.g.*, the gene encoding D-alanine racemase for *Bacilli*.

An example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the nucleic acid encoding the PRO polypeptide such as DHFR or thymidine kinase. An appropriate host cell when wild-type DHFR is employed is the CHO cell line deficient in DHFR activity, prepared and propagated as described by Urlaub *et al.*, Proc. Natl. Acad. Sci. USA, 77: 4216 (1980). A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid YRp7. Stinchcomb *et al.*, Nature, 282: 39 (1979); Kingsman *et al.*, Gene, 7: 141 (1979); Tschemper *et al.*, Gene, 10: 157 (1980). The *trp1* gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1. Jones, Genetics, 85: 12 (1977).

Expression and cloning vectors usually contain a promoter operably linked to the nucleic acid sequence encoding the PRO polypeptide to direct mRNA synthesis. Promoters recognized by a variety of potential host cells are well known. Promoters suitable for use with prokaryotic hosts include the β -lactamase and lactose promoter systems (Chang *et al.*, Nature, 275: 615 (1978); Goeddel *et al.*, Nature, 281: 544 (1979)), alkaline phosphatase, a tryptophan (*trp*) promoter system (Goeddel, Nucleic Acids Res., 8: 4057 (1980); EP 36,776), and hybrid promoters such as the *tac* promoter (deBoer *et al.*, Proc. Natl. Acad. Sci. USA, 80: 21-25 (1983)). Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding the PRO polypeptide.

Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase (Hitzeman *et al.*, J. Biol. Chem., 255: 2073 (1980)) or other glycolytic enzymes (Hess *et al.*, J. Adv. Enzyme Reg., 7: 149 (1968); Holland, Biochemistry, 17: 4900 (1978)), such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

Other yeast promoters that are inducible promoters having the additional advantage of transcription controlled by growth conditions are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657.

PRO nucleic acid transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus (UK 2,211,504 published 5 July 1989), adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus, and Simian Virus 40 (SV40); by heterologous mammalian promoters, *e.g.*, the actin

promoter or an immunoglobulin promoter; and by heat-shock promoters, provided such promoters are compatible with the host cell systems.

Transcription of a DNA encoding the PRO polypeptide by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp, that act on a promoter to increase its transcription. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, α -fetoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. The enhancer may be spliced into the vector at a position 5' or 3' to the sequence coding for PRO polypeptides, but is preferably located at a site 5' from the promoter.

Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding the PRO polypeptide.

Still other methods, vectors, and host cells suitable for adaptation to the synthesis of the PRO polypeptide in recombinant vertebrate cell culture are described in Gething *et al.*, Nature, 293: 620-625 (1981); Mantei *et al.*, Nature, 281: 40-46 (1979); EP 117,060; and EP 117,058.

5.2.3.4. Detecting Gene Amplification/Expression

Gene amplification and/or expression may be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA (Thomas, Proc. Natl. Acad. Sci. USA, 77:5201-5205 (1980)), dot blotting (DNA analysis), or *in situ* hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn may be labeled and the assay may be carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

Gene expression, alternatively, may be measured by immunological methods, such as immunohistochemical staining of cells or tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a native-sequence PRO polypeptide or against a synthetic peptide based on the DNA sequences provided herein or against exogenous sequence fused to DNA encoding the PRO polypeptide and encoding a specific antibody epitope.

5.2.3.5. Purification of PRO Polypeptides

Forms of PRO polypeptides may be recovered from culture medium or from host cell lysates. If

membrane-bound, it can be released from the membrane using a suitable detergent solution (e.g., TRITON-X™ 100) or by enzymatic cleavage. Cells employed in expression of nucleic acid encoding the PRO polypeptide can be disrupted by various physical or chemical means, such as freeze-thaw cycling, sonication, mechanical disruption, or cell-lysing agents. It may be desired to purify the PRO polypeptide from recombinant cell proteins or polypeptides. The following procedures are exemplary of suitable purification procedures: by fractionation on an ion-exchange column; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; protein A Sepharose columns to remove contaminants such as IgG; and metal chelating columns to bind epitope-tagged forms of the PRO polypeptide. Various methods of protein purification may be employed and such methods are known in the art and described, for example, in Deutscher, Methods in Enzymology, 182 (1990); Scopes, Protein Purification: Principles and Practice (Springer-Verlag: New York, 1982). The purification step(s) selected will depend, for example, on the nature of the production process used and the particular PRO polypeptide produced.

5.2.4. Uses of PRO Polypeptides

5.2.4.1. Assays for Cardiovascular, Endothelial, and Angiogenic Activity

Various assays can be used to test the polypeptide herein for cardiovascular, endothelial, and angiogenic activity. Such assays include those provided in the Examples below.

Assays for testing for endothelin antagonist activity, as disclosed in U.S. Pat. No. 5,773,414, include a rat heart ventricle binding assay where the polypeptide is tested for its ability to inhibit iodinated endothelin-1 binding in a receptor assay, an endothelin receptor binding assay testing for intact cell binding of radiolabeled endothelin-1 using rabbit renal artery vascular smooth muscle cells, an inositol phosphate accumulation assay where functional activity is determined in Rat-1 cells by measuring intra-cellular levels of second messengers, an arachidonic acid release assay that measures the ability of added compounds to reduce endothelin-stimulated arachidonic acid release in cultured vascular smooth muscles, *in vitro* (isolated vessel) studies using endothelium from male New Zealand rabbits, and *in vivo* studies using male Sprague-Dawley rats.

Assays for tissue generation activity include, without limitation, those described in WO 95/16035 (bone, cartilage, tendon); WO 95/05846 (nerve, neuronal), and WO 91/07491 (skin, endothelium).

Assays for wound-healing activity include, for example, those described in Winter, Epidermal Wound Healing, Maibach, HI and Rovee, DT, eds. (Year Book Medical Publishers, Inc., Chicago), pp. 71-112, as modified by the article of Eaglstein and Mertz, J. Invest. Dermatol., 71: 382-384 (1978).

An assay to screen for a test molecule relating to a PRO polypeptide that binds an endothelin B₁ (ETB₁) receptor polypeptide and modulates signal transduction activity involves providing a host cell transformed with a DNA encoding endothelin B₁ receptor polypeptide, exposing the cells to the test candidate, and measuring endothelin B₁ receptor signal transduction activity, as described, e.g., in U.S. Pat. No. 5,773,223.

There are several cardiac hypertrophy assays. *In vitro* assays include induction of spreading of adult rat cardiac myocytes. In this assay, ventricular myocytes are isolated from a single (male Sprague-Dawley) rat,

essentially following a modification of the procedure described in detail by Piper *et al.*, "Adult ventricular rat heart muscle cells" in Cell Culture Techniques in Heart and Vessel Research, H.M. Piper, ed. (Berlin: Springer-Verlag, 1990), pp. 36-60. This procedure permits the isolation of adult ventricular myocytes and the long-term culture of these cells in the rod-shaped phenotype. Phenylephrine and Prostaglandin $F_{2\alpha}$ ($PGF_{2\alpha}$) have been shown to induce a spreading response in these adult cells. The inhibition of myocyte spreading induced by $PGF_{2\alpha}$ or $PGF_{2\alpha}$ analogs (e.g., fluprostenol) and phenylephrine by various potential inhibitors of cardiac hypertrophy is then tested.

One example of an *in vivo* assay is a test for inhibiting cardiac hypertrophy induced by fluprostenol *in vivo*. This pharmacological model tests the ability of the PRO polypeptide to inhibit cardiac hypertrophy induced in rats (e.g., male Wistar or Sprague-Dawley) by subcutaneous injection of fluprostenol (an agonist analog of $PGF_{2\alpha}$). It is known that rats with pathologic cardiac hypertrophy induced by myocardial infarction have chronically elevated levels of extractable $PGF_{2\alpha}$ in their myocardium. Lai *et al.*, Am. J. Physiol. (Heart Circ. Physiol.), 271: H2197-H2208 (1996). Accordingly, factors that can inhibit the effects of fluprostenol on myocardial growth *in vivo* are potentially useful for treating cardiac hypertrophy. The effects of the PRO polypeptide on cardiac hypertrophy are determined by measuring the weight of heart, ventricles, and left ventricle (normalized by body weight) relative to fluprostenol-treated rats not receiving the PRO polypeptide.

Another example of an *in vivo* assay is the pressure-overload cardiac hypertrophy assay. For *in vivo* testing it is common to induce pressure-overload cardiac hypertrophy by constriction of the abdominal aorta of test animals. In a typical protocol, rats (e.g., male Wistar or Sprague-Dawley) are treated under anesthesia, and the abdominal aorta of each rat is narrowed down just below the diaphragm. Beznak M., Can. J. Biochem. Physiol., 33: 985-94 (1955). The aorta is exposed through a surgical incision, and a blunted needle is placed next to the vessel. The aorta is constricted with a ligature of silk thread around the needle, which is immediately removed and which reduces the lumen of the aorta to the diameter of the needle. This approach is described, for example, in Rossi *et al.*, Am. Heart J., 124: 700-709 (1992) and O'Rourke and Reibel, P.S.E.M.B., 200: 95-100 (1992).

In yet another *in vivo* assay, the effect on cardiac hypertrophy following experimentally induced myocardial infarction (MI) is measured. Acute MI is induced in rats by left coronary artery ligation and confirmed by electrocardiographic examination. A sham-operated group of animals is also prepared as control animals. Earlier data have shown that cardiac hypertrophy is present in the group of animals with MI, as evidenced by an 18% increase in heart weight-to-body weight ratio. Lai *et al.*, *supra*. Treatment of these animals with candidate blockers of cardiac hypertrophy, e.g., the PRO polypeptide, provides valuable information about the therapeutic potential of the candidates tested. One further such assay test for induction of cardiac hypertrophy is disclosed in U.S. Pat. No. 5,773,415, using Sprague-Dawley rats.

For cancer, a variety of well-known animal models can be used to further understand the role of the genes identified herein in the development and pathogenesis of tumors, and to test the efficacy of candidate therapeutic agents, including antibodies and other antagonists of native PRO polypeptides, such as small-molecule antagonists. The *in vivo* nature of such models makes them particularly predictive of responses in human patients. Animal models of tumors and cancers (e.g., breast cancer, colon cancer, prostate cancer, lung cancer, etc.) include both non-recombinant and recombinant (transgenic) animals. Non-recombinant animal models include, for example,

rodent, *e.g.*, murine models. Such models can be generated by introducing tumor cells into syngeneic mice using standard techniques, *e.g.*, subcutaneous injection, tail vein injection, spleen implantation, intraperitoneal implantation, implantation under the renal capsule, or orthotopic implantation, *e.g.*, colon cancer cells implanted in colonic tissue. See, *e.g.*, PCT publication No. WO 97/33551, published September 18, 1997. Probably the most often used animal species in oncological studies are immunodeficient mice and, in particular, nude mice. The observation that the nude mouse with thymic hypo/aplasia could successfully act as a host for human tumor xenografts has lead to its widespread use for this purpose. The autosomal recessive *nu* gene has been introduced into a very large number of distinct congenic strains of nude mouse, including, for example, ASW, A/He, AKR, BALB/c, B10.LP, C17, C3H, C57BL, C57, CBA, DBA, DDD, I/st, NC, NFR, NFS, NFS/N, NZB, NZC, NZW, P, RIII, and SJL. In addition, a wide variety of other animals with inherited immunological defects other than the nude mouse have been bred and used as recipients of tumor xenografts. For further details see, *e.g.*, The Nude Mouse in Oncology Research, E. Boven and B. Winograd, eds. (CRC Press, Inc., 1991).

The cells introduced into such animals can be derived from known tumor/cancer cell lines, such as any of the above-listed tumor cell lines, and, for example, the B104-1-1 cell line (stable NIH-3T3 cell line transfected with the *neu* protooncogene); *ras*-transfected NIH-3T3 cells; Caco-2 (ATCC HTB-37); or a moderately well-differentiated grade II human colon adenocarcinoma cell line, HT-29 (ATCC HTB-38); or from tumors and cancers. Samples of tumor or cancer cells can be obtained from patients undergoing surgery, using standard conditions involving freezing and storing in liquid nitrogen. Karmali *et al.*, Br. J. Cancer, 48: 689-696 (1983).

Tumor cells can be introduced into animals such as nude mice by a variety of procedures. The subcutaneous (s.c.) space in mice is very suitable for tumor implantation. Tumors can be transplanted s.c. as solid blocks, as needle biopsies by use of a trochar, or as cell suspensions. For solid-block or trochar implantation, tumor tissue fragments of suitable size are introduced into the s.c. space. Cell suspensions are freshly prepared from primary tumors or stable tumor cell lines, and injected subcutaneously. Tumor cells can also be injected as subdermal implants. In this location, the inoculum is deposited between the lower part of the dermal connective tissue and the s.c. tissue.

Animal models of breast cancer can be generated, for example, by implanting rat neuroblastoma cells (from which the *neu* oncogene was initially isolated), or *neu*-transformed NIH-3T3 cells into nude mice, essentially as described by Drebin *et al.* Proc. Nat. Acad. Sci. USA, 83: 9129-9133 (1986).

Similarly, animal models of colon cancer can be generated by passaging colon cancer cells in animals, *e.g.*, nude mice, leading to the appearance of tumors in these animals. An orthotopic transplant model of human colon cancer in nude mice has been described, for example, by Wang *et al.*, Cancer Research, 54: 4726-4728 (1994) and Too *et al.*, Cancer Research, 55: 681-684 (1995). This model is based on the so-called "METAMOUSETM" sold by AntiCancer, Inc., (San Diego, California).

Tumors that arise in animals can be removed and cultured *in vitro*. Cells from the *in vitro* cultures can then be passaged to animals. Such tumors can serve as targets for further testing or drug screening. Alternatively, the tumors resulting from the passage can be isolated and RNA from pre-passage cells and cells isolated after one or more rounds of passage analyzed for differential expression of genes of interest. Such passaging techniques can

be performed with any known tumor or cancer cell lines.

For example, Meth A, CMS4, CMS5, CMS21, and WEHI-164 are chemically induced fibrosarcomas of BALB/c female mice (DeLeo *et al.*, J. Exp. Med., **146**: 720 (1977)), which provide a highly controllable model system for studying the anti-tumor activities of various agents. Palladino *et al.*, J. Immunol., **138**: 4023-4032 (1987). Briefly, tumor cells are propagated *in vitro* in cell culture. Prior to injection into the animals, the cell lines are washed and suspended in buffer, at a cell density of about 10×10^6 to 10×10^7 cells/ml. The animals are then infected subcutaneously with 10 to 100 μ l of the cell suspension, allowing one to three weeks for a tumor to appear.

In addition, the Lewis lung (3LL) carcinoma of mice, which is one of the most thoroughly studied experimental tumors, can be used as an investigational tumor model. Efficacy in this tumor model has been correlated with beneficial effects in the treatment of human patients diagnosed with small-cell carcinoma of the lung (SCCL). This tumor can be introduced in normal mice upon injection of tumor fragments from an affected mouse or of cells maintained in culture. Zupi *et al.*, Br. J. Cancer, **41**: suppl. 4, 30 (1980). Evidence indicates that tumors can be started from injection of even a single cell and that a very high proportion of infected tumor cells survive. For further information about this tumor model see, Zacharski, Haemostasis, **16**: 300-320 (1986).

One way of evaluating the efficacy of a test compound in an animal model with an implanted tumor is to measure the size of the tumor before and after treatment. Traditionally, the size of implanted tumors has been measured with a slide caliper in two or three dimensions. The measure limited to two dimensions does not accurately reflect the size of the tumor; therefore, it is usually converted into the corresponding volume by using a mathematical formula. However, the measurement of tumor size is very inaccurate. The therapeutic effects of a drug candidate can be better described as treatment-induced growth delay and specific growth delay. Another important variable in the description of tumor growth is the tumor volume doubling time. Computer programs for the calculation and description of tumor growth are also available, such as the program reported by Rygaard and Spang-Thomsen, Proc. 6th Int. Workshop on Immune-Deficient Animals, Wu and Sheng eds. (Basel, 1989), p. 301. It is noted, however, that necrosis and inflammatory responses following treatment may actually result in an increase in tumor size, at least initially. Therefore, these changes need to be carefully monitored, by a combination of a morphometric method and flow cytometric analysis.

Further, recombinant (transgenic) animal models can be engineered by introducing the coding portion of the PRO gene identified herein into the genome of animals of interest, using standard techniques for producing transgenic animals. Animals that can serve as a target for transgenic manipulation include, without limitation, mice, rats, rabbits, guinea pigs, sheep, goats, pigs, and non-human primates, *e.g.*, baboons, chimpanzees and monkeys. Techniques known in the art to introduce a transgene into such animals include pronucleic microinjection (U.S. Patent No. 4,873,191); retrovirus-mediated gene transfer into germ lines (*e.g.*, Van der Putten *et al.*, Proc. Natl. Acad. Sci. USA, **82**: 6148-615 (1985)); gene targeting in embryonic stem cells (Thompson *et al.*, Cell, **56**: 313-321 (1989)); electroporation of embryos (Lo, Mol. Cell. Biol., **3**: 1803-1814 (1983)); and sperm-mediated gene transfer. Lavitrano *et al.*, Cell, **57**: 717-73 (1989). For a review, see for example, U.S. Patent No. 4,736,866.

For the purpose of the present invention, transgenic animals include those that carry the transgene only in part of their cells ("mosaic animals"). The transgene can be integrated either as a single transgene, or in

concatamers, *e.g.*, head-to-head or head-to-tail tandems. Selective introduction of a transgene into a particular cell type is also possible by following, for example, the technique of Lasko *et al.*, Proc. Natl. Acad. Sci. USA, **89**: 6232-636 (1992). The expression of the transgene in transgenic animals can be monitored by standard techniques. For example, Southern blot analysis or PCR amplification can be used to verify the integration of the transgene. The level of mRNA expression can then be analyzed using techniques such as *in situ* hybridization, Northern blot analysis, PCR, or immunocytochemistry. The animals are further examined for signs of tumor or cancer development.

Alternatively, "knock-out" animals can be constructed that have a defective or altered gene encoding a PRO polypeptide identified herein, as a result of homologous recombination between the endogenous gene encoding the PRO polypeptide and altered genomic DNA encoding the same polypeptide introduced into an embryonic cell of the animal. For example, cDNA encoding a particular PRO polypeptide can be used to clone genomic DNA encoding that polypeptide in accordance with established techniques. A portion of the genomic DNA encoding a particular PRO polypeptide can be deleted or replaced with another gene, such as a gene encoding a selectable marker that can be used to monitor integration. Typically, several kilobases of unaltered flanking DNA (both at the 5' and 3' ends) are included in the vector. *See, e.g.*, Thomas and Capecchi, Cell, **51**: 503 (1987) for a description of homologous recombination vectors. The vector is introduced into an embryonic stem cell line (*e.g.*, by electroporation) and cells in which the introduced DNA has homologously recombined with the endogenous DNA are selected. *See, e.g.*, Li *et al.*, Cell, **69**: 915 (1992). The selected cells are then injected into a blastocyst of an animal (*e.g.*, a mouse or rat) to form aggregation chimeras. *See, e.g.*, Bradley, in Teratocarcinomas and Embryonic Stem Cells: A Practical Approach, E. J. Robertson, ed. (IRL: Oxford, 1987), pp. 113-152. A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term to create a "knock-out" animal. Progeny harboring the homologously recombined DNA in their germ cells can be identified by standard techniques and used to breed animals in which all cells of the animal contain the homologously recombined DNA. Knockout animals can be characterized, for instance, by their ability to defend against certain pathological conditions and by their development of pathological conditions due to absence of the PRO polypeptide.

The efficacy of antibodies specifically binding the PRO polypeptides identified herein, and other drug candidates, can be tested also in the treatment of spontaneous animal tumors. A suitable target for such studies is the feline oral squamous cell carcinoma (SCC). Feline oral SCC is a highly invasive, malignant tumor that is the most common oral malignancy of cats, accounting for over 60% of the oral tumors reported in this species. It rarely metastasizes to distant sites, although this low incidence of metastasis may merely be a reflection of the short survival times for cats with this tumor. These tumors are usually not amenable to surgery, primarily because of the anatomy of the feline oral cavity. At present, there is no effective treatment for this tumor. Prior to entry into the study, each cat undergoes complete clinical examination and biopsy, and is scanned by computed tomography (CT). Cats diagnosed with sublingual oral squamous cell tumors are excluded from the study. The tongue can become paralyzed as a result of such tumor, and even if the treatment kills the tumor, the animals may not be able to feed themselves. Each cat is treated repeatedly, over a longer period of time. Photographs of the tumors will be taken daily during the treatment period, and at each subsequent recheck. After treatment, each cat undergoes another CT

scan. CT scans and thoracic radiograms are evaluated every 8 weeks thereafter. The data are evaluated for differences in survival, response, and toxicity as compared to control groups. Positive response may require evidence of tumor regression, preferably with improvement of quality of life and/or increased life span.

In addition, other spontaneous animal tumors, such as fibrosarcoma, adenocarcinoma, lymphoma, chondroma, or leiomyosarcoma of dogs, cats, and baboons can also be tested. Of these, mammary adenocarcinoma in dogs and cats is a preferred model as its appearance and behavior are very similar to those in humans. However, the use of this model is limited by the rare occurrence of this type of tumor in animals.

Other *in vitro* and *in vivo* cardiovascular, endothelial, and angiogenic tests known in the art are also suitable herein.

5.2.4.2. Tissue Distribution

The results of the cardiovascular, endothelial, and angiogenic assays herein can be verified by further studies, such as by determining mRNA expression in various human tissues.

As noted before, gene amplification and/or gene expression in various tissues may be measured by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA (Thomas, Proc. Natl. Acad. Sci. USA, 77:5201-5205 (1980)), dot blotting (DNA analysis), or *in situ* hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes.

Gene expression in various tissues, alternatively, may be measured by immunological methods, such as immunohistochemical staining of tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a native-sequence PRO polypeptide or against a synthetic peptide based on the DNA sequences provided herein or against exogenous sequence fused to PRO DNA and encoding a specific antibody epitope. General techniques for generating antibodies, and special protocols for *in situ* hybridization are provided hereinbelow.

5.2.4.3. Antibody Binding Studies

The results of the cardiovascular, endothelial, and angiogenic study can be further verified by antibody binding studies, in which the ability of anti-PRO antibodies to inhibit the effect of the PRO polypeptides on endothelial cells or other cells used in the cardiovascular, endothelial, and angiogenic assays is tested. Exemplary antibodies include polyclonal, monoclonal, humanized, bispecific, and heteroconjugate antibodies, the preparation of which will be described hereinbelow.

Antibody binding studies may be carried out in any known assay method, such as competitive binding assays, direct and indirect sandwich assays, and immunoprecipitation assays. Zola, Monoclonal Antibodies: A Manual of Techniques (CRC Press, Inc., 1987), pp.147-158.

Competitive binding assays rely on the ability of a labeled standard to compete with the test sample analyte for binding with a limited amount of antibody. The amount of target protein in the test sample is inversely proportional to the amount of standard that becomes bound to the antibodies. To facilitate determining the amount of standard that becomes bound, the antibodies preferably are insolubilized before or after the competition, so that the standard and analyte that are bound to the antibodies may conveniently be separated from the standard and analyte that remain unbound.

Sandwich assays involve the use of two antibodies, each capable of binding to a different immunogenic portion, or epitope, of the protein to be detected. In a sandwich assay, the test sample analyte is bound by a first antibody that is immobilized on a solid support, and thereafter a second antibody binds to the analyte, thus forming an insoluble three-part complex. *See, e.g.,* U.S. Pat. No. 4,376,110. The second antibody may itself be labeled with a detectable moiety (direct sandwich assays) or may be measured using an anti-immunoglobulin antibody that is labeled with a detectable moiety (indirect sandwich assay). For example, one type of sandwich assay is an ELISA assay, in which case the detectable moiety is an enzyme.

For immunohistochemistry, the tissue sample may be fresh or frozen or may be embedded in paraffin and fixed with a preservative such as formalin, for example.

5.2.4.4. Cell-Based Tumor Assays

Cell-based assays and animal models for cardiovascular, endothelial, and angiogenic disorders, such as tumors, can be used to verify the findings of a cardiovascular, endothelial, and angiogenic assay herein, and further to understand the relationship between the genes identified herein and the development and pathogenesis of undesirable cardiovascular, endothelial, and angiogenic cell growth. The role of gene products identified herein in the development and pathology of undesirable cardiovascular, endothelial, and angiogenic cell growth, *e.g.,* tumor cells, can be tested by using cells or cells lines that have been identified as being stimulated or inhibited by the PRO polypeptide herein. Such cells include, for example, those set forth in the Examples below.

In a different approach, cells of a cell type known to be involved in a particular cardiovascular, endothelial, and angiogenic disorder are transfected with the cDNAs herein, and the ability of these cDNAs to induce excessive growth or inhibit growth is analyzed. If the cardiovascular, endothelial, and angiogenic disorder is cancer, suitable tumor cells include, for example, stable tumor cell lines such as the B104-1-1 cell line (stable NIH-3T3 cell line transfected with the *neu* protooncogene) and *ras*-transfected NIH-3T3 cells, which can be transfected with the desired gene and monitored for tumorigenic growth. Such transfected cell lines can then be used to test the ability of poly- or monoclonal antibodies or antibody compositions to inhibit tumorigenic cell growth by exerting cytostatic or cytotoxic activity on the growth of the transformed cells, or by mediating antibody-dependent cellular cytotoxicity (ADCC). Cells transfected with the coding sequences of the genes identified herein can further be used to identify drug candidates for the treatment of cardiovascular, endothelial, and angiogenic disorders such as cancer.

In addition, primary cultures derived from tumors in transgenic animals (as described above) can be used in the cell-based assays herein, although stable cell lines are preferred. Techniques to derive continuous cell lines from transgenic animals are well known in the art. *See, e.g.,* Small *et al.*, Mol. Cell. Biol., **5**: 642-648 (1985).

5.2.4.5. Gene Therapy

Described below are methods and compositions whereby disease symptoms may be ameliorated. Certain diseases are brought about, at least in part, by an excessive level of gene product, or by the presence of a gene product exhibiting an abnormal or excessive activity. As such, the reduction in the level and/or activity of such gene products would bring about the amelioration of such disease symptoms.

Alternatively, certain other diseases are brought about, at least in part, by the absence or reduction of the level of gene expression, or a reduction in the level of a gene product's activity. As such, an increase in the level of gene expression and/or the activity of such gene products would bring about the amelioration of such disease symptoms.

In some cases, the up-regulation of a gene in a disease state reflects a protective role for that gene product in responding to the disease condition. Enhancement of such a target gene's expression, or the activity of the target gene product, will reinforce the protective effect it exerts. Some disease states may result from an abnormally low level of activity of such a protective gene. In these cases also, an increase in the level of gene expression and/or the activity of such gene products would bring about the amelioration of such disease symptoms.

The PRO polypeptides described herein and polypeptidyl agonists and antagonists may be employed in accordance with the present invention by expression of such polypeptides *in vivo*, which is often referred to as gene therapy.

There are two major approaches to getting the nucleic acid (optionally contained in a vector) into the patient's cells: *in vivo* and *ex vivo*. For *in vivo* delivery the nucleic acid is injected directly into the patient, usually at the sites where the PRO polypeptide is required, *i.e.*, the site of synthesis of the PRO polypeptide, if known, and the site (*e.g.*, wound) where biological activity of the PRO polypeptide is needed. For *ex vivo* treatment, the patient's cells are removed, the nucleic acid is introduced into these isolated cells, and the modified cells are administered to the patient either directly or, for example, encapsulated within porous membranes that are implanted into the patient (*see, e.g.*, U.S. Pat. Nos. 4,892,538 and 5,283,187). There are a variety of techniques available for introducing nucleic acids into viable cells. The techniques vary depending upon whether the nucleic acid is transferred into cultured cells *in vitro*, or transferred *in vivo* in the cells of the intended host. Techniques suitable for the transfer of nucleic acid into mammalian cells *in vitro* include the use of liposomes, electroporation, microinjection, transduction, cell fusion, DEAE-dextran, the calcium phosphate precipitation method, etc. Transduction involves the association of a replication-defective, recombinant viral (preferably retroviral) particle with a cellular receptor, followed by introduction of the nucleic acids contained by the particle into the cell. A commonly used vector for *ex vivo* delivery of the gene is a retrovirus.

The currently preferred *in vivo* nucleic acid transfer techniques include transfection with viral or non-viral vectors (such as adenovirus, lentivirus, Herpes simplex I virus, or adeno-associated virus (AAV)) and lipid-based systems (useful lipids for lipid-mediated transfer of the gene are, for example, DOTMA, DOPE, and DC-Chol; *see, e.g.*, Tonkinson *et al.*, Cancer Investigation, 14(1): 54-65 (1996)). The most preferred vectors for use in gene therapy are viruses, most preferably adenoviruses, AAV, lentiviruses, or retroviruses. A viral vector such as a retroviral vector includes at least one transcriptional promoter/enhancer or locus-defining element(s), or other

elements that control gene expression by other means such as alternate splicing, nuclear RNA export, or post-translational modification of messenger. In addition, a viral vector such as a retroviral vector includes a nucleic acid molecule that, when transcribed in the presence of a gene encoding the PRO polypeptide, is operably linked thereto and acts as a translation initiation sequence. Such vector constructs also include a packaging signal, long terminal repeats (LTRs) or portions thereof, and positive and negative strand primer binding sites appropriate to the virus used (if these are not already present in the viral vector). In addition, such vector typically includes a signal sequence for secretion of the PRO polypeptide from a host cell in which it is placed. Preferably the signal sequence for this purpose is a mammalian signal sequence, most preferably the native signal sequence for the PRO polypeptide. Optionally, the vector construct may also include a signal that directs polyadenylation, as well as one or more restriction sites and a translation termination sequence. By way of example, such vectors will typically include a 5' LTR, a tRNA binding site, a packaging signal, an origin of second-strand DNA synthesis, and a 3' LTR or a portion thereof. Other vectors can be used that are non-viral, such as cationic lipids, polylysine, and dendrimers.

In some situations, it is desirable to provide the nucleic acid source with an agent that targets the target cells, such as an antibody specific for a cell-surface membrane protein or the target cell, a ligand for a receptor on the target cell, etc. Where liposomes are employed, proteins that bind to a cell-surface membrane protein associated with endocytosis may be used for targeting and/or to facilitate uptake, *e.g.*, capsid proteins or fragments thereof tropic for a particular cell type, antibodies for proteins that undergo internalization in cycling, and proteins that target intracellular localization and enhance intracellular half-life. The technique of receptor-mediated endocytosis is described, for example, by Wu *et al.*, J. Biol. Chem., **262**: 4429-4432 (1987); and Wagner *et al.*, Proc. Natl. Acad. Sci. USA, **87**: 3410-3414 (1990). For a review of the currently known gene marking and gene therapy protocols, see, Anderson *et al.*, Science, **256**: 808-813 (1992). See also WO 93/25673 and the references cited therein.

Suitable gene therapy and methods for making retroviral particles and structural proteins can be found in, *e.g.*, U.S. Pat. No. 5,681,746.

5.2.4.6. Use of Gene as a Diagnostic

This invention is also related to the use of the gene encoding the PRO polypeptide as a diagnostic. Detection of a mutated form of the PRO polypeptide will allow a diagnosis of a cardiovascular, endothelial, and angiogenic disease or a susceptibility to a cardiovascular, endothelial, and angiogenic disease, such as a tumor, since mutations in the PRO polypeptide may cause tumors.

Individuals carrying mutations in the genes encoding a human PRO polypeptide may be detected at the DNA level by a variety of techniques. Nucleic acids for diagnosis may be obtained from a patient's cells, such as from blood, urine, saliva, tissue biopsy, and autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR (Saiki *et al.*, Nature, **324**: 163-166 (1986)) prior to analysis. RNA or cDNA may also be used for the same purpose. As an example, PCR primers complementary to the nucleic acid encoding the PRO polypeptide can be used to identify and analyze the PRO polypeptide mutations. For example, deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal

genotype. Point mutations can be identified by hybridizing amplified DNA to radiolabeled RNA encoding the PRO polypeptide, or alternatively, radiolabeled antisense DNA sequences encoding the PRO polypeptide. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase A digestion or by differences in melting temperatures.

5 Genetic testing based on DNA sequence differences may be achieved by detection of alteration in electrophoretic mobility of DNA fragments in gels with or without denaturing agents. Small sequence deletions and insertions can be visualized by high resolution gel electrophoresis. DNA fragments of different sequences may be distinguished on denaturing formamide gradient gels in which the mobilities of different DNA fragments are retarded in the gel at different positions according to their specific melting or partial melting temperatures. See, 10 *e.g.*, Myers *et al.*, Science, 230: 1242 (1985).

Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method, for example, Cotton *et al.*, Proc. Natl. Acad. Sci. USA, 85: 4397-4401 (1985).

15 Thus, the detection of a specific DNA sequence may be achieved by methods such as hybridization, RNase protection, chemical cleavage, direct DNA sequencing, or the use of restriction enzymes, *e.g.*, restriction fragment length polymorphisms (RFLP), and Southern blotting of genomic DNA.

5.2.4.7. Use to Detect PRO Polypeptide Levels

In addition to more conventional gel-electrophoresis and DNA sequencing, mutations can also be detected by *in situ* analysis.

20 Expression of nucleic acid encoding the PRO polypeptide may be linked to vascular disease or neovascularization associated with tumor formation. If the PRO polypeptide has a signal sequence and the mRNA is highly expressed in endothelial cells and to a lesser extent in smooth muscle cells, this indicates that the PRO polypeptide is present in serum. Accordingly, an anti-PRO polypeptide antibody could be used to diagnose vascular disease or neovascularization associated with tumor formation, since an altered level of this PRO polypeptide may 25 be indicative of such disorders.

A competition assay may be employed wherein antibodies specific to the PRO polypeptide are attached to a solid support and the labeled PRO polypeptide and a sample derived from the host are passed over the solid support and the amount of label detected attached to the solid support can be correlated to a quantity of the PRO polypeptide in the sample.

30 5.2.4.8. Chromosome Mapping

The sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. Moreover, there is a current need for identifying particular sites on the chromosome. Few chromosome marking reagents based on actual sequence data (repeat polymorphisms) are presently available for marking chromosomal 35 location. The mapping of DNAs to chromosomes according to the present invention is an important first step in

correlating those sequences with genes associated with disease.

Briefly, sequences can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp) from the cDNA. Computer analysis for the 3'- untranslated region is used to rapidly select primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers are then used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the primer will yield an amplified fragment.

PCR mapping of somatic cell hybrids is a rapid procedure for assigning a particular DNA to a particular chromosome. Using the present invention with the same oligonucleotide primers, sublocalization can be achieved with panels of fragments from specific chromosomes or pools of large genomic clones in an analogous manner. Other mapping strategies that can similarly be used to map to its chromosome include *in situ* hybridization, prescreening with labeled flow-sorted chromosomes, and preselection by hybridization to construct chromosome-specific cDNA libraries.

Fluorescence *in situ* hybridization (FISH) of a cDNA clone to a metaphase chromosomal spread can be used to provide a precise chromosomal location in one step. This technique can be used with cDNA as short as 500 or 600 bases; however, clones larger than 2,000 bp have a higher likelihood of binding to a unique chromosomal location with sufficient signal intensity for simple detection. FISH requires use of the clones from which the gene encoding the PRO polypeptide was derived, and the longer the better. For example, 2,000 bp is good, 4,000 bp is better, and more than 4,000 is probably not necessary to get good results a reasonable percentage of the time. For a review of this technique, see, Verma *et al.*, Human Chromosomes: a Manual of Basic Techniques (Pergamon Press, New York, 1988).

Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available online through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region is then identified through linkage analysis (coinheritance of physically adjacent genes).

Next, it is necessary to determine the differences in the cDNA or genomic sequence between affected and unaffected individuals. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

With current resolution of physical mapping and genetic mapping techniques, a cDNA precisely localized to a chromosomal region associated with the disease could be one of between 50 and 500 potential causative genes. (This assumes 1 megabase mapping resolution and one gene per 20 kb).

5.2.4.9. Screening Assays for Drug Candidates

This invention encompasses methods of screening compounds to identify those that mimic the PRO polypeptide (agonists) or prevent the effect of the PRO polypeptide (antagonists). Screening assays for antagonist drug candidates are designed to identify compounds that bind or complex with the PRO polypeptide encoded by the genes identified herein, or otherwise interfere with the interaction of the encoded polypeptides with other

cellular proteins. Such screening assays will include assays amenable to high-throughput screening of chemical libraries, making them particularly suitable for identifying small molecule drug candidates.

The assays can be performed in a variety of formats, including protein-protein binding assays, biochemical screening assays, immunoassays, and cell-based assays, which are well characterized in the art.

5 All assays for antagonists are common in that they call for contacting the drug candidate with a PRO polypeptide encoded by a nucleic acid identified herein under conditions and for a time sufficient to allow these two components to interact.

10 In binding assays, the interaction is binding and the complex formed can be isolated or detected in the reaction mixture. In a particular embodiment, the PRO polypeptide encoded by the gene identified herein or the drug candidate is immobilized on a solid phase, *e.g.*, on a microtiter plate, by covalent or non-covalent attachments. Non-covalent attachment generally is accomplished by coating the solid surface with a solution of the PRO polypeptide and drying. Alternatively, an immobilized antibody, *e.g.*, a monoclonal antibody, specific for the PRO polypeptide to be immobilized can be used to anchor it to a solid surface. The assay is performed by adding the non-immobilized component, which may be labeled by a detectable label, to the immobilized component, *e.g.*, the coated surface containing the anchored component. When the reaction is complete, the non-reacted components are removed, *e.g.*, by washing, and complexes anchored on the solid surface are detected. When the originally non-immobilized component carries a detectable label, the detection of label immobilized on the surface indicates that complexing occurred. Where the originally non-immobilized component does not carry a label, complexing can be detected, for example, by using a labeled antibody specifically binding the immobilized complex.

20 If the candidate compound interacts with but does not bind to a particular PRO polypeptide encoded by a gene identified herein, its interaction with that polypeptide can be assayed by methods well known for detecting protein-protein interactions. Such assays include traditional approaches, such as, *e.g.*, cross-linking, co-immunoprecipitation, and co-purification through gradients or chromatographic columns. In addition, protein-protein interactions can be monitored by using a yeast-based genetic system described by Fields and co-workers (Fields and Song, Nature (London), 340: 245-246 (1989); Chien *et al.*, Proc. Natl. Acad. Sci. USA, 88: 9578-9582 (1991)) as disclosed by Chevray and Nathans, Proc. Natl. Acad. Sci. USA, 89: 5789-5793 (1991). Many transcriptional activators, such as yeast GAL4, consist of two physically discrete modular domains, one acting as the DNA-binding domain, the other one functioning as the transcription-activation domain. The yeast expression system described in the foregoing publications (generally referred to as the "two-hybrid system") takes advantage of this property, and employs two hybrid proteins, one in which the target protein is fused to the DNA-binding domain of GAL4, and another, in which candidate activating proteins are fused to the activation domain. The expression of a GAL1-*lacZ* reporter gene under control of a GAL4-activated promoter depends on reconstitution of GAL4 activity via protein-protein interaction. Colonies containing interacting polypeptides are detected with a chromogenic substrate for β -galactosidase. A complete kit (MATCHMAKER™) for identifying protein-protein interactions between two specific proteins using the two-hybrid technique is commercially available from Clontech. This system can also be extended to map protein domains involved in specific protein interactions as well as to pinpoint amino acid residues that are crucial for these interactions.

Compounds that interfere with the interaction of a gene encoding a PRO polypeptide identified herein and other intra- or extracellular components can be tested as follows: usually a reaction mixture is prepared containing the product of the gene and the intra- or extracellular component under conditions and for a time allowing for the interaction and binding of the two products. To test the ability of a candidate compound to inhibit binding, the reaction is run in the absence and in the presence of the test compound. In addition, a placebo may be added to a third reaction mixture, to serve as positive control. The binding (complex formation) between the test compound and the intra- or extracellular component present in the mixture is monitored as described hereinabove. The formation of a complex in the control reaction(s) but not in the reaction mixture containing the test compound indicates that the test compound interferes with the interaction of the test compound and its reaction partner.

If the PRO polypeptide has the ability to stimulate the proliferation of endothelial cells in the presence of the co-mitogen ConA, then one example of a screening method takes advantage of this ability. Specifically, in the proliferation assay, human umbilical vein endothelial cells are obtained and cultured in 96-well flat-bottomed culture plates (Costar, Cambridge, MA) and supplemented with a reaction mixture appropriate for facilitating proliferation of the cells, the mixture containing Con-A (Calbiochem, La Jolla, CA). Con-A and the compound to be screened are added and after incubation at 37°C, cultures are pulsed with ³H-thymidine and harvested onto glass fiber filters (phD; Cambridge Technology, Watertown, MA). Mean ³H-thymidine incorporation (cpm) of triplicate cultures is determined using a liquid scintillation counter (Beckman Instruments, Irvine, CA). Significant ³(H)-thymidine incorporation indicates stimulation of endothelial cell proliferation.

To assay for antagonists, the assay described above is performed; however, in this assay the PRO polypeptide is added along with the compound to be screened and the ability of the compound to inhibit ³(H)thymidine incorporation in the presence of the PRO polypeptide indicates that the compound is an antagonist to the PRO polypeptide. Alternatively, antagonists may be detected by combining the PRO polypeptide and a potential antagonist with membrane-bound PRO polypeptide receptors or recombinant receptors under appropriate conditions for a competitive inhibition assay. The PRO polypeptide can be labeled, such as by radioactivity, such that the number of PRO polypeptide molecules bound to the receptor can be used to determine the effectiveness of the potential antagonist. The gene encoding the receptor can be identified by numerous methods known to those of skill in the art, for example, ligand panning and FACS sorting. Coligan *et al.*, Current Protocols in Immun., 1(2): Chapter 5 (1991). Preferably, expression cloning is employed wherein polyadenylated RNA is prepared from a cell responsive to the PRO polypeptide and a cDNA library created from this RNA is divided into pools and used to transfect COS cells or other cells that are not responsive to the PRO polypeptide. Transfected cells that are grown on glass slides are exposed to the labeled PRO polypeptide. The PRO polypeptide can be labeled by a variety of means including iodination or inclusion of a recognition site for a site-specific protein kinase. Following fixation and incubation, the slides are subjected to autoradiographic analysis. Positive pools are identified and sub-pools are prepared and re-transfected using an interactive sub-pooling and re-screening process, eventually yielding a single clone that encodes the putative receptor.

As an alternative approach for receptor identification, the labeled PRO polypeptide can be photoaffinity-linked with cell membrane or extract preparations that express the receptor molecule. Cross-linked material is

resolved by PAGE and exposed to X-ray film. The labeled complex containing the receptor can be excised, resolved into peptide fragments, and subjected to protein micro-sequencing. The amino acid sequence obtained from micro-sequencing would be used to design a set of degenerate oligonucleotide probes to screen a cDNA library to identify the gene encoding the putative receptor.

5 In another assay for antagonists, mammalian cells or a membrane preparation expressing the receptor would be incubated with the labeled PRO polypeptide in the presence of the candidate compound. The ability of the compound to enhance or block this interaction could then be measured.

The compositions useful in the treatment of cardiovascular, endothelial, and angiogenic disorders include, without limitation, antibodies, small organic and inorganic molecules, peptides, phosphopeptides, antisense and
10 ribozyme molecules, triple-helix molecules, etc., that inhibit the expression and/or activity of the target gene product.

More specific examples of potential antagonists include an oligonucleotide that binds to the fusions of immunoglobulin with a PRO polypeptide, and, in particular, antibodies including, without limitation, poly- and monoclonal antibodies and antibody fragments, single-chain antibodies, anti-idiotypic antibodies, and chimeric or
15 humanized versions of such antibodies or fragments, as well as human antibodies and antibody fragments. Alternatively, a potential antagonist may be a closely related protein, for example, a mutated form of the PRO polypeptide that recognizes the receptor but imparts no effect, thereby competitively inhibiting the action of the PRO polypeptide.

Another potential PRO polypeptide antagonist is an antisense RNA or DNA construct prepared using
20 antisense technology, where, *e.g.*, an antisense RNA or DNA molecule acts to block directly the translation of mRNA by hybridizing to targeted mRNA and preventing protein translation. Antisense technology can be used to control gene expression through triple-helix formation or antisense DNA or RNA, both of which methods are based on binding of a polynucleotide to DNA or RNA. For example, the 5' coding portion of the polynucleotide sequence, which encodes the mature PRO polypeptides herein, is used to design an antisense RNA oligonucleotide
25 of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription (triple helix - *see, Lee et al., Nucl. Acids Res.*, 6:3073 (1979); Cooney *et al., Science*, 241: 456 (1988); Dervan *et al., Science*, 251:1360 (1991)), thereby preventing transcription and the production of the PRO polypeptide. A sequence "complementary" to a portion of an RNA, as referred to herein, means a sequence having sufficient complementarity to be able to hybridize with the RNA, forming a stable duplex;
30 in the case of double-stranded antisense nucleic acids, a single strand of the duplex DNA may thus be tested, or triplex helix formation may be assayed. The ability to hybridize will depend on both the degree of complementarity and the length of the antisense nucleic acid. Generally, the longer the hybridizing nucleic acid, the more base mismatches with an RNA it may contain and still form a stable duplex (or triplex, as the case may be). One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point
35 of the hybridized complex. The antisense RNA oligonucleotide hybridizes to the mRNA *in vivo* and blocks translation of the mRNA molecule into the PRO polypeptide (antisense - Okano, Neurochem., 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression (CRC Press: Boca Raton, FL, 1988).

The antisense oligonucleotides can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone, for example, to improve stability of the molecule, hybridization, etc. The oligonucleotide may include other appended groups such as peptides (*e.g.*, for targeting host cell receptors *in vivo*),
5 or agents facilitating transport across the cell membrane (see, *e.g.*, Letsinger, *et al.*, 1989, *Proc. Natl. Acad. Sci. U.S.A.* 86:6553-6556; Lemaitre, *et al.*, 1987, *Proc. Natl. Acad. Sci. U.S.A.* 84:648-652; PCT Publication No. WO88/09810, published December 15, 1988) or the blood-brain barrier (see, *e.g.*, PCT Publication No. WO89/10134, published April 25, 1988), hybridization-triggered cleavage agents (see, *e.g.*, Krol *et al.*, 1988, *BioTechniques* 6:958-976) or intercalating agents (see, *e.g.*, Zon, 1988, *Pharm. Res.* 5:539-549). To this end, the
10 oligonucleotide may be conjugated to another molecule, *e.g.*, a peptide, hybridization triggered cross-linking agent, transport agent, hybridization-triggered cleavage agent, etc.

The antisense oligonucleotide may comprise at least one modified base moiety which is selected from the group including but not limited to 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine,
15 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil,
20 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

The antisense oligonucleotide may also comprise at least one modified sugar moiety selected from the group including but not limited to arabinose, 2-fluoroarabinose, xylulose, and hexose.

In yet another embodiment, the antisense oligonucleotide comprises at least one modified phosphate backbone selected from the group consisting of a phosphorothioate, a phosphorodithioate, a phosphoramidothioate,
25 a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

In yet another embodiment, the antisense oligonucleotide is an α -anomeric oligonucleotide. An α -anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to
30 the usual β -units, the strands run parallel to each other (Gautier, *et al.*, 1987, *Nucl. Acids Res.* 15:6625-6641). The oligonucleotide is a 2'-O-methylribonucleotide (Inoue, *et al.*, 1987, *Nucl. Acids Res.* 15:6131-6148), or a chimeric RNA-DNA analogue (Inoue, *et al.*, 1987, *FEBS Lett.* 215:327-330).

Oligonucleotides of the invention may be synthesized by standard methods known in the art, *e.g.*, by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.).
35 As examples, phosphorothioate oligonucleotides may be synthesized by the method of Stein, *et al.* (1988, *Nucl. Acids Res.* 16:3209), methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin, *et al.*, 1988, *Proc. Natl. Acad. Sci. U.S.A.* 85:7448-7451), etc.

The oligonucleotides described above can also be delivered to cells such that the antisense RNA or DNA may be expressed *in vivo* to inhibit production of the PRO polypeptide. When antisense DNA is used, oligodeoxyribonucleotides derived from the translation-initiation site, *e.g.*, between about -10 and +10 positions of the target gene nucleotide sequence, are preferred.

Antisense RNA or DNA molecules are generally at least about 5 bases in length, about 10 bases in length, about 15 bases in length, about 20 bases in length, about 25 bases in length, about 30 bases in length, about 35 bases in length, about 40 bases in length, about 45 bases in length, about 50 bases in length, about 55 bases in length, about 60 bases in length, about 65 bases in length, about 70 bases in length, about 75 bases in length, about 80 bases in length, about 85 bases in length, about 90 bases in length, about 95 bases in length, about 100 bases in length, or more.

Potential antagonists further include small molecules that bind to the active site, the receptor binding site, or growth factor or other relevant binding site of the PRO polypeptide, thereby blocking the normal biological activity of the PRO polypeptide. Examples of small molecules include, but are not limited to, small peptides or peptide-like molecules, preferably soluble peptides, and synthetic non-peptidyl organic or inorganic compounds.

Additional potential antagonists are ribozymes, which are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. Ribozymes act by sequence-specific hybridization to the complementary target RNA, followed by endonucleolytic cleavage. Specific ribozyme cleavage sites within a potential RNA target can be identified by known techniques. For further details *see, e.g.*, Rossi, Current Biology, 4: 469-471 (1994), and PCT publication No. WO 97/33551 (published September 18, 1997).

While ribozymes that cleave mRNA at site specific recognition sequences can be used to destroy target gene mRNAs, the use of hammerhead ribozymes is preferred. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions which form complementary base pairs with the target mRNA. The sole requirement is that the target mRNA have the following sequence of two bases: 5'-UG-3'. The construction and production of hammerhead ribozymes is well known in the art and is described more fully in Myers, 1995, *Molecular Biology and Biotechnology: A Comprehensive Desk Reference*, VCH Publishers, New York, (see especially Figure 4, page 833) and in Haseloff and Gerlach, 1988, *Nature*, 334:585-591, which is incorporated herein by reference in its entirety.

Preferably the ribozyme is engineered so that the cleavage recognition site is located near the 5' end of the target gene mRNA, *i.e.*, to increase efficiency and minimize the intracellular accumulation of non-functional mRNA transcripts.

The ribozymes of the present invention also include RNA endoribonucleases (hereinafter "Cech-type ribozymes") such as the one which occurs naturally in *Tetrahymena thermophila* (known as the IVS, or L-19 IVS RNA) and which has been extensively described by Thomas Cech and collaborators (Zaug, *et al.*, 1984, *Science*, 224:574-578; Zaug and Cech, 1986, *Science*, 231:470-475; Zaug, *et al.*, 1986, *Nature*, 324:429-433; published International patent application No. WO 88/04300 by University Patents Inc.; Been and Cech, 1986, *Cell*, 47:207-216). The Cech-type ribozymes have an eight base pair active site that hybridizes to a target RNA sequence

whereafter cleavage of the target RNA takes place. The invention encompasses those Cech-type ribozymes that target eight base-pair active site sequences that are present in the target gene.

As in the antisense approach, the ribozymes can be composed of modified oligonucleotides (*e.g.*, for improved stability, targeting, *etc.*) and should be delivered to cells that express the target gene *in vivo*. A preferred method of delivery involves using a DNA construct "encoding" the ribozyme under the control of a strong constitutive pol III or pol II promoter, so that transfected cells will produce sufficient quantities of the ribozyme to destroy endogenous target gene messages and inhibit translation. Because ribozymes, unlike antisense molecules, are catalytic, a lower intracellular concentration is required for efficiency.

Nucleic acid molecules in triple-helix formation used to inhibit transcription should be single-stranded and composed of deoxynucleotides. The base composition of these oligonucleotides is designed such that it promotes triple-helix formation via Hoogsteen base-pairing rules, which generally require sizeable stretches of purines or pyrimidines on one strand of a duplex. For further details *see, e.g.*, PCT publication No. WO 97/33551, *supra*.

These small molecules can be identified by any one or more of the screening assays discussed hereinabove and/or by any other screening techniques well known for those skilled in the art.

5.2.4.10. Types of Cardiovascular, Endothelial, and Angiogenic Disorders to be Treated

The PRO polypeptides, or agonists or antagonists thereto, that have activity in the cardiovascular, angiogenic, and endothelial assays described herein, and/or whose gene product has been found to be localized to the cardiovascular system, are likely to have therapeutic uses in a variety of cardiovascular, endothelial, and angiogenic disorders, including systemic disorders that affect vessels, such as diabetes mellitus. Their therapeutic utility could include diseases of the arteries, capillaries, veins, and/or lymphatics. Examples of treatments hereunder include treating muscle wasting disease, treating osteoporosis, aiding in implant fixation to stimulate the growth of cells around the implant and therefore facilitate its attachment to its intended site, increasing IGF stability in tissues or in serum, if applicable, and increasing binding to the IGF receptor (since IGF has been shown *in vitro* to enhance human marrow erythroid and granulocytic progenitor cell growth).

The PRO polypeptides or agonists or antagonists thereto may also be employed to stimulate erythropoiesis or granulopoiesis, to stimulate wound healing or tissue regeneration and associated therapies concerned with re-growth of tissue, such as connective tissue, skin, bone, cartilage, muscle, lung, or kidney, to promote angiogenesis, to stimulate or inhibit migration of endothelial cells, and to proliferate the growth of vascular smooth muscle and endothelial cell production. The increase in angiogenesis mediated by the PRO polypeptide or agonist would be beneficial to ischemic tissues and to collateral coronary development in the heart subsequent to coronary stenosis. Antagonists are used to inhibit the action of such polypeptides, for example, to limit the production of excess connective tissue during wound healing or pulmonary fibrosis if the PRO polypeptide promotes such production. This would include treatment of acute myocardial infarction and heart failure.

Moreover, the present invention provides the treatment of cardiac hypertrophy, regardless of the underlying cause, by administering a therapeutically effective dose of the PRO polypeptide, or agonist or antagonist thereto. If the objective is the treatment of human patients, the PRO polypeptide preferably is recombinant human PRO

polypeptide (rhPRO polypeptide). The treatment for cardiac hypertrophy can be performed at any of its various stages, which may result from a variety of diverse pathologic conditions, including myocardial infarction, hypertension, hypertrophic cardiomyopathy, and valvular regurgitation. The treatment extends to all stages of the progression of cardiac hypertrophy, with or without structural damage of the heart muscle, regardless of the underlying cardiac disorder.

The decision of whether to use the molecule itself or an agonist thereof for any particular indication, as opposed to an antagonist to the molecule, would depend mainly on whether the molecule herein promotes cardiovascularization, genesis of endothelial cells, or angiogenesis or inhibits these conditions. For example, if the molecule promotes angiogenesis, an antagonist thereof would be useful for treatment of disorders where it is desired to limit or prevent angiogenesis. Examples of such disorders include vascular tumors such as haemangioma, tumor angiogenesis, neovascularization in the retina, choroid, or cornea, associated with diabetic retinopathy or premature infant retinopathy or macular degeneration and proliferative vitreoretinopathy, rheumatoid arthritis, Crohn's disease, atherosclerosis, ovarian hyperstimulation, psoriasis, endometriosis associated with neovascularization, restenosis subsequent to balloon angioplasty, scar tissue overproduction, for example, that seen in a keloid that forms after surgery, fibrosis after myocardial infarction, or fibrotic lesions associated with pulmonary fibrosis.

If, however, the molecule inhibits angiogenesis, it would be expected to be used directly for treatment of the above conditions.

On the other hand, if the molecule stimulates angiogenesis it would be used itself (or an agonist thereof) for indications where angiogenesis is desired such as peripheral vascular disease, hypertension, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, arterial restenosis, thrombophlebitis, lymphangitis, lymphedema, wound healing and tissue repair, ischemia reperfusion injury, angina, myocardial infarctions such as acute myocardial infarctions, chronic heart conditions, heart failure such as congestive heart failure, and osteoporosis.

If, however, the molecule inhibits angiogenesis, an antagonist thereof would be used for treatment of those conditions where angiogenesis is desired.

Specific types of diseases are described below, where the PRO polypeptide herein or agonists or antagonists thereof may serve as useful for vascular-related drug targeting or as therapeutic targets for the treatment or prevention of the disorders. Atherosclerosis is a disease characterized by accumulation of plaques of intimal thickening in arteries, due to accumulation of lipids, proliferation of smooth muscle cells, and formation of fibrous tissue within the arterial wall. The disease can affect large, medium, and small arteries in any organ. Changes in endothelial and vascular smooth muscle cell function are known to play an important role in modulating the accumulation and regression of these plaques.

Hypertension is characterized by raised vascular pressure in the systemic arterial, pulmonary arterial, or portal venous systems. Elevated pressure may result from or result in impaired endothelial function and/or vascular disease.

Inflammatory vasculitides include giant cell arteritis, Takayasu's arteritis, polyarteritis nodosa (including the microangiopathic form), Kawasaki's disease, microscopic polyangiitis, Wegener's granulomatosis, and a variety

of infectious-related vascular disorders (including Henoch-Schonlein purpura). Altered endothelial cell function has been shown to be important in these diseases.

5 Reynaud's disease and Reynaud's phenomenon are characterized by intermittent abnormal impairment of the circulation through the extremities on exposure to cold. Altered endothelial cell function has been shown to be important in this disease.

Aneurysms are saccular or fusiform dilatations of the arterial or venous tree that are associated with altered endothelial cell and/or vascular smooth muscle cells.

Arterial restenosis (restenosis of the arterial wall) may occur following angioplasty as a result of alteration in the function and proliferation of endothelial and vascular smooth muscle cells.

10 Thrombophlebitis and lymphangitis are inflammatory disorders of veins and lymphatics, respectively, that may result from, and/or in, altered endothelial cell function. Similarly, lymphedema is a condition involving impaired lymphatic vessels resulting from endothelial cell function.

The family of benign and malignant vascular tumors are characterized by abnormal proliferation and growth of cellular elements of the vascular system. For example, lymphangiomas are benign tumors of the lymphatic system that are congenital, often cystic, malformations of the lymphatics that usually occur in newborns. Cystic tumors tend to grow into the adjacent tissue. Cystic tumors usually occur in the cervical and axillary region. They can also occur in the soft tissue of the extremities. The main symptoms are dilated, sometimes reticular, structured lymphatics and lymphocysts surrounded by connective tissue. Lymphangiomas are assumed to be caused by improperly connected embryonic lymphatics or their deficiency. The result is impaired local lymph drainage.

15 Griener *et al.*, Lymphology, 4: 140-144 (1971).

Another use for the PRO polypeptides herein or agonists or antagonists thereto is in the prevention of tumor angiogenesis, which involves vascularization of a tumor to enable it to grow and/or metastasize. This process is dependent on the growth of new blood vessels. Examples of neoplasms and related conditions that involve tumor angiogenesis include breast carcinomas, lung carcinomas, gastric carcinomas, esophageal carcinomas, colorectal carcinomas, liver carcinomas, ovarian carcinomas, thecomas, arrhenoblastomas, cervical carcinomas, endometrial carcinoma, endometrial hyperplasia, endometriosis, fibrosarcomas, choriocarcinoma, head and neck cancer, nasopharyngeal carcinoma, laryngeal carcinomas, hepatoblastoma, Kaposi's sarcoma, melanoma, skin carcinomas, hemangioma, cavernous hemangioma, hemangioblastoma, pancreas carcinomas, retinoblastoma, astrocytoma, glioblastoma, Schwannoma, oligodendroglioma, medulloblastoma, neuroblastomas, rhabdomyosarcoma, osteogenic sarcoma, leiomyosarcomas, urinary tract carcinomas, thyroid carcinomas, Wilm's tumor, renal cell carcinoma, prostate carcinoma, abnormal vascular proliferation associated with phakomatoses, edema (such as that associated with brain tumors), and Meigs' syndrome.

25

30

Age-related macular degeneration (AMD) is a leading cause of severe visual loss in the elderly population. The exudative form of AMD is characterized by choroidal neovascularization and retinal pigment epithelial cell detachment. Because choroidal neovascularization is associated with a dramatic worsening in prognosis, the PRO polypeptide or agonist or antagonist thereto is expected to be useful in reducing the severity of AMD.

35

Healing of trauma such as wound healing and tissue repair is also a targeted use for the PRO polypeptides herein or their agonists or antagonists. Formation and regression of new blood vessels is essential for tissue healing and repair. This category includes bone, cartilage, tendon, ligament, and/or nerve tissue growth or regeneration, as well as wound healing and tissue repair and replacement, and in the treatment of burns, incisions, and ulcers.

5 A PRO polypeptide or agonist or antagonist thereof that induces cartilage and/or bone growth in circumstances where bone is not normally formed has application in the healing of bone fractures and cartilage damage or defects in humans and other animals. Such a preparation employing a PRO polypeptide or agonist or antagonist thereof may have prophylactic use in closed as well as open fracture reduction and also in the improved fixation of artificial joints. *De novo* bone formation induced by an osteogenic agent contributes to the repair of congenital, trauma-

10 induced, or oncologic, resection-induced craniofacial defects, and also is useful in cosmetic plastic surgery.

PRO polypeptides or agonists or antagonists thereto may also be useful to promote better or faster closure of non-healing wounds, including without limitation pressure ulcers, ulcers associated with vascular insufficiency, surgical and traumatic wounds, and the like.

It is expected that a PRO polypeptide or agonist or antagonist thereto may also exhibit activity for

15 generation or regeneration of other tissues, such as organs (including, for example, pancreas, liver, intestine, kidney, skin, or endothelium), muscle (smooth, skeletal, or cardiac), and vascular (including vascular endothelium) tissue, or for promoting the growth of cells comprising such tissues. Part of the desired effects may be by inhibition or modulation of fibrotic scarring to allow normal tissue to regenerate.

A PRO polypeptide herein or agonist or antagonist thereto may also be useful for gut protection or

20 regeneration and treatment of lung or liver fibrosis, reperfusion injury in various tissues, and conditions resulting from systemic cytokine damage. Also, the PRO polypeptide or agonist or antagonist thereto may be useful for promoting or inhibiting differentiation of tissues described above from precursor tissues or cells, or for inhibiting the growth of tissues described above.

A PRO polypeptide or agonist or antagonist thereto may also be used in the treatment of periodontal

25 diseases and in other tooth-repair processes. Such agents may provide an environment to attract bone-forming cells, stimulate growth of bone-forming cells, or induce differentiation of progenitors of bone-forming cells. A PRO polypeptide herein or an agonist or an antagonist thereto may also be useful in the treatment of osteoporosis or osteoarthritis, such as through stimulation of bone and/or cartilage repair or by blocking inflammation or processes of tissue destruction (collagenase activity, osteoclast activity, etc.) mediated by inflammatory processes, since blood

30 vessels play an important role in the regulation of bone turnover and growth.

Another category of tissue regeneration activity that may be attributable to the PRO polypeptide herein or agonist or antagonist thereto is tendon/ligament formation. A protein that induces tendon/ligament-like tissue or other tissue formation in circumstances where such tissue is not normally formed has application in the healing of tendon or ligament tears, deformities, and other tendon or ligament defects in humans and other animals. Such

35 a preparation may have prophylactic use in preventing damage to tendon or ligament tissue, as well as use in the improved fixation of tendon or ligament to bone or other tissues, and in repairing defects to tendon or ligament tissue. *De novo* tendon/ligament-like tissue formation induced by a composition of the PRO polypeptide herein or

agonist or antagonist thereto contributes to the repair of congenital, trauma-induced, or other tendon or ligament defects of other origin, and is also useful in cosmetic plastic surgery for attachment or repair of tendons or ligaments. The compositions herein may provide an environment to attract tendon- or ligament-forming cells, stimulate growth of tendon- or ligament-forming cells, induce differentiation of progenitors of tendon- or ligament-forming cells, or induce growth of tendon/ligament cells or progenitors *ex vivo* for return *in vivo* to effect tissue repair. The compositions herein may also be useful in the treatment of tendinitis, carpal tunnel syndrome, and other tendon or ligament defects. The compositions may also include an appropriate matrix and/or sequestering agent as a carrier as is well known in the art.

The PRO polypeptide or its agonist or antagonist may also be useful for proliferation of neural cells and for regeneration of nerve and brain tissue, *i.e.*, for the treatment of central and peripheral nervous system disease and neuropathies, as well as mechanical and traumatic disorders, that involve degeneration, death, or trauma to neural cells or nerve tissue. More specifically, a PRO polypeptide or its agonist or antagonist may be used in the treatment of diseases of the peripheral nervous system, such as peripheral nerve injuries, peripheral neuropathy and localized neuropathies, and central nervous system diseases, such as Alzheimer's, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome. Further conditions that may be treated in accordance with the present invention include mechanical and traumatic disorders, such as spinal cord disorders, head trauma, and cerebrovascular diseases such as stroke. Peripheral neuropathies resulting from chemotherapy or other medical therapies may also be treatable using a PRO polypeptide herein or agonist or antagonist thereto.

Ischemia-reperfusion injury is another indication. Endothelial cell dysfunction may be important in both the initiation of, and in regulation of the sequelae of events that occur following ischemia-reperfusion injury.

Rheumatoid arthritis is a further indication. Blood vessel growth and targeting of inflammatory cells through the vasculature is an important component in the pathogenesis of rheumatoid and sero-negative forms of arthritis.

A PRO polypeptide or its agonist or antagonist may also be administered prophylactically to patients with cardiac hypertrophy, to prevent the progression of the condition, and avoid sudden death, including death of asymptomatic patients. Such preventative therapy is particularly warranted in the case of patients diagnosed with massive left ventricular cardiac hypertrophy (a maximal wall thickness of 35 mm or more in adults, or a comparable value in children), or in instances when the hemodynamic burden on the heart is particularly strong.

A PRO polypeptide or its agonist or antagonist may also be useful in the management of atrial fibrillation, which develops in a substantial portion of patients diagnosed with hypertrophic cardiomyopathy.

Further indications include angina, myocardial infarctions such as acute myocardial infarctions, and heart failure such as congestive heart failure. Additional non-neoplastic conditions include psoriasis, diabetic and other proliferative retinopathies including retinopathy of prematurity, retrolental fibroplasia, neovascular glaucoma, thyroid hyperplasias (including Grave's disease), corneal and other tissue transplantation, chronic inflammation, lung inflammation, nephrotic syndrome, preeclampsia, ascites, pericardial effusion (such as that associated with pericarditis), and pleural effusion.

In view of the above, the PRO polypeptides or agonists or antagonists thereof described herein, which are shown to alter or impact endothelial cell function, proliferation, and/or form, are likely to play an important role in the etiology and pathogenesis of many or all of the disorders noted above, and as such can serve as therapeutic targets to augment or inhibit these processes or for vascular-related drug targeting in these disorders.

5

5.2.4.11. Administration Protocols, Schedules, Doses, and Formulations

The molecules herein and agonists and antagonists thereto are pharmaceutically useful as a prophylactic and therapeutic agent for various disorders and diseases as set forth above.

Therapeutic compositions of the PRO polypeptides or agonists or antagonists are prepared for storage by mixing the desired molecule having the appropriate degree of purity with optional pharmaceutically acceptable carriers, excipients, or stabilizers (Remington's Pharmaceutical Sciences, 16th edition, Osol, A. ed. (1980)), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (*e.g.*, Zn-protein complexes); and/or non-ionic surfactants such as TWEEN™, PLURONICS™ or polyethylene glycol (PEG).

Additional examples of such carriers include ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts, or electrolytes such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, and polyethylene glycol. Carriers for topical or gel-based forms of agonist or antagonist include polysaccharides such as sodium carboxymethylcellulose or methylcellulose, polyvinylpyrrolidone, polyacrylates, polyoxyethylene-polyoxypropylene-block polymers, polyethylene glycol, and wood wax alcohols. For all administrations, conventional depot forms are suitably used. Such forms include, for example, microcapsules, nano-capsules, liposomes, plasters, inhalation forms, nose sprays, sublingual tablets, and sustained-release preparations. The PRO polypeptides or agonists or antagonists will typically be formulated in such vehicles at a concentration of about 0.1 mg/ml to 100 mg/ml.

Another formulation comprises incorporating a PRO polypeptide or agonist or antagonist thereof into formed articles. Such articles can be used in modulating endothelial cell growth and angiogenesis. In addition, tumor invasion and metastasis may be modulated with these articles.

PRO polypeptides or agonists or antagonists to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes, prior to or following lyophilization and reconstitution. PRO polypeptides ordinarily will be stored in lyophilized form or in solution if administered systemically. If in lyophilized form, the PRO polypeptide or agonist or antagonist thereto is typically formulated in combination with other ingredients for reconstitution with an appropriate diluent at the time for use. An example of a liquid formulation of a PRO polypeptide or agonist or antagonist is a sterile, clear, colorless unpreserved solution filled in a single-dose vial for subcutaneous injection. Preserved pharmaceutical compositions suitable for repeated use may contain, for example, depending mainly on the indication and type of polypeptide:

- a) PRO polypeptide or agonist or antagonist thereto;
- b) a buffer capable of maintaining the pH in a range of maximum stability of the polypeptide or other molecule in solution, preferably about 4-8;
- c) a detergent/surfactant primarily to stabilize the polypeptide or molecule against agitation-induced aggregation;
- d) an isotonicifier;
- e) a preservative selected from the group of phenol, benzyl alcohol and a benzethonium halide, *e.g.*, chloride; and
- f) water.

If the detergent employed is non-ionic, it may, for example, be polysorbates (*e.g.*, POLYSORBATE™ (TWEEN™) 20, 80, etc.) or poloxamers (*e.g.*, POLOXAMER™ 188). The use of non-ionic surfactants permits the formulation to be exposed to shear surface stresses without causing denaturation of the polypeptide. Further, such surfactant-containing formulations may be employed in aerosol devices such as those used in a pulmonary dosing, and needleless jet injector guns (*see, e.g.*, EP 257,956).

An isotonicifier may be present to ensure isotonicity of a liquid composition of the PRO polypeptide or agonist or antagonist thereto, and includes polyhydric sugar alcohols, preferably trihydric or higher sugar alcohols, such as glycerin, erythritol, arabitol, xylitol, sorbitol, and mannitol. These sugar alcohols can be used alone or in combination. Alternatively, sodium chloride or other appropriate inorganic salts may be used to render the solutions isotonic.

The buffer may, for example, be an acetate, citrate, succinate, or phosphate buffer depending on the pH desired. The pH of one type of liquid formulation of this invention is buffered in the range of about 4 to 8, preferably about physiological pH.

The preservatives phenol, benzyl alcohol and benzethonium halides, *e.g.*, chloride, are known antimicrobial agents that may be employed.

Therapeutic PRO polypeptide compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle. The formulations are preferably administered as repeated intravenous (i.v.), subcutaneous (s.c.), or intramuscular (i.m.) injections, or as aerosol formulations suitable for intranasal or intrapulmonary delivery (for intrapulmonary delivery *see, e.g.*, EP 257,956).

PRO polypeptides can also be administered in the form of sustained-released preparations. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the protein, which matrices are in the form of shaped articles, *e.g.*, films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (*e.g.*, poly(2-hydroxyethyl-methacrylate) as described by Langer *et al.*, J. Biomed. Mater. Res., 15: 167-277 (1981) and Langer, Chem. Tech., 12: 98-105 (1982) or poly(vinylalcohol)), polylactides (U.S. Patent No. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma ethyl-L-glutamate (Sidman *et al.*, Biopolymers, 22: 547-556 (1983)), non-degradable ethylene-vinyl acetate (Langer *et al.*, *supra*), degradable lactic acid-glycolic acid copolymers such as the Lupron Depot™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid (EP 133,988).

While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated proteins remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37°C, resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for protein stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S-S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

Sustained-release PRO polypeptide compositions also include liposomally entrapped PRO polypeptides. Liposomes containing the PRO polypeptide are prepared by methods known *per se*: DE 3,218,121; Epstein *et al.*, Proc. Natl. Acad. Sci. USA, 82: 3688-3692 (1985); Hwang *et al.*, Proc. Natl. Acad. Sci. USA, 77: 4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese patent application 83-118008; U.S. Patent Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. % cholesterol, the selected proportion being adjusted for the optimal therapy.

The therapeutically effective dose of a PRO polypeptide or agonist or antagonist thereto will, of course, vary depending on such factors as the pathological condition to be treated (including prevention), the method of administration, the type of compound being used for treatment, any co-therapy involved, the patient's age, weight, general medical condition, medical history, etc., and its determination is well within the skill of a practicing physician. Accordingly, it will be necessary for the therapist to titer the dosage and modify the route of administration as required to obtain the maximal therapeutic effect. If the PRO polypeptide has a narrow host range, for the treatment of human patients formulations comprising human PRO polypeptide, more preferably native-sequence human PRO polypeptide, are preferred. The clinician will administer the PRO polypeptide until a dosage is reached that achieves the desired effect for treatment of the condition in question. For example, if the objective is the treatment of CHF, the amount would be one that inhibits the progressive cardiac hypertrophy associated with this condition. The progress of this therapy is easily monitored by echo cardiography. Similarly, in patients with hypertrophic cardiomyopathy, the PRO polypeptide can be administered on an empirical basis.

With the above guidelines, the effective dose generally is within the range of from about 0.001 to about 1.0 mg/kg, more preferably about 0.01-1.0 mg/kg, most preferably about 0.01-0.1 mg/kg.

For non-oral use in treating human adult hypertension, it is advantageous to administer the PRO polypeptide in the form of an injection at about 0.01 to 50 mg, preferably about 0.05 to 20 mg, most preferably 1 to 20 mg, per kg body weight, 1 to 3 times daily by intravenous injection. For oral administration, a molecule based on the PRO polypeptide is preferably administered at about 5 mg to 1 g, preferably about 10 to 100 mg, per kg body weight, 1 to 3 times daily. It should be appreciated that endotoxin contamination should be kept minimally at a safe level, for example, less than 0.5 ng/mg protein. Moreover, for human administration, the formulations preferably meet sterility, pyrogenicity, general safety, and purity as required by FDA Office and Biologics standards.

The dosage regimen of a pharmaceutical composition containing the PRO polypeptide to be used in tissue regeneration will be determined by the attending physician considering various factors that modify the action of the polypeptides, *e.g.*, amount of tissue weight desired to be formed, the site of damage, the condition of the damaged tissue, the size of a wound, type of damaged tissue (*e.g.*, bone), the patient's age, sex, and diet, the severity of any infection, time of administration, and other clinical factors. The dosage may vary with the type of matrix used in the reconstitution and with inclusion of other proteins in the pharmaceutical composition. For example, the addition of other known growth factors, such as IGF-I, to the final composition may also affect the dosage. Progress can be monitored by periodic assessment of tissue/bone growth and/or repair, for example, X-rays, histomorphometric determinations, and tetracycline labeling.

The route of PRO polypeptide or antagonist or agonist administration is in accord with known methods, *e.g.*, by injection or infusion by intravenous, intramuscular, intracerebral, intraperitoneal, intracerebrospinal, subcutaneous, intraocular, intraarticular, intrasynovial, intrathecal, oral, topical, or inhalation routes, or by sustained-release systems as noted below. The PRO polypeptide or agonist or antagonists thereof also are suitably administered by intratumoral, peritumoral, intralesional, or perilesional routes, to exert local as well as systemic therapeutic effects. The intraperitoneal route is expected to be particularly useful, for example, in the treatment of ovarian tumors.

If a peptide or small molecule is employed as an antagonist or agonist, it is preferably administered orally or non-orally in the form of a liquid or solid to mammals.

Examples of pharmacologically acceptable salts of molecules that form salts and are useful hereunder include alkali metal salts (*e.g.*, sodium salt, potassium salt), alkaline earth metal salts (*e.g.*, calcium salt, magnesium salt), ammonium salts, organic base salts (*e.g.*, pyridine salt, triethylamine salt), inorganic acid salts (*e.g.*, hydrochloride, sulfate, nitrate), and salts of organic acid (*e.g.*, acetate, oxalate, p-toluenesulfonate).

For compositions herein that are useful for bone, cartilage, tendon, or ligament regeneration, the therapeutic method includes administering the composition topically, systemically, or locally as an implant or device. When administered, the therapeutic composition for use is in a pyrogen-free, physiologically acceptable form. Further, the composition may desirably be encapsulated or injected in a viscous form for delivery to the site of bone, cartilage, or tissue damage. Topical administration may be suitable for wound healing and tissue repair. Preferably, for bone and/or cartilage formation, the composition would include a matrix capable of delivering the

protein-containing composition to the site of bone and/or cartilage damage, providing a structure for the developing bone and cartilage and preferably capable of being resorbed into the body. Such matrices may be formed of materials presently in use for other implanted medical applications.

5 The choice of matrix material is based on biocompatibility, biodegradability, mechanical properties, cosmetic appearance, and interface properties. The particular application of the compositions will define the appropriate formulation. Potential matrices for the compositions may be biodegradable and chemically defined calcium sulfate, tricalcium phosphate, hydroxyapatite, polylactic acid, polyglycolic acid, and polyanhydrides. Other potential materials are biodegradable and biologically well-defined, such as bone or dermal collagen. Further matrices are comprised of pure proteins or extracellular matrix components. Other potential matrices are
10 nonbiodegradable and chemically defined, such as sintered hydroxyapatite, bioglass, aluminates, or other ceramics. Matrices may be comprised of combinations of any of the above-mentioned types of material, such as polylactic acid and hydroxyapatite or collagen and tricalcium phosphate. The bioceramics may be altered in composition, such as in calcium-aluminate-phosphate and processing to alter pore size, particle size, particle shape, and biodegradability.

15 One specific embodiment is a 50:50 (mole weight) copolymer of lactic acid and glycolic acid in the form of porous particles having diameters ranging from 150 to 800 microns. In some applications, it will be useful to utilize a sequestering agent, such as carboxymethyl cellulose or autologous blood clot, to prevent the polypeptide compositions from disassociating from the matrix.

20 One suitable family of sequestering agents is cellulosic materials such as alkylcelluloses (including hydroxyalkylcelluloses), including methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose, and carboxymethylcellulose, one preferred being cationic salts of carboxymethylcellulose (CMC). Other preferred sequestering agents include hyaluronic acid, sodium alginate, poly(ethylene glycol), polyoxyethylene oxide, carboxyvinyl polymer, and poly(vinyl alcohol). The amount of sequestering agent useful herein is 0.5-20 wt%, preferably 1-10 wt%, based on total formulation weight, which
25 represents the amount necessary to prevent desorption of the polypeptide (or its antagonist) from the polymer matrix and to provide appropriate handling of the composition, yet not so much that the progenitor cells are prevented from infiltrating the matrix, thereby providing the polypeptide (or its antagonist) the opportunity to assist the osteogenic activity of the progenitor cells.

5.2.4.12. Combination Therapies

30 The effectiveness of the PRO polypeptide or an agonist or antagonist thereof in preventing or treating the disorder in question may be improved by administering the active agent serially or in combination with another agent that is effective for those purposes, either in the same composition or as separate compositions.

For example, for treatment of cardiac hypertrophy, PRO polypeptide therapy can be combined with the administration of inhibitors of known cardiac myocyte hypertrophy factors, *e.g.*, inhibitors of α -adrenergic agonists
35 such as phenylephrine; endothelin-1 inhibitors such as BOSENTANTM and MOXONODINTM; inhibitors to CT-1

(U.S. Pat. No. 5,679,545); inhibitors to LIF; ACE inhibitors; des-aspartate-angiotensin I inhibitors (U.S. Pat. No. 5,773,415), and angiotensin II inhibitors.

For treatment of cardiac hypertrophy associated with hypertension, the PRO polypeptide can be administered in combination with β -adrenergic receptor blocking agents, *e.g.*, propranolol, timolol, tertalolol, carteolol, nadolol, betaxolol, penbutolol, acetobutolol, atenolol, metoprolol, or carvedilol; ACE inhibitors, *e.g.*, quinapril, captopril, enalapril, ramipril, benazepril, fosinopril, or lisinopril; diuretics, *e.g.*, chlorothiazide, hydrochlorothiazide, hydroflumethazide, methylchlorothiazide, benzthiazide, dichlorphenamide, acetazolamide, or indapamide; and/or calcium channel blockers, *e.g.*, diltiazem, nifedipine, verapamil, or nocardipine. Pharmaceutical compositions comprising the therapeutic agents identified herein by their generic names are commercially available, and are to be administered following the manufacturers' instructions for dosage, administration, adverse effects, contraindications, etc. *See, e.g., Physicians' Desk Reference* (Medical Economics Data Production Co.: Montvale, N.J., 1997), 51th Edition.

Preferred candidates for combination therapy in the treatment of hypertrophic cardiomyopathy are β -adrenergic-blocking drugs (*e.g.*, propranolol, timolol, tertalolol, carteolol, nadolol, betaxolol, penbutolol, acetobutolol, atenolol, metoprolol, or carvedilol), verapamil, difedipine, or diltiazem. Treatment of hypertrophy associated with high blood pressure may require the use of antihypertensive drug therapy, using calcium channel blockers, *e.g.*, diltiazem, nifedipine, verapamil, or nocardipine; β -adrenergic blocking agents; diuretics, *e.g.*, chlorothiazide, hydrochlorothiazide, hydroflumethazide, methylchlorothiazide, benzthiazide, dichlorphenamide, acetazolamide, or indapamide; and/or ACE-inhibitors, *e.g.*, quinapril, captopril, enalapril, ramipril, benazepril, fosinopril, or lisinopril.

For other indications, PRO polypeptides or their agonists or antagonists may be combined with other agents beneficial to the treatment of the bone and/or cartilage defect, wound, or tissue in question. These agents include various growth factors such as EGF, PDGF, TGF- α or TGF- β , IGF, FGF, and CTGF.

In addition, PRO polypeptides or their agonists or antagonists used to treat cancer may be combined with cytotoxic, chemotherapeutic, or growth-inhibitory agents as identified above. Also, for cancer treatment, the PRO polypeptide or agonist or antagonist thereof is suitably administered serially or in combination with radiological treatments, whether involving irradiation or administration of radioactive substances.

The effective amounts of the therapeutic agents administered in combination with the PRO polypeptide or agonist or antagonist thereof will be at the physician's or veterinarian's discretion. Dosage administration and adjustment is done to achieve maximal management of the conditions to be treated. For example, for treating hypertension, these amounts ideally take into account use of diuretics or digitalis, and conditions such as hyper- or hypotension, renal impairment, etc. The dose will additionally depend on such factors as the type of the therapeutic agent to be used and the specific patient being treated. Typically, the amount employed will be the same dose as that used, if the given therapeutic agent is administered without the PRO polypeptide.

5.2.4.13. Articles of Manufacture

An article of manufacture such as a kit containing the PRO polypeptide or agonists or antagonists thereof useful for the diagnosis or treatment of the disorders described above comprises at least a container and a label. Suitable containers include, for example, bottles, vials, syringes, and test tubes. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition that is effective for diagnosing or treating the condition and may have a sterile access port (for example, the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). The active agent in the composition is the PRO polypeptide or an agonist or antagonist thereto. The label on, or associated with, the container indicates that the composition is used for diagnosing or treating the condition of choice. The article of manufacture may further comprise a second container comprising a pharmaceutically-acceptable buffer, such as phosphate-buffered saline, Ringer's solution, and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, syringes, and package inserts with instructions for use. The article of manufacture may also comprise a second or third container with another active agent as described above.

5.2.5. Antibodies

Some of the most promising drug candidates according to the present invention are antibodies and antibody fragments that may inhibit the production or the gene product of the genes identified herein and/or reduce the activity of the gene products.

5.2.5.1. Polyclonal Antibodies

Methods of preparing polyclonal antibodies are known to the skilled artisan. Polyclonal antibodies can be raised in a mammal, for example, by one or more injections of an immunizing agent and, if desired, an adjuvant. Typically, the immunizing agent and/or adjuvant will be injected in the mammal by multiple subcutaneous or intraperitoneal injections. The immunizing agent may include the PRO polypeptide or a fusion protein thereof. It may be useful to conjugate the immunizing agent to a protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include, but are not limited to, keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. Examples of adjuvants that may be employed include Freund's complete adjuvant and MPL-TDM adjuvant (monophosphoryl Lipid A or synthetic trehalose dicorynomycolate). The immunization protocol may be selected by one skilled in the art without undue experimentation.

5.2.5.2. Monoclonal Antibodies

The anti-PRO antibodies may, alternatively, be monoclonal antibodies. Monoclonal antibodies may be prepared using hybridoma methods, such as those described by Kohler and Milstein, *Nature*, 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal is typically immunized with an

immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes may be immunized *in vitro*.

5 The immunizing agent will typically include the PRO polypeptide or a fusion protein thereof. Generally, either peripheral blood lymphocytes ("PBLs") are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell. Goding, Monoclonal Antibodies: Principles and Practice (New York: Academic Press, 1986), pp. 59-103. Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine, and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells may be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, 10 immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells:

15 Preferred immortalized cell lines are those that fuse efficiently, support stable high-level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, California and the American Type Culture Collection, Manassas, Virginia. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies. Kozbor, J. Immunol., 133:3001 (1984); Brodeur *et al.*, Monoclonal Antibody Production Techniques and Applications (Marcel Dekker, Inc.: New York, 1987) pp. 51-63. 20

The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal antibodies directed against the PRO polypeptide. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson and Pollard, Anal. Biochem., 107:220 (1980). 25

After the desired hybridoma cells are identified, the clones may be subcloned by limiting dilution procedures and grown by standard methods. Goding, *supra*. Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium. Alternatively, the hybridoma cells may be grown *in vivo* as ascites in a mammal. 30

The monoclonal antibodies secreted by the subclones may be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

35 The monoclonal antibodies may also be made by recombinant DNA methods, such as those described in U.S. Patent No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the

invention serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also may be modified, for example, by substituting the coding sequence for human heavy- and light-chain constant domains in place of the homologous murine sequences (U.S. Patent No. 4,816,567; Morrison *et al.*, *supra*) or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

The antibodies may be monovalent antibodies. Methods for preparing monovalent antibodies are well known in the art. For example, one method involves recombinant expression of immunoglobulin light chain and modified heavy chain. The heavy chain is truncated generally at any point in the Fc region so as to prevent heavy-chain crosslinking. Alternatively, the relevant cysteine residues are substituted with another amino acid residue or are deleted so as to prevent crosslinking.

In vitro methods are also suitable for preparing monovalent antibodies. Digestion of antibodies to produce fragments thereof, particularly Fab fragments, can be accomplished using routine techniques known in the art.

5.2.5.3. Human and Humanized Antibodies

The anti-PRO antibodies may further comprise humanized antibodies or human antibodies. Humanized forms of non-human (*e.g.*, murine) antibodies are chimeric immunoglobulins, immunoglobulin chains, or fragments thereof (such as Fv, Fab, Fab', F(ab')₂, or other antigen-binding subsequences of antibodies) that contain minimal sequence derived from non-human immunoglobulin. Humanized antibodies include human immunoglobulins (recipient antibody) in which residues from a CDR of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat, or rabbit having the desired specificity, affinity, and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies may also comprise residues that are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin, and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody preferably also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. Jones *et al.*, Nature, 321: 522-525 (1986); Riechmann *et al.*, Nature, 332: 323-329 (1988); Presta, Curr. Op. Struct. Biol., 2:593-596 (1992).

Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source that is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers (Jones *et al.*, Nature, 321: 522-525 (1986); Riechmann *et al.*, Nature, 332: 323-327 (1988); Verhoeyen *et al.*, Science, 239: 1534-

1536 (1988)), by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Patent No. 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

Human antibodies can also be produced using various techniques known in the art, including phage display libraries. Hoogenboom and Winter, *J. Mol. Biol.*, 227: 381 (1991); Marks *et al.*, *J. Mol. Biol.*, 222: 581 (1991). The techniques of Cole *et al.* and Boerner *et al.* are also available for the preparation of human monoclonal antibodies. Cole *et al.*, Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, p. 77 (1985) and Boerner *et al.*, *J. Immunol.*, 147(1): 86-95 (1991). Similarly, human antibodies can be made by introducing human immunoglobulin loci into transgenic animals, *e.g.*, mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed that closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; and 5,661,016, and in the following scientific publications: Marks *et al.*, *Bio/Technology*, 10: 779-783 (1992); Lonberg *et al.*, *Nature*, 368: 856-859 (1994); Morrison, *Nature*, 368: 812-813 (1994); Fishwild *et al.*, *Nature Biotechnology*, 14: 845-851 (1996); Neuberger, *Nature Biotechnology*, 14: 826 (1996); Lonberg and Huszar, *Intern. Rev. Immunol.*, 13: 65-93 (1995).

5.2.5.4. Bispecific Antibodies

Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for the PRO polypeptide, the other one is for any other antigen, and preferably for a cell-surface protein or receptor or receptor subunit.

Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain pairs, where the two heavy chains have different specificities. Milstein and Cuello, *Nature*, 305: 537-539 (1983). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule is usually accomplished by affinity chromatography steps. Similar procedures are disclosed in WO 93/08829, published 13 May 1993, and in Traunecker *et al.*, *EMBO J.*, 10: 3655-3659 (1991).

Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant-domain sequences. The fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least one of the fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. For further

details of generating bispecific antibodies, *see*, for example, Suresh *et al.*, Methods in Enzymology, 121: 210 (1986).

5.2.5.5. Heteroconjugate Antibodies

Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune-system cells to unwanted cells (U.S. Patent No. 4,676,980), and for treatment of HIV infection. WO 91/00360; WO 92/200373; EP 03089. It is contemplated that the antibodies may be prepared *in vitro* using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins may be constructed using a disulfide-exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate and those disclosed, for example, in U.S. Patent No. 4,676,980.

5.2.5.6. Effector Function Engineering

It may be desirable to modify the antibody of the invention with respect to effector function, so as to enhance, *e.g.*, the effectiveness of the antibody in treating cancer. For example, cysteine residue(s) may be introduced into the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated may have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). *See*, Caron *et al.*, J. Exp. Med., 176: 1191-1195 (1992) and Shopes, J. Immunol., 148: 2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity may also be prepared using heterobifunctional cross-linkers as described in Wolff *et al.*, Cancer Research, 53: 2560-2565 (1993). Alternatively, an antibody can be engineered that has dual Fc regions and may thereby have enhanced complement lysis and ADCC capabilities. *See*, Stevenson *et al.*, Anti-Cancer Drug Design, 3: 219-230 (1989).

5.2.5.7. Immunoconjugates

The invention also pertains to immunoconjugates comprising an antibody conjugated to a cytotoxic agent such as a chemotherapeutic agent, toxin (*e.g.*, an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (*i.e.*, a radioconjugate).

Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above. Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolaca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcun, croton, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated antibodies. Examples include ^{212}Bi , ^{131}I , ^{131}In , ^{90}Y , and ^{186}Re .

Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives

of imidoesters (such as dimethyl adipimidate HCl), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta *et al.*, Science, 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See, WO94/11026.

In another embodiment, the antibody may be conjugated to a "receptor" (such as streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (e.g., avidin) that is conjugated to a cytotoxic agent (e.g., a radionucleotide).

5.2.5.8. Immunoliposomes

The antibodies disclosed herein may also be formulated as immunoliposomes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein *et al.*, Proc. Natl. Acad. Sci. USA, 82: 3688 (1985); Hwang *et al.*, Proc. Natl. Acad. Sci. USA, 77: 4030 (1980); and U.S. Pat. Nos. 4,485,045 and 4,544,545. Liposomes with enhanced circulation time are disclosed in U.S. Patent No. 5,013,556.

Particularly useful liposomes can be generated by the reverse-phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol, and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present invention can be conjugated to the liposomes as described in Martin *et al.*, J. Biol. Chem., 257: 286-288 (1982) via a disulfide-interchange reaction. A chemotherapeutic agent (such as Doxorubicin) is optionally contained within the liposome. See, Gabizon *et al.*, J. National Cancer Inst., 81(19): 1484 (1989).

5.2.5.9. Pharmaceutical Compositions of Antibodies

Antibodies specifically binding a PRO polypeptide identified herein, as well as other molecules identified by the screening assays disclosed hereinbefore, can be administered for the treatment of various disorders as noted above and below in the form of pharmaceutical compositions.

If the PRO polypeptide is intracellular and whole antibodies are used as inhibitors, internalizing antibodies are preferred. However, lipofections or liposomes can also be used to deliver the antibody, or an antibody fragment, into cells. Where antibody fragments are used, the smallest inhibitory fragment that specifically binds to the binding domain of the target protein is preferred. For example, based upon the variable-region sequences of an antibody, peptide molecules can be designed that retain the ability to bind the target protein sequence. Such peptides can be synthesized chemically and/or produced by recombinant DNA technology. See, e.g., Marasco *et al.*, Proc. Natl. Acad. Sci. USA, 90: 7889-7893 (1993).

The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. Alternatively, or in addition, the composition may comprise an agent that enhances its function, such as, for example, a cytotoxic agent, cytokine, chemotherapeutic agent, or growth-inhibitory agent. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

The active ingredients may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles, and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences, *supra*.

The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, *e.g.*, films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOTTM (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated antibodies remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37°C, resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S-S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

5.2.5.10. Methods of Treatment using the Antibody

It is contemplated that the antibodies to a PRO polypeptide may be used to treat various cardiovascular, endothelial, and angiogenic conditions as noted above.

The antibodies are administered to a mammal, preferably a human, in accord with known methods, such as intravenous administration as a bolus or by continuous infusion over a period of time, by intramuscular, intraperitoneal, intracerebrospinal, subcutaneous, intra-articular, intrasynovial, intrathecal, oral, topical, or inhalation routes. Intravenous administration of the antibody is preferred.

Other therapeutic regimens may be combined with the administration of the antibodies of the instant invention as noted above. For example, if the antibodies are to treat cancer, the patient to be treated with such

antibodies may also receive radiation therapy. Alternatively, or in addition, a chemotherapeutic agent may be administered to the patient. Preparation and dosing schedules for such chemotherapeutic agents may be used according to manufacturers' instructions or as determined empirically by the skilled practitioner. Preparation and dosing schedules for such chemotherapy are also described in Chemotherapy Service, Ed., M.C. Perry (Williams & Wilkins: Baltimore, MD, 1992). The chemotherapeutic agent may precede, or follow administration of the antibody, or may be given simultaneously therewith. The antibody may be combined with an anti-estrogen compound such as tamoxifen or EVISTA™ or an anti-progesterone such as onapristone (*see*, EP 616812) in dosages known for such molecules.

If the antibodies are used for treating cancer, it may be desirable also to administer antibodies against other tumor-associated antigens, such as antibodies that bind to one or more of the ErbB2, EGFR, ErbB3, ErbB4, or VEGF receptor(s). These also include the agents set forth above. Also, the antibody is suitably administered serially or in combination with radiological treatments, whether involving irradiation or administration of radioactive substances. Alternatively, or in addition, two or more antibodies binding the same or two or more different antigens disclosed herein may be co-administered to the patient. Sometimes, it may be beneficial also to administer one or more cytokines to the patient. In a preferred embodiment, the antibodies herein are co-administered with a growth-inhibitory agent. For example, the growth-inhibitory agent may be administered first, followed by an antibody of the present invention. However, simultaneous administration or administration of the antibody of the present invention first is also contemplated. Suitable dosages for the growth-inhibitory agent are those presently used and may be lowered due to the combined action (synergy) of the growth-inhibitory agent and the antibody herein.

In one embodiment, vascularization of tumors is attacked in combination therapy. The anti-PRO polypeptide antibody and another antibody (*e.g.*, anti-VEGF) are administered to tumor-bearing patients at therapeutically effective doses as determined, for example, by observing necrosis of the tumor or its metastatic foci, if any. This therapy is continued until such time as no further beneficial effect is observed or clinical examination shows no trace of the tumor or any metastatic foci. Then TNF is administered, alone or in combination with an auxiliary agent such as alpha-, beta-, or gamma-interferon, anti-HER2 antibody, heregulin, anti-heregulin antibody, D-factor, interleukin-1 (IL-1), interleukin-2 (IL-2), granulocyte-macrophage colony stimulating factor (GM-CSF), or agents that promote microvascular coagulation in tumors, such as anti-protein C antibody, anti-protein S antibody, or C4b binding protein (*see*, WO 91/01753, published 21 February 1991), or heat or radiation.

Since the auxiliary agents will vary in their effectiveness, it is desirable to compare their impact on the tumor by matrix screening in conventional fashion. The administration of anti-PRO polypeptide antibody and TNF is repeated until the desired clinical effect is achieved. Alternatively, the anti-PRO polypeptide antibody is administered together with TNF and, optionally, auxiliary agent(s). In instances where solid tumors are found in the limbs or in other locations susceptible to isolation from the general circulation, the therapeutic agents described herein are administered to the isolated tumor or organ. In other embodiments, a FGF or PDGF antagonist, such as an anti-FGF or an anti-PDGF neutralizing antibody, is administered to the patient in conjunction with the anti-PRO

polypeptide antibody. Treatment with anti-PRO polypeptide antibodies preferably may be suspended during periods of wound healing or desirable neovascularization.

For the prevention or treatment of cardiovascular, endothelial, and angiogenic disorder, the appropriate dosage of an antibody herein will depend on the type of disorder to be treated, as defined above, the severity and course of the disease, whether the antibody is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the antibody, and the discretion of the attending physician. The antibody is suitably administered to the patient at one time or over a series of treatments.

For example, depending on the type and severity of the disorder, about 1 $\mu\text{g/kg}$ to 50 mg/kg (e.g., 0.1-20 mg/kg) of antibody is an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. A typical daily or weekly dosage might range from about 1 $\mu\text{g/kg}$ to 100 mg/kg or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment is repeated or sustained until a desired suppression of disorder symptoms occurs. However, other dosage regimens may be useful. The progress of this therapy is easily monitored by conventional techniques and assays, including, for example, radiographic tumor imaging.

5.2.5.11. Articles of Manufacture with Antibodies

An article of manufacture containing a container with the antibody and a label is also provided. Such articles are described above, wherein the active agent is an anti-PRO antibody.

5.2.5.12. Diagnosis and Prognosis of Tumors using Antibodies

If the indication for which the antibodies are used is cancer, while cell-surface proteins, such as growth receptors over expressed in certain tumors, are excellent targets for drug candidates or tumor (e.g., cancer) treatment, the same proteins along with PRO polypeptides find additional use in the diagnosis and prognosis of tumors. For example, antibodies directed against the PRO polypeptides may be used as tumor diagnostics or prognostics.

For example, antibodies, including antibody fragments, can be used qualitatively or quantitatively to detect the expression of genes including the gene encoding the PRO polypeptide. The antibody preferably is equipped with a detectable, e.g., fluorescent label, and binding can be monitored by light microscopy, flow cytometry, fluorimetry, or other techniques known in the art. Such binding assays are performed essentially as described above.

In situ detection of antibody binding to the marker gene products can be performed, for example, by immunofluorescence or immunoelectron microscopy. For this purpose, a histological specimen is removed from the patient, and a labeled antibody is applied to it, preferably by overlaying the antibody on a biological sample. This procedure also allows for determining the distribution of the marker gene product in the tissue examined. It will be apparent to those skilled in the art that a wide variety of histological methods are readily available for *in situ* detection.

The following Examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

The disclosures of all patent and literature references cited in the present specification are hereby incorporated by reference in their entirety.

5 6. EXAMPLES

Commercially available reagents referred to in the Examples were used according to manufacturer's instructions unless otherwise indicated. The source of those cells identified in the following Examples, and throughout the specification, by ATCC accession numbers is the American Type Culture Collection, Manassas, VA. Unless otherwise noted, the present invention uses standard procedures of recombinant DNA technology, such as those described hereinabove and in the following textbooks: Sambrook *et al.*, supra; Ausubel *et al.*, Current Protocols in Molecular Biology (Green Publishing Associates and Wiley Interscience, N.Y., 1989); Innis *et al.*, PCR Protocols: A Guide to Methods and Applications (Academic Press, Inc.: N.Y., 1990); Harlow *et al.*, Antibodies: A Laboratory Manual (Cold Spring Harbor Press: Cold Spring Harbor, 1988); Gait, Oligonucleotide Synthesis (IRL Press: Oxford, 1984); Freshney, Animal Cell Culture, 1987; Coligan *et al.*, Current Protocols in Immunology, 1991.

15 6.1. EXAMPLE 1: Extracellular Domain Homology Screening to Identify Novel Polypeptides and cDNA Encoding Therefor

The extracellular domain (ECD) sequences (including the secretion signal sequence, if any) from about 950 known secreted proteins from the Swiss-Prot public database were used to search EST databases. The EST databases included public databases (*e.g.*, Dayhoff, GenBank), and proprietary databases (*e.g.* LIFESEQ®, Incyte Pharmaceuticals, Palo Alto, CA). The search was performed using the computer program BLAST or BLAST-2 (Altschul *et al.*, Methods in Enzymology, 266:460-480 (1996)) as a comparison of the ECD protein sequences to a 6 frame translation of the EST sequences. Those comparisons with a BLAST score of 70 (or in some cases, 90) or greater that did not encode known proteins were clustered and assembled into consensus DNA sequences with the program "phrap" (Phil Green, University of Washington, Seattle, WA).

25 Using this extracellular domain homology screen, consensus DNA sequences were assembled relative to the other identified EST sequences using phrap. In addition, the consensus DNA sequences obtained were often (but not always) extended using repeated cycles of BLAST or BLAST-2 and phrap to extend the consensus sequence as far as possible using the sources of EST sequences discussed above.

30 Based upon the consensus sequences obtained as described above, oligonucleotides were then synthesized and used to identify by PCR a cDNA library that contained the sequence of interest and for use as probes to isolate a clone of the full-length coding sequence for a PRO polypeptide. Forward and reverse PCR primers generally range from 20 to 30 nucleotides and are often designed to give a PCR product of about 100-1000 bp in length. The probe sequences are typically 40-55 bp in length. In some cases, additional oligonucleotides are synthesized when the consensus sequence is greater than about 1-1.5 kbp. In order to screen several libraries for a full-length clone, 35 DNA from the libraries was screened by PCR amplification, as per Ausubel *et al.*, Current Protocols in Molecular

Biology, with the PCR primer pair. A positive library was then used to isolate clones encoding the gene of interest using the probe oligonucleotide and one of the primer pairs.

The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with oligo dT containing a NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; *see*, Holmes *et al.*, Science, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

6.2. EXAMPLE 2: Isolation of cDNA Clones by Amylase Screening

6.2.1. Preparation of oligo dT primed cDNA library

mRNA was isolated from a human tissue of interest using reagents and protocols from Invitrogen, San Diego, CA (Fast Track 2). This RNA was used to generate an oligo dT primed cDNA library in the vector pRK5D using reagents and protocols from Life Technologies, Gaithersburg, MD (Super Script Plasmid System). In this procedure, the double stranded cDNA was sized to greater than 1000 bp and the SalI/NotI linked cDNA was cloned into XhoI/NotI cleaved vector. pRK5D is a cloning vector that has an sp6 transcription initiation site followed by an SfiI restriction enzyme site preceding the XhoI/NotI cDNA cloning sites.

6.2.2. Preparation of random primed cDNA library

A secondary cDNA library was generated in order to preferentially represent the 5' ends of the primary cDNA clones. Sp6 RNA was generated from the primary library (described above), and this RNA was used to generate a random primed cDNA library in the vector pSST-AMY.0 using reagents and protocols from Life Technologies (Super Script Plasmid System, referenced above). In this procedure the double stranded cDNA was sized to 500-1000 bp, linked with blunt to NotI adaptors, cleaved with SfiI, and cloned into SfiI/NotI cleaved vector. pSST-AMY.0 is a cloning vector that has a yeast alcohol dehydrogenase promoter preceding the cDNA cloning sites and the mouse amylase sequence (the mature sequence without the secretion signal) followed by the yeast alcohol dehydrogenase terminator, after the cloning sites. Thus, cDNAs cloned into this vector that are fused in frame with amylase sequence will lead to the secretion of amylase from appropriately transfected yeast colonies.

6.2.3. Transformation and Detection

DNA from the library described in paragraph 2 above was chilled on ice to which was added electrocompetent DH10B bacteria (Life Technologies, 20 ml). The bacteria and vector mixture was then electroporated as recommended by the manufacturer. Subsequently, SOC media (Life Technologies, 1 ml) was added and the mixture was incubated at 37°C for 30 minutes. The transformants were then plated onto 20 standard 150 mm LB plates containing ampicillin and incubated for 16 hours (37°C). Positive colonies were scraped off the plates and the DNA was isolated from the bacterial pellet using standard protocols, *e.g.*, CsCl-gradient. The purified DNA was then carried on to the yeast protocols below.

The yeast methods were divided into three categories: (1) Transformation of yeast with the plasmid/cDNA combined vector; (2) Detection and isolation of yeast clones secreting amylase; and (3) PCR amplification of the insert directly from the yeast colony and purification of the DNA for sequencing and further analysis.

5 The yeast strain used was HD56-5A (ATCC-90785). This strain has the following genotype: MAT alpha, ura3-52, leu2-3, leu2-112, his3-11, his3-15, MAL⁺, SUC⁺, GAL⁺. Preferably, yeast mutants can be employed that have deficient post-translational pathways. Such mutants may have translocation deficient alleles in *sec71*, *sec72*, *sec62*, with truncated *sec71* being most preferred. Alternatively, antagonists (including antisense nucleotides and/or ligands) which interfere with the normal operation of these genes, other proteins implicated in this post translation pathway (e.g., SEC61p, SEC72p, SEC62p, SEC63p, TDJ1p or SSA1p-4p) or the complex formation
10 of these proteins may also be preferably employed in combination with the amylase-expressing yeast.

Transformation was performed based on the protocol outlined by Gietz *et al.*, Nucl. Acid. Res., 20:1425 (1992). Transformed cells were then inoculated from agar into YEPD complex media broth (100 ml) and grown overnight at 30°C. The YEPD broth was prepared as described in Kaiser *et al.*, Methods in Yeast Genetics, Cold Spring Harbor Press, Cold Spring Harbor, NY, p. 207 (1994). The overnight culture was then diluted to about 2
15 x 10⁶ cells/ml (approx. OD₆₀₀=0.1) into fresh YEPD broth (500 ml) and regrown to 1 x 10⁷ cells/ml (approx. OD₆₀₀=0.4-0.5).

The cells were then harvested and prepared for transformation by transfer into GS3 rotor bottles in a Sorval GS3 rotor at 5,000 rpm for 5 minutes, the supernatant discarded, and then resuspended into sterile water, and centrifuged again in 50 ml falcon tubes at 3,500 rpm in a Beckman GS-6KR centrifuge. The supernatant was
20 discarded and the cells were subsequently washed with LiAc/TE (10 ml, 10 mM Tris-HCl, 1 mM EDTA pH 7.5, 100 mM Li₂OOCCH₃), and resuspended into LiAc/TE (2.5 ml).

Transformation took place by mixing the prepared cells (100 µl) with freshly denatured single stranded salmon testes DNA (Lofstrand Labs, Gaithersburg, MD) and transforming DNA (1 µg, vol. < 10 µl) in microfuge tubes. The mixture was mixed briefly by vortexing, then 40% PEG/TE (600 µl, 40% polyethylene glycol-4000,
25 10 mM Tris-HCl, 1 mM EDTA, 100 mM Li₂OOCCH₃, pH 7.5) was added. This mixture was gently mixed and incubated at 30°C while agitating for 30 minutes. The cells were then heat shocked at 42°C for 15 minutes, and the reaction vessel centrifuged in a microfuge at 12,000 rpm for 5-10 seconds, decanted and resuspended into TE (500 µl, 10 mM Tris-HCl, 1 mM EDTA pH 7.5) followed by recentrifugation. The cells were then diluted into TE (1 ml) and aliquots (200 µl) were spread onto the selective media previously prepared in 150 mm growth plates
30 (VWR).

Alternatively, instead of multiple small reactions, the transformation was performed using a single, large scale reaction, wherein reagent amounts were scaled up accordingly.

The selective media used was a synthetic complete dextrose agar lacking uracil (SCD-Ura) prepared as described in Kaiser *et al.*, Methods in Yeast Genetics, Cold Spring Harbor Press, Cold Spring Harbor, NY, p. 208-
35 210 (1994). Transformants were grown at 30°C for 2-3 days.

The detection of colonies secreting amylase was performed by including red starch in the selective growth media. Starch was coupled to the red dye (Reactive Red-120, Sigma) as per the procedure described by Biely *et*

al., Anal. Biochem., 172:176-179 (1988). The coupled starch was incorporated into the SCD-Ura agar plates at a final concentration of 0.15% (w/v), and was buffered with potassium phosphate to a pH of 7.0 (50-100 mM final concentration).

5 The positive colonies were picked and streaked across fresh selective media (onto 150 mm plates) in order to obtain well isolated and identifiable single colonies. Well isolated single colonies positive for amylase secretion were detected by direct incorporation of red starch into buffered SCD-Ura agar. Positive colonies were determined by their ability to break down starch resulting in a clear halo around the positive colony visualized directly.

6.2.4. Isolation of DNA by PCR Amplification

10 When a positive colony was isolated, a portion of it was picked by a toothpick and diluted into sterile water (30 μ l) in a 96 well plate. At this time, the positive colonies were either frozen and stored for subsequent analysis or immediately amplified. An aliquot of cells (5 μ l) was used as a template for the PCR reaction in a 25 μ l volume containing: 0.5 μ l KlenTaq (Clontech, Palo Alto, CA); 4.0 μ l 10 mM dNTP's (Perkin Elmer-Cetus); 2.5 μ l KlenTaq buffer (Clontech); 0.25 μ l forward oligo 1; 0.25 μ l reverse oligo 2; 12.5 μ l distilled water. The sequence of the forward oligonucleotide 1 was:

15 5'-TGTAACGACGGCCAGTTAAATAGACCTGCAATTATTAATCT-3' (SEQ ID NO:382)

The sequence of reverse oligonucleotide 2 was:

5'-CAGGAAACAGCTATGACCACCTGCACACCTGCAAATCCATT-3' (SEQ ID NO:383)

PCR was then performed as follows:

	a.	Denature	92°C, 5 minutes
20	b.	3 cycles of:	Denature 92°C, 30 seconds
			Anneal 59°C, 30 seconds
			Extend 72°C, 60 seconds
	c.	3 cycles of:	Denature 92°C, 30 seconds
25			Anneal 57°C, 30 seconds
			Extend 72°C, 60 seconds
	d.	25 cycles of:	Denature 92°C, 30 seconds
			Anneal 55°C, 30 seconds
			Extend 72°C, 60 seconds
	e.	Hold	4°C

30 The underlined regions of the oligonucleotides annealed to the ADH promoter region and the amylase region, respectively, and amplified a 307 bp region from vector pSST-AMY.0 when no insert was present. Typically, the first 18 nucleotides of the 5' end of these oligonucleotides contained annealing sites for the sequencing primers. Thus, the total product of the PCR reaction from an empty vector was 343 bp. However, signal sequence-fused cDNA resulted in considerably longer nucleotide sequences.

Following the PCR, an aliquot of the reaction (5 µl) was examined by agarose gel electrophoresis in a 1% agarose gel using a Tris-Borate-EDTA (TBE) buffering system as described by Sambrook *et al.*, *supra*. Clones resulting in a single strong PCR product larger than 400 bp were further analyzed by DNA sequencing after purification with a 96 Qiaquick PCR clean-up column (Qiagen Inc., Chatsworth, CA).

5 6.3. EXAMPLE 3: Isolation of cDNA Clones Using Signal Algorithm Analysis

Various polypeptide-encoding nucleic acid sequences were identified by applying a proprietary signal sequence finding algorithm developed by Genentech, Inc., (South San Francisco, CA) upon ESTs as well as clustered and assembled EST fragments from public (e.g., GenBank) and/or private (LIFESEQ®, Incyte Pharmaceuticals, Inc., Palo Alto, CA) databases. The signal sequence algorithm computes a secretion signal score based on the character of the DNA nucleotides surrounding the first and optionally the second methionine codon(s) (ATG) at the 5'-end of the sequence or sequence fragment under consideration. The nucleotides following the first ATG must code for at least 35 unambiguous amino acids without any stop codons. If the first ATG has the required amino acids, the second is not examined. If neither meets the requirement, the candidate sequence is not scored. In order to determine whether the EST sequence contains an authentic signal sequence, the DNA and corresponding amino acid sequences surrounding the ATG codon are scored using a set of seven sensors (evaluation parameters) known to be associated with secretion signals. Use of this algorithm resulted in the identification of numerous polypeptide-encoding nucleic acid sequences.

6.4. EXAMPLE 4: Isolation of cDNA clones Encoding Human PRO Polypeptides

Using the techniques described in Examples 1 to 3 above, numerous full-length cDNA clones were identified as encoding PRO polypeptides as disclosed herein. These cDNAs were then deposited under the terms of the Budapest Treaty with the American Type Culture Collection, 10801 University Blvd., Manassas, VA 20110-2209, USA (ATCC) as shown in Table 7 below.

Table 7

<u>Material</u>	<u>ATCC Dep. No.</u>	<u>Deposit Date</u>
25 23330-1390	209775	4/14/1998
23339-1130	209282	9/18/1997
26846-1397	203406	10/27/1998
26847-1395	209772	4/14/1998
27865-1091	209296	9/23/1997
30 30868-1156	1437-PTA	3/2/2000
30871-1157	209380	10/16/1997
32286-1191	209385	10/16/1997
33089-1132	209262	9/16/1997
33092-1202	209420	10/28/1997

	33100-1159	209377	10/16/1997
	33223-1136	209264	9/16/1997
	34392-1170	209526	12/10/1997
	34431-1177	209399	10/17/1997
5	34433-1308	209719	3/31/1998
	34434-1139	209252	9/16/1997
	35600-1162	209370	10/16/1997
	35673-1201	209418	10/28/1997
	35880-1160	209379	10/16/1997
10	35918-1174	209402	10/17/1997
	36350-1158	209378	10/16/1997
	36638-1056	209456	11/12/1997
	38268-1188	209421	10/28/1997
	40370-1217	209485	11/21/1997
15	40628-1216	209432	11/7/1997
	43316-1237	209487	11/21/1997
	44196-1353	209847	5/6/1998
	45409-2511	203579	1/12/1999
	45419-1252	209616	2/5/1998
20	46777-1253	209619	2/5/1998
	48336-1309	209669	3/11/1998
	48606-1479	203040	7/1/1998
	49435-1219	209480	11/21/1997
	49631-1328	209806	4/28/1998
25	50919-1361	209848	5/6/1998
	50920-1325	209700	3/26/1998
	50921-1458	209859	5/12/1998
	52758-1399	209773	4/14/1998
	53517-1366-1	209802	4/23/1998
30	53915-1258	209593	1/21/1998
	53974-1401	209774	4/14/1998
	53987-1438	209858	5/12/1998
	56047-1456	209948	6/9/1998
	56050-1455	203011	6/23/1998
35	56110-1437	203113	8/11/1998
	56405-1357	209849	5/6/1998
	56433-1406	209857	5/12/1998

	56439-1376	209864	5/14/1998
	56529-1647	203293	9/29/1998
	56865-1491	203022	6/23/1998
	56965-1356	209842	5/6/1998
5	57033-1403-1	209905	5/27/1998
	57037-1444	209903	5/27/1998
	57039-1402	209777	4/14/1998
	57689-1385	209869	5/14/1998
	57690-1374	209950	6/9/1998
10	57694-1341	203017	6/23/1998
	57695-1340	203006	6/23/1998
	57699-1412	203020	6/23/1998
	57700-1408	203583	1/12/1999
	57708-1411	203021	6/23/1998
15	57838-1337	203014	6/23/1998
	58847-1383	209879	5/20/1998
	58852-1637	203271	9/22/1998
	58853-1423	203016	6/23/1998
	59212-1627	203245	9/9/1998
20	59220-1514	209962	6/9/1998
	59493-1420	203050	7/1/1998
	59497-1496	209941	6/4/1998
	59586-1520	203288	9/29/1998
	59588-1571	203106	8/11/1998
25	59620-1463	209989	6/16/1998
	59622-1334	209984	6/16/1998
	59777-1480	203111	8/11/1998
	59848-1512	203088	8/4/1998
	59849-1504	209986	6/16/1998
30	60621-1516	203091	8/4/1998
	60622-1525	203090	8/4/1998
	60764-1533	203452	11/10/1998
	60783-1611	203130	8/18/1998
	61755-1554	203112	8/11/1998
35	62306-1570	203254	9/9/1998
	62312-2558	203836	3/9/1999
	62814-1521	203093	8/4/1998

	62872-1509	203100	8/4/1998
	64883-1526	203253	9/9/1998
	64886-1601	203241	9/9/1998
	64889-1541	203250	9/9/1998
5	64896-1539	203238	9/9/1998
	64897-1628	203216	9/15/1998
	64903-1553	203223	9/15/1998
	64908-1163-1	203243	9/9/1998
	64950-1590	203224	9/15/1998
10	65402-1540	203252	9/9/1998
	65404-1551	203244	9/9/1998
	65405-1547	203476	11/17/1998
	65410-1569	203231	9/15/1998
	65412-1523	203094	8/4/1998
15	66307-2661	431-PTA	7/27/1999
	66526-1616	203246	9/9/1998
	66659-1593	203269	9/22/1998
	66660-1585	203279	9/22/1998
	66667-1596	203267	9/22/1998
20	66672-1586	203265	9/22/1998
	66675-1587	203282	9/22/1998
	67300-1605	203163	8/25/1998
	68818-2536	203657	2/9/1999
	68862-2546	203652	2/9/1999
25	68872-1620	203160	8/25/1998
	71290-1630	203275	9/22/1998
	73736-1657	203466	11/17/1998
	73739-1645	203270	9/22/1998
	73742-1662	203316	10/6/1998
30	76385-1692	203664	2/9/1999
	76393-1664	203323	10/6/1998
	76399-1700	203472	11/17/1998
	76400-2528	203573	1/12/1999
	76510-2504	203477	11/17/1998
35	76529-1666	203315	10/6/1998
	76532-1702	203473	11/17/1998
	76541-1675	203409	10/27/1998

	77503-1686	203362	10/20/1998
	77624-2515	203553	12/22/1998
	79230-2525	203549	12/22/1998
	79862-2522	203550	12/22/1998
5	80145-2594	204-PTA	6/8/1999
	80899-2501	203539	12/15/1998
	81754-2532	203542	12/15/1998
	81757-2512	203543	12/15/1998
	81761-2583	203862	3/23/1999
10	82358-2738	510-PTA	8/10/1999
	82364-2538	203603	1/20/1999
	82403-2959	2317-PTA	8/1/2000
	83500-2506	203391	10/29/1998
	83560-2569	203816	3/2/1999
15	84210-2576	203818	3/2/1999
	84920-2614	203966	4/27/1999
	86576-2595	203868	3/23/1999
	92218-2554	203834	3/9/1999
	92233-2599	134-PTA	5/25/1999
20	92256-2596	203891	3/30/1999
	92265-2669	256-PTA	6/22/1999
	92274-2617	203971	4/27/1999
	92929-2534-1	203586	1/12/1999
	93011-2637	20-PTA	5/4/1999
25	94854-2586	203864	3/23/1999
	96787-2534-1	203589	1/12/1999
	96867-2620	203972	4/27/1999
	96872-2674	550-PTA	8/17/1999
	96878-2626	23-PTA	5/4/1999
30	96889-2641	119-PTA	5/25/1999
	100312-2645	44-PTA	5/11/1999
	105782-2693	387-PTA	7/20/1999
	105849-2704	473-PTA	8/3/1999
	108725-2766	863-PTA	10/19/1999
35	108769-2765	861-PTA	10/19/1999
	119498-2965	2298-PTA	7/25/2000
	119535-2756	613-PTA	8/31/1999

	125185-2806	1031-PTA	12/7/1999
	131639-2874	1784-PTA	4/25/2000
	139623-2893	1670-PTA	4/11/2000
	143076-2787	1028-PTA	12/7/1999
5	143276-2975	2387-PTA	8/8/2000
	164625-2890	1535-PTA	3/21/2000
	167678-2963	2302-PTA	7/25/2000
	170021-2923	1906-PTA	5/23/2000
	170212-3000	2583-PTA	10/10/2000
10	177313-2982	2251-PTA	7/19/2000

These deposits were made under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure and the Regulations thereunder (Budapest Treaty). This assures maintenance of a viable culture of the deposit for 30 years from the date of deposit. The deposits will be made available by ATCC under the terms of the Budapest Treaty, and subject to an agreement between Genentech, Inc. and ATCC, which assures permanent and unrestricted availability of the progeny of the culture of the deposit to the public upon issuance of the pertinent U.S. patent or upon laying open to the public of any U.S. or foreign patent application, whichever comes first, and assures availability of the progeny to one determined by the U.S. Commissioner of Patents and Trademarks to be entitled thereto according to 35 USC § 122 and the Commissioner's rules pursuant thereto (including 37 CFR § 1.14 with particular reference to 886 OG 638).

The assignee of the present application has agreed that if a culture of the materials on deposit should die or be lost or destroyed when cultivated under suitable conditions, the materials will be promptly replaced on notification with another of the same. Availability of the deposited material is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any government in accordance with its patent laws.

6.5 EXAMPLE 5: Isolation of cDNA clones Encoding Human PRO1873, PRO7223, PRO7248, PRO730, PRO532, PRO7261, PRO734, PRO771, PRO2010, PRO5723, PRO3444, PRO9940, PRO3562, PRO10008, PRO5730, PRO6008, PRO4527, PRO4538 and PRO4553

DNA molecules encoding the PRO1873, PRO7223, PRO7248, PRO730, PRO532, PRO7261, PRO734, PRO771, PRO2010, PRO5723, PRO3444, PRO9940, PRO3562, PRO10008, PRO5730, PRO6008, PRO4527, PRO4538 and PRO4553 polypeptides shown in the accompanying figures were obtained through GenBank.

6.6. EXAMPLE 6: Use of PRO as a Hybridization Probe

The following method describes use of a nucleotide sequence encoding PRO as a hybridization probe.

DNA comprising the coding sequence of full-length or mature PRO (as shown in accompanying figures) or a fragment thereof is employed as a probe to screen for homologous DNAs (such as those encoding naturally-occurring variants of PRO) in human tissue cDNA libraries or human tissue genomic libraries.

5 Hybridization and washing of filters containing either library DNAs is performed under the following high-stringency conditions. Hybridization of radiolabeled probe derived from the gene encoding PRO polypeptide to the filters is performed in a solution of 50% formamide, 5x SSC, 0.1% SDS, 0.1% sodium pyrophosphate, 50 mM sodium phosphate, pH 6.8, 2x Denhardt's solution, and 10% dextran sulfate at 42°C for 20 hours. Washing of the filters is performed in an aqueous solution of 0.1x SSC and 0.1% SDS at 42°C.

10 DNAs having a desired sequence identity with the DNA encoding full-length native sequence can then be identified using standard techniques known in the art.

6.7. EXAMPLE 7: Expression of PRO in *E. coli*

This example illustrates preparation of an unglycosylated form of PRO by recombinant expression in *E. coli*.

15 The DNA sequence encoding PRO is initially amplified using selected PCR primers. The primers should contain restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector. A variety of expression vectors may be employed. An example of a suitable vector is pBR322 (derived from *E. coli*; see, Bolivar *et al.*, Gene, 2:95 (1977)) which contains genes for ampicillin and tetracycline resistance. The vector is digested with restriction enzyme and dephosphorylated. The PCR amplified sequences are then ligated into the vector. The vector will preferably include sequences which encode for an antibiotic resistance gene, a trp promoter, a poly-His leader (including the first six STII codons, poly-His sequence, and enterokinase cleavage site),
20 the PRO coding region, lambda transcriptional terminator, and an argU gene.

The ligation mixture is then used to transform a selected *E. coli* strain using the methods described in Sambrook *et al.*, *supra*. Transformants are identified by their ability to grow on LB plates and antibiotic resistant colonies are then selected. Plasmid DNA can be isolated and confirmed by restriction analysis and DNA
25 sequencing.

Selected clones can be grown overnight in liquid culture medium such as LB broth supplemented with antibiotics. The overnight culture may subsequently be used to inoculate a larger scale culture. The cells are then grown to a desired optical density, during which the expression promoter is turned on.

30 After culturing the cells for several more hours, the cells can be harvested by centrifugation. The cell pellet obtained by the centrifugation can be solubilized using various agents known in the art, and the solubilized PRO protein can then be purified using a metal chelating column under conditions that allow tight binding of the protein.

PRO may be expressed in *E. coli* in a poly-His tagged form, using the following procedure. The DNA encoding PRO is initially amplified using selected PCR primers. The primers will contain restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector, and other useful sequences
35 providing for efficient and reliable translation initiation, rapid purification on a metal chelation column, and proteolytic removal with enterokinase. The PCR-amplified, poly-His tagged sequences are then ligated into an

expression vector, which is used to transform an *E. coli* host based on strain 52 (W3110 fuhA(tonA) lon galE rpoHts(htpRts) clpP(lacIq). Transformants are first grown in LB containing 50 mg/ml carbenicillin at 30°C with shaking until an OD₆₀₀ of 3-5 is reached. Cultures are then diluted 50-100 fold into CRAP media (prepared by mixing 3.57 g (NH₄)₂SO₄, 0.71 g sodium citrate·2H₂O, 1.07 g KCl, 5.36 g Difco yeast extract, 5.36 g Sheffield hycase SF in 500 ml water, as well as 110 mM MPOS, pH 7.3, 0.55% (w/v) glucose and 7 mM MgSO₄) and grown for approximately 20-30 hours at 30°C with shaking. Samples are removed to verify expression by SDS-PAGE analysis, and the bulk culture is centrifuged to pellet the cells. Cell pellets are frozen until purification and refolding.

E. coli paste from 0.5 to 1 L fermentations (6-10 g pellets) is resuspended in 10 volumes (w/v) in 7 M guanidine, 20 mM Tris, pH 8 buffer. Solid sodium sulfite and sodium tetrathionate is added to make final concentrations of 0.1M and 0.02 M, respectively, and the solution is stirred overnight at 4°C. This step results in a denatured protein with all cysteine residues blocked by sulfitolization. The solution is centrifuged at 40,000 rpm in a Beckman Ultracentrifuge for 30 min. The supernatant is diluted with 3-5 volumes of metal chelate column buffer (6 M guanidine, 20 mM Tris, pH 7.4) and filtered through 0.22 micron filters to clarify. The clarified extract is loaded onto a 5 ml Qiagen Ni²⁺-NTA metal chelate column equilibrated in the metal chelate column buffer. The column is washed with additional buffer containing 50 mM imidazole (Calbiochem, Utrol grade), pH 7.4. The protein is eluted with buffer containing 250 mM imidazole. Fractions containing the desired protein are pooled and stored at 4°C. Protein concentration is estimated by its absorbance at 280 nm using the calculated extinction coefficient based on its amino acid sequence.

The proteins are refolded by diluting the sample slowly into freshly prepared refolding buffer consisting of: 20 mM Tris, pH 8.6, 0.3 M NaCl, 2.5 M urea, 5 mM cysteine, 20 mM glycine and 1 mM EDTA. Refolding volumes are chosen so that the final protein concentration is between 50 to 100 micrograms/ml. The refolding solution is stirred gently at 4°C for 12-36 hours. The refolding reaction is quenched by the addition of TFA to a final concentration of 0.4% (pH of approximately 3). Before further purification of the protein, the solution is filtered through a 0.22 micron filter and acetonitrile is added to 2-10% final concentration. The refolded protein is chromatographed on a Poros R1/H reversed phase column using a mobile buffer of 0.1% TFA with elution with a gradient of acetonitrile from 10 to 80%. Aliquots of fractions with A₂₈₀ absorbance are analyzed on SDS polyacrylamide gels and fractions containing homogeneous refolded protein are pooled. Generally, the properly refolded species of most proteins are eluted at the lowest concentrations of acetonitrile since those species are the most compact with their hydrophobic interiors shielded from interaction with the reversed phase resin. Aggregated species are usually eluted at higher acetonitrile concentrations. In addition to resolving misfolded forms of proteins from the desired form, the reversed phase step also removes endotoxin from the samples.

Fractions containing the desired folded PRO polypeptide are pooled and the acetonitrile removed using a gentle stream of nitrogen directed at the solution. Proteins are formulated into 20 mM Hepes, pH 6.8 with 0.14 M sodium chloride and 4% mannitol by dialysis or by gel filtration using G25 Superfine (Pharmacia) resins equilibrated in the formulation buffer and sterile filtered.

Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

6.8. EXAMPLE 8: Expression of PRO in mammalian cells

This example illustrates preparation of a potentially glycosylated form of PRO by recombinant expression in mammalian cells.

5 The vector, pRK5 (see EP 307,247, published March 15, 1989), is employed as the expression vector. Optionally, the PRO DNA is ligated into pRK5 with selected restriction enzymes to allow insertion of the PRO DNA using ligation methods such as described in Sambrook *et al.*, *supra*. The resulting vector is called pRK5-PRO.

10 In one embodiment, the selected host cells may be 293 cells. Human 293 cells (ATCC CCL 1573) are grown to confluence in tissue culture plates in medium such as DMEM supplemented with fetal calf serum and optionally, nutrient components and/or antibiotics. About 10 μ g pRK5-PRO DNA is mixed with about 1 μ g DNA encoding the VA RNA gene [Thimmappaya *et al.*, *Cell*, 31:543 (1982)] and dissolved in 500 μ l of 1 mM Tris-HCl, 0.1 mM EDTA, 0.227 M CaCl_2 . To this mixture is added, dropwise, 500 μ l of 50 mM HEPES (pH 7.35), 280 mM NaCl, 1.5 mM NaPO_4 , and a precipitate is allowed to form for 10 minutes at 25°C. The precipitate is suspended and added to the 293 cells and allowed to settle for about four hours at 37°C. The culture medium is aspirated off and 2 ml of 20% glycerol in PBS is added for 30 seconds. The 293 cells are then washed with serum free medium, 15 fresh medium is added and the cells are incubated for about 5 days.

Approximately 24 hours after the transfections, the culture medium is removed and replaced with culture medium (alone) or culture medium containing 200 μ Ci/ml ^{35}S -cysteine and 200 μ Ci/ml ^{35}S -methionine. After a 12 hour incubation, the conditioned medium is collected, concentrated on a spin filter, and loaded onto a 15% SDS gel. The processed gel may be dried and exposed to film for a selected period of time to reveal the presence of the PRO polypeptide. The cultures containing transfected cells may undergo further incubation (in serum free medium) and the medium is tested in selected bioassays. 20

In an alternative technique, PRO may be introduced into 293 cells transiently using the dextran sulfate method described by Sompanyrac *et al.*, *Proc. Natl. Acad. Sci.*, 12:7575 (1981). 293 cells are grown to maximal density in a spinner flask and 700 μ g pRK5-PRO DNA is added. The cells are first concentrated from the spinner flask by centrifugation and washed with PBS. The DNA-dextran precipitate is incubated on the cell pellet for four hours. The cells are treated with 20% glycerol for 90 seconds, washed with tissue culture medium, and re-introduced into the spinner flask containing tissue culture medium, 5 μ g/ml bovine insulin and 0.1 μ g/ml bovine transferrin. After about four days, the conditioned media is centrifuged and filtered to remove cells and debris. The sample containing expressed PRO can then be concentrated and purified by any selected method, such as dialysis and/or column chromatography. 25 30

In another embodiment, PRO can be expressed in CHO cells. The pRK5-PRO can be transfected into CHO cells using known reagents such as CaPO_4 or DEAE-dextran. As described above, the cell cultures can be incubated, and the medium replaced with culture medium (alone) or medium containing a radiolabel such as ^{35}S -methionine. After determining the presence of a PRO polypeptide, the culture medium may be replaced with serum free medium. Preferably, the cultures are incubated for about 6 days, and then the conditioned medium is harvested. 35

The medium containing the expressed PRO polypeptide can then be concentrated and purified by any selected method.

Epitope-tagged PRO may also be expressed in host CHO cells. The PRO may be subcloned out of the pRK5 vector. The subclone insert can undergo PCR to fuse in frame with a selected epitope tag such as a poly-His tag into a Baculovirus expression vector. The poly-His tagged PRO insert can then be subcloned into a SV40 driven vector containing a selection marker such as DHFR for selection of stable clones. Finally, the CHO cells can be transfected (as described above) with the SV40 driven vector. Labeling may be performed, as described above, to verify expression. The culture medium containing the expressed poly-His tagged PRO can then be concentrated and purified by any selected method, such as by Ni²⁺-chelate affinity chromatography.

PRO may also be expressed in CHO and/or COS cells by a transient expression procedure or in CHO cells by another stable expression procedure.

Stable expression in CHO cells is performed using the following procedure. The proteins are expressed as an IgG construct (immunoadhesin), in which the coding sequences for the soluble forms (*e.g.*, extracellular domains) of the respective proteins are fused to an IgG1 constant region sequence containing the hinge, CH2 and CH2 domains and/or as a poly-His tagged form.

Following PCR amplification, the respective DNAs are subcloned in a CHO expression vector using standard techniques as described in Ausubel *et al.*, Current Protocols of Molecular Biology, Unit 3.16, John Wiley and Sons (1997). CHO expression vectors are constructed to have compatible restriction sites 5' and 3' of the DNA of interest to allow the convenient shuttling of cDNA's. The vector used in expression in CHO cells is as described in Lucas *et al.*, Nucl. Acids Res., 24:9 (1774-1779 (1996), and uses the SV40 early promoter/enhancer to drive expression of the cDNA of interest and dihydrofolate reductase (DHFR). DHFR expression permits selection for stable maintenance of the plasmid following transfection.

Twelve micrograms of the desired plasmid DNA is introduced into approximately 10 million CHO cells using commercially available transfection reagents Superfect® (Qiagen), Dosper® or Eugene® (Boehringer Mannheim). The cells are grown as described in Lucas *et al.*, *supra*. Approximately 3×10^7 cells are frozen in an ampule for further growth and production as described below.

The ampules containing the plasmid DNA are thawed by placement into a water bath and mixed by vortexing. The contents are pipetted into a centrifuge tube containing 10 ml of media and centrifuged at 1000 rpm for 5 minutes. The supernatant is aspirated and the cells are resuspended in 10 ml of selective media (0.2 µm filtered PS20 with 5% 0.2 µm diafiltered fetal bovine serum). The cells are then aliquoted into a 100 ml spinner containing 90 ml of selective media. After 1-2 days, the cells are transferred into a 250 ml spinner filled with 150 ml selective growth medium and incubated at 37°C. After another 2-3 days, 250 ml, 500 ml and 2000 ml spinners are seeded with 3×10^5 cells/ml. The cell media is exchanged with fresh media by centrifugation and resuspension in production medium. Although any suitable CHO media may be employed, a production medium described in U.S. Patent No. 5,122,469, issued June 16, 1992 may actually be used. A 3L production spinner is seeded at 1.2×10^6 cells/ml. On day 0, the cell number and pH is determined. On day 1, the spinner is sampled and sparging with filtered air is commenced. On day 2, the spinner is sampled, the temperature shifted to 33°C, and 30 ml of 500 g/L

glucose and 0.6 ml of 10% antifoam (e.g., 35% polydimethylsiloxane emulsion, Dow Corning 365 Medical Grade Emulsion) taken. Throughout the production, the pH is adjusted as necessary to keep it at around 7.2. After 10 days, or until the viability drops below 70%, the cell culture is harvested by centrifugation and filtering through a 0.22 μ m filter. The filtrate is either stored at 4°C or immediately loaded onto columns for purification.

5 For the poly-His tagged constructs, the proteins are purified using a Ni²⁺-NTA column (Qiagen). Before purification, imidazole is added to the conditioned media to a concentration of 5 mM. The conditioned media is pumped onto a 6 ml Ni²⁺-NTA column equilibrated in 20 mM Hepes, pH 7.4, buffer containing 0.3 M NaCl and 5 mM imidazole at a flow rate of 4-5 ml/min. at 4°C. After loading, the column is washed with additional equilibration buffer and the protein eluted with equilibration buffer containing 0.25 M imidazole. The highly
10 purified protein is subsequently desalted into a storage buffer containing 10 mM Hepes, 0.14 M NaCl and 4% mannitol, pH 6.8, with a 25 ml G25 Superfine (Pharmacia) column and stored at -80°C.

Immunoadhesin (Fc-containing) constructs are purified from the conditioned media as follows. The conditioned medium is pumped onto a 5 ml Protein A column (Pharmacia) which has been equilibrated in 20 mM Na phosphate buffer, pH 6.8. After loading, the column is washed extensively with equilibration buffer before
15 elution with 100 mM citric acid, pH 3.5. The eluted protein is immediately neutralized by collecting 1 ml fractions into tubes containing 275 μ l of 1 M Tris buffer, pH 9. The highly purified protein is subsequently desalted into storage buffer as described above for the poly-His tagged proteins. The homogeneity is assessed by SDS polyacrylamide gels and by N-terminal amino acid sequencing by Edman degradation.

Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

20 6.9. EXAMPLE 9: Expression of PRO in Yeast

The following method describes recombinant expression of PRO in yeast.

First, yeast expression vectors are constructed for intracellular production or secretion of PRO from the ADH2/GAPDH promoter. DNA encoding PRO and the promoter is inserted into suitable restriction enzyme sites in the selected plasmid to direct intracellular expression of PRO. For secretion, DNA encoding PRO can be cloned
25 into the selected plasmid, together with DNA encoding the ADH2/GAPDH promoter, a native PRO signal peptide or other mammalian signal peptide, or, for example, a yeast alpha-factor or invertase secretory signal/leader sequence, and linker sequences (if needed) for expression of PRO.

Yeast cells, such as yeast strain AB110, can then be transformed with the expression plasmids described above and cultured in selected fermentation media. The transformed yeast supernatants can be analyzed by
30 precipitation with 10% trichloroacetic acid and separation by SDS-PAGE, followed by staining of the gels with Coomassie Blue stain.

Recombinant PRO can subsequently be isolated and purified by removing the yeast cells from the fermentation medium by centrifugation and then concentrating the medium using selected cartridge filters. The concentrate containing PRO may further be purified using selected column chromatography resins.

35 Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

6.10. EXAMPLE 10: Expression of PRO in Baculovirus-Infected Insect Cells

The following method describes recombinant expression in Baculovirus-infected insect cells.

The sequence coding for PRO is fused upstream of an epitope tag contained within a baculovirus expression vector. Such epitope tags include poly-His tags and immunoglobulin tags (like Fc regions of IgG). A variety of plasmids may be employed, including plasmids derived from commercially available plasmids such as pVL1393 (Novagen). Briefly, the sequence encoding PRO or the desired portion of the coding sequence of PRO (such as the sequence encoding the extracellular domain of a transmembrane protein or the sequence encoding the mature protein if the protein is extracellular) is amplified by PCR with primers complementary to the 5' and 3' regions. The 5' primer may incorporate flanking (selected) restriction enzyme sites. The product is then digested with those selected restriction enzymes and subcloned into the expression vector.

Recombinant baculovirus is generated by co-transfecting the above plasmid and BaculoGold™ virus DNA (Pharmingen) into *Spodoptera frugiperda* ("Sf9") cells (ATCC CRL 1711) using lipofectin (commercially available from GIBCO-BRL). After 4 - 5 days of incubation at 28°C, the released viruses are harvested and used for further amplifications. Viral infection and protein expression are performed as described by O'Reilley *et al.*, Baculovirus expression vectors: A Laboratory Manual, Oxford: Oxford University Press (1994).

Expressed poly-His tagged PRO can then be purified, for example, by Ni²⁺-chelate affinity chromatography as follows. Extracts are prepared from recombinant virus-infected Sf9 cells as described by Rupert *et al.*, Nature, **362**:175-179 (1993). Briefly, Sf9 cells are washed, resuspended in sonication buffer (25 ml Hepes, pH 7.9; 12.5 mM MgCl₂; 0.1 mM EDTA; 10% glycerol; 0.1% NP-40; 0.4 M KCl), and sonicated twice for 20 seconds on ice. The sonicates are cleared by centrifugation, and the supernatant is diluted 50-fold in loading buffer (50 mM phosphate, 300 mM NaCl, 10% glycerol, pH 7.8) and filtered through a 0.45 µm filter. A Ni²⁺-NTA agarose column (commercially available from Qiagen) is prepared with a bed volume of 5 ml, washed with 25 ml of water and equilibrated with 25 ml of loading buffer. The filtered cell extract is loaded onto the column at 0.5 ml per minute. The column is washed to baseline A₂₈₀ with loading buffer, at which point fraction collection is started. Next, the column is washed with a secondary wash buffer (50 mM phosphate; 300 mM NaCl, 10% glycerol, pH 6.0), which elutes nonspecifically bound protein. After reaching A₂₈₀ baseline again, the column is developed with a 0 to 500 mM imidazole gradient in the secondary wash buffer. One ml fractions are collected and analyzed by SDS-PAGE and silver staining or Western blot with Ni²⁺-NTA-conjugated to alkaline phosphatase (Qiagen). Fractions containing the eluted His₁₀-tagged PRO are pooled and dialyzed against loading buffer.

Alternatively, purification of the IgG tagged (or Fc tagged) PRO can be performed using known chromatography techniques, including for instance, Protein A or protein G column chromatography.

Following PCR amplification, the respective coding sequences are subcloned into a baculovirus expression vector (pb.PH.IgG for IgG fusions and pb.PH.His.c for poly-His tagged proteins), and the vector and Baculogold® baculovirus DNA (Pharmingen) are co-transfected into 105 *Spodoptera frugiperda* ("Sf9") cells (ATCC CRL 1711), using Lipofectin (Gibco BRL). pb.PH.IgG and pb.PH.His are modifications of the commercially available baculovirus expression vector pVL1393 (Pharmingen), with modified polylinker regions to include the His or Fc tag sequences. The cells are grown in Hink's TNM-FH medium supplemented with 10% FBS (Hyclone). Cells are

incubated for 5 days at 28°C. The supernatant is harvested and subsequently used for the first viral amplification by infecting Sf9 cells in Hink's TNM-FH medium supplemented with 10% FBS at an approximate multiplicity of infection (MOI) of 10. Cells are incubated for 3 days at 28°C. The supernatant is harvested and the expression of the constructs in the baculovirus expression vector is determined by batch binding of 1 ml of supernatant to 25 ml of Ni²⁺-NTA beads (QIAGEN) for histidine tagged proteins or Protein-A Sepharose CL-4B beads (Pharmacia) for IgG tagged proteins followed by SDS-PAGE analysis comparing to a known concentration of protein standard by Coomassie blue staining.

The first viral amplification supernatant is used to infect a spinner culture (500 ml) of Sf9 cells grown in ESF-921 medium (Expression Systems LLC) at an approximate MOI of 0.1. Cells are incubated for 3 days at 28°C. The supernatant is harvested and filtered. Batch binding and SDS-PAGE analysis is repeated, as necessary, until expression of the spinner culture is confirmed.

The conditioned medium from the transfected cells (0.5 to 3 L) is harvested by centrifugation to remove the cells and filtered through 0.22 micron filters. For the poly-His tagged constructs, the protein construct is purified using a Ni²⁺-NTA column (Qiagen). Before purification, imidazole is added to the conditioned media to a concentration of 5 mM. The conditioned media is pumped onto a 6 ml Ni²⁺-NTA column equilibrated in 20 mM Hepes, pH 7.4, buffer containing 0.3 M NaCl and 5 mM imidazole at a flow rate of 4-5 ml/min. at 4°C. After loading, the column is washed with additional equilibration buffer and the protein eluted with equilibration buffer containing 0.25 M imidazole. The highly purified protein is subsequently desalted into a storage buffer containing 10 mM Hepes, 0.14 M NaCl and 4% mannitol, pH 6.8, with a 25 ml G25 Superfine (Pharmacia) column and stored at -80°C.

Immunoadhesin (Fc containing) constructs of proteins are purified from the conditioned media as follows. The conditioned media is pumped onto a 5 ml Protein A column (Pharmacia) which has been equilibrated in 20 mM Na phosphate buffer, pH 6.8. After loading, the column is washed extensively with equilibration buffer before elution with 100 mM citric acid, pH 3.5. The eluted protein is immediately neutralized by collecting 1 ml fractions into tubes containing 275 ml of 1 M Tris buffer, pH 9. The highly purified protein is subsequently desalted into storage buffer as described above for the poly-His tagged proteins. The homogeneity of the proteins is verified by SDS polyacrylamide gel (PAGE) electrophoresis and N-terminal amino acid sequencing by Edman degradation.

Alternatively, a modified baculovirus procedure may be used incorporating high-5 cells. In this procedure, the DNA encoding the desired sequence is amplified with suitable systems, such as Pfu (Stratagene), or fused upstream (5'-of) of an epitope tag contained with a baculovirus expression vector. Such epitope tags include poly-His tags and immunoglobulin tags (like Fc regions of IgG). A variety of plasmids may be employed, including plasmids derived from commercially available plasmids such as pIE1-1 (Novagen). The pIE1-1 and pIE1-2 vectors are designed for constitutive expression of recombinant proteins from the baculovirus ie1 promoter in stably-transformed insect cells (1). The plasmids differ only in the orientation of the multiple cloning sites and contain all promoter sequences known to be important for ie1-mediated gene expression in uninfected insect cells as well as the hr5 enhancer element. pIE1-1 and pIE1-2 include the translation initiation site and can be used to produce fusion proteins. Briefly, the desired sequence or the desired portion of the sequence (such as the sequence encoding the

extracellular domain of a transmembrane protein) is amplified by PCR with primers complementary to the 5' and 3' regions. The 5' primer may incorporate flanking (selected) restriction enzyme sites. The product is then digested with those selected restriction enzymes and subcloned into the expression vector. For example, derivatives of pIE1-1 can include the Fc region of human IgG (pb.PH.IgG) or an 8 histidine (pb.PH.His) tag downstream (3'-of) the desired sequence. Preferably, the vector construct is sequenced for confirmation.

High-5 cells are grown to a confluency of 50% under the conditions of, 27°C, no CO₂, NO pen/strep. For each 150 mm plate, 30 µg of pIE based vector containing the sequence is mixed with 1 ml Ex-Cell medium (Media: Ex-Cell 401 + 1/100 L-Glu JRH Biosciences #14401-78P (note: this media is light sensitive)), and in a separate tube, 100 µl of CellFectin (CellFECTIN (GibcoBRL #10362-010) (vortexed to mix)) is mixed with 1 ml of Ex-Cell medium. The two solutions are combined and allowed to incubate at room temperature for 15 minutes. 8 ml of Ex-Cell media is added to the 2 ml of DNA/CellFECTIN mix and this is layered on high-5 cells that have been washed once with Ex-Cell media. The plate is then incubated in darkness for 1 hour at room temperature. The DNA/CellFECTIN mix is then aspirated, and the cells are washed once with Ex-Cell to remove excess CellFECTIN, 30 ml of fresh Ex-Cell media is added and the cells are incubated for 3 days at 28°C. The supernatant is harvested and the expression of the sequence in the baculovirus expression vector is determined by batch binding of 1 ml of supernatant to 25 ml of Ni²⁺-NTA beads (QIAGEN) for histidine tagged proteins or Protein-A Sepharose CL-4B beads (Pharmacia) for IgG tagged proteins followed by SDS-PAGE analysis comparing to a known concentration of protein standard by Coomassie blue staining.

The conditioned media from the transfected cells (0.5 to 3 L) is harvested by centrifugation to remove the cells and filtered through 0.22 micron filters. For the poly-His tagged constructs, the protein comprising the sequence is purified using a Ni²⁺-NTA column (Qiagen). Before purification, imidazole is added to the conditioned media to a concentration of 5 mM. The conditioned media is pumped onto a 6 ml Ni²⁺-NTA column equilibrated in 20 mM Hepes, pH 7.4, buffer containing 0.3 M NaCl and 5 mM imidazole at a flow rate of 4-5 ml/min. at 48°C. After loading, the column is washed with additional equilibration buffer and the protein eluted with equilibration buffer containing 0.25 M imidazole. The highly purified protein is then subsequently desalted into a storage buffer containing 10 mM Hepes, 0.14 M NaCl and 4% mannitol, pH 6.8, with a 25 ml G25 Superfine (Pharmacia) column and stored at -80°C.

Immunoadhesin (Fc containing) constructs of proteins are purified from the conditioned media as follows. The conditioned media is pumped onto a 5 ml Protein A column (Pharmacia) which had been equilibrated in 20 mM Na phosphate buffer, pH 6.8. After loading, the column is washed extensively with equilibration buffer before elution with 100 mM citric acid, pH 3.5. The eluted protein is immediately neutralized by collecting 1 ml fractions into tubes containing 275 ml of 1 M Tris buffer, pH 9. The highly purified protein is subsequently desalted into storage buffer as described above for the poly-His tagged proteins. The homogeneity of the sequence is assessed by SDS polyacrylamide gels and by N-terminal amino acid sequencing by Edman degradation and other analytical procedures as desired or necessary.

Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

6.11. EXAMPLE 11: Preparation of Antibodies that Bind PRO

This example illustrates preparation of monoclonal antibodies which can specifically bind the PRO polypeptide or an epitope on the PRO polypeptide without substantially binding to any other polypeptide or polypeptide epitope.

5 Techniques for producing the monoclonal antibodies are known in the art and are described, for instance, in Goding, *supra*. Immunogens that may be employed include purified PRO, fusion proteins containing PRO, and cells expressing recombinant PRO on the cell surface. Selection of the immunogen can be made by the skilled artisan without undue experimentation.

10 Mice, such as Balb/c, are immunized with the PRO immunogen emulsified in complete Freund's adjuvant and injected subcutaneously or intraperitoneally in an amount from 1-100 micrograms. Alternatively, the immunogen is emulsified in MPL-TDM adjuvant (Ribi Immunochemical Research, Hamilton, MT) and injected into the animal's hind foot pads. The immunized mice are then boosted 10 to 12 days later with additional immunogen emulsified in the selected adjuvant. Thereafter, for several weeks, the mice may also be boosted with additional immunization injections. Serum samples may be periodically obtained from the mice by retro-orbital
15 bleeding for testing in ELISA assays to detect anti-PRO antibodies.

After a suitable antibody titer has been detected, the animals "positive" for antibodies can be injected with a final intravenous injection of PRO. Three to four days later, the mice are sacrificed and the spleen cells are harvested. The spleen cells are then fused (using 35% polyethylene glycol) to a selected murine myeloma cell line such as P3X63AgU.1, available from ATCC, No. CRL 1597. The fusions generate hybridoma cells which can then
20 be plated in 96 well tissue culture plates containing HAT (hypoxanthine, aminopterin, and thymidine) medium to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

The hybridoma cells will be screened in an ELISA for reactivity against PRO. Determination of "positive" hybridoma cells secreting the desired monoclonal antibodies against PRO is within the skill in the art.

25 The positive hybridoma cells can be injected intraperitoneally into syngeneic Balb/c mice to produce ascites containing the anti-PRO monoclonal antibodies. Alternatively, the hybridoma cells can be grown in tissue culture flasks or roller bottles. Purification of the monoclonal antibodies produced in the ascites can be accomplished using ammonium sulfate precipitation, followed by gel exclusion chromatography. Alternatively, affinity chromatography based upon binding of antibody to protein A or protein G can be employed.

6.12. EXAMPLE 12: Purification of PRO Polypeptides Using Specific Antibodies

30 Native or recombinant PRO polypeptides may be purified by a variety of standard techniques in the art of protein purification. For example, pro-PRO polypeptide, mature PRO polypeptide, or pre-PRO polypeptide is purified by immunoaffinity chromatography using antibodies specific for the PRO polypeptide of interest. In general, an immunoaffinity column is constructed by covalently coupling the anti-PRO polypeptide antibody to an activated chromatographic resin.

35 Polyclonal immunoglobulins are prepared from immune sera either by precipitation with ammonium sulfate or by purification on immobilized Protein A (Pharmacia LKB Biotechnology, Piscataway, N.J.). Likewise,

monoclonal antibodies are prepared from mouse ascites fluid by ammonium sulfate precipitation or chromatography on immobilized Protein A. Partially purified immunoglobulin is covalently attached to a chromatographic resin such as CnBr-activated SEPHAROSE™ (Pharmacia LKB Biotechnology). The antibody is coupled to the resin, the resin is blocked, and the derivative resin is washed according to the manufacturer's instructions.

Such an immunoaffinity column is utilized in the purification of PRO polypeptide by preparing a fraction from cells containing PRO polypeptide in a soluble form. This preparation is derived by solubilization of the whole cell or of a subcellular fraction obtained via differential centrifugation by the addition of detergent or by other methods well known in the art. Alternatively, soluble PRO polypeptide containing a signal sequence may be secreted in useful quantity into the medium in which the cells are grown.

A soluble PRO polypeptide-containing preparation is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of PRO polypeptide (*e.g.*, high ionic strength buffers in the presence of detergent). Then, the column is eluted under conditions that disrupt antibody/PRO polypeptide binding (*e.g.*, a low pH buffer such as approximately pH 2-3, or a high concentration of a chaotrope such as urea or thiocyanate ion), and PRO polypeptide is collected.

6.13. EXAMPLE 13: Drug Screening

This invention is particularly useful for screening compounds by using PRO polypeptides or binding fragment thereof in any of a variety of drug screening techniques. The PRO polypeptide or fragment employed in such a test may either be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the PRO polypeptide or fragment. Drugs are screened against such transformed cells in competitive binding assays. Such cells, either in viable or fixed form, can be used for standard binding assays. One may measure, for example, the formation of complexes between PRO polypeptide or a fragment and the agent being tested. Alternatively, one can examine the diminution in complex formation between the PRO polypeptide and its target cell or target receptors caused by the agent being tested.

Thus, the present invention provides methods of screening for drugs or any other agents which can affect a PRO polypeptide-associated disease or disorder. These methods comprise contacting such an agent with an PRO polypeptide or fragment thereof and assaying (i) for the presence of a complex between the agent and the PRO polypeptide or fragment, or (ii) for the presence of a complex between the PRO polypeptide or fragment and the cell, by methods well known in the art. In such competitive binding assays, the PRO polypeptide or fragment is typically labeled. After suitable incubation, free PRO polypeptide or fragment is separated from that present in bound form, and the amount of free or uncomplexed label is a measure of the ability of the particular agent to bind to PRO polypeptide or to interfere with the PRO polypeptide/cell complex.

Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to a polypeptide and is described in detail in WO 84/03564, published on September 13, 1984. Briefly stated, large numbers of different small peptide test compounds are synthesized on a solid substrate, such

as plastic pins or some other surface. As applied to a PRO polypeptide, the peptide test compounds are reacted with PRO polypeptide and washed. Bound PRO polypeptide is detected by methods well known in the art. Purified PRO polypeptide can also be coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies can be used to capture the peptide and immobilize it on the solid support.

This invention also contemplates the use of competitive drug screening assays in which neutralizing antibodies capable of binding PRO polypeptide specifically compete with a test compound for binding to PRO polypeptide or fragments thereof. In this manner, the antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with PRO polypeptide.

6.14. EXAMPLE 14: Rational Drug Design

The goal of rational drug design is to produce structural analogs of biologically active polypeptide of interest (*i.e.*, a PRO polypeptide) or of small molecules with which they interact, *e.g.*, agonists, antagonists, or inhibitors. Any of these examples can be used to fashion drugs which are more active or stable forms of the PRO polypeptide or which enhance or interfere with the function of the PRO polypeptide *in vivo* (*c.f.*, Hodgson, Bio/Technology, 9: 19-21 (1991)).

In one approach, the three-dimensional structure of the PRO polypeptide, or of an PRO polypeptide-inhibitor complex, is determined by x-ray crystallography, by computer modeling or, most typically, by a combination of the two approaches. Both the shape and charges of the PRO polypeptide must be ascertained to elucidate the structure and to determine active site(s) of the molecule. Less often, useful information regarding the structure of the PRO polypeptide may be gained by modeling based on the structure of homologous proteins. In both cases, relevant structural information is used to design analogous PRO polypeptide-like molecules or to identify efficient inhibitors. Useful examples of rational drug design may include molecules which have improved activity or stability as shown by Braxton and Wells, Biochemistry, 31:7796-7801 (1992) or which act as inhibitors, agonists, or antagonists of native peptides as shown by Athauda *et al.*, J. Biochem., 113:742-746 (1993).

It is also possible to isolate a target-specific antibody, selected by functional assay, as described above, and then to solve its crystal structure. This approach, in principle, yields a pharmacore upon which subsequent drug design can be based. It is possible to bypass protein crystallography altogether by generating anti-idiotypic antibodies (anti-ids) to a functional, pharmacologically active antibody. As a mirror image of a mirror image, the binding site of the anti-ids would be expected to be an analog of the original receptor. The anti-id could then be used to identify and isolate peptides from banks of chemically or biologically produced peptides. The isolated peptides would then act as the pharmacore.

By virtue of the present invention, sufficient amounts of the PRO polypeptide may be made available to perform such analytical studies as X-ray crystallography. In addition, knowledge of the PRO polypeptide amino acid sequence provided herein will provide guidance to those employing computer modeling techniques in place of or in addition to x-ray crystallography.

6.15. EXAMPLE 15: Stimulation of Endothelial Cell Proliferation (Assay 8)

This assay is designed to determine whether PRO polypeptides of the present invention show the ability to stimulate adrenal cortical capillary endothelial cell (ACE) growth. PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of conditions or disorders where angiogenesis would be beneficial including, for example, wound healing, and the like (as would agonists of these PRO polypeptides). Antagonists of the PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of cancerous tumors.

Bovine adrenal cortical capillary endothelial (ACE) cells (from primary culture, maximum of 12-14 passages) were plated in 96-well plates at 500 cells/well per 100 microliter. Assay media included low glucose DMEM, 10% calf serum, 2 mM glutamine, and 1X penicillin/streptomycin/fungizone. Control wells included the following: (1) no ACE cells added; (2) ACE cells alone; (3) ACE cells plus VEGF (5 ng/ml); and (4) ACE cells plus FGF (5ng/ml). The control or test sample, (in 100 microliter volumes), was then added to the wells (at dilutions of 1%, 0.1% and 0.01%, respectively). The cell cultures were incubated for 6-7 days at 37°C/5% CO₂. After the incubation, the media in the wells was aspirated, and the cells were washed 1X with PBS. An acid phosphatase reaction mixture (100 microliter; 0.1M sodium acetate, pH 5.5, 0.1% Triton X-100, 10 mM p-nitrophenyl phosphate) was then added to each well. After a 2 hour incubation at 37°C, the reaction was stopped by addition of 10 microliters 1N NaOH. Optical density (OD) was measured on a microplate reader at 405 nm.

The activity of a PRO polypeptide was calculated as the fold increase in proliferation (as determined by the acid phosphatase activity, OD 405 nm) relative to (1) cell only background, and (2) relative to maximum stimulation by VEGF. VEGF (at 3-10 ng/ml) and FGF (at 1-5 ng/ml) were employed as an activity reference for maximum stimulation. Results of the assay were considered "positive" if the observed stimulation was \geq 50% increase over background. VEGF (5 ng/ml) control at 1% dilution gave 1.24 fold stimulation; FGF (5 ng/ml) control at 1% dilution gave 1.46 fold stimulation.

PRO21 tested positive in this assay.

6.16. EXAMPLE 16: Inhibition of Vascular Endothelial Growth Factor (VEGF) Stimulated Proliferation of Endothelial Cell Growth (Assay 9)

The ability of various PRO polypeptides to inhibit VEGF stimulated proliferation of endothelial cells was tested. Polypeptides testing positive in this assay are useful for inhibiting endothelial cell growth in mammals where such an effect would be beneficial, e.g., for inhibiting tumor growth.

Specifically, bovine adrenal cortical capillary endothelial cells (ACE) (from primary culture, maximum of 12-14 passages) were plated in 96-well plates at 500 cells/well per 100 microliter. Assay media included low glucose DMEM, 10% calf serum, 2 mM glutamine, and 1X penicillin/streptomycin/fungizone. Control wells included the following: (1) no ACE cells added; (2) ACE cells alone; (3) ACE cells plus 5 ng/ml FGF; (4) ACE cells plus 3 ng/ml VEGF; (5) ACE cells plus 3 ng/ml VEGF plus 1 ng/ml TGF-beta; and (6) ACE cells plus 3 ng/ml VEGF plus 5 ng/ml LIF. The test samples, poly-his tagged PRO polypeptides (in 100 microliter volumes), were then added to the wells (at dilutions of 1%, 0.1% and 0.01%, respectively). The cell cultures were

incubated for 6-7 days at 37°C/5% CO₂. After the incubation, the media in the wells was aspirated, and the cells were washed 1X with PBS. An acid phosphatase reaction mixture (100 microliter; 0.1M sodium acetate, pH 5.5, 0.1% Triton X-100, 10 mM p-nitrophenyl phosphate) was then added to each well. After a 2 hour incubation at 37°C, the reaction was stopped by addition of 10 microliters 1N NaOH. Optical density (OD) was measured on a microplate reader at 405 nm.

The activity of PRO polypeptides was calculated as the percent inhibition of VEGF (3 ng/ml) stimulated proliferation (as determined by measuring acid phosphatase activity at OD 405 nm) relative to the cells without stimulation. TGF-beta was employed as an activity reference at 1 ng/ml, since TGF-beta blocks 70-90% of VEGF-stimulated ACE cell proliferation. The results are indicative of the utility of the PRO polypeptides in cancer therapy and specifically in inhibiting tumor angiogenesis. Numerical values (relative inhibition) are determined by calculating the percent inhibition of VEGF stimulated proliferation by the PRO polypeptides relative to cells without stimulation and then dividing that percentage into the percent inhibition obtained by TGF-β at 1 ng/ml which is known to block 70-90% of VEGF stimulated cell proliferation. The results are considered positive if the PRO polypeptide exhibits 30% or greater inhibition of VEGF stimulation of endothelial cell growth (relative inhibition 30% or greater).

PRO247, PRO720 and PRO4302 tested positive in this assay.

6.17. EXAMPLE 17: Enhancement of Heart Neonatal Hypertrophy Induced by LIF+ET-1 (Assay 75)

This assay is designed to determine whether PRO polypeptides of the present invention show the ability to enhance neonatal heart hypertrophy induced by LIF and endothelin-1 (ET-1). A test compound that provides a positive response in the present assay would be useful for the therapeutic treatment of cardiac insufficiency diseases or disorders characterized or associated with an undesired level of hypertrophy of the cardiac muscle.

Cardiac myocytes from 1-day old Harlan Sprague Dawley rats (180 μl at 7.5 x 10⁴/ml, serum <0.1, freshly isolated) are introduced on day 1 to 96-well plates previously coated with DMEM/F12 + 4%FCS. Test PRO polypeptide samples or growth medium alone (negative control) are then added directly to the wells on day 2 in 20 μl volume. LIF + ET-1 are then added to the wells on day 3. The cells are stained after an additional 2 days in culture and are then scored visually the next day. A positive in the assay occurs when the PRO polypeptide treated myocytes obtain a score greater than zero. A score of zero represents non-responsive cells whereas scores of 1 or 2 represent enhancement (*i.e.* they are visually larger on the average or more numerous than the untreated myocytes).

PRO21 polypeptides tested positive in this assay.

6.18. EXAMPLE 18: Detection of Endothelial Cell Apoptosis (FACS) (Assay 96)

The ability of PRO polypeptides of the present invention to induce apoptosis in endothelial cells was tested in human venous umbilical vein endothelial cells (HUVEC, Cell Systems) in gelatinized T175 flasks using HUVEC cells below passage 10. PRO polypeptides testing positive in this assay are expected to be useful for

therapeutically treating conditions where apoptosis of endothelial cells would be beneficial including, for example, the therapeutic treatment of tumors.

On day one, the cells were split [420,000 cells per gelatinized 6 cm dishes - (11×10^3 cells/cm² Falcon, Primaria)] and grown in media containing serum (CS-C, Cell System) overnight or for 16 hours to 24 hours.

On day 2, the cells were washed 1x with 5 ml PBS ; 3 ml of 0% serum medium was added with VEGF (100 ng/ml); and 30 μ l of the PRO test compound (final dilution 1%) or 0% serum medium (negative control) was added. The mixtures were incubated for 48 hours before harvesting.

The cells were then harvested for FACS analysis. The medium was aspirated and the cells washed once with PBS. 5 ml of 1 x trypsin was added to the cells in a T-175 flask, and the cells were allowed to stand until they were released from the plate (about 5-10 minutes). Trypsinization was stopped by adding 5 ml of growth media. The cells were spun at 1000 rpm for 5 minutes at 4°C. The media was aspirated and the cells were resuspended in 10 ml of 10% serum complemented medium (Cell Systems), 5 μ l of Annexin-FITC (BioVison) added and chilled tubes were submitted for FACS. A positive result was determined to be enhanced apoptosis in the PRO polypeptide treated samples as compared to the negative control.

PRO4302 polypeptide tested positive in this assay.

6.19. EXAMPLE 19: Induction of c-fos in HUVEC Cells (Assay 123)

This assay is designed to determine whether PRO polypeptides show the ability to induce c-fos in HUVEC cells. PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of conditions or disorders where angiogenesis would be beneficial including, for example, wound healing, and the like (as would agonists of these PRO polypeptides). Antagonists of the PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of cancerous tumors.

Human venous umbilical vein endothelial cells (HUVEC, Cell Systems) in growth media (50% Ham's F12 w/o GHT: low glucose, and 50% DMEM without glycine: with NaHCO₃, 1% glutamine, 10 mM HEPES, 10% FBS, 10 ng/ml bFGF) were plated on 96-well microtiter plates at a cell density of 5×10^3 cells/well. The day after plating (day 2), the cells were starved for 24 hours by removing the growth media and replacing with serum free media. On day 3, the cells are treated with 100 μ l/well test samples and controls (positive control = growth media; negative control = Protein 32 buffer = 10 mM HEPES, 140 mM NaCl, 4% (w/v) mannitol, pH 6.8). One plate of cells was incubated for 30 minutes at 37°C, in 5% CO₂. Another plate of cells was incubated for 60 minutes at 37°C, in 5% CO₂. The samples were removed, and RNA was harvested using the RNeasy 96 kit (Qiagen). Next, the RNA was assayed for c-fos, egr-1 and GAPDH induction using Taqman.

The measure of activity of the fold increase over the negative control (Protein 32/HEPES buffer described above) value was by obtained by calculating the fold increase of the ratio of c-fos to GAPDH in test samples as compared to the negative control. The results are considered positive if the PRO polypeptide exhibits at least a two-fold value over the negative buffer control.

PRO1376 polypeptide tested positive in this assay.

6.20. EXAMPLE 20: Normal Human Iliac Artery Endothelial Cell Proliferation (Assay 138)

This assay is designed to determine whether PRO polypeptides of the present invention show the ability to induce proliferation of human iliac artery endothelial cells in culture and, therefore, function as useful growth factors.

On day 0, human iliac artery endothelial cells (from cell lines, maximum of 12-14 passages) were plated in 96-well plates at 1000 cells/well per 100 microliter and incubated overnight in complete media [epithelial cell growth media (EGM, Clonetics), plus supplements: human epithelial growth factor (hEGF), bovine brain extract (BBE), hydrocortisone, GA-1000, and fetal bovine serum (FBS, Clonetics)]. On day 1, complete media was replaced by basal media [EGM plus 1% FBS] and addition of PRO polypeptides at 1%, 0.1% and 0.01%. On day 7, an assessment of cell proliferation was performed by Alamar Blue assay followed by Crystal Violet. Results are expressed as % of the cell growth observed with control buffer.

The following PRO polypeptides tested positive in this assay: PRO214, PRO238, PRO256, PRO363, PRO365, PRO791, PRO836, PRO1025, PRO1029, PRO1186, PRO1192, PRO1272, PRO1274, PRO1279, PRO1306, PRO1325, PRO1329, PRO1376, PRO1411, PRO1419, PRO1508, PRO1787, PRO1868, PRO1890, PRO4324, PRO4333, PRO4408, PRO4499, PRO5725, PRO6006, PRO9821, PRO9873, PRO10008, PRO10096, PRO19670, PRO20040, PRO20044, PRO21384 and PRO28631.

6.21. EXAMPLE 21: Pooled Human Umbilical Vein Endothelial Cell Proliferation (Assay 139)

This assay is designed to determine whether PRO polypeptides of the present invention show the ability to induce proliferation of pooled human umbilical vein endothelial cells in culture and, therefore, function as useful growth factors.

On day 0, pooled human umbilical vein endothelial cells (from cell lines, maximum of 12-14 passages) were plated in 96-well plates at 1000 cells/well per 100 microliter and incubated overnight in complete media [epithelial cell growth media (EGM, Clonetics), plus supplements: human epithelial growth factor (hEGF), bovine brain extract (BBE), hydrocortisone, GA-1000, and fetal bovine serum (FBS, Clonetics)]. On day 1, complete media was replaced by basal media [EGM plus 1% FBS] and addition of PRO polypeptides at 1%, 0.1% and 0.01%. On day 7, an assessment of cell proliferation was performed by Alamar Blue assay followed by Crystal Violet. Results are expressed as % of the cell growth observed with control buffer.

The following PRO polypeptides tested positive in this assay: PRO181, PRO205, PRO221, PRO229, PRO231, PRO238, PRO241, PRO247, PRO256, PRO258, PRO263, PRO265, PRO295, PRO321, PRO322, PRO337, PRO363, PRO444, PRO533, PRO697, PRO725, PRO771, PRO788, PRO819, PRO827, PRO828, PRO846, PRO865, PRO1005, PRO1006, PRO1007, PRO1025, PRO1054, PRO1071, PRO1075, PRO1079, PRO1080, PRO1114, PRO1131, PRO1155, PRO1160, PRO1184, PRO1190, PRO1192, PRO1195, PRO1244, PRO1272, PRO1273, PRO1279, PRO1283, PRO1286, PRO1306, PRO1309, PRO1325, PRO1329, PRO1347, PRO1356, PRO1376, PRO1382, PRO1412, PRO1419, PRO1474, PRO1477, PRO1488, PRO1550, PRO1556, PRO1760, PRO1782, PRO1787, PRO1801, PRO1868, PRO1887, PRO3438, PRO3444, PRO4302, PRO4324, PRO4341, PRO4342, PRO4353, PRO4354, PRO4356, PRO4371, PRO4405, PRO4422, PRO4425, PRO5723,

PRO5725, PRO5737, PRO5776, PRO6029, PRO6071, PRO7436, PRO9771, PRO10008, PRO10096, PRO21055 and PRO21384.

6.22. EXAMPLE 22: Human Coronary Artery Smooth Muscle Cell Proliferation (Assay 140)

This assay is designed to determine whether PRO polypeptides of the present invention show the ability to induce proliferation of human coronary artery smooth muscle cells in culture and, therefore, function as useful growth factors.

On day 0, human coronary artery smooth muscle cells (from cell lines, maximum of 12-14 passages) were plated in 96-well plates at 1000 cells/well per 100 microliter and incubated overnight in complete media [smooth muscle growth media (SmGM, Clonetics), plus supplements: insulin, human epithelial growth factor (hEGF), human fibroblast growth factor (hFGF), GA-1000, and fetal bovine serum (FBS, Clonetics)]. On day 1, complete media was replaced by basal media [SmGM plus 1% FBS] and addition of PRO polypeptides at 1%, 0.1% and 0.01%. On day 7, an assessment of cell proliferation was performed by Alamar Blue assay followed by Crystal Violet. Results are expressed as % of the cell growth observed with control buffer.

The following PRO polypeptides tested positive in this assay: PRO162, PRO181, PRO182, PRO195, PRO204, PRO221, PRO230, PRO256, PRO258, PRO533, PRO697, PRO725, PRO738, PRO826, PRO836, PRO840, PRO846, PRO865, PRO982, PRO1025, PRO1029, PRO1071, PRO1080, PRO1083, PRO1134, PRO1160, PRO1182, PRO1184, PRO1186, PRO1192, PRO1265, PRO1274, PRO1279, PRO1283, PRO1306, PRO1308, PRO1309, PRO1325, PRO1337, PRO1338, PRO1343, PRO1376, PRO1387, PRO1411, PRO1412, PRO1415, PRO1434, PRO1474, PRO1488, PRO1550, PRO1556, PRO1567, PRO1600, PRO1754, PRO1758, PRO1760, PRO1787, PRO1865, PRO1868, PRO1917, PRO1928, PRO3438, PRO3562, PRO4302, PRO4333, PRO4345, PRO4353, PRO4354, PRO4405, PRO4408, PRO4430, PRO4503, PRO5725, PRO6714, PRO9771, PRO9820, PRO9940, PRO10096, PRO21055, PRO21184 and PRO21366.

6.23. EXAMPLE 23: Microarray Analysis to Detect Overexpression of PRO Polypeptides in HUVEC Cells Treated with Growth Factors

This assay is designed to determine whether PRO polypeptides of the present invention show the ability to induce angiogenesis by stimulating endothelial cell tube formation in HUVEC cells.

Nucleic acid microarrays, often containing thousands of gene sequences, are useful for identifying differentially expressed genes in tissues exposed to various stimuli (*e.g.*, growth factors) as compared to their normal, unexposed counterparts. Using nucleic acid microarrays, test and control mRNA samples from test and control tissue samples are reverse transcribed and labeled to generate cDNA probes. The cDNA probes are then hybridized to an array of nucleic acids immobilized on a solid support. The array is configured such that the sequence and position of each member of the array is known. Hybridization of a labeled probe with a particular array member indicates that the sample from which the probe was derived expresses that gene. If the hybridization signal of a probe from a test (exposed tissue) sample is greater than hybridization signal of a probe from a control (normal, unexposed tissue) sample, the gene or genes overexpressed in the exposed tissue are identified. The

implication of this result is that an overexpressed protein in an exposed tissue may be involved in the functional changes within the tissue following exposure to the stimuli (*e.g.*, tube formation).

The methodology of hybridization of nucleic acids and microarray technology is well known in the art. In the present example, the specific preparation of nucleic acids for hybridization and probes, slides, and hybridization conditions are all detailed in U.S. Provisional Patent Application Serial No. 60/193,767, filed on March 31, 2000 and which is herein incorporated by reference.

In the present example, HUVEC cells grown in either collagen gels or fibrin gels were induced to form tubes by the addition of various growth factors. Specifically, collagen gels were prepared as described previously in Yang *et al.*, *American J. Pathology*, 1999, 155(3):887-895 and Xin *et al.*, *American J. Pathology*, 2001, 158(3):1111-1120. Following gelation of the HUVEC cells, 1X basal medium containing M199 supplemented with 1%FBS, 1X ITS, 2 mM L-glutamine, 50 µg/ml ascorbic acid, 26.5 mM NaHCO₃, 100U/ml penicillin and 100 U/ml streptomycin was added. Tube formation was elicited by the inclusion in the culture media of either a mixture of phorbol myrsitate acetate (50 nM), vascular endothelial cell growth factor (40 ng/ml) and basic fibroblast growth factor (40 ng/ml) ("PMA growth factor mix") or hepatocyte growth factor (40 ng/ml) and vascular endothelial cell growth factor (40 ng/ml) (HGF/VEGF mix) for the indicated period of time. Fibrin Gels were prepared by suspending Huvec (4 x 10⁵ cells/ml) in M199 containing 1% fetal bovine serum (Hyclone) and human fibrinogen (2.5mg/ml). Thrombin (50U/ml) was then added to the fibrinogen suspension at a ratio of 1 part thrombin solution:30 parts fibrinogen suspension. The solution was then layered onto 10 cm tissue culture plates (total volume: 15 ml/plate) and allowed to solidify at 37°C for 20 min. Tissue culture media (10 ml of BM containing PMA (50 nM), bFGF (40ng/ml) and VEGF (40 ng/ml)) was then added and the cells incubated at 37°C in 5%CO₂ in air for the indicated period of time.

Total RNA was extracted from the HUVEC cells incubated for 0, 4, 8, 24, 40 and 50 hours in the different matrix and media combinations using a TRIzol extraction followed by a second purification using RNAeasy Mini Kit (Qiagen). The total RNA was used to prepare cRNA which was then hybridized to the microarrays.

In the present experiments, nucleic acid probes derived from the herein described PRO polypeptide-encoding nucleic acid sequences were used in the creation of the microarray and RNA from the HUVEC cells described above were used for the hybridization thereto. Pairwise comparisons were made using time 0 chips as a baseline. Three replicate samples were analyzed for each experimental condition and time. Hence there were 3 time 0 samples for each treatment and 3 replicates of each successive time point. Therefore, a 3 by 3 comparison was performed for each time point compared against each time 0 point. This resulted in 9 comparisons per time point. Only those genes that had increased expression in all three non-time-0 replicates in each of the different matrix and media combinations as compared to any of the three time zero replicates were considered positive. Although this stringent method of data analysis does allow for false negatives, it minimizes false positives.

PRO178, PRO195, PRO228, PRO301, PRO302, PRO532, PRO724, PRO730, PRO734, PRO793, PRO871, PRO938, PRO1012, PRO1120, PRO1139, PRO1198, PRO1287, PRO1361, PRO1864, PRO1873,

PRO2010, PRO3579, PRO4313, PRO4527, PRO4538, PRO4553, PRO4995, PRO5730, PRO6008, PRO7223, PRO7248 and PRO7261 tested positive in this assay.

6.24. EXAMPLE 24: *In situ* Hybridization

In situ hybridization is a powerful and versatile technique for the detection and localization of nucleic acid sequences within cell or tissue preparations. It may be useful, for example, to identify sites of gene expression, analyze the tissue distribution of transcription, identify and localize viral infection, follow changes in specific mRNA synthesis, and aid in chromosome mapping.

In situ hybridization was performed following an optimized version of the protocol by Lu and Gillett, Cell Vision, 1: 169-176 (1994), using PCR-generated ^{32}P -labeled riboprobes. Briefly, formalin-fixed, paraffin-embedded human tissues were sectioned, deparaffinized, deproteinized in proteinase K (20 g/ml) for 15 minutes at 37°C, and further processed for *in situ* hybridization as described by Lu and Gillett, *supra*. A (^{32}P -UTP)-labeled antisense riboprobe was generated from a PCR product and hybridized at 55°C overnight. The slides were dipped in Kodak NTB2™ nuclear track emulsion and exposed for 4 weeks.

6.24.1. ^{32}P -Riboprobe synthesis

6.0 μl (125 mCi) of ^{32}P -UTP (Amersham BF 1002, SA < 2000 Ci/mmol) were speed-vacuum dried. To each tube containing dried ^{32}P -UTP, the following ingredients were added:

2.0 μl 5x transcription buffer

1.0 μl DTT (100 mM)

2.0 μl NTP mix (2.5 mM: 10 μl each of 10 mM GTP, CTP & ATP + 10 μl H₂O)

1.0 μl UTP (50 μM)

1.0 μl RNAsin

1.0 μl DNA template (1 μg)

1.0 μl H₂O

1.0 μl RNA polymerase (for PCR products T3 = AS, T7 = S, usually)

The tubes were incubated at 37°C for one hour. A total of 1.0 μl RQ1 DNase was added, followed by incubation at 37°C for 15 minutes. A total of 90 μl TE (10 mM Tris pH 7.6/1 mM EDTA pH 8.0) was added, and the mixture was pipetted onto DE81 paper. The remaining solution was loaded in a MICROCON-50™ ultrafiltration unit, and spun using program 10 (6 minutes). The filtration unit was inverted over a second tube and spun using program 2 (3 minutes). After the final recovery spin, a total of 100 μl TE was added, then 1 μl of the final product was pipetted on DE81 paper and counted in 6 ml of BIOFLUOR II™.

The probe was run on a TBE/urea gel. A total of 1-3 μl of the probe or 5 μl of RNA Mrk III was added to 3 μl of loading buffer. After heating on a 95°C heat block for three minutes, the gel was immediately placed on ice. The wells of gel were flushed, and the sample was loaded and run at 180-250 volts for 45 minutes. The gel was wrapped in plastic wrap (SARAN™ brand) and exposed to XAR film with an intensifying screen in a -70°C freezer one hour to overnight.

6.24.2. ³²P-Hybridization6.24.2.1. *Pretreatment of frozen sections*

5 The slides were removed from the freezer, placed on aluminum trays, and thawed at room temperature for 5 minutes. The trays were placed in a 55°C incubator for five minutes to reduce condensation. The slides were fixed for 10 minutes in 4% paraformaldehyde on ice in the fume hood, and washed in 0.5 x SSC for 5 minutes, at room temperature (25 ml 20 x SSC + 975 ml SQ H₂O). After deproteinization in 0.5 µg/ml proteinase K for 10 minutes at 37°C (12.5 µl of 10 mg/ml stock in 250 ml prewarmed RNase-free RNase buffer), the sections were washed in 0.5 x SSC for 10 minutes at room temperature. The sections were dehydrated in 70%, 95%, and 100% ethanol, 2 minutes each.

6.24.2.2. *Pretreatment of paraffin-embedded sections*

10 The slides were deparaffinized, placed in SQ H₂O, and rinsed twice in 2 x SSC at room temperature, for 5 minutes each time. The sections were deproteinized in 20 µg/ml proteinase K (500 µl of 10 mg/ml in 250 ml RNase-free RNase buffer; 37°C, 15 minutes) for human embryo tissue, or 8 x proteinase K (100 µl in 250 ml RNase buffer, 37°C, 30 minutes) for formalin tissues. Subsequent rinsing in 0.5 x SSC and dehydration were performed as described above.

6.24.2.3. *Prehybridization*

20 The slides were laid out in a plastic box lined with Box buffer (4 x SSC, 50% formamide) - saturated filter paper. The tissue was covered with 50 µl of hybridization buffer (3.75 g dextran sulfate + 6 ml SQ H₂O), vortexed, and heated in the microwave for 2 minutes with the cap loosened. After cooling on ice, 18.75 ml formamide, 3.75 ml 20 x SSC, and 9 ml SQ H₂O were added, and the tissue was vortexed well and incubated at 42°C for 1-4 hours.

6.24.2.4. *Hybridization*

25 1.0 x 10⁶ cpm probe and 1.0 µl tRNA (50 mg/ml stock) per slide were heated at 95°C for 3 minutes. The slides were cooled on ice, and 48 µl hybridization buffer was added per slide. After vortexing, 50 µl ³²P mix was added to 50 µl prehybridization on the slide. The slides were incubated overnight at 55°C.

6.24.2.5. *Washes*

30 Washing was done for 2x10 minutes with 2xSSC, EDTA at room temperature (400 ml 20 x SSC + 16 ml 0.25 M EDTA, V_f=4L), followed by RNaseA treatment at 37°C for 30 minutes (500 µl of 10 mg/ml in 250 ml RNase buffer = 20 µg/ml). The slides were washed 2 x10 minutes with 2 x SSC, EDTA at room temperature. The stringency wash conditions were as follows: 2 hours at 55°C, 0.1 x SSC, EDTA (20 ml 20 x SSC + 16 ml EDTA, V_f=4L).

6.24.2.6. *Oligonucleotides*

In situ analysis was performed on three of the DNA sequences disclosed herein. The primers used to generate the probes and/or the probes employed for these analyses are as follows:

- 5 DNA33100-p1: 5'GGA TTC TAA TAC GAC TCA CTA TAG GGC CGG GTG GAG GTG GAA CAG AAA
3' (SEQ ID NO:375)
- DNA33100-p2: 5' CTA TGA AAT TAA CCC TCA CTA AAG GGA CAC AGA CAG AGC CCC ATA CGC
3' (SEQ ID NO:376)
- DNA34431-p1: 5'GGA TTC TAA TAC GAC TCA CTA TAG GGC CAG GGA AAT CCG GAT GTC TC
3' (SEQ ID NO:377)
- 10 DNA34431-p2: 5' CTA TGA AAT TAA CCC TCA CTA AAG GGA GTA AGG GGA TGC CAC CGA GTA
3' (SEQ ID NO:378)
- DNA38268-p1: 5'GGA TTC TAA TAC GAC TCA CTA TAG GGC CAG CTA CCC GCA GGA GGA GG
3' (SEQ ID NO:379)
- DNA38268-p2: 5'CTA TGA AAT TAA CCC TCA CTA AAG GGA TCC CAG GTG ATG AGG TCC AGA
15 3' (SEQ ID NO:380)
- DNA64908 probe: 5'CCATCTCGGAGACCTTTGTGCAGCGTGTATACCAGCCTTACCTCACCA
CTTGCGACGGACACAGAGCCTGCAGCACCTACCGAACCATCTACCGGAC
TGCCTATCGCCGTAGCCCTGGGGTGACTCCCGCAAGCCTCGCTATGCTTG
CTGCCCTGGTTGGAAGAGGACCAGTGGGCTCCCTGGGGCTTGTGGAGCA
20 GCAATATGCCAGCCTCCATGTGGGAATGGAGGGAGTTGCATCCGCCCAG
GACACTGCCGCTGCCCTGTGGGATGGCAGGGAGATACTTGCCAGACAGA
TGTTGATGAATGCAGTACAGGAGAGGCCAGTTGTCCCCAGCGCTGTGTC
AATACTGTGGGAAGTTACTGGTGCCAGGGATGGGAGGGACAAAGCCCAT
CTGCAGATGGGACGCGCTGCCTGTCTAAGGAGGGGCCCTCCCGGTGGCC
25 CCAACCCACAGCAGGAGTGGACAGCA3' (SEQ ID NO:381)

6.24.2.7. *Results*

In situ analysis was performed and the results from these analyses are as follows:

6.24.2.7.1. DNA33100-1159 (PRO229) (Scavenger-R/CD6
homologTNF motif)

A specific positive signal was observed in mononuclear phagocytes (macrophages) of fetal and adult spleen, liver, lymph node and thymus. All other tissues were negative.

5

6.24.2.7.2. DNA34431-1177 (PRO263) (CD44)

A specific positive signal was observed in human fetal tissues and placenta over mononuclear cells, with strong expression in epithelial cells of the adrenal cortex. All adult tissues were negative.

6.24.2.7.3. DNA38268-1188 (PRO295) (Integrin)

10

A specific positive signal was observed in human fetal ganglion cells, fetal neurons, adult adrenal medulla and adult neurons. All other tissues were negative.

6.24.2.7.4. DNA64908-1163-1 (PRO1449)

15

A specific positive signal was observed in the developing vasculature (from E7-E11), in endothelial cells and in progenitors of endothelial cells in wholemount in situ hybridizations of mouse embryos (Figure 375). Specific expression was also observed in a subset of blood vessels and epidermis from E12 onward. A mouse orthologue of PRO1449 which has about 78 % amino acid identity with PRO1449 was used as the probe.

In normal adult tissues, expression was low to absent. When present, expression was confined to the vasculature (Figure 376). Figure 376 further shows that highest expression in adult tissues was observed regionally in vessels running within the white matter of the brain. Elevated expression was also observed in vasculature of many inflamed and diseased tissues, including, but not limited to, tumor vasculature.

20

Following electroporation of the mouse orthologue of PRO1449 into the choroid layer in the eyes of chicken embryos, new vessel formation was observed in the electroporated eye (top right), but not in the control side from the same embryo (top left), or an embryo that was electroporated with a control cDNA (bottom right) (Figure 377).

25

The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the invention. The present invention is not to be limited in scope by the construct(s) deposited, since the deposited embodiment(s) is/are intended as single illustration(s) of certain aspects of the invention and any constructs that are functionally equivalent are within the scope of this invention. The deposit of material(s) herein does not constitute an admission that the written description herein contained is inadequate to enable the practice of any aspect of the invention, including the best mode thereof, nor is it to be construed as limiting the scope of the claims to the specific illustrations that it represents. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the appended claims.

30

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule having at least 80% nucleic acid sequence identity to a nucleotide sequence that encodes an amino acid sequence selected from the group consisting of the amino acid sequence shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:4), Figure 6 (SEQ ID NO:6), Figure 8 (SEQ ID NO:8), Figure 10 (SEQ ID NO:10), Figure 12 (SEQ ID NO:12), Figure 14 (SEQ ID NO:14), Figure 16 (SEQ ID NO:16), Figure 18 (SEQ ID NO:18), Figure 20 (SEQ ID NO:20), Figure 22 (SEQ ID NO:22), Figure 24 (SEQ ID NO:24), Figure 26 (SEQ ID NO:26), Figure 28 (SEQ ID NO:28), Figure 30 (SEQ ID NO:30), Figure 32 (SEQ ID NO:32), Figure 34 (SEQ ID NO:34), Figure 36 (SEQ ID NO:36), Figure 38 (SEQ ID NO:38), Figure 40 (SEQ ID NO:40), Figure 42 (SEQ ID NO:42), Figure 44 (SEQ ID NO:44), Figure 46 (SEQ ID NO:46), Figure 48 (SEQ ID NO:48), Figure 50 (SEQ ID NO:50), Figure 52 (SEQ ID NO:52), Figure 54 (SEQ ID NO:54), Figure 56 (SEQ ID NO:56), Figure 58 (SEQ ID NO:58), Figure 60 (SEQ ID NO:60), Figure 62 (SEQ ID NO:62), Figure 64 (SEQ ID NO:64), Figure 66 (SEQ ID NO:66), Figure 68 (SEQ ID NO:68), Figure 70 (SEQ ID NO:70), Figure 72 (SEQ ID NO:72), Figure 74 (SEQ ID NO:74), Figure 76 (SEQ ID NO:76), Figure 78 (SEQ ID NO:78), Figure 80 (SEQ ID NO:80), Figure 82 (SEQ ID NO:82), Figure 84 (SEQ ID NO:84), Figure 86 (SEQ ID NO:86), Figure 88 (SEQ ID NO:88), Figure 90 (SEQ ID NO:90), Figure 92 (SEQ ID NO:92), Figure 94 (SEQ ID NO:94), Figure 96 (SEQ ID NO:96), Figure 98 (SEQ ID NO:98), Figure 100 (SEQ ID NO:100), Figure 102 (SEQ ID NO:102), Figure 104 (SEQ ID NO:104), Figure 106 (SEQ ID NO:106), Figure 108 (SEQ ID NO:108), Figure 110 (SEQ ID NO:110), Figure 112 (SEQ ID NO:112), Figure 114 (SEQ ID NO:114), Figure 116 (SEQ ID NO:116), Figure 118 (SEQ ID NO:118), Figure 120 (SEQ ID NO:120), Figure 122 (SEQ ID NO:122), Figure 124 (SEQ ID NO:124), Figure 126 (SEQ ID NO:126), Figure 128 (SEQ ID NO:128), Figure 130 (SEQ ID NO:130), Figure 132 (SEQ ID NO:132), Figure 134 (SEQ ID NO:134), Figure 136 (SEQ ID NO:136), Figure 138 (SEQ ID NO:138), Figure 140 (SEQ ID NO:140), Figure 142 (SEQ ID NO:142), Figure 144 (SEQ ID NO:144), Figure 146 (SEQ ID NO:146), Figure 148 (SEQ ID NO:148), Figure 150 (SEQ ID NO:150), Figure 152 (SEQ ID NO:152), Figure 154 (SEQ ID NO:154), Figure 156 (SEQ ID NO:156), Figure 158 (SEQ ID NO:158), Figure 160 (SEQ ID NO:160), Figure 162 (SEQ ID NO:162), Figure 164 (SEQ ID NO:164), Figure 166 (SEQ ID NO:166), Figure 168 (SEQ ID NO:168), Figure 170 (SEQ ID NO:170), Figure 172 (SEQ ID NO:172), Figure 174 (SEQ ID NO:174), Figure 176 (SEQ ID NO:176), Figure 178 (SEQ ID NO:178), Figure 180 (SEQ ID NO:180), Figure 182 (SEQ ID NO:182), Figure 184 (SEQ ID NO:184), Figure 186 (SEQ ID NO:186), Figure 188 (SEQ ID NO:188), Figure 190 (SEQ ID NO:190), Figure 192 (SEQ ID NO:192), Figure 194 (SEQ ID NO:194), Figure 196 (SEQ ID NO:196), Figure 198 (SEQ ID NO:198), Figure 200 (SEQ ID NO:200), Figure 202 (SEQ ID NO:202), Figure 204 (SEQ ID NO:204), Figure 206 (SEQ ID NO:206), Figure 208 (SEQ ID NO:208), Figure 210 (SEQ ID NO:210), Figure 212 (SEQ ID NO:212), Figure 214 (SEQ ID NO:214), Figure 216 (SEQ ID NO:216), Figure 218 (SEQ ID NO:218), Figure 220 (SEQ ID NO:220), Figure 222 (SEQ ID NO:222), Figure 224 (SEQ ID NO:224), Figure 226 (SEQ ID NO:226), Figure 228 (SEQ ID NO:228), Figure 230 (SEQ ID NO:230), Figure 232 (SEQ ID NO:232), Figure 234 (SEQ ID NO:234), Figure 236 (SEQ ID NO:236), Figure 238 (SEQ ID NO:238), Figure 240 (SEQ ID NO:240), Figure 242 (SEQ ID NO:242), Figure 244 (SEQ ID NO:244), Figure 246 (SEQ ID NO:246), Figure

248 (SEQ ID NO:248), Figure 250 (SEQ ID NO:250), Figure 252 (SEQ ID NO:252), Figure 254 (SEQ ID NO:254), Figure 256 (SEQ ID NO:256), Figure 258 (SEQ ID NO:258), Figure 260 (SEQ ID NO:260), Figure 262 (SEQ ID NO:262), Figure 264 (SEQ ID NO:264), Figure 266 (SEQ ID NO:266), Figure 268 (SEQ ID NO:268), Figure 270 (SEQ ID NO:270), Figure 272 (SEQ ID NO:272), Figure 274 (SEQ ID NO:274), Figure 276 (SEQ ID NO:276), Figure 278 (SEQ ID NO:278), Figure 280 (SEQ ID NO:280), Figure 282 (SEQ ID NO:282), Figure 284 (SEQ ID NO:284), Figure 286 (SEQ ID NO:286), Figure 288 (SEQ ID NO:288), Figure 290 (SEQ ID NO:290), Figure 292 (SEQ ID NO:292), Figure 294 (SEQ ID NO:294), Figure 296 (SEQ ID NO:296), Figure 298 (SEQ ID NO:298), Figure 300 (SEQ ID NO:300), Figure 302 (SEQ ID NO:302), Figure 304 (SEQ ID NO:304), Figure 306 (SEQ ID NO:306), Figure 308 (SEQ ID NO:308), Figure 310 (SEQ ID NO:310), Figure 312 (SEQ ID NO:312), Figure 314 (SEQ ID NO:314), Figure 316 (SEQ ID NO:316), Figure 318 (SEQ ID NO:318), Figure 320 (SEQ ID NO:320), Figure 322 (SEQ ID NO:322), Figure 324 (SEQ ID NO:324), Figure 326 (SEQ ID NO:326), Figure 328 (SEQ ID NO:328), Figure 330 (SEQ ID NO:330), Figure 332 (SEQ ID NO:332), Figure 334 (SEQ ID NO:334), Figure 336 (SEQ ID NO:336), Figure 338 (SEQ ID NO:338), Figure 340 (SEQ ID NO:340), Figure 342 (SEQ ID NO:342), Figure 344 (SEQ ID NO:344), Figure 346 (SEQ ID NO:346), Figure 348 (SEQ ID NO:348), Figure 350 (SEQ ID NO:350), Figure 352 (SEQ ID NO:352), Figure 354 (SEQ ID NO:354), Figure 356 (SEQ ID NO:356), Figure 358 (SEQ ID NO:358), Figure 360 (SEQ ID NO:360), Figure 362 (SEQ ID NO:362), Figure 364 (SEQ ID NO:364), Figure 366 (SEQ ID NO:366), Figure 368 (SEQ ID NO:368), Figure 370 (SEQ ID NO:370), Figure 372 (SEQ ID NO:372) and Figure 374 (SEQ ID NO:374).

2. An isolated nucleic acid molecule having at least 80% nucleic acid sequence identity to a nucleotide sequence selected from the group consisting of the nucleotide sequence shown in Figure 1 (SEQ ID NO:1), Figure 3 (SEQ ID NO:3), Figure 5 (SEQ ID NO:5), Figure 7 (SEQ ID NO:7), Figure 9 (SEQ ID NO:9), Figure 11 (SEQ ID NO:11), Figure 13 (SEQ ID NO:13), Figure 15 (SEQ ID NO:15), Figure 17 (SEQ ID NO:17), Figure 19 (SEQ ID NO:19), Figure 21 (SEQ ID NO:21), Figure 23 (SEQ ID NO:23), Figure 25 (SEQ ID NO:25), Figure 27 (SEQ ID NO:27), Figure 29 (SEQ ID NO:29), Figure 31 (SEQ ID NO:31), Figure 33 (SEQ ID NO:33), Figure 35 (SEQ ID NO:35), Figure 37 (SEQ ID NO:37), Figure 39 (SEQ ID NO:39), Figure 41 (SEQ ID NO:41), Figure 43 (SEQ ID NO:43), Figure 45 (SEQ ID NO:45), Figure 47 (SEQ ID NO:47), Figure 49 (SEQ ID NO:49), Figure 51 (SEQ ID NO:51), Figure 53 (SEQ ID NO:53), Figure 55 (SEQ ID NO:55), Figure 57 (SEQ ID NO:57), Figure 59 (SEQ ID NO:59), Figure 61 (SEQ ID NO:61), Figure 63 (SEQ ID NO:63), Figure 65 (SEQ ID NO:65), Figure 67 (SEQ ID NO:67), Figure 69 (SEQ ID NO:69), Figure 71 (SEQ ID NO:71), Figure 73 (SEQ ID NO:73), Figures 75A-75B (SEQ ID NO:75), Figure 77 (SEQ ID NO:77), Figure 79 (SEQ ID NO:79), Figure 81 (SEQ ID NO:81), Figure 83 (SEQ ID NO:83), Figure 85 (SEQ ID NO:85), Figure 87 (SEQ ID NO:87), Figure 89 (SEQ ID NO:89), Figure 91 (SEQ ID NO:91), Figure 93 (SEQ ID NO:93), Figure 95 (SEQ ID NO:95), Figure 97 (SEQ ID NO:97), Figure 99 (SEQ ID NO:99), Figure 101 (SEQ ID NO:101), Figure 103 (SEQ ID NO:103), Figure 105 (SEQ ID NO:105), Figure 107 (SEQ ID NO:107), Figure 109 (SEQ ID NO:109), Figure 111 (SEQ ID NO:111), Figure 113 (SEQ ID NO:113), Figure 115 (SEQ

5 ID NO:115), Figure 117 (SEQ ID NO:117), Figure 119 (SEQ ID NO:119), Figure 121 (SEQ ID NO:121), Figure
123 (SEQ ID NO:123), Figure 125 (SEQ ID NO:125), Figure 127 (SEQ ID NO:127), Figure 129 (SEQ ID
NO:129), Figure 131 (SEQ ID NO:131), Figure 133 (SEQ ID NO:133), Figure 135 (SEQ ID NO:135), Figure
137 (SEQ ID NO:137), Figure 139 (SEQ ID NO:139), Figure 141 (SEQ ID NO:141), Figure 143 (SEQ ID
NO:143), Figure 145 (SEQ ID NO:145), Figure 147 (SEQ ID NO:147), Figure 149 (SEQ ID NO:149), Figure
151 (SEQ ID NO:151), Figure 153 (SEQ ID NO:153), Figure 155 (SEQ ID NO:155), Figure 157 (SEQ ID
NO:157), Figure 159 (SEQ ID NO:159), Figure 161 (SEQ ID NO:161), Figure 163 (SEQ ID NO:163), Figure
165 (SEQ ID NO:165), Figure 167 (SEQ ID NO:167), Figure 169 (SEQ ID NO:169), Figure 171 (SEQ ID
NO:171), Figure 173 (SEQ ID NO:173), Figure 175 (SEQ ID NO:175), Figure 177 (SEQ ID NO:177), Figure
10 179 (SEQ ID NO:179), Figure 181 (SEQ ID NO:181), Figure 183 (SEQ ID NO:183), Figure 185 (SEQ ID
NO:185), Figure 187 (SEQ ID NO:187), Figure 189 (SEQ ID NO:189), Figure 191 (SEQ ID NO:191), Figure
193 (SEQ ID NO:193), Figure 195 (SEQ ID NO:195), Figure 197 (SEQ ID NO:197), Figure 199 (SEQ ID
NO:199), Figure 201 (SEQ ID NO:201), Figure 203 (SEQ ID NO:203), Figure 205 (SEQ ID NO:205), Figure
207 (SEQ ID NO:207), Figure 209 (SEQ ID NO:209), Figure 211 (SEQ ID NO:211), Figure 213 (SEQ ID
15 NO:213), Figure 215 (SEQ ID NO:215), Figure 217 (SEQ ID NO:217), Figure 219 (SEQ ID NO:219), Figure
221 (SEQ ID NO:221), Figure 223 (SEQ ID NO:223), Figure 225 (SEQ ID NO:225), Figure 227 (SEQ ID
NO:227), Figure 229 (SEQ ID NO:229), Figure 231 (SEQ ID NO:231), Figure 233 (SEQ ID NO:233), Figure
235 (SEQ ID NO:235), Figure 237 (SEQ ID NO:237), Figure 239 (SEQ ID NO:239), Figure 241 (SEQ ID
NO:241), Figure 243 (SEQ ID NO:243), Figure 245 (SEQ ID NO:245), Figure 247 (SEQ ID NO:247), Figure
20 249 (SEQ ID NO:249), Figure 251 (SEQ ID NO:251), Figure 253 (SEQ ID NO:253), Figure 255 (SEQ ID
NO:255), Figure 257 (SEQ ID NO:257), Figure 259 (SEQ ID NO:259), Figure 261 (SEQ ID NO:261), Figure
263 (SEQ ID NO:263), Figure 265 (SEQ ID NO:265), Figure 267 (SEQ ID NO:267), Figure 269 (SEQ ID
NO:269), Figure 271 (SEQ ID NO:271), Figure 273 (SEQ ID NO:273), Figure 275 (SEQ ID NO:275), Figure
277 (SEQ ID NO:277), Figure 279 (SEQ ID NO:279), Figure 281 (SEQ ID NO:281), Figure 283 (SEQ ID
25 NO:283), Figure 285 (SEQ ID NO:285), Figure 287 (SEQ ID NO:287), Figures 289A-289B (SEQ ID NO:289),
Figure 291 (SEQ ID NO:291), Figure 293 (SEQ ID NO:293), Figure 295 (SEQ ID NO:295), Figure 297 (SEQ
ID NO:297), Figure 299 (SEQ ID NO:299), Figure 301 (SEQ ID NO:301), Figure 303 (SEQ ID NO:303), Figure
305 (SEQ ID NO:305), Figure 307 (SEQ ID NO:307), Figure 309 (SEQ ID NO:309), Figures 311A-311B (SEQ
ID NO:311), Figure 313 (SEQ ID NO:313), Figure 315 (SEQ ID NO:315), Figure 317 (SEQ ID NO:317), Figure
30 319 (SEQ ID NO:319), Figure 321 (SEQ ID NO:321), Figure 323 (SEQ ID NO:323), Figure 325 (SEQ ID
NO:325), Figure 327 (SEQ ID NO:327), Figure 329 (SEQ ID NO:329), Figure 331 (SEQ ID NO:331), Figure
333 (SEQ ID NO:333), Figure 335 (SEQ ID NO:335), Figure 337 (SEQ ID NO:337), Figure 339 (SEQ ID
NO:339), Figure 341 (SEQ ID NO:341), Figure 343 (SEQ ID NO:343), Figure 345 (SEQ ID NO:345), Figure
347 (SEQ ID NO:347), Figure 349 (SEQ ID NO:349), Figures 351A-351B (SEQ ID NO:351), Figure 353 (SEQ
35 ID NO:353), Figure 355 (SEQ ID NO:355), Figure 357 (SEQ ID NO:357), Figure 359 (SEQ ID NO:359), Figure
361 (SEQ ID NO:361), Figure 363 (SEQ ID NO:363), Figure 365 (SEQ ID NO:365), Figure 367 (SEQ ID
NO:367), Figure 369 (SEQ ID NO:369), Figure 371 (SEQ ID NO:371) and Figure 373 (SEQ ID NO:373).

3. An isolated nucleic acid molecule having at least 80% nucleic acid sequence identity to a nucleotide sequence selected from the group consisting of the full-length coding sequence of the nucleotide sequence shown in Figure 1 (SEQ ID NO:1), Figure 3 (SEQ ID NO:3), Figure 5 (SEQ ID NO:5), Figure 7 (SEQ ID NO:7), Figure 9 (SEQ ID NO:9), Figure 11 (SEQ ID NO:11), Figure 13 (SEQ ID NO:13), Figure 15 (SEQ ID NO:15), Figure 17 (SEQ ID NO:17), Figure 19 (SEQ ID NO:19), Figure 21 (SEQ ID NO:21), Figure 23 (SEQ ID NO:23), Figure 25 (SEQ ID NO:25), Figure 27 (SEQ ID NO:27), Figure 29 (SEQ ID NO:29), Figure 31 (SEQ ID NO:31), Figure 33 (SEQ ID NO:33), Figure 35 (SEQ ID NO:35), Figure 37 (SEQ ID NO:37), Figure 39 (SEQ ID NO:39), Figure 41 (SEQ ID NO:41), Figure 43 (SEQ ID NO:43), Figure 45 (SEQ ID NO:45), Figure 47 (SEQ ID NO:47), Figure 49 (SEQ ID NO:49), Figure 51 (SEQ ID NO:51), Figure 53 (SEQ ID NO:53), Figure 55 (SEQ ID NO:55), Figure 57 (SEQ ID NO:57), Figure 59 (SEQ ID NO:59), Figure 61 (SEQ ID NO:61), Figure 63 (SEQ ID NO:63), Figure 65 (SEQ ID NO:65), Figure 67 (SEQ ID NO:67), Figure 69 (SEQ ID NO:69), Figure 71 (SEQ ID NO:71), Figure 73 (SEQ ID NO:73), Figures 75A-75B (SEQ ID NO:75), Figure 77 (SEQ ID NO:77), Figure 79 (SEQ ID NO:79), Figure 81 (SEQ ID NO:81), Figure 83 (SEQ ID NO:83), Figure 85 (SEQ ID NO:85), Figure 87 (SEQ ID NO:87), Figure 89 (SEQ ID NO:89), Figure 91 (SEQ ID NO:91), Figure 93 (SEQ ID NO:93), Figure 95 (SEQ ID NO:95), Figure 97 (SEQ ID NO:97), Figure 99 (SEQ ID NO:99), Figure 101 (SEQ ID NO:101), Figure 103 (SEQ ID NO:103), Figure 105 (SEQ ID NO:105), Figure 107 (SEQ ID NO:107), Figure 109 (SEQ ID NO:109), Figure 111 (SEQ ID NO:111), Figure 113 (SEQ ID NO:113), Figure 115 (SEQ ID NO:115), Figure 117 (SEQ ID NO:117), Figure 119 (SEQ ID NO:119), Figure 121 (SEQ ID NO:121), Figure 123 (SEQ ID NO:123), Figure 125 (SEQ ID NO:125), Figure 127 (SEQ ID NO:127), Figure 129 (SEQ ID NO:129), Figure 131 (SEQ ID NO:131), Figure 133 (SEQ ID NO:133), Figure 135 (SEQ ID NO:135), Figure 137 (SEQ ID NO:137), Figure 139 (SEQ ID NO:139), Figure 141 (SEQ ID NO:141), Figure 143 (SEQ ID NO:143), Figure 145 (SEQ ID NO:145), Figure 147 (SEQ ID NO:147), Figure 149 (SEQ ID NO:149), Figure 151 (SEQ ID NO:151), Figure 153 (SEQ ID NO:153), Figure 155 (SEQ ID NO:155), Figure 157 (SEQ ID NO:157), Figure 159 (SEQ ID NO:159), Figure 161 (SEQ ID NO:161), Figure 163 (SEQ ID NO:163), Figure 165 (SEQ ID NO:165), Figure 167 (SEQ ID NO:167), Figure 169 (SEQ ID NO:169), Figure 171 (SEQ ID NO:171), Figure 173 (SEQ ID NO:173), Figure 175 (SEQ ID NO:175), Figure 177 (SEQ ID NO:177), Figure 179 (SEQ ID NO:179), Figure 181 (SEQ ID NO:181), Figure 183 (SEQ ID NO:183), Figure 185 (SEQ ID NO:185), Figure 187 (SEQ ID NO:187), Figure 189 (SEQ ID NO:189), Figure 191 (SEQ ID NO:191), Figure 193 (SEQ ID NO:193), Figure 195 (SEQ ID NO:195), Figure 197 (SEQ ID NO:197), Figure 199 (SEQ ID NO:199), Figure 201 (SEQ ID NO:201), Figure 203 (SEQ ID NO:203), Figure 205 (SEQ ID NO:205), Figure 207 (SEQ ID NO:207), Figure 209 (SEQ ID NO:209), Figure 211 (SEQ ID NO:211), Figure 213 (SEQ ID NO:213), Figure 215 (SEQ ID NO:215), Figure 217 (SEQ ID NO:217), Figure 219 (SEQ ID NO:219), Figure 221 (SEQ ID NO:221), Figure 223 (SEQ ID NO:223), Figure 225 (SEQ ID NO:225), Figure 227 (SEQ ID NO:227), Figure 229 (SEQ ID NO:229), Figure 231 (SEQ ID NO:231), Figure 233 (SEQ ID NO:233), Figure 235 (SEQ ID NO:235), Figure 237 (SEQ ID NO:237), Figure 239 (SEQ ID NO:239), Figure 241 (SEQ ID NO:241), Figure 243 (SEQ ID NO:243), Figure 245 (SEQ ID NO:245), Figure 247 (SEQ ID NO:247), Figure 249 (SEQ ID NO:249), Figure 251 (SEQ ID NO:251), Figure

253 (SEQ ID NO:253), Figure 255 (SEQ ID NO:255), Figure 257 (SEQ ID NO:257), Figure 259 (SEQ ID NO:259), Figure 261 (SEQ ID NO:261), Figure 263 (SEQ ID NO:263), Figure 265 (SEQ ID NO:265), Figure 267 (SEQ ID NO:267), Figure 269 (SEQ ID NO:269), Figure 271 (SEQ ID NO:271), Figure 273 (SEQ ID NO:273), Figure 275 (SEQ ID NO:275), Figure 277 (SEQ ID NO:277), Figure 279 (SEQ ID NO:279), Figure 281 (SEQ ID NO:281), Figure 283 (SEQ ID NO:283), Figure 285 (SEQ ID NO:285), Figure 287 (SEQ ID NO:287), Figures 289A-289B (SEQ ID NO:289), Figure 291 (SEQ ID NO:291), Figure 293 (SEQ ID NO:293), Figure 295 (SEQ ID NO:295), Figure 297 (SEQ ID NO:297), Figure 299 (SEQ ID NO:299), Figure 301 (SEQ ID NO:301), Figure 303 (SEQ ID NO:303), Figure 305 (SEQ ID NO:305), Figure 307 (SEQ ID NO:307), Figure 309 (SEQ ID NO:309), Figures 311A-311B (SEQ ID NO:311), Figure 313 (SEQ ID NO:313), Figure 315 (SEQ ID NO:315), Figure 317 (SEQ ID NO:317), Figure 319 (SEQ ID NO:319), Figure 321 (SEQ ID NO:321), Figure 323 (SEQ ID NO:323), Figure 325 (SEQ ID NO:325), Figure 327 (SEQ ID NO:327), Figure 329 (SEQ ID NO:329), Figure 331 (SEQ ID NO:331), Figure 333 (SEQ ID NO:333), Figure 335 (SEQ ID NO:335), Figure 337 (SEQ ID NO:337), Figure 339 (SEQ ID NO:339), Figure 341 (SEQ ID NO:341), Figure 343 (SEQ ID NO:343), Figure 345 (SEQ ID NO:345), Figure 347 (SEQ ID NO:347), Figure 349 (SEQ ID NO:349), Figures 351A-351B (SEQ ID NO:351), Figure 353 (SEQ ID NO:353), Figure 355 (SEQ ID NO:355), Figure 357 (SEQ ID NO:357), Figure 359 (SEQ ID NO:359), Figure 361 (SEQ ID NO:361), Figure 363 (SEQ ID NO:363), Figure 365 (SEQ ID NO:365), Figure 367 (SEQ ID NO:367), Figure 369 (SEQ ID NO:369), Figure 371 (SEQ ID NO:371) and Figure 373 (SEQ ID NO:373).

4. An isolated nucleic acid molecule having at least 80% nucleic acid sequence identity to the full-length coding sequence of the DNA deposited under any ATCC accession number shown in Table 7.

5. A vector comprising the nucleic acid of Claim 1.

6. A host cell comprising the vector of Claim 5.

7. The host cell of Claim 6, wherein said cell is a CHO cell.

8. The host cell of Claim 6, wherein said cell is an *E. coli*.

9. The host cell of Claim 6, wherein said cell is a yeast cell.

10. A process for producing a PRO polypeptide comprising culturing the host cell of Claim 6 under conditions suitable for expression of said PRO polypeptide and recovering said PRO polypeptide from the cell culture.

11. An isolated polypeptide having at least 80% amino acid sequence identity to an amino acid sequence selected from the group consisting of the amino acid sequence shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:4), Figure 6 (SEQ ID NO:6), Figure 8 (SEQ ID NO:8), Figure 10 (SEQ ID NO:10), Figure 12 (SEQ ID NO:12), Figure 14 (SEQ ID NO:14), Figure 16 (SEQ ID NO:16), Figure 18 (SEQ ID NO:18), Figure 20 (SEQ ID NO:20), Figure 22 (SEQ ID NO:22), Figure 24 (SEQ ID NO:24), Figure 26 (SEQ ID NO:26), Figure 28 (SEQ ID NO:28), Figure 30 (SEQ ID NO:30), Figure 32 (SEQ ID NO:32), Figure 34 (SEQ ID NO:34), Figure 36 (SEQ ID NO:36), Figure 38 (SEQ ID NO:38), Figure 40 (SEQ ID NO:40), Figure 42 (SEQ ID NO:42), Figure 44 (SEQ ID NO:44), Figure 46 (SEQ ID NO:46), Figure 48 (SEQ ID NO:48), Figure 50 (SEQ ID NO:50), Figure 52 (SEQ ID NO:52), Figure 54 (SEQ ID NO:54), Figure 56 (SEQ ID NO:56), Figure 58 (SEQ ID NO:58), Figure 60 (SEQ ID NO:60), Figure 62 (SEQ ID NO:62), Figure 64 (SEQ ID NO:64), Figure 66 (SEQ ID NO:66), Figure 68 (SEQ ID NO:68), Figure 70 (SEQ ID NO:70), Figure 72 (SEQ ID NO:72), Figure 74 (SEQ ID NO:74), Figure 76 (SEQ ID NO:76), Figure 78 (SEQ ID NO:78), Figure 80 (SEQ ID NO:80), Figure 82 (SEQ ID NO:82), Figure 84 (SEQ ID NO:84), Figure 86 (SEQ ID NO:86), Figure 88 (SEQ ID NO:88), Figure 90 (SEQ ID NO:90), Figure 92 (SEQ ID NO:92), Figure 94 (SEQ ID NO:94), Figure 96 (SEQ ID NO:96), Figure 98 (SEQ ID NO:98), Figure 100 (SEQ ID NO:100), Figure 102 (SEQ ID NO:102), Figure 104 (SEQ ID NO:104), Figure 106 (SEQ ID NO:106), Figure 108 (SEQ ID NO:108), Figure 110 (SEQ ID NO:110), Figure 112 (SEQ ID NO:112), Figure 114 (SEQ ID NO:114), Figure 116 (SEQ ID NO:116), Figure 118 (SEQ ID NO:118), Figure 120 (SEQ ID NO:120), Figure 122 (SEQ ID NO:122), Figure 124 (SEQ ID NO:124), Figure 126 (SEQ ID NO:126), Figure 128 (SEQ ID NO:128), Figure 130 (SEQ ID NO:130), Figure 132 (SEQ ID NO:132), Figure 134 (SEQ ID NO:134), Figure 136 (SEQ ID NO:136), Figure 138 (SEQ ID NO:138), Figure 140 (SEQ ID NO:140), Figure 142 (SEQ ID NO:142), Figure 144 (SEQ ID NO:144), Figure 146 (SEQ ID NO:146), Figure 148 (SEQ ID NO:148), Figure 150 (SEQ ID NO:150), Figure 152 (SEQ ID NO:152), Figure 154 (SEQ ID NO:154), Figure 156 (SEQ ID NO:156), Figure 158 (SEQ ID NO:158), Figure 160 (SEQ ID NO:160), Figure 162 (SEQ ID NO:162), Figure 164 (SEQ ID NO:164), Figure 166 (SEQ ID NO:166), Figure 168 (SEQ ID NO:168), Figure 170 (SEQ ID NO:170), Figure 172 (SEQ ID NO:172), Figure 174 (SEQ ID NO:174), Figure 176 (SEQ ID NO:176), Figure 178 (SEQ ID NO:178), Figure 180 (SEQ ID NO:180), Figure 182 (SEQ ID NO:182), Figure 184 (SEQ ID NO:184), Figure 186 (SEQ ID NO:186), Figure 188 (SEQ ID NO:188), Figure 190 (SEQ ID NO:190), Figure 192 (SEQ ID NO:192), Figure 194 (SEQ ID NO:194), Figure 196 (SEQ ID NO:196), Figure 198 (SEQ ID NO:198), Figure 200 (SEQ ID NO:200), Figure 202 (SEQ ID NO:202), Figure 204 (SEQ ID NO:204), Figure 206 (SEQ ID NO:206), Figure 208 (SEQ ID NO:208), Figure 210 (SEQ ID NO:210), Figure 212 (SEQ ID NO:212), Figure 214 (SEQ ID NO:214), Figure 216 (SEQ ID NO:216), Figure 218 (SEQ ID NO:218), Figure 220 (SEQ ID NO:220), Figure 222 (SEQ ID NO:222), Figure 224 (SEQ ID NO:224), Figure 226 (SEQ ID NO:226), Figure 228 (SEQ ID NO:228), Figure 230 (SEQ ID NO:230), Figure 232 (SEQ ID NO:232), Figure 234 (SEQ ID NO:234), Figure 236 (SEQ ID NO:236), Figure 238 (SEQ ID NO:238), Figure 240 (SEQ ID NO:240), Figure 242 (SEQ ID NO:242), Figure 244 (SEQ ID NO:244), Figure 246 (SEQ ID NO:246), Figure 248 (SEQ ID NO:248), Figure 250 (SEQ ID NO:250), Figure 252 (SEQ ID NO:252), Figure 254 (SEQ ID NO:254), Figure 256 (SEQ ID

NO:256), Figure 258 (SEQ ID NO:258), Figure 260 (SEQ ID NO:260), Figure 262 (SEQ ID NO:262), Figure 264 (SEQ ID NO:264), Figure 266 (SEQ ID NO:266), Figure 268 (SEQ ID NO:268), Figure 270 (SEQ ID NO:270), Figure 272 (SEQ ID NO:272), Figure 274 (SEQ ID NO:274), Figure 276 (SEQ ID NO:276), Figure 278 (SEQ ID NO:278), Figure 280 (SEQ ID NO:280), Figure 282 (SEQ ID NO:282), Figure 284 (SEQ ID NO:284), Figure 286 (SEQ ID NO:286), Figure 288 (SEQ ID NO:288), Figure 290 (SEQ ID NO:290), Figure 292 (SEQ ID NO:292), Figure 294 (SEQ ID NO:294), Figure 296 (SEQ ID NO:296), Figure 298 (SEQ ID NO:298), Figure 300 (SEQ ID NO:300), Figure 302 (SEQ ID NO:302), Figure 304 (SEQ ID NO:304), Figure 306 (SEQ ID NO:306), Figure 308 (SEQ ID NO:308), Figure 310 (SEQ ID NO:310), Figure 312 (SEQ ID NO:312), Figure 314 (SEQ ID NO:314), Figure 316 (SEQ ID NO:316), Figure 318 (SEQ ID NO:318), Figure 320 (SEQ ID NO:320), Figure 322 (SEQ ID NO:322), Figure 324 (SEQ ID NO:324), Figure 326 (SEQ ID NO:326), Figure 328 (SEQ ID NO:328), Figure 330 (SEQ ID NO:330), Figure 332 (SEQ ID NO:332), Figure 334 (SEQ ID NO:334), Figure 336 (SEQ ID NO:336), Figure 338 (SEQ ID NO:338), Figure 340 (SEQ ID NO:340), Figure 342 (SEQ ID NO:342), Figure 344 (SEQ ID NO:344), Figure 346 (SEQ ID NO:346), Figure 348 (SEQ ID NO:348), Figure 350 (SEQ ID NO:350), Figure 352 (SEQ ID NO:352), Figure 354 (SEQ ID NO:354), Figure 356 (SEQ ID NO:356), Figure 358 (SEQ ID NO:358), Figure 360 (SEQ ID NO:360), Figure 362 (SEQ ID NO:362), Figure 364 (SEQ ID NO:364), Figure 366 (SEQ ID NO:366), Figure 368 (SEQ ID NO:368), Figure 370 (SEQ ID NO:370), Figure 372 (SEQ ID NO:372) and Figure 374 (SEQ ID NO:374).

12. An isolated polypeptide having at least 80% amino acid sequence identity to an amino acid sequence encoded by the full-length coding sequence of the DNA deposited under any ATCC accession number shown in Table 7.

13. A chimeric molecule comprising a polypeptide according to Claim 11 fused to a heterologous amino acid sequence.

14. The chimeric molecule of Claim 13, wherein said heterologous amino acid sequence is an epitope tag sequence.

15. The chimeric molecule of Claim 13, wherein said heterologous amino acid sequence is a Fc region of an immunoglobulin.

16. An antibody which specifically binds to a polypeptide according to Claim 11.

17. The antibody of Claim 16, wherein said antibody is a monoclonal antibody, a humanized antibody or a single-chain antibody.

18. An isolated nucleic acid molecule having at least 80% nucleic acid sequence identity to:

(a) a nucleotide sequence encoding the polypeptide shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:4), Figure 6 (SEQ ID NO:6), Figure 8 (SEQ ID NO:8), Figure 10 (SEQ ID NO:10), Figure 12 (SEQ ID NO:12), Figure 14 (SEQ ID NO:14), Figure 16 (SEQ ID NO:16), Figure 18 (SEQ ID NO:18), Figure 20 (SEQ ID NO:20), Figure 22 (SEQ ID NO:22), Figure 24 (SEQ ID NO:24), Figure 26 (SEQ ID NO:26),
5 Figure 28 (SEQ ID NO:28), Figure 30 (SEQ ID NO:30), Figure 32 (SEQ ID NO:32), Figure 34 (SEQ ID NO:34), Figure 36 (SEQ ID NO:36), Figure 38 (SEQ ID NO:38), Figure 40 (SEQ ID NO:40), Figure 42 (SEQ ID NO:42), Figure 44 (SEQ ID NO:44), Figure 46 (SEQ ID NO:46), Figure 48 (SEQ ID NO:48), Figure 50 (SEQ ID NO:50), Figure 52 (SEQ ID NO:52), Figure 54 (SEQ ID NO:54), Figure 56 (SEQ ID NO:56), Figure 58 (SEQ ID NO:58), Figure 60 (SEQ ID NO:60), Figure 62 (SEQ ID NO:62), Figure 64 (SEQ ID NO:64),
10 Figure 66 (SEQ ID NO:66), Figure 68 (SEQ ID NO:68), Figure 70 (SEQ ID NO:70), Figure 72 (SEQ ID NO:72), Figure 74 (SEQ ID NO:74), Figure 76 (SEQ ID NO:76), Figure 78 (SEQ ID NO:78), Figure 80 (SEQ ID NO:80), Figure 82 (SEQ ID NO:82), Figure 84 (SEQ ID NO:84), Figure 86 (SEQ ID NO:86), Figure 88 (SEQ ID NO:88), Figure 90 (SEQ ID NO:90), Figure 92 (SEQ ID NO:92), Figure 94 (SEQ ID NO:94), Figure 96 (SEQ ID NO:96), Figure 98 (SEQ ID NO:98), Figure 100 (SEQ ID NO:100), Figure 102 (SEQ ID NO:102),
15 Figure 104 (SEQ ID NO:104), Figure 106 (SEQ ID NO:106), Figure 108 (SEQ ID NO:108), Figure 110 (SEQ ID NO:110), Figure 112 (SEQ ID NO:112), Figure 114 (SEQ ID NO:114), Figure 116 (SEQ ID NO:116), Figure 118 (SEQ ID NO:118), Figure 120 (SEQ ID NO:120), Figure 122 (SEQ ID NO:122), Figure 124 (SEQ ID NO:124), Figure 126 (SEQ ID NO:126), Figure 128 (SEQ ID NO:128), Figure 130 (SEQ ID NO:130), Figure 132 (SEQ ID NO:132), Figure 134 (SEQ ID NO:134), Figure 136 (SEQ ID NO:136), Figure 138 (SEQ ID NO:138), Figure 140 (SEQ ID NO:140), Figure 142 (SEQ ID NO:142), Figure 144 (SEQ ID NO:144), Figure 146 (SEQ ID NO:146), Figure 148 (SEQ ID NO:148), Figure 150 (SEQ ID NO:150), Figure 152 (SEQ ID NO:152), Figure 154 (SEQ ID NO:154), Figure 156 (SEQ ID NO:156), Figure 158 (SEQ ID NO:158), Figure 160 (SEQ ID NO:160), Figure 162 (SEQ ID NO:162), Figure 164 (SEQ ID NO:164), Figure 166 (SEQ ID NO:166), Figure 168 (SEQ ID NO:168), Figure 170 (SEQ ID NO:170), Figure 172 (SEQ ID NO:172), Figure 174 (SEQ ID NO:174), Figure 176 (SEQ ID NO:176), Figure 178 (SEQ ID NO:178), Figure 180 (SEQ ID NO:180), Figure 182 (SEQ ID NO:182), Figure 184 (SEQ ID NO:184), Figure 186 (SEQ ID NO:186), Figure 188 (SEQ ID NO:188), Figure 190 (SEQ ID NO:190), Figure 192 (SEQ ID NO:192), Figure 194 (SEQ ID NO:194), Figure 196 (SEQ ID NO:196), Figure 198 (SEQ ID NO:198), Figure 200 (SEQ ID NO:200), Figure 202 (SEQ ID NO:202), Figure 204 (SEQ ID NO:204), Figure 206 (SEQ ID NO:206), Figure 208 (SEQ ID NO:208), Figure 210 (SEQ ID NO:210), Figure 212 (SEQ ID NO:212), Figure 214 (SEQ ID NO:214), Figure 216 (SEQ ID NO:216), Figure 218 (SEQ ID NO:218), Figure 220 (SEQ ID NO:220), Figure 222 (SEQ ID NO:222), Figure 224 (SEQ ID NO:224), Figure 226 (SEQ ID NO:226), Figure 228 (SEQ ID NO:228), Figure 230 (SEQ ID NO:230), Figure 232 (SEQ ID NO:232), Figure 234 (SEQ ID NO:234), Figure 236 (SEQ ID NO:236), Figure 238 (SEQ ID NO:238), Figure 240 (SEQ ID NO:240), Figure 242 (SEQ ID NO:242), Figure 244 (SEQ ID NO:244), Figure 246 (SEQ ID NO:246), Figure 248 (SEQ ID NO:248), Figure 250 (SEQ ID NO:250), Figure 252 (SEQ ID NO:252), Figure 254 (SEQ ID NO:254), Figure 256 (SEQ ID NO:256), Figure 258 (SEQ ID NO:258), Figure 260 (SEQ ID NO:260), Figure 262 (SEQ ID NO:262), Figure 264 (SEQ ID NO:264)

NO:264), Figure 266 (SEQ ID NO:266), Figure 268 (SEQ ID NO:268), Figure 270 (SEQ ID NO:270), Figure 272 (SEQ ID NO:272), Figure 274 (SEQ ID NO:274), Figure 276 (SEQ ID NO:276), Figure 278 (SEQ ID NO:278), Figure 280 (SEQ ID NO:280), Figure 282 (SEQ ID NO:282), Figure 284 (SEQ ID NO:284), Figure 286 (SEQ ID NO:286), Figure 288 (SEQ ID NO:288), Figure 290 (SEQ ID NO:290), Figure 292 (SEQ ID NO:292), Figure 294 (SEQ ID NO:294), Figure 296 (SEQ ID NO:296), Figure 298 (SEQ ID NO:298), Figure 300 (SEQ ID NO:300), Figure 302 (SEQ ID NO:302), Figure 304 (SEQ ID NO:304), Figure 306 (SEQ ID NO:306), Figure 308 (SEQ ID NO:308), Figure 310 (SEQ ID NO:310), Figure 312 (SEQ ID NO:312), Figure 314 (SEQ ID NO:314), Figure 316 (SEQ ID NO:316), Figure 318 (SEQ ID NO:318), Figure 320 (SEQ ID NO:320), Figure 322 (SEQ ID NO:322), Figure 324 (SEQ ID NO:324), Figure 326 (SEQ ID NO:326), Figure 328 (SEQ ID NO:328), Figure 330 (SEQ ID NO:330), Figure 332 (SEQ ID NO:332), Figure 334 (SEQ ID NO:334), Figure 336 (SEQ ID NO:336), Figure 338 (SEQ ID NO:338), Figure 340 (SEQ ID NO:340), Figure 342 (SEQ ID NO:342), Figure 344 (SEQ ID NO:344), Figure 346 (SEQ ID NO:346), Figure 348 (SEQ ID NO:348), Figure 350 (SEQ ID NO:350), Figure 352 (SEQ ID NO:352), Figure 354 (SEQ ID NO:354), Figure 356 (SEQ ID NO:356), Figure 358 (SEQ ID NO:358), Figure 360 (SEQ ID NO:360), Figure 362 (SEQ ID NO:362), Figure 364 (SEQ ID NO:364), Figure 366 (SEQ ID NO:366), Figure 368 (SEQ ID NO:368), Figure 370 (SEQ ID NO:370), Figure 372 (SEQ ID NO:372) or Figure 374 (SEQ ID NO:374), lacking its associated signal peptide;

(b) a nucleotide sequence encoding an extracellular domain of the polypeptide shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:4), Figure 6 (SEQ ID NO:6), Figure 8 (SEQ ID NO:8), Figure 10 (SEQ ID NO:10), Figure 12 (SEQ ID NO:12), Figure 14 (SEQ ID NO:14), Figure 16 (SEQ ID NO:16), Figure 18 (SEQ ID NO:18), Figure 20 (SEQ ID NO:20), Figure 22 (SEQ ID NO:22), Figure 24 (SEQ ID NO:24), Figure 26 (SEQ ID NO:26), Figure 28 (SEQ ID NO:28), Figure 30 (SEQ ID NO:30), Figure 32 (SEQ ID NO:32), Figure 34 (SEQ ID NO:34), Figure 36 (SEQ ID NO:36), Figure 38 (SEQ ID NO:38), Figure 40 (SEQ ID NO:40), Figure 42 (SEQ ID NO:42), Figure 44 (SEQ ID NO:44), Figure 46 (SEQ ID NO:46), Figure 48 (SEQ ID NO:48), Figure 50 (SEQ ID NO:50), Figure 52 (SEQ ID NO:52), Figure 54 (SEQ ID NO:54), Figure 56 (SEQ ID NO:56), Figure 58 (SEQ ID NO:58), Figure 60 (SEQ ID NO:60), Figure 62 (SEQ ID NO:62), Figure 64 (SEQ ID NO:64), Figure 66 (SEQ ID NO:66), Figure 68 (SEQ ID NO:68), Figure 70 (SEQ ID NO:70), Figure 72 (SEQ ID NO:72), Figure 74 (SEQ ID NO:74), Figure 76 (SEQ ID NO:76), Figure 78 (SEQ ID NO:78), Figure 80 (SEQ ID NO:80), Figure 82 (SEQ ID NO:82), Figure 84 (SEQ ID NO:84), Figure 86 (SEQ ID NO:86), Figure 88 (SEQ ID NO:88), Figure 90 (SEQ ID NO:90), Figure 92 (SEQ ID NO:92), Figure 94 (SEQ ID NO:94), Figure 96 (SEQ ID NO:96), Figure 98 (SEQ ID NO:98), Figure 100 (SEQ ID NO:100), Figure 102 (SEQ ID NO:102), Figure 104 (SEQ ID NO:104), Figure 106 (SEQ ID NO:106), Figure 108 (SEQ ID NO:108), Figure 110 (SEQ ID NO:110), Figure 112 (SEQ ID NO:112), Figure 114 (SEQ ID NO:114), Figure 116 (SEQ ID NO:116), Figure 118 (SEQ ID NO:118), Figure 120 (SEQ ID NO:120), Figure 122 (SEQ ID NO:122), Figure 124 (SEQ ID NO:124), Figure 126 (SEQ ID NO:126), Figure 128 (SEQ ID NO:128), Figure 130 (SEQ ID NO:130), Figure 132 (SEQ ID NO:132), Figure 134 (SEQ ID NO:134), Figure 136 (SEQ ID NO:136), Figure 138 (SEQ ID NO:138), Figure 140 (SEQ ID NO:140), Figure 142 (SEQ ID NO:142), Figure

144 (SEQ ID NO:144), Figure 146 (SEQ ID NO:146), Figure 148 (SEQ ID NO:148), Figure 150 (SEQ ID NO:150), Figure 152 (SEQ ID NO:152), Figure 154 (SEQ ID NO:154), Figure 156 (SEQ ID NO:156), Figure 158 (SEQ ID NO:158), Figure 160 (SEQ ID NO:160), Figure 162 (SEQ ID NO:162), Figure 164 (SEQ ID NO:164), Figure 166 (SEQ ID NO:166), Figure 168 (SEQ ID NO:168), Figure 170 (SEQ ID NO:170), Figure 172 (SEQ ID NO:172), Figure 174 (SEQ ID NO:174), Figure 176 (SEQ ID NO:176), Figure 178 (SEQ ID NO:178), Figure 180 (SEQ ID NO:180), Figure 182 (SEQ ID NO:182), Figure 184 (SEQ ID NO:184), Figure 186 (SEQ ID NO:186), Figure 188 (SEQ ID NO:188), Figure 190 (SEQ ID NO:190), Figure 192 (SEQ ID NO:192), Figure 194 (SEQ ID NO:194), Figure 196 (SEQ ID NO:196), Figure 198 (SEQ ID NO:198), Figure 200 (SEQ ID NO:200), Figure 202 (SEQ ID NO:202), Figure 204 (SEQ ID NO:204), Figure 206 (SEQ ID NO:206), Figure 208 (SEQ ID NO:208), Figure 210 (SEQ ID NO:210), Figure 212 (SEQ ID NO:212), Figure 214 (SEQ ID NO:214), Figure 216 (SEQ ID NO:216), Figure 218 (SEQ ID NO:218), Figure 220 (SEQ ID NO:220), Figure 222 (SEQ ID NO:222), Figure 224 (SEQ ID NO:224), Figure 226 (SEQ ID NO:226), Figure 228 (SEQ ID NO:228), Figure 230 (SEQ ID NO:230), Figure 232 (SEQ ID NO:232), Figure 234 (SEQ ID NO:234), Figure 236 (SEQ ID NO:236), Figure 238 (SEQ ID NO:238), Figure 240 (SEQ ID NO:240), Figure 242 (SEQ ID NO:242), Figure 244 (SEQ ID NO:244), Figure 246 (SEQ ID NO:246), Figure 248 (SEQ ID NO:248), Figure 250 (SEQ ID NO:250), Figure 252 (SEQ ID NO:252), Figure 254 (SEQ ID NO:254), Figure 256 (SEQ ID NO:256), Figure 258 (SEQ ID NO:258), Figure 260 (SEQ ID NO:260), Figure 262 (SEQ ID NO:262), Figure 264 (SEQ ID NO:264), Figure 266 (SEQ ID NO:266), Figure 268 (SEQ ID NO:268), Figure 270 (SEQ ID NO:270), Figure 272 (SEQ ID NO:272), Figure 274 (SEQ ID NO:274), Figure 276 (SEQ ID NO:276), Figure 278 (SEQ ID NO:278), Figure 280 (SEQ ID NO:280), Figure 282 (SEQ ID NO:282), Figure 284 (SEQ ID NO:284), Figure 286 (SEQ ID NO:286), Figure 288 (SEQ ID NO:288), Figure 290 (SEQ ID NO:290), Figure 292 (SEQ ID NO:292), Figure 294 (SEQ ID NO:294), Figure 296 (SEQ ID NO:296), Figure 298 (SEQ ID NO:298), Figure 300 (SEQ ID NO:300), Figure 302 (SEQ ID NO:302), Figure 304 (SEQ ID NO:304), Figure 306 (SEQ ID NO:306), Figure 308 (SEQ ID NO:308), Figure 310 (SEQ ID NO:310), Figure 312 (SEQ ID NO:312), Figure 314 (SEQ ID NO:314), Figure 316 (SEQ ID NO:316), Figure 318 (SEQ ID NO:318), Figure 320 (SEQ ID NO:320), Figure 322 (SEQ ID NO:322), Figure 324 (SEQ ID NO:324), Figure 326 (SEQ ID NO:326), Figure 328 (SEQ ID NO:328), Figure 330 (SEQ ID NO:330), Figure 332 (SEQ ID NO:332), Figure 334 (SEQ ID NO:334), Figure 336 (SEQ ID NO:336), Figure 338 (SEQ ID NO:338), Figure 340 (SEQ ID NO:340), Figure 342 (SEQ ID NO:342), Figure 344 (SEQ ID NO:344), Figure 346 (SEQ ID NO:346), Figure 348 (SEQ ID NO:348), Figure 350 (SEQ ID NO:350), Figure 352 (SEQ ID NO:352), Figure 354 (SEQ ID NO:354), Figure 356 (SEQ ID NO:356), Figure 358 (SEQ ID NO:358), Figure 360 (SEQ ID NO:360), Figure 362 (SEQ ID NO:362), Figure 364 (SEQ ID NO:364), Figure 366 (SEQ ID NO:366), Figure 368 (SEQ ID NO:368), Figure 370 (SEQ ID NO:370), Figure 372 (SEQ ID NO:372) or Figure 374 (SEQ ID NO:374), with its associated signal peptide; or

(c) a nucleotide sequence encoding an extracellular domain of the polypeptide shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:4), Figure 6 (SEQ ID NO:6), Figure 8 (SEQ ID NO:8), Figure 10 (SEQ ID NO:10), Figure 12 (SEQ ID NO:12), Figure 14 (SEQ ID NO:14), Figure 16 (SEQ ID NO:16), Figure 18

(SEQ ID NO:18), Figure 20 (SEQ ID NO:20), Figure 22 (SEQ ID NO:22), Figure 24 (SEQ ID NO:24), Figure 26 (SEQ ID NO:26), Figure 28 (SEQ ID NO:28), Figure 30 (SEQ ID NO:30), Figure 32 (SEQ ID NO:32), Figure 34 (SEQ ID NO:34), Figure 36 (SEQ ID NO:36), Figure 38 (SEQ ID NO:38), Figure 40 (SEQ ID NO:40), Figure 42 (SEQ ID NO:42), Figure 44 (SEQ ID NO:44), Figure 46 (SEQ ID NO:46), Figure 48 (SEQ ID NO:48), Figure 50 (SEQ ID NO:50), Figure 52 (SEQ ID NO:52), Figure 54 (SEQ ID NO:54), Figure 56 (SEQ ID NO:56), Figure 58 (SEQ ID NO:58), Figure 60 (SEQ ID NO:60), Figure 62 (SEQ ID NO:62), Figure 64 (SEQ ID NO:64), Figure 66 (SEQ ID NO:66), Figure 68 (SEQ ID NO:68), Figure 70 (SEQ ID NO:70), Figure 72 (SEQ ID NO:72), Figure 74 (SEQ ID NO:74), Figure 76 (SEQ ID NO:76), Figure 78 (SEQ ID NO:78), Figure 80 (SEQ ID NO:80), Figure 82 (SEQ ID NO:82), Figure 84 (SEQ ID NO:84), Figure 86 (SEQ ID NO:86), Figure 88 (SEQ ID NO:88), Figure 90 (SEQ ID NO:90), Figure 92 (SEQ ID NO:92), Figure 94 (SEQ ID NO:94), Figure 96 (SEQ ID NO:96), Figure 98 (SEQ ID NO:98), Figure 100 (SEQ ID NO:100), Figure 102 (SEQ ID NO:102), Figure 104 (SEQ ID NO:104), Figure 106 (SEQ ID NO:106), Figure 108 (SEQ ID NO:108), Figure 110 (SEQ ID NO:110), Figure 112 (SEQ ID NO:112), Figure 114 (SEQ ID NO:114), Figure 116 (SEQ ID NO:116), Figure 118 (SEQ ID NO:118), Figure 120 (SEQ ID NO:120), Figure 122 (SEQ ID NO:122), Figure 124 (SEQ ID NO:124), Figure 126 (SEQ ID NO:126), Figure 128 (SEQ ID NO:128), Figure 130 (SEQ ID NO:130), Figure 132 (SEQ ID NO:132), Figure 134 (SEQ ID NO:134), Figure 136 (SEQ ID NO:136), Figure 138 (SEQ ID NO:138), Figure 140 (SEQ ID NO:140), Figure 142 (SEQ ID NO:142), Figure 144 (SEQ ID NO:144), Figure 146 (SEQ ID NO:146), Figure 148 (SEQ ID NO:148), Figure 150 (SEQ ID NO:150), Figure 152 (SEQ ID NO:152), Figure 154 (SEQ ID NO:154), Figure 156 (SEQ ID NO:156), Figure 158 (SEQ ID NO:158), Figure 160 (SEQ ID NO:160), Figure 162 (SEQ ID NO:162), Figure 164 (SEQ ID NO:164), Figure 166 (SEQ ID NO:166), Figure 168 (SEQ ID NO:168), Figure 170 (SEQ ID NO:170), Figure 172 (SEQ ID NO:172), Figure 174 (SEQ ID NO:174), Figure 176 (SEQ ID NO:176), Figure 178 (SEQ ID NO:178), Figure 180 (SEQ ID NO:180), Figure 182 (SEQ ID NO:182), Figure 184 (SEQ ID NO:184), Figure 186 (SEQ ID NO:186), Figure 188 (SEQ ID NO:188), Figure 190 (SEQ ID NO:190), Figure 192 (SEQ ID NO:192), Figure 194 (SEQ ID NO:194), Figure 196 (SEQ ID NO:196), Figure 198 (SEQ ID NO:198), Figure 200 (SEQ ID NO:200), Figure 202 (SEQ ID NO:202), Figure 204 (SEQ ID NO:204), Figure 206 (SEQ ID NO:206), Figure 208 (SEQ ID NO:208), Figure 210 (SEQ ID NO:210), Figure 212 (SEQ ID NO:212), Figure 214 (SEQ ID NO:214), Figure 216 (SEQ ID NO:216), Figure 218 (SEQ ID NO:218), Figure 220 (SEQ ID NO:220), Figure 222 (SEQ ID NO:222), Figure 224 (SEQ ID NO:224), Figure 226 (SEQ ID NO:226), Figure 228 (SEQ ID NO:228), Figure 230 (SEQ ID NO:230), Figure 232 (SEQ ID NO:232), Figure 234 (SEQ ID NO:234), Figure 236 (SEQ ID NO:236), Figure 238 (SEQ ID NO:238), Figure 240 (SEQ ID NO:240), Figure 242 (SEQ ID NO:242), Figure 244 (SEQ ID NO:244), Figure 246 (SEQ ID NO:246), Figure 248 (SEQ ID NO:248), Figure 250 (SEQ ID NO:250), Figure 252 (SEQ ID NO:252), Figure 254 (SEQ ID NO:254), Figure 256 (SEQ ID NO:256), Figure 258 (SEQ ID NO:258), Figure 260 (SEQ ID NO:260), Figure 262 (SEQ ID NO:262), Figure 264 (SEQ ID NO:264), Figure 266 (SEQ ID NO:266), Figure 268 (SEQ ID NO:268), Figure 270 (SEQ ID NO:270), Figure 272 (SEQ ID NO:272), Figure 274 (SEQ ID NO:274), Figure 276 (SEQ ID NO:276), Figure 278 (SEQ ID NO:278), Figure 280 (SEQ ID NO:280), Figure 282 (SEQ ID NO:282), Figure

284 (SEQ ID NO:284), Figure 286 (SEQ ID NO:286), Figure 288 (SEQ ID NO:288), Figure 290 (SEQ ID NO:290), Figure 292 (SEQ ID NO:292), Figure 294 (SEQ ID NO:294), Figure 296 (SEQ ID NO:296), Figure 298 (SEQ ID NO:298), Figure 300 (SEQ ID NO:300), Figure 302 (SEQ ID NO:302), Figure 304 (SEQ ID NO:304), Figure 306 (SEQ ID NO:306), Figure 308 (SEQ ID NO:308), Figure 310 (SEQ ID NO:310), Figure 312 (SEQ ID NO:312), Figure 314 (SEQ ID NO:314), Figure 316 (SEQ ID NO:316), Figure 318 (SEQ ID NO:318), Figure 320 (SEQ ID NO:320), Figure 322 (SEQ ID NO:322), Figure 324 (SEQ ID NO:324), Figure 326 (SEQ ID NO:326), Figure 328 (SEQ ID NO:328), Figure 330 (SEQ ID NO:330), Figure 332 (SEQ ID NO:332), Figure 334 (SEQ ID NO:334), Figure 336 (SEQ ID NO:336), Figure 338 (SEQ ID NO:338), Figure 340 (SEQ ID NO:340), Figure 342 (SEQ ID NO:342), Figure 344 (SEQ ID NO:344), Figure 346 (SEQ ID NO:346), Figure 348 (SEQ ID NO:348), Figure 350 (SEQ ID NO:350), Figure 352 (SEQ ID NO:352), Figure 354 (SEQ ID NO:354), Figure 356 (SEQ ID NO:356), Figure 358 (SEQ ID NO:358), Figure 360 (SEQ ID NO:360), Figure 362 (SEQ ID NO:362), Figure 364 (SEQ ID NO:364), Figure 366 (SEQ ID NO:366), Figure 368 (SEQ ID NO:368), Figure 370 (SEQ ID NO:370), Figure 372 (SEQ ID NO:372) or Figure 374 (SEQ ID NO:374), lacking its associated signal peptide.

19. An isolated polypeptide having at least 80% amino acid sequence identity to:

(a) an amino acid sequence of the polypeptide shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:4), Figure 6 (SEQ ID NO:6), Figure 8 (SEQ ID NO:8), Figure 10 (SEQ ID NO:10), Figure 12 (SEQ ID NO:12), Figure 14 (SEQ ID NO:14), Figure 16 (SEQ ID NO:16), Figure 18 (SEQ ID NO:18), Figure 20 (SEQ ID NO:20), Figure 22 (SEQ ID NO:22), Figure 24 (SEQ ID NO:24), Figure 26 (SEQ ID NO:26), Figure 28 (SEQ ID NO:28), Figure 30 (SEQ ID NO:30), Figure 32 (SEQ ID NO:32), Figure 34 (SEQ ID NO:34), Figure 36 (SEQ ID NO:36), Figure 38 (SEQ ID NO:38), Figure 40 (SEQ ID NO:40), Figure 42 (SEQ ID NO:42), Figure 44 (SEQ ID NO:44), Figure 46 (SEQ ID NO:46), Figure 48 (SEQ ID NO:48), Figure 50 (SEQ ID NO:50), Figure 52 (SEQ ID NO:52), Figure 54 (SEQ ID NO:54), Figure 56 (SEQ ID NO:56), Figure 58 (SEQ ID NO:58), Figure 60 (SEQ ID NO:60), Figure 62 (SEQ ID NO:62), Figure 64 (SEQ ID NO:64), Figure 66 (SEQ ID NO:66), Figure 68 (SEQ ID NO:68), Figure 70 (SEQ ID NO:70), Figure 72 (SEQ ID NO:72), Figure 74 (SEQ ID NO:74), Figure 76 (SEQ ID NO:76), Figure 78 (SEQ ID NO:78), Figure 80 (SEQ ID NO:80), Figure 82 (SEQ ID NO:82), Figure 84 (SEQ ID NO:84), Figure 86 (SEQ ID NO:86), Figure 88 (SEQ ID NO:88), Figure 90 (SEQ ID NO:90), Figure 92 (SEQ ID NO:92), Figure 94 (SEQ ID NO:94), Figure 96 (SEQ ID NO:96), Figure 98 (SEQ ID NO:98), Figure 100 (SEQ ID NO:100), Figure 102 (SEQ ID NO:102), Figure 104 (SEQ ID NO:104), Figure 106 (SEQ ID NO:106), Figure 108 (SEQ ID NO:108), Figure 110 (SEQ ID NO:110), Figure 112 (SEQ ID NO:112), Figure 114 (SEQ ID NO:114), Figure 116 (SEQ ID NO:116), Figure 118 (SEQ ID NO:118), Figure 120 (SEQ ID NO:120), Figure 122 (SEQ ID NO:122), Figure 124 (SEQ ID NO:124), Figure 126 (SEQ ID NO:126), Figure 128 (SEQ ID NO:128), Figure 130 (SEQ ID NO:130), Figure 132 (SEQ ID NO:132), Figure 134 (SEQ ID NO:134), Figure 136 (SEQ ID NO:136), Figure 138 (SEQ ID NO:138), Figure 140 (SEQ ID NO:140), Figure 142 (SEQ ID NO:142), Figure 144 (SEQ ID NO:144), Figure 146 (SEQ ID NO:146), Figure 148 (SEQ ID NO:148), Figure 150 (SEQ ID NO:150), Figure 152 (SEQ ID

NO:152), Figure 154 (SEQ ID NO:154), Figure 156 (SEQ ID NO:156), Figure 158 (SEQ ID NO:158), Figure 160 (SEQ ID NO:160), Figure 162 (SEQ ID NO:162), Figure 164 (SEQ ID NO:164), Figure 166 (SEQ ID NO:166), Figure 168 (SEQ ID NO:168), Figure 170 (SEQ ID NO:170), Figure 172 (SEQ ID NO:172), Figure 174 (SEQ ID NO:174), Figure 176 (SEQ ID NO:176), Figure 178 (SEQ ID NO:178), Figure 180 (SEQ ID NO:180), Figure 182 (SEQ ID NO:182), Figure 184 (SEQ ID NO:184), Figure 186 (SEQ ID NO:186), Figure 188 (SEQ ID NO:188), Figure 190 (SEQ ID NO:190), Figure 192 (SEQ ID NO:192), Figure 194 (SEQ ID NO:194), Figure 196 (SEQ ID NO:196), Figure 198 (SEQ ID NO:198), Figure 200 (SEQ ID NO:200), Figure 202 (SEQ ID NO:202), Figure 204 (SEQ ID NO:204), Figure 206 (SEQ ID NO:206), Figure 208 (SEQ ID NO:208), Figure 210 (SEQ ID NO:210), Figure 212 (SEQ ID NO:212), Figure 214 (SEQ ID NO:214), Figure 216 (SEQ ID NO:216), Figure 218 (SEQ ID NO:218), Figure 220 (SEQ ID NO:220), Figure 222 (SEQ ID NO:222), Figure 224 (SEQ ID NO:224), Figure 226 (SEQ ID NO:226), Figure 228 (SEQ ID NO:228), Figure 230 (SEQ ID NO:230), Figure 232 (SEQ ID NO:232), Figure 234 (SEQ ID NO:234), Figure 236 (SEQ ID NO:236), Figure 238 (SEQ ID NO:238), Figure 240 (SEQ ID NO:240), Figure 242 (SEQ ID NO:242), Figure 244 (SEQ ID NO:244), Figure 246 (SEQ ID NO:246), Figure 248 (SEQ ID NO:248), Figure 250 (SEQ ID NO:250), Figure 252 (SEQ ID NO:252), Figure 254 (SEQ ID NO:254), Figure 256 (SEQ ID NO:256), Figure 258 (SEQ ID NO:258), Figure 260 (SEQ ID NO:260), Figure 262 (SEQ ID NO:262), Figure 264 (SEQ ID NO:264), Figure 266 (SEQ ID NO:266), Figure 268 (SEQ ID NO:268), Figure 270 (SEQ ID NO:270), Figure 272 (SEQ ID NO:272), Figure 274 (SEQ ID NO:274), Figure 276 (SEQ ID NO:276), Figure 278 (SEQ ID NO:278), Figure 280 (SEQ ID NO:280), Figure 282 (SEQ ID NO:282), Figure 284 (SEQ ID NO:284), Figure 286 (SEQ ID NO:286), Figure 288 (SEQ ID NO:288), Figure 290 (SEQ ID NO:290), Figure 292 (SEQ ID NO:292), Figure 294 (SEQ ID NO:294), Figure 296 (SEQ ID NO:296), Figure 298 (SEQ ID NO:298), Figure 300 (SEQ ID NO:300), Figure 302 (SEQ ID NO:302), Figure 304 (SEQ ID NO:304), Figure 306 (SEQ ID NO:306), Figure 308 (SEQ ID NO:308), Figure 310 (SEQ ID NO:310), Figure 312 (SEQ ID NO:312), Figure 314 (SEQ ID NO:314), Figure 316 (SEQ ID NO:316), Figure 318 (SEQ ID NO:318), Figure 320 (SEQ ID NO:320), Figure 322 (SEQ ID NO:322), Figure 324 (SEQ ID NO:324), Figure 326 (SEQ ID NO:326), Figure 328 (SEQ ID NO:328), Figure 330 (SEQ ID NO:330), Figure 332 (SEQ ID NO:332), Figure 334 (SEQ ID NO:334), Figure 336 (SEQ ID NO:336), Figure 338 (SEQ ID NO:338), Figure 340 (SEQ ID NO:340), Figure 342 (SEQ ID NO:342), Figure 344 (SEQ ID NO:344), Figure 346 (SEQ ID NO:346), Figure 348 (SEQ ID NO:348), Figure 350 (SEQ ID NO:350), Figure 352 (SEQ ID NO:352), Figure 354 (SEQ ID NO:354), Figure 356 (SEQ ID NO:356), Figure 358 (SEQ ID NO:358), Figure 360 (SEQ ID NO:360), Figure 362 (SEQ ID NO:362), Figure 364 (SEQ ID NO:364), Figure 366 (SEQ ID NO:366), Figure 368 (SEQ ID NO:368), Figure 370 (SEQ ID NO:370), Figure 372 (SEQ ID NO:372) or Figure 374 (SEQ ID NO:374), lacking its associated signal peptide;

(b) an amino acid sequence of an extracellular domain of the polypeptide shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:4), Figure 6 (SEQ ID NO:6), Figure 8 (SEQ ID NO:8), Figure 10 (SEQ ID NO:10), Figure 12 (SEQ ID NO:12), Figure 14 (SEQ ID NO:14), Figure 16 (SEQ ID NO:16), Figure 18 (SEQ ID NO:18), Figure 20 (SEQ ID NO:20), Figure 22 (SEQ ID NO:22), Figure 24 (SEQ ID NO:24), Figure 26

(SEQ ID NO:26), Figure 28 (SEQ ID NO:28), Figure 30 (SEQ ID NO:30), Figure 32 (SEQ ID NO:32), Figure 34 (SEQ ID NO:34), Figure 36 (SEQ ID NO:36), Figure 38 (SEQ ID NO:38), Figure 40 (SEQ ID NO:40), Figure 42 (SEQ ID NO:42), Figure 44 (SEQ ID NO:44), Figure 46 (SEQ ID NO:46), Figure 48 (SEQ ID NO:48), Figure 50 (SEQ ID NO:50), Figure 52 (SEQ ID NO:52), Figure 54 (SEQ ID NO:54), Figure 56 (SEQ ID NO:56), Figure 58 (SEQ ID NO:58), Figure 60 (SEQ ID NO:60), Figure 62 (SEQ ID NO:62), Figure 64 (SEQ ID NO:64), Figure 66 (SEQ ID NO:66), Figure 68 (SEQ ID NO:68), Figure 70 (SEQ ID NO:70), Figure 72 (SEQ ID NO:72), Figure 74 (SEQ ID NO:74), Figure 76 (SEQ ID NO:76), Figure 78 (SEQ ID NO:78), Figure 80 (SEQ ID NO:80), Figure 82 (SEQ ID NO:82), Figure 84 (SEQ ID NO:84), Figure 86 (SEQ ID NO:86), Figure 88 (SEQ ID NO:88), Figure 90 (SEQ ID NO:90), Figure 92 (SEQ ID NO:92), Figure 94 (SEQ ID NO:94), Figure 96 (SEQ ID NO:96), Figure 98 (SEQ ID NO:98), Figure 100 (SEQ ID NO:100), Figure 102 (SEQ ID NO:102), Figure 104 (SEQ ID NO:104), Figure 106 (SEQ ID NO:106), Figure 108 (SEQ ID NO:108), Figure 110 (SEQ ID NO:110), Figure 112 (SEQ ID NO:112), Figure 114 (SEQ ID NO:114), Figure 116 (SEQ ID NO:116), Figure 118 (SEQ ID NO:118), Figure 120 (SEQ ID NO:120), Figure 122 (SEQ ID NO:122), Figure 124 (SEQ ID NO:124), Figure 126 (SEQ ID NO:126), Figure 128 (SEQ ID NO:128), Figure 130 (SEQ ID NO:130), Figure 132 (SEQ ID NO:132), Figure 134 (SEQ ID NO:134), Figure 136 (SEQ ID NO:136), Figure 138 (SEQ ID NO:138), Figure 140 (SEQ ID NO:140), Figure 142 (SEQ ID NO:142), Figure 144 (SEQ ID NO:144), Figure 146 (SEQ ID NO:146), Figure 148 (SEQ ID NO:148), Figure 150 (SEQ ID NO:150), Figure 152 (SEQ ID NO:152), Figure 154 (SEQ ID NO:154), Figure 156 (SEQ ID NO:156), Figure 158 (SEQ ID NO:158), Figure 160 (SEQ ID NO:160), Figure 162 (SEQ ID NO:162), Figure 164 (SEQ ID NO:164), Figure 166 (SEQ ID NO:166), Figure 168 (SEQ ID NO:168), Figure 170 (SEQ ID NO:170), Figure 172 (SEQ ID NO:172), Figure 174 (SEQ ID NO:174), Figure 176 (SEQ ID NO:176), Figure 178 (SEQ ID NO:178), Figure 180 (SEQ ID NO:180), Figure 182 (SEQ ID NO:182), Figure 184 (SEQ ID NO:184), Figure 186 (SEQ ID NO:186), Figure 188 (SEQ ID NO:188), Figure 190 (SEQ ID NO:190), Figure 192 (SEQ ID NO:192), Figure 194 (SEQ ID NO:194), Figure 196 (SEQ ID NO:196), Figure 198 (SEQ ID NO:198), Figure 200 (SEQ ID NO:200), Figure 202 (SEQ ID NO:202), Figure 204 (SEQ ID NO:204), Figure 206 (SEQ ID NO:206), Figure 208 (SEQ ID NO:208), Figure 210 (SEQ ID NO:210), Figure 212 (SEQ ID NO:212), Figure 214 (SEQ ID NO:214), Figure 216 (SEQ ID NO:216), Figure 218 (SEQ ID NO:218), Figure 220 (SEQ ID NO:220), Figure 222 (SEQ ID NO:222), Figure 224 (SEQ ID NO:224), Figure 226 (SEQ ID NO:226), Figure 228 (SEQ ID NO:228), Figure 230 (SEQ ID NO:230), Figure 232 (SEQ ID NO:232), Figure 234 (SEQ ID NO:234), Figure 236 (SEQ ID NO:236), Figure 238 (SEQ ID NO:238), Figure 240 (SEQ ID NO:240), Figure 242 (SEQ ID NO:242), Figure 244 (SEQ ID NO:244), Figure 246 (SEQ ID NO:246), Figure 248 (SEQ ID NO:248), Figure 250 (SEQ ID NO:250), Figure 252 (SEQ ID NO:252), Figure 254 (SEQ ID NO:254), Figure 256 (SEQ ID NO:256), Figure 258 (SEQ ID NO:258), Figure 260 (SEQ ID NO:260), Figure 262 (SEQ ID NO:262), Figure 264 (SEQ ID NO:264), Figure 266 (SEQ ID NO:266), Figure 268 (SEQ ID NO:268), Figure 270 (SEQ ID NO:270), Figure 272 (SEQ ID NO:272), Figure 274 (SEQ ID NO:274), Figure 276 (SEQ ID NO:276), Figure 278 (SEQ ID NO:278), Figure 280 (SEQ ID NO:280), Figure 282 (SEQ ID NO:282), Figure 284 (SEQ ID NO:284), Figure 286 (SEQ ID NO:286), Figure 288 (SEQ ID NO:288), Figure 290 (SEQ ID NO:290), Figure

292 (SEQ ID NO:292), Figure 294 (SEQ ID NO:294), Figure 296 (SEQ ID NO:296), Figure 298 (SEQ ID NO:298), Figure 300 (SEQ ID NO:300), Figure 302 (SEQ ID NO:302), Figure 304 (SEQ ID NO:304), Figure 306 (SEQ ID NO:306), Figure 308 (SEQ ID NO:308), Figure 310 (SEQ ID NO:310), Figure 312 (SEQ ID NO:312), Figure 314 (SEQ ID NO:314), Figure 316 (SEQ ID NO:316), Figure 318 (SEQ ID NO:318), Figure 320 (SEQ ID NO:320), Figure 322 (SEQ ID NO:322), Figure 324 (SEQ ID NO:324), Figure 326 (SEQ ID NO:326), Figure 328 (SEQ ID NO:328), Figure 330 (SEQ ID NO:330), Figure 332 (SEQ ID NO:332), Figure 334 (SEQ ID NO:334), Figure 336 (SEQ ID NO:336), Figure 338 (SEQ ID NO:338), Figure 340 (SEQ ID NO:340), Figure 342 (SEQ ID NO:342), Figure 344 (SEQ ID NO:344), Figure 346 (SEQ ID NO:346), Figure 348 (SEQ ID NO:348), Figure 350 (SEQ ID NO:350), Figure 352 (SEQ ID NO:352), Figure 354 (SEQ ID NO:354), Figure 356 (SEQ ID NO:356), Figure 358 (SEQ ID NO:358), Figure 360 (SEQ ID NO:360), Figure 362 (SEQ ID NO:362), Figure 364 (SEQ ID NO:364), Figure 366 (SEQ ID NO:366), Figure 368 (SEQ ID NO:368), Figure 370 (SEQ ID NO:370), Figure 372 (SEQ ID NO:372) or Figure 374 (SEQ ID NO:374), with its associated signal peptide; or

(c) an amino acid sequence of an extracellular domain of the polypeptide shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:4), Figure 6 (SEQ ID NO:6), Figure 8 (SEQ ID NO:8), Figure 10 (SEQ ID NO:10), Figure 12 (SEQ ID NO:12), Figure 14 (SEQ ID NO:14), Figure 16 (SEQ ID NO:16), Figure 18 (SEQ ID NO:18), Figure 20 (SEQ ID NO:20), Figure 22 (SEQ ID NO:22), Figure 24 (SEQ ID NO:24), Figure 26 (SEQ ID NO:26), Figure 28 (SEQ ID NO:28), Figure 30 (SEQ ID NO:30), Figure 32 (SEQ ID NO:32), Figure 34 (SEQ ID NO:34), Figure 36 (SEQ ID NO:36), Figure 38 (SEQ ID NO:38), Figure 40 (SEQ ID NO:40), Figure 42 (SEQ ID NO:42), Figure 44 (SEQ ID NO:44), Figure 46 (SEQ ID NO:46), Figure 48 (SEQ ID NO:48), Figure 50 (SEQ ID NO:50), Figure 52 (SEQ ID NO:52), Figure 54 (SEQ ID NO:54), Figure 56 (SEQ ID NO:56), Figure 58 (SEQ ID NO:58), Figure 60 (SEQ ID NO:60), Figure 62 (SEQ ID NO:62), Figure 64 (SEQ ID NO:64), Figure 66 (SEQ ID NO:66), Figure 68 (SEQ ID NO:68), Figure 70 (SEQ ID NO:70), Figure 72 (SEQ ID NO:72), Figure 74 (SEQ ID NO:74), Figure 76 (SEQ ID NO:76), Figure 78 (SEQ ID NO:78), Figure 80 (SEQ ID NO:80), Figure 82 (SEQ ID NO:82), Figure 84 (SEQ ID NO:84), Figure 86 (SEQ ID NO:86), Figure 88 (SEQ ID NO:88), Figure 90 (SEQ ID NO:90), Figure 92 (SEQ ID NO:92), Figure 94 (SEQ ID NO:94), Figure 96 (SEQ ID NO:96), Figure 98 (SEQ ID NO:98), Figure 100 (SEQ ID NO:100), Figure 102 (SEQ ID NO:102), Figure 104 (SEQ ID NO:104), Figure 106 (SEQ ID NO:106), Figure 108 (SEQ ID NO:108), Figure 110 (SEQ ID NO:110), Figure 112 (SEQ ID NO:112), Figure 114 (SEQ ID NO:114), Figure 116 (SEQ ID NO:116), Figure 118 (SEQ ID NO:118), Figure 120 (SEQ ID NO:120), Figure 122 (SEQ ID NO:122), Figure 124 (SEQ ID NO:124), Figure 126 (SEQ ID NO:126), Figure 128 (SEQ ID NO:128), Figure 130 (SEQ ID NO:130), Figure 132 (SEQ ID NO:132), Figure 134 (SEQ ID NO:134), Figure 136 (SEQ ID NO:136), Figure 138 (SEQ ID NO:138), Figure 140 (SEQ ID NO:140), Figure 142 (SEQ ID NO:142), Figure 144 (SEQ ID NO:144), Figure 146 (SEQ ID NO:146), Figure 148 (SEQ ID NO:148), Figure 150 (SEQ ID NO:150), Figure 152 (SEQ ID NO:152), Figure 154 (SEQ ID NO:154), Figure 156 (SEQ ID NO:156), Figure 158 (SEQ ID NO:158), Figure 160 (SEQ ID NO:160), Figure 162 (SEQ ID NO:162), Figure 164 (SEQ ID NO:164), Figure 166 (SEQ ID NO:166), Figure 168 (SEQ ID NO:168), Figure 170 (SEQ ID NO:170), Figure 172 (SEQ ID

NO:172), Figure 174 (SEQ ID NO:174), Figure 176 (SEQ ID NO:176), Figure 178 (SEQ ID NO:178), Figure
 180 (SEQ ID NO:180), Figure 182 (SEQ ID NO:182), Figure 184 (SEQ ID NO:184), Figure 186 (SEQ ID
 NO:186), Figure 188 (SEQ ID NO:188), Figure 190 (SEQ ID NO:190), Figure 192 (SEQ ID NO:192), Figure
 194 (SEQ ID NO:194), Figure 196 (SEQ ID NO:196), Figure 198 (SEQ ID NO:198), Figure 200 (SEQ ID
 5 NO:200), Figure 202 (SEQ ID NO:202), Figure 204 (SEQ ID NO:204), Figure 206 (SEQ ID NO:206), Figure
 208 (SEQ ID NO:208), Figure 210 (SEQ ID NO:210), Figure 212 (SEQ ID NO:212), Figure 214 (SEQ ID
 NO:214), Figure 216 (SEQ ID NO:216), Figure 218 (SEQ ID NO:218), Figure 220 (SEQ ID NO:220), Figure
 222 (SEQ ID NO:222), Figure 224 (SEQ ID NO:224), Figure 226 (SEQ ID NO:226), Figure 228 (SEQ ID
 NO:228), Figure 230 (SEQ ID NO:230), Figure 232 (SEQ ID NO:232), Figure 234 (SEQ ID NO:234), Figure
 10 236 (SEQ ID NO:236), Figure 238 (SEQ ID NO:238), Figure 240 (SEQ ID NO:240), Figure 242 (SEQ ID
 NO:242), Figure 244 (SEQ ID NO:244), Figure 246 (SEQ ID NO:246), Figure 248 (SEQ ID NO:248), Figure
 250 (SEQ ID NO:250), Figure 252 (SEQ ID NO:252), Figure 254 (SEQ ID NO:254), Figure 256 (SEQ ID
 NO:256), Figure 258 (SEQ ID NO:258), Figure 260 (SEQ ID NO:260), Figure 262 (SEQ ID NO:262), Figure
 264 (SEQ ID NO:264), Figure 266 (SEQ ID NO:266), Figure 268 (SEQ ID NO:268), Figure 270 (SEQ ID
 15 NO:270), Figure 272 (SEQ ID NO:272), Figure 274 (SEQ ID NO:274), Figure 276 (SEQ ID NO:276), Figure
 278 (SEQ ID NO:278), Figure 280 (SEQ ID NO:280), Figure 282 (SEQ ID NO:282), Figure 284 (SEQ ID
 NO:284), Figure 286 (SEQ ID NO:286), Figure 288 (SEQ ID NO:288), Figure 290 (SEQ ID NO:290), Figure
 292 (SEQ ID NO:292), Figure 294 (SEQ ID NO:294), Figure 296 (SEQ ID NO:296), Figure 298 (SEQ ID
 NO:298), Figure 300 (SEQ ID NO:300), Figure 302 (SEQ ID NO:302), Figure 304 (SEQ ID NO:304), Figure
 20 306 (SEQ ID NO:306), Figure 308 (SEQ ID NO:308), Figure 310 (SEQ ID NO:310), Figure 312 (SEQ ID
 NO:312), Figure 314 (SEQ ID NO:314), Figure 316 (SEQ ID NO:316), Figure 318 (SEQ ID NO:318), Figure
 320 (SEQ ID NO:320), Figure 322 (SEQ ID NO:322), Figure 324 (SEQ ID NO:324), Figure 326 (SEQ ID
 NO:326), Figure 328 (SEQ ID NO:328), Figure 330 (SEQ ID NO:330), Figure 332 (SEQ ID NO:332), Figure
 334 (SEQ ID NO:334), Figure 336 (SEQ ID NO:336), Figure 338 (SEQ ID NO:338), Figure 340 (SEQ ID
 25 NO:340), Figure 342 (SEQ ID NO:342), Figure 344 (SEQ ID NO:344), Figure 346 (SEQ ID NO:346), Figure
 348 (SEQ ID NO:348), Figure 350 (SEQ ID NO:350), Figure 352 (SEQ ID NO:352), Figure 354 (SEQ ID
 NO:354), Figure 356 (SEQ ID NO:356), Figure 358 (SEQ ID NO:358), Figure 360 (SEQ ID NO:360), Figure
 362 (SEQ ID NO:362), Figure 364 (SEQ ID NO:364), Figure 366 (SEQ ID NO:366), Figure 368 (SEQ ID
 NO:368), Figure 370 (SEQ ID NO:370), Figure 372 (SEQ ID NO:372) or Figure 374 (SEQ ID NO:374), lacking
 30 its associated signal peptide.

20. A method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal
 comprising administering to the mammal a therapeutically effective amount of a polypeptide shown in Figure 2
 (SEQ ID NO:2), Figure 4 (SEQ ID NO:4), Figure 6 (SEQ ID NO:6), Figure 8 (SEQ ID NO:8), Figure 10 (SEQ
 35 ID NO:10), Figure 12 (SEQ ID NO:12), Figure 14 (SEQ ID NO:14), Figure 16 (SEQ ID NO:16), Figure 18
 (SEQ ID NO:18), Figure 20 (SEQ ID NO:20), Figure 22 (SEQ ID NO:22), Figure 24 (SEQ ID NO:24), Figure
 26 (SEQ ID NO:26), Figure 28 (SEQ ID NO:28), Figure 30 (SEQ ID NO:30), Figure 32 (SEQ ID NO:32),

Figure 34 (SEQ ID NO:34), Figure 36 (SEQ ID NO:36), Figure 38 (SEQ ID NO:38), Figure 40 (SEQ ID NO:40), Figure 42 (SEQ ID NO:42), Figure 44 (SEQ ID NO:44), Figure 46 (SEQ ID NO:46), Figure 48 (SEQ ID NO:48), Figure 50 (SEQ ID NO:50), Figure 52 (SEQ ID NO:52), Figure 54 (SEQ ID NO:54), Figure 56 (SEQ ID NO:56), Figure 58 (SEQ ID NO:58), Figure 60 (SEQ ID NO:60), Figure 62 (SEQ ID NO:62), Figure 64 (SEQ ID NO:64), Figure 66 (SEQ ID NO:66), Figure 68 (SEQ ID NO:68), Figure 70 (SEQ ID NO:70), Figure 72 (SEQ ID NO:72), Figure 74 (SEQ ID NO:74), Figure 76 (SEQ ID NO:76), Figure 78 (SEQ ID NO:78), Figure 80 (SEQ ID NO:80), Figure 82 (SEQ ID NO:82), Figure 84 (SEQ ID NO:84), Figure 86 (SEQ ID NO:86), Figure 88 (SEQ ID NO:88), Figure 90 (SEQ ID NO:90), Figure 92 (SEQ ID NO:92), Figure 94 (SEQ ID NO:94), Figure 96 (SEQ ID NO:96), Figure 98 (SEQ ID NO:98), Figure 100 (SEQ ID NO:100), Figure 102 (SEQ ID NO:102), Figure 104 (SEQ ID NO:104), Figure 106 (SEQ ID NO:106), Figure 108 (SEQ ID NO:108), Figure 110 (SEQ ID NO:110), Figure 112 (SEQ ID NO:112), Figure 114 (SEQ ID NO:114), Figure 116 (SEQ ID NO:116), Figure 118 (SEQ ID NO:118), Figure 120 (SEQ ID NO:120), Figure 122 (SEQ ID NO:122), Figure 124 (SEQ ID NO:124), Figure 126 (SEQ ID NO:126), Figure 128 (SEQ ID NO:128), Figure 130 (SEQ ID NO:130), Figure 132 (SEQ ID NO:132), Figure 134 (SEQ ID NO:134), Figure 136 (SEQ ID NO:136), Figure 138 (SEQ ID NO:138), Figure 140 (SEQ ID NO:140), Figure 142 (SEQ ID NO:142), Figure 144 (SEQ ID NO:144), Figure 146 (SEQ ID NO:146), Figure 148 (SEQ ID NO:148), Figure 150 (SEQ ID NO:150), Figure 152 (SEQ ID NO:152), Figure 154 (SEQ ID NO:154), Figure 156 (SEQ ID NO:156), Figure 158 (SEQ ID NO:158), Figure 160 (SEQ ID NO:160), Figure 162 (SEQ ID NO:162), Figure 164 (SEQ ID NO:164), Figure 166 (SEQ ID NO:166), Figure 168 (SEQ ID NO:168), Figure 170 (SEQ ID NO:170), Figure 172 (SEQ ID NO:172), Figure 174 (SEQ ID NO:174), Figure 176 (SEQ ID NO:176), Figure 178 (SEQ ID NO:178), Figure 180 (SEQ ID NO:180), Figure 182 (SEQ ID NO:182), Figure 184 (SEQ ID NO:184), Figure 186 (SEQ ID NO:186), Figure 188 (SEQ ID NO:188), Figure 190 (SEQ ID NO:190), Figure 192 (SEQ ID NO:192), Figure 194 (SEQ ID NO:194), Figure 196 (SEQ ID NO:196), Figure 198 (SEQ ID NO:198), Figure 200 (SEQ ID NO:200), Figure 202 (SEQ ID NO:202), Figure 204 (SEQ ID NO:204), Figure 206 (SEQ ID NO:206), Figure 208 (SEQ ID NO:208), Figure 210 (SEQ ID NO:210), Figure 212 (SEQ ID NO:212), Figure 214 (SEQ ID NO:214), Figure 216 (SEQ ID NO:216), Figure 218 (SEQ ID NO:218), Figure 220 (SEQ ID NO:220), Figure 222 (SEQ ID NO:222), Figure 224 (SEQ ID NO:224), Figure 226 (SEQ ID NO:226), Figure 228 (SEQ ID NO:228), Figure 230 (SEQ ID NO:230), Figure 232 (SEQ ID NO:232), Figure 234 (SEQ ID NO:234), Figure 236 (SEQ ID NO:236), Figure 238 (SEQ ID NO:238), Figure 240 (SEQ ID NO:240), Figure 242 (SEQ ID NO:242), Figure 244 (SEQ ID NO:244), Figure 246 (SEQ ID NO:246), Figure 248 (SEQ ID NO:248), Figure 250 (SEQ ID NO:250), Figure 252 (SEQ ID NO:252), Figure 254 (SEQ ID NO:254), Figure 256 (SEQ ID NO:256), Figure 258 (SEQ ID NO:258), Figure 260 (SEQ ID NO:260), Figure 262 (SEQ ID NO:262), Figure 264 (SEQ ID NO:264), Figure 266 (SEQ ID NO:266), Figure 268 (SEQ ID NO:268), Figure 270 (SEQ ID NO:270), Figure 272 (SEQ ID NO:272), Figure 274 (SEQ ID NO:274), Figure 276 (SEQ ID NO:276), Figure 278 (SEQ ID NO:278), Figure 280 (SEQ ID NO:280), Figure 282 (SEQ ID NO:282), Figure 284 (SEQ ID NO:284), Figure 286 (SEQ ID NO:286), Figure 288 (SEQ ID NO:288), Figure 290 (SEQ ID NO:290), Figure 292 (SEQ ID NO:292), Figure 294 (SEQ ID NO:294), Figure 296 (SEQ ID NO:296), Figure

298 (SEQ ID NO:298), Figure 300 (SEQ ID NO:300), Figure 302 (SEQ ID NO:302), Figure 304 (SEQ ID NO:304), Figure 306 (SEQ ID NO:306), Figure 308 (SEQ ID NO:308), Figure 310 (SEQ ID NO:310), Figure 312 (SEQ ID NO:312), Figure 314 (SEQ ID NO:314), Figure 316 (SEQ ID NO:316), Figure 318 (SEQ ID NO:318), Figure 320 (SEQ ID NO:320), Figure 322 (SEQ ID NO:322), Figure 324 (SEQ ID NO:324), Figure 326 (SEQ ID NO:326), Figure 328 (SEQ ID NO:328), Figure 330 (SEQ ID NO:330), Figure 332 (SEQ ID NO:332), Figure 334 (SEQ ID NO:334), Figure 336 (SEQ ID NO:336), Figure 338 (SEQ ID NO:338), Figure 340 (SEQ ID NO:340), Figure 342 (SEQ ID NO:342), Figure 344 (SEQ ID NO:344), Figure 346 (SEQ ID NO:346), Figure 348 (SEQ ID NO:348), Figure 350 (SEQ ID NO:350), Figure 352 (SEQ ID NO:352), Figure 354 (SEQ ID NO:354), Figure 356 (SEQ ID NO:356), Figure 358 (SEQ ID NO:358), Figure 360 (SEQ ID NO:360), Figure 362 (SEQ ID NO:362), Figure 364 (SEQ ID NO:364), Figure 366 (SEQ ID NO:366), Figure 368 (SEQ ID NO:368), Figure 370 (SEQ ID NO:370), Figure 372 (SEQ ID NO:372) or Figure 374 (SEQ ID NO:374), or agonist or antagonist thereof.

21. The method according to Claim 20, wherein the mammal is human.
22. The method of Claim 21, wherein the human has suffered myocardial infarction.
23. The method of Claim 21, wherein the human has cardiac hypertrophy, trauma, a cancer, or age-related macular degeneration.
24. The method of Claim 23, wherein the cardiac hypertrophy is characterized by the presence of an elevated level of $\text{PGF}_{2\alpha}$.
25. The method of Claim 20, wherein the polypeptide is administered together with a cardiovascular, endothelial or angiogenic agent.
26. The method of Claim 23, wherein the polypeptide is administered following primary angioplasty.
27. The method of Claim 20, wherein the cardiovascular, endothelial or angiogenic disorder is cancer.
28. The method of Claim 27, wherein the polypeptide is administered in combination with a chemotherapeutic agent, a growth inhibitory agent or a cytotoxic agent.
29. The method of Claim 20, wherein said agonist is an antibody to said polypeptide.
30. The method of Claim 20, wherein said antagonist is an antibody to said polypeptide.

31. A method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal comprising administering to the mammal a nucleic acid molecule that encodes a polypeptide shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:4), Figure 6 (SEQ ID NO:6), Figure 8 (SEQ ID NO:8), Figure 10 (SEQ ID NO:10), Figure 12 (SEQ ID NO:12), Figure 14 (SEQ ID NO:14), Figure 16 (SEQ ID NO:16), Figure 18 (SEQ ID NO:18), Figure 20 (SEQ ID NO:20), Figure 22 (SEQ ID NO:22), Figure 24 (SEQ ID NO:24), Figure 26 (SEQ ID NO:26), Figure 28 (SEQ ID NO:28), Figure 30 (SEQ ID NO:30), Figure 32 (SEQ ID NO:32), Figure 34 (SEQ ID NO:34), Figure 36 (SEQ ID NO:36), Figure 38 (SEQ ID NO:38), Figure 40 (SEQ ID NO:40), Figure 42 (SEQ ID NO:42), Figure 44 (SEQ ID NO:44), Figure 46 (SEQ ID NO:46), Figure 48 (SEQ ID NO:48), Figure 50 (SEQ ID NO:50), Figure 52 (SEQ ID NO:52), Figure 54 (SEQ ID NO:54), Figure 56 (SEQ ID NO:56), Figure 58 (SEQ ID NO:58), Figure 60 (SEQ ID NO:60), Figure 62 (SEQ ID NO:62), Figure 64 (SEQ ID NO:64), Figure 66 (SEQ ID NO:66), Figure 68 (SEQ ID NO:68), Figure 70 (SEQ ID NO:70), Figure 72 (SEQ ID NO:72), Figure 74 (SEQ ID NO:74), Figure 76 (SEQ ID NO:76), Figure 78 (SEQ ID NO:78), Figure 80 (SEQ ID NO:80), Figure 82 (SEQ ID NO:82), Figure 84 (SEQ ID NO:84), Figure 86 (SEQ ID NO:86), Figure 88 (SEQ ID NO:88), Figure 90 (SEQ ID NO:90), Figure 92 (SEQ ID NO:92), Figure 94 (SEQ ID NO:94), Figure 96 (SEQ ID NO:96), Figure 98 (SEQ ID NO:98), Figure 100 (SEQ ID NO:100), Figure 102 (SEQ ID NO:102), Figure 104 (SEQ ID NO:104), Figure 106 (SEQ ID NO:106), Figure 108 (SEQ ID NO:108), Figure 110 (SEQ ID NO:110), Figure 112 (SEQ ID NO:112), Figure 114 (SEQ ID NO:114), Figure 116 (SEQ ID NO:116), Figure 118 (SEQ ID NO:118), Figure 120 (SEQ ID NO:120), Figure 122 (SEQ ID NO:122), Figure 124 (SEQ ID NO:124), Figure 126 (SEQ ID NO:126), Figure 128 (SEQ ID NO:128), Figure 130 (SEQ ID NO:130), Figure 132 (SEQ ID NO:132), Figure 134 (SEQ ID NO:134), Figure 136 (SEQ ID NO:136), Figure 138 (SEQ ID NO:138), Figure 140 (SEQ ID NO:140), Figure 142 (SEQ ID NO:142), Figure 144 (SEQ ID NO:144), Figure 146 (SEQ ID NO:146), Figure 148 (SEQ ID NO:148), Figure 150 (SEQ ID NO:150), Figure 152 (SEQ ID NO:152), Figure 154 (SEQ ID NO:154), Figure 156 (SEQ ID NO:156), Figure 158 (SEQ ID NO:158), Figure 160 (SEQ ID NO:160), Figure 162 (SEQ ID NO:162), Figure 164 (SEQ ID NO:164), Figure 166 (SEQ ID NO:166), Figure 168 (SEQ ID NO:168), Figure 170 (SEQ ID NO:170), Figure 172 (SEQ ID NO:172), Figure 174 (SEQ ID NO:174), Figure 176 (SEQ ID NO:176), Figure 178 (SEQ ID NO:178), Figure 180 (SEQ ID NO:180), Figure 182 (SEQ ID NO:182), Figure 184 (SEQ ID NO:184), Figure 186 (SEQ ID NO:186), Figure 188 (SEQ ID NO:188), Figure 190 (SEQ ID NO:190), Figure 192 (SEQ ID NO:192), Figure 194 (SEQ ID NO:194), Figure 196 (SEQ ID NO:196), Figure 198 (SEQ ID NO:198), Figure 200 (SEQ ID NO:200), Figure 202 (SEQ ID NO:202), Figure 204 (SEQ ID NO:204), Figure 206 (SEQ ID NO:206), Figure 208 (SEQ ID NO:208), Figure 210 (SEQ ID NO:210), Figure 212 (SEQ ID NO:212), Figure 214 (SEQ ID NO:214), Figure 216 (SEQ ID NO:216), Figure 218 (SEQ ID NO:218), Figure 220 (SEQ ID NO:220), Figure 222 (SEQ ID NO:222), Figure 224 (SEQ ID NO:224), Figure 226 (SEQ ID NO:226), Figure 228 (SEQ ID NO:228), Figure 230 (SEQ ID NO:230), Figure 232 (SEQ ID NO:232), Figure 234 (SEQ ID NO:234), Figure 236 (SEQ ID NO:236), Figure 238 (SEQ ID NO:238), Figure 240 (SEQ ID NO:240), Figure 242 (SEQ ID NO:242), Figure 244 (SEQ ID NO:244), Figure 246 (SEQ ID NO:246), Figure 248 (SEQ ID NO:248), Figure 250 (SEQ ID NO:250), Figure 252 (SEQ ID NO:252), Figure 254 (SEQ ID NO:254), Figure

256 (SEQ ID NO:256), Figure 258 (SEQ ID NO:258), Figure 260 (SEQ ID NO:260), Figure 262 (SEQ ID NO:262), Figure 264 (SEQ ID NO:264), Figure 266 (SEQ ID NO:266), Figure 268 (SEQ ID NO:268), Figure 270 (SEQ ID NO:270), Figure 272 (SEQ ID NO:272), Figure 274 (SEQ ID NO:274), Figure 276 (SEQ ID NO:276), Figure 278 (SEQ ID NO:278), Figure 280 (SEQ ID NO:280), Figure 282 (SEQ ID NO:282), Figure 284 (SEQ ID NO:284), Figure 286 (SEQ ID NO:286), Figure 288 (SEQ ID NO:288), Figure 290 (SEQ ID NO:290), Figure 292 (SEQ ID NO:292), Figure 294 (SEQ ID NO:294), Figure 296 (SEQ ID NO:296), Figure 298 (SEQ ID NO:298), Figure 300 (SEQ ID NO:300), Figure 302 (SEQ ID NO:302), Figure 304 (SEQ ID NO:304), Figure 306 (SEQ ID NO:306), Figure 308 (SEQ ID NO:308), Figure 310 (SEQ ID NO:310), Figure 312 (SEQ ID NO:312), Figure 314 (SEQ ID NO:314), Figure 316 (SEQ ID NO:316), Figure 318 (SEQ ID NO:318), Figure 320 (SEQ ID NO:320), Figure 322 (SEQ ID NO:322), Figure 324 (SEQ ID NO:324), Figure 326 (SEQ ID NO:326), Figure 328 (SEQ ID NO:328), Figure 330 (SEQ ID NO:330), Figure 332 (SEQ ID NO:332), Figure 334 (SEQ ID NO:334), Figure 336 (SEQ ID NO:336), Figure 338 (SEQ ID NO:338), Figure 340 (SEQ ID NO:340), Figure 342 (SEQ ID NO:342), Figure 344 (SEQ ID NO:344), Figure 346 (SEQ ID NO:346), Figure 348 (SEQ ID NO:348), Figure 350 (SEQ ID NO:350), Figure 352 (SEQ ID NO:352), Figure 354 (SEQ ID NO:354), Figure 356 (SEQ ID NO:356), Figure 358 (SEQ ID NO:358), Figure 360 (SEQ ID NO:360), Figure 362 (SEQ ID NO:362), Figure 364 (SEQ ID NO:364), Figure 366 (SEQ ID NO:366), Figure 368 (SEQ ID NO:368), Figure 370 (SEQ ID NO:370), Figure 372 (SEQ ID NO:372) or Figure 374 (SEQ ID NO:374), or agonist or antagonist thereof.

- 20 32. The method of Claim 31, wherein said agonist is an antibody to said polypeptide.
33. The method of Claim 31, wherein said antagonist is an antibody to said polypeptide.
34. The method of Claim 31, wherein the mammal is human.
- 25 35. The method of Claim 31, wherein the nucleic acid molecule is administered via *ex vivo* gene therapy.
36. A method for inhibiting endothelial cell growth in a mammal comprising administering to the mammal a PRO247, PRO720 or PRO4302 polypeptide or agonist thereof, wherein endothelial cell growth in said mammal is inhibited.
- 30 37. A method for stimulating endothelial cell growth in a mammal comprising administering to the mammal a PRO21, PRO181, PRO205, PRO214, PRO221, PRO229, PRO231, PRO238, PRO241, PRO247, PRO256, PRO258, PRO263, PRO265, PRO295, PRO321, PRO322, PRO337, PRO363, PRO365, PRO444, PRO533, PRO697, PRO725, PRO771, PRO788, PRO791, PRO819, PRO827, PRO828, PRO836, PRO846, PRO865, PRO1005, PRO1006, PRO1007, PRO1025, PRO1029, PRO1054, PRO1071, PRO1075, PRO1079,
- 35

PRO1080, PRO1114, PRO1131, PRO1155, PRO1160, PRO1184, PRO1186, PRO1190, PRO1192, PRO1195, PRO1244, PRO1272, PRO1273, PRO1274, PRO1279, PRO1283, PRO1286, PRO1306, PRO1309, PRO1325, PRO1329, PRO1347, PRO1356, PRO1376, PRO1382, PRO1411, PRO1412, PRO1419, PRO1474, PRO1477, PRO1488, PRO1508, PRO1550, PRO1556, PRO1760, PRO1782, PRO1787, PRO1801, PRO1868, PRO1887, PRO1890, PRO3438, PRO3444, PRO4302, PRO4324, PRO4333, PRO4341, PRO4342, PRO4353, PRO4354, PRO4356, PRO4371, PRO4405, PRO4408, PRO4422, PRO4425, PRO4499, PRO5723, PRO5725, PRO5737, PRO5776, PRO6006, PRO6029, PRO6071, PRO7436, PRO9771, PRO9821, PRO9873, PRO10008, PRO10096, PRO19670, PRO20040, PRO20044, PRO21055, PRO21384 or PRO28631 polypeptide, or agonist thereof, wherein endothelial cell growth in said mammal is stimulated.

38. A method for inducing cardiac hypertrophy in a mammal comprising administering to the mammal a PRO21 polypeptide or agonist thereof, wherein cardiac hypertrophy in said mammal is induced.

39. A method for stimulating angiogenesis induced by a PRO1376 or PRO1449 polypeptide in a mammal comprising administering a therapeutically effective amount of said polypeptide to the mammal, wherein said angiogenesis is stimulated.

40. A method for inducing endothelial cell apoptosis comprising administering to the endothelial cell a PRO4302 polypeptide or agonist thereof, wherein apoptosis in said endothelial cell is induced.

41. A method for stimulating smooth muscle cell growth comprising administering to the smooth muscle cell a PRO162, PRO181, PRO182, PRO195, PRO204, PRO221, PRO230, PRO256, PRO258, PRO533, PRO697, PRO725, PRO738, PRO826, PRO836, PRO840, PRO846, PRO865, PRO982, PRO1025, PRO1029, PRO1071, PRO1080, PRO1083, PRO1134, PRO1160, PRO1182, PRO1184, PRO1186, PRO1192, PRO1265, PRO1274, PRO1279, PRO1283, PRO1306, PRO1308, PRO1309, PRO1325, PRO1337, PRO1338, PRO1343, PRO1376, PRO1387, PRO1411, PRO1412, PRO1415, PRO1434, PRO1474, PRO1488, PRO1550, PRO1556, PRO1567, PRO1600, PRO1754, PRO1758, PRO1760, PRO1787, PRO1865, PRO1868, PRO1917, PRO1928, PRO3438, PRO3562, PRO4302, PRO4333, PRO4345, PRO4353, PRO4354, PRO4405, PRO4408, PRO4430, PRO4503, PRO5725, PRO6714, PRO9771, PRO9820, PRO9940, PRO10096, PRO21055, PRO21184 or PRO21366 polypeptide, or agonist thereof, wherein smooth muscle cell growth in said smooth muscle cell is stimulated.

42. A method for inducing endothelial cell tube formation comprising administering to the endothelial cell a PRO178, PRO195, PRO228, PRO301, PRO302, PRO532, PRO724, PRO730, PRO734, PRO793, PRO871, PRO938, PRO1012, PRO1120, PRO1139, PRO1198, PRO1287, PRO1361, PRO1864, PRO1873, PRO2010, PRO3579, PRO4313, PRO4527, PRO4538, PRO4553, PRO4995, PRO5730, PRO6008, PRO7223, PRO7248 or PRO7261 polypeptide, or agonist thereof, wherein tube formation in said endothelial cell is induced.

1/392

FIGURE 1

GCCCACGCGTCCGATGGCGTTCACGTTTCGCGGCCTTCTGCTACATGCTGGCGCTGCTGCT
CACTGCCGCGCTCATCTTCTTCGCCATTTGGCACATTATAGCATTGATGAGCTGAAGAC
TGATTACAAGAATCCTATAGACCAGTGTAATACCCTGAATCCCCCTTGTAAGTCCCAGAGTA
CCTCATCCACGCTTTCTTCTGTGTCATGTTTCTTTGTGCAGCAGAGTGGCTTACACTGGG
TCTCAATATGCCCCCTCTTGGCATATCATATTTGGAGGTATATGAGTAGACCAGTGATGAG
TGGCCCAGGACTCTATGACCCCTACAACCATCATGAATGCAGATATTCTAGCATATTGTCA
GAAGGAAGGATGGTGCAAATTAGCTTTTTATCTTCTAGCATTTTTTTTACTACCTATATGG
CATGATCTATGTTTTGGTGAGCTCTTAGAACAACACACAGAAGAATTGGTCCAGTTAAGT
GCATGCAAAAAGCCACCAAATGAAGGGATTCTATCCAGCAAGATCCTGTCCAAGAGTAGC
CTGTGGAATCTGATCAGTTACTTTAAAAAATGACTCCTTATTTTTTAAATGTTTCCACAT
TTTTGCTTGTGGAAAGACTGTTTTCATATGTTTATACTCAGATAAAGATTTTTAAATGGTAT
TACGTATAAATTAATATAAAATGATTACCTCTGGTGTTGACAGGTTTGAAGTTGCACTTC
TTAAGGAACAGCCATAATCCTCTGAATGATGCATTAATTACTGACTGTCCTAGTACATTG
GAAGCTTTTGTATTATAGGAAGTTGTAGGGCTCATTTTGGTTTCATTGAAACAGTATCTAA
TTATAAATTAGCTGTAGATATCAGGTGCTTCTGATGAAGTGAAAATGTATATCTGACTAG
TGGGAAACTTCATGGGTTTCCCTCATCTGTGATGTCGATGATTATATATGGATACATTTAC
AAAAATAAAAAGCGGGAATTTTCCCTTCGCTTGAATATTATCCCTGTATATTGCATGAAT
GAGAGATTTCCCATATTTCCATCAGAGTAATAAATATACTTGCTTTAATTCTTAAGCATA
AGTAAACATGATATAAAAATATATGCTGAATTACTTGTGAAGAATGCATTTAAAGCTATT
TTAAATGTGTTTTTATTTGTAAGACATTACTTATTAAGAAATTGGTTATTATGCTTACTG
TTCTAATCTGGTGGTAAAGGTATTCTTAAGAAATTGCAGGTACTACAGATTTTCAAACT
GAATGAGAGAAAATTGTATAACCATCCTGCTGTTCTTTAGTGCAATACAATAAACTCT
GAAATTAAGACTC

2/392

FIGURE 2

MAFTFAAFCYMLALLLTAALIFFAIWHIIAFDELKTDYKNPIDQCNTLNPLVLPEYLIHA
FFCVMFLCAAEWLTLGLNMPLLAYHIWRYMSRPVMSGPLYDPTTIMNADILAYCQKEGW
CKLAFYLLAFFYYLYGMIYVLVSS

Important features:

Signal peptide:

amino acids 1-20

Type II transmembrane domain:

amino acids 11-31

Other transmembrane domain:

amino acids 57-77 and 123-143

3/392

FIGURE 3

GGCTCAGAGGCCCCACTGGACCCTCGGCTCTTCCTTGGACTTCTTGTGTGTTCTGTGAGC
TTCGCTGGATTCAAGGTCTTGGGCATCAGAGGTGAGAGGGTGGGAAGGTCCGCCGCGATG
GGGAAGCCCTGGCTGCGTGCGCTACAGCTGCTGCTCCTGCTGGGCGCGTCTGTGGGCGCGG
GCGGGCGCCCCGCGCTGCACCTACACCTTCGTGCTGCCCCCGCAGAAGTTCACGGGCGCT
GTGTGCTGGAGCGGCCCCGCATCCACGCGGGCGACGCCCCGAGGCCGCAACGCCAGCGAG
CTGGCGGCGCTGCGCATGCGCGTGGGCCGCCACGAGGAGCTGTTACGCGAGCTGCAGAGG
CTGGCGGCGGCCGACGGCGCCGTGGCCGGCGAGGTGCGCGCGCTGCGCAAGGAGAGCCGC
GGCCTGAGCGCGCGCCTGGGCCAGTTGCGCGCGCAGCTGCAGCACGAGGCGGGGCCCGGG
GCGGGCCCCGGGGCGGATCTGGGGGGCGGAGCCTGCCGCGGCGCTGGCGCTGCTCGGGGAG
CGCGTGCTCAACGCGTCCGCCGAGGCTCAGCGCGCAGCCGCCCGGTTCCACCAGCTGGAC
GTCAAGTTCGCGAGCTGGCGCAGCTCGTCACCCAGCAGAGCAGTCTCATCGCCCGCCTG
GAGCGCCTGTGCCCCGGGAGGCGGGCGGGCAGCAGCAGGTCTGCGGCCACCCCCACTG
GTGCCTGTGGTTCCGGTCCGTCTTGTGGGTAGCACACAGTGACACCAAGTAGGATGCTGGAC
CCAGCCCCAGAGCCCCAGAGAGACCAGACCCAGAGACAGCAGGAGCCCATGGCTTCTCCC
ATGCCTGCAGGTACCCCTGCGGTCCCCACCAAGCCTGTGGGCCCCGTGGCAGGATTGTGCA
GAGGCCCGCCAGGCAGGCCATGAACAGAGTGAGTGTATGAACTGCGAGTGGGCCGTGAC
GTAGTGTGAGTATGGTGTGAGCAGCAACTGGAGGGTGGAGGCTGGACTGTGATCCAGCGG
AGGCAAGATGGTTCAAGTCAACTTCTTCACTACCTGGCAGCACTATAAGGCGGGCTTTGGG
CGGCCAGACGGAGAATACTGGCTGGGCCTTGAACCCGTGTATCAGCTGACCAGCCGTGGG
GACCATGAGCTGCTGGTTCTCCTGGAGGACTGGGGGGGCGTGGAGCACGTGCCCACTAT
GATGGCTTCTCCCTGGAACCCGAGAGCGACCACTACCGCCTGCGGCTTGGCCAGTACCAT
GGTGATGCTGGAGACTCTCTTCTCCTGGCACAATGACAAGCCCTTCAGCACCGTGGATAGG
GACCGAGACTCCTATTCTGGTAACTGTGCCCTGTACCAGCGGGGAGGCTGGTGGTACCAT
GCCTGTGCCCCACTCCAACCTCAACGGTGTGTGGCACCACGGCGGCCACTACCGAAGCCGC
TACCAGGATGGTGTCTACTGGGCTGAGTTTCGTGGTGGGGCATATTCTCTCAGGAAGGCC
GCCATGCTCATTGCGCCCCCTGAAGCTGTGACTCTGTGTTCTCTGTCCCCTAGGCCCTAG
AGGACATTGGTCAGCAGGAGCCCCAAGTTGTTCTGGCCACACCTTCTTTGTGGCTCAGTGC
CAATGTGTCCACAGAACTTCCCACTGTGGATCTGTGACCCTGGGCGCTGAAAATGGGAC
CCAGGAATCCCCCGGTCAATATCTTGGCCTCAGATGGCTCCCCAAGGTCAATTCATATCT
CGGTTTGAGCTCATATCTTATAATAACACAAAGTAGCCAC

4/392

FIGURE 4

MGKPWLRALQLLLLLGASWARAGAPRCTYTFVLPPQKFTGAVCWSGPASTRATPEANAS
ELAALRMVRGRHEELLRELQRLAAADGAVAGEVRALRKESRGLSARLGQLRAQLQHEAGP
GAGPGADLGAEPAAALALLGERVLNASAEQAARFQHQLDVKFRELAQLVTQQSSLIAR
LERLCPPGAGGQQQVLPPPPLVPVVPVRLVGSTSDTSRMLDPAPEPQRDQTQRQQEPMAS
PMPAGHPAVPTKPVGPWQDCAEARQAGHEQSGVYELRVGRHVSVWCEQQLEGGGWTVIQ
RRQDGSVNFFTTWQHYKAGFGRPDGEYWLGLEPVYQLTSRGDHELLVLLEDWGGRGARAH
YDGFSLEPESDHYRLRLGQYHGDAGDSLWHNDKPFSTVDRDRDSYSGNCALYQRGGWY
HACAHSNLNGVWHHGGHYRSRYQDGVYWAEFRGGAYSLRKAAMLIRPLKL

Signal peptide:

Amino acids 1-20

N-glycosylation sites:

Amino acids 58-62;145-149

cAMP- and cGMP-dependent protein kinase phosphorylation site:

Amino acids 97-101

Tyrosine kinase phosphorylation site:

Amino acids 441-448

N-myristoylation sites:

Amino acids

16-22;23-29;87-93;108-114;121-127;125-131;129-135;187-193;293-299;353-359;378-384;445-451;453-459

Cell attachment sequence:

Amino acids 340-343

Fibrinogen beta and gamma chains C-terminal domain signature:

Amino acids 418-431

5/392

FIGURE 5

CCCACGCGTCCGGCGCCGTGGCCTCGCGTCCATCTTTGCCGTTCTCTCGGACCTGTCACA
AAGGAGTCGCGCCGCCGCCGCCGCCCTCCCTCCGGTGGGCCCCGGGAGGTAGAGAAAGT
CAGTGCCACAGCCCGACCGCGCTGCTCTGAGCCCTGGGCACGCGGAACGGGAGGGAGTCT
GAGGGTTGGGGACGTCTGTGAGGGAGGGGAACAGCCGCTCGAGCCTGGGGCGGGCGGACC
GGACTGGGGCCGGGGTAGGCTCTGGAAAGGGCCCCGGGAGAGAGGTGGCGTTGGTCAGAAC
CTGAGAAAACAGCCGAGAGGTTTTCCACCGAGGCCCGCGCTTGAGGGATCTGAAGAGGTTC
CTAGAAGAGGGTGTTCCCTCTTTCCGGGGGTCTCACCAGAAGAGGTTCTTGGGGGTCCGC
CTTCTGAGGAGGCTGCGGCTAACAGGGCCCAGAACTGCCATTGGATGTCCAGAATCCCCCT
GTAGTTGATAATGTTGGGAATAAGCTCTGCAACTTTCTTTGGCATTTCAGTTGTTAAAAAC
AAATAGGATGCAAATTCCTCAACTCCAGGTTATGAAAACAGTACTTGGAAAACGAAAAC
TACCTAAATGATCGTCTTTGGTTGGGCCGTGTTCTTAGCGAGCAGAAGCCTTGGCCAGGG
TCTGTTGTTGACTCTCGAAGAGCACATAGCCCACTTCCTAGGGACTGGAGGTGCCGCTAC
TACCATGGGTAATTCCTGTATCTGCCGAGATGACAGTGGAACAGATGACAGTGTGACAC
CCAACAGCAACAGGCCGAGAACAGTGCAGTACCCACTGCTGACACAAGGAGCCAACCACG
GGACCCTGTTCCGGCCACCAAGGAGGGGCGGAGGACCTCATGAGCCAAGGAGAAAGAAACA
AAATGTGGATGGGCTAGTGTTGGACACACTGGCAGTAATACGGACTCTTGTAAGATAAGTA
AGTATCTGACTCACGGTCACCTCCAGTGGAATGAAAAGTGTTCTGCCCGGAACCATGACT
TTAGGACTCCTTCAGTTCCTTTAGGACATACTCGCCAAGCCTTGTTGCTCACAGGGCAAAG
GAGAATATTTTAATGCTCCGCTGATGGCAGAGTAAATGATAAGATTTGATGTTTTTGCTT
GCTGTCATCTACTTTGTCTGGAAATGTCTAAATGTTTCTGTAGCAGAAAACACGATAAAG
CTATGATCTTTATTAGAG

6/392

FIGURE 6

MIVFGWAVFLASRSLGQGLLLTLEEHIAHFLGTGGAATTMGNSCICRDDSGTDDSVDTQQ
QQAENSAVPTADTRSQPRDPVRPPRRGRGPHEPRRKKQNV DGLVLDTLAVIRTLVDKO

Signal peptide:
amino acids 1-16

Casein kinase II phosphorylation site:
amino acids 22-26, 50-54, 113-117

N-myristoylation site:
amino acids 18-24, 32-38, 34-40, 35-41, 51-57

7/392

FIGURE 7

CGGACGCGTGGGGGAAACCCCTTCCGAGAAAACAGCAACAAGCTGAGCTGCTGTGACAGAG
GGGAACAAGATGGCGGCGCCGAAGGGGAGCCTCTGGGTGAGGACCCAACTGGGGCTCCCG
CCGCTGCTGCTGCTGACCATGGCCTTGGCCGGAGGTTCTGGGGACCGCTTCGGCTGAAGCA
TTTGAATCGGTCTTGGGTGATACGGCGTCTTGCCACCGGGCCTGTCAGTTGACCTACCCC
TTGCACACCTACCCTAAGGAAGAGGAGTTGTACGCATGTCAGAGAGGTTGCAGGCTGTTT
TCAATTTGTCAGTTTGTGGATGATGGAATTGACTTAAATCGAACTAAATTGGAATGTGAA
TCTGCATGTACAGAAGCATATTCCTAATCTGATGAGCAATATGCTTGCCATCTTGTTGC
CAGAATCAGCTGCCATTCGCTGAACTGAGACAAGAACAACCTTATGTCCCTGATGCCAAAA
ATGCACCTACTCTTTCTCTAATCTGGTGAGGTCATTCTGGAGTGACATGATGGACTCC
GCACAGAGCTTCATAACCTCTTCATGGACTTTTTATCTTCAAGCCGATGACGGAAAAATA
GTTATATTCCAGTCTAAGCCAGAAATCCAGTACGCACCACATTTGGAGCAGGAGCCTACA
AATTTGAGAGAATCATCTCTAAGCAAAATGTCCTATCTGCAAATGAGAAATTCACAAGCG
CACAGGAATTTTCTTGAAGATGGAGAAAGTGATGGCTTTTTAAGATGCCTCTCTCTTAAC
TCTGGGTGGATTTTAACTACAACCTCTTGTCCTCTCGGTGATGGTATTGCTTTGGATTTGT
TGTGCAACTGTTGCTACAGCTGTGGAGCAGTATGTTCCCTCTGAGAAGCTGAGTATCTAT
GGTGACTTGGAGTTTATGAATGAACAAAAGCTAAACAGATATCCAGCTTCTTCTCTTG
GTTGTTAGATCTAAAAGTGAAGATCATGAAGAAGCAGGGCCTCTACCTACAAAAGTGAAT
CTTGCTCATTCTGAAATTTAAGCATTTTTCTTTTAAAAGACAAGTGTAATAGACATCTAA
AATTCCTCCTCATAGAGCTTTTAAATGGTTTCATTGGATATAGGCCTTAAGAAATCA
CTATAAAATGCAAATAAAGTTACTCAAATCTGTG

8/392

FIGURE 8

MAAPKGSLWVRTQLGLPPLLLLTMALAGGSGTASAEAFDSVLGDTASCHRAQCLTYPLHT
YPKEEELYACQRCRLFSICQFVDDGIDLNRKLECESECTEAYSQSDEQYACHLGCONQ
LPFAELRQEQQLMSLMPKMHLFFPLTLVRSFWSMDMSAQSFITSSWTFYLQADDGKIVIF
QSKPEIQYAPHLEQEPTNLRESSLSKMSYLMRNSQAHRNFLEDGESDGFLRCLSLNSGW
ILTTTLVLVSMVLLWICCATVATAVEQYVPSEKLSIYGDLEFMNEQKLNRYPASSLVVVR
SKTEDHEEAGPLPTKVNLAHSEI

Important features:

Signal peptide:

amino acids 1-31

Transmembrane domain:

amino acids 241-260

N-glycosylation site:

amino acids 90-93

9/392

FIGURE 9

TATTTACCATATCAGATTCACATTCAGTCCTCAGCAAAATGAAGGGCTCCATTTTCACTC
TGTTTTTATTCTCTGTCTTATTTGCCATCTCAGAAGTGCGGAGCAAGGAGTCTGTGAGAC
TCTGTGGGCTAGAATACATACGGACAGTCATCTATATCTGTGCTAGCTCCAGGTGGAGAA
GGCATCTGGAGGGGATCCCTCAAGCTCAGCAAGCTGAGACAGGAAACTCCTTCCAGCTCC
CACATAAACGTGAGTTTTCTGAGGAAAATCCAGCGCAAAACCTTCCGAAGGTGGATGCCT
CAGGGGAAGACCGTCTTTGGGGTGGACAGATGCCCACTGAAGAGCTTTGGAAGTCAAAGA
AGCATTCAAGTATGTCAAGACAAGATTTACAAACTTTGTGTTGCACTGATGGCTGTTCCA
TGAATGATTTGAGTGCTCTTTGCTAAGACAAGAGCAAATACCCAATGGGTGGCAGAGCTT
TATCACATGTTTAATTACAGTGTTTTACTGCCTGGTAGAACACTAATATTGTGTTATTAA
AATGATGGCTTTTGGGTAGGCAAACTTCTTTTCTAAAAGGTATAGCTGAGCGGTTGAAA
CCACAGTGATCTCTATTTTCTCCCTTTGCCAAGGTTAATGAACTGTTCTTTCAAATTCT
ACTAATGCTTTGAAATTTCAAATGCTGCGCAAAATTGCAATAAAAATGCTATAAA

10/392

FIGURE 10

MKGSIFTLFLFSVLFAISEVRSKESVRLCGLEYIRTVIYICASSRWRRLHLEGIPQAQQAE
TGNSFQLPHKREFSEENPAQNLPKVDASGEDRLWGGQMPTEELWKSCKHVSMSRQDLQTL
CCTDGCSMTDLSALC

Important features:

Signal sequence:

amino acids 1-18

**cAMP- and cGMP-dependent protein kinase phosphorylation
site:**

amino acids 107-111

N-myristoylation sites:

amino acids 3-9, 52-58, 96-102, 125-131

Insulin family signature:

amino acids 121-136

Insulin family proteins:

amino acids 28-46

11/392

FIGURE 11

CCCACGCGTCCGGACAAACTGGAGGTGAAAGGAGCTGGTACTGTCCACTGTGCTGTCCGGT
GCTGAACCTGAGACGCGAGCGGACCAGTTGCTCCAGCACCTGAAGGCAACGCCCTCTTGC
ACCCCTCTGTGCCCTGTGGGACCCGCTTCACCAACAGGACCCATATCAACTTGACAAAGGA
GTGTGGTATCGGACGTGGGAGAGAGTCTCTGTTTGCCACCTGGGCGCTCATTTCAGGCGT
GACTTTGGAGATTTCTATAGTTTTAGACCAAATATTTTTTTTTTCCCAGCTAAGACGAT
CTTTTGAGAGTTTTTTTTTTTTTATTGTGATTTATATTTCCACAGCGTTTAGGAATCTTCT
GGGGGACTTTTGTGACTGTTAAAATAAGGTGAAAAGCAATAAGGATGTTTAAGTGCTGGT
CAGTTGTCTTGGTTCTCGGATTCATTTTTCTGGAGTCGGAAGGAAGGCCAACCAAAGAAG
GAGGATATGGCCTTAAATCCTATCAGCCTCTAATGAGATTGCGACATAAGCAGGAAAAAA
ATCAAGAAAGTTCAAGAGTCAAAGGATTTATGATTTCAGGATGGCCCTTTTGGATCTTGTG
AAAATAAGTACTGTGGTTTGGGAAGACACTGTGTTACCAGCAGAGAGACAGGGCAAGCAG
AATGTGCCTGTATGGACCTTTGCAAACGTCACCTACAAACCTGTGTGTGGATCTGACGGAG
AATTCTATGAAAACCACTGTGAAGTGCACAGAGCTGCTTGCCCTGAAAAAACAAAAGATTA
CCATTGTTTACAATGAAGACTGCTTCTTTAAAGGAGATAAGTGCAAGACTACTGAATACA
GCAAGATGAAAAATATGCTATTAGATTTACAAAATCAAAAATATATTATGCAAGAAAATG
AAAATCCTAATGGCGACGACATATCTCGGAAGAAGCTATTGGTGGATCAAATGTTTAAAT
ATTTTGATGCAGACAGTAATGGACTTGTAGATATTAATGAACTAACTCAGGTGATAAAAC
AGGAAGAACTTGGCAAGGATCTCTTTGATTGTACTTTGTATGTTCTATTGAAATATGATG
ATTTTAATGCTGACAAGCACCTGGCTCTTGAAGAATTTTATAGAGCATTCCAAGTGATCC
AGTTGAGTCTGCCAGAAGATCAGAACTAAGCATCACTGCAGCAACTGTGGGACAAAGTG
CTGTTCTGAGCTGTGCCATTCAAGGAACCTGAGACCTCCCATTATCTGGAAAAGGAACA
ATATTATTCTAAATAATTTAGATTTGGAAGACATCAATGACTTTGGAGATGATGGGTCCT
TGTATATTACTAAGGTTACCACAACCTCACGTTGGCAATTACACCTGCTATGCAGATGGCT
ATGAACAAGTCTATCAGACTCACATCTTCCAAGTGAATGTTCTCCAGTCATCC

12/392

FIGURE 12

MFKCWSVVLVLGFIFLESEGRPTKEGGYGLKSYQPLMRLRHKQEKQESSRVKGFMIQDG
PFGSCENKYCGLGRHCVTSRETGQAECACMDLCKRHYKPVCSDGEFYENHCEVHRAACL
KKQKITIVHNEDCFFKGDCKTTEYSKMKNMLLDLQNQKYIMQENENPNGDDISRKKLLV
DQMFKYFDADSNGLVDINELTQVIKQEELGKDLFDCTLYVLLKYDDFNADKHLALEEFYR
AFQVIQLSLPEDQKLSITAATVGQSAVLSCAIQGTLRPPIIWKRNNIILNLDLEDINDF
GDDGSLYITKVTTHVGNVTCYADGYEQVYQTHIFQVNVPPVI

Signal sequence:

Amino acids 1-20

N-glycosylation site:

Amino acids 318-322

Tyrosine kinase phosphorylation sites:

Amino acids 21-29;211-220

N-myristoylation sites:

Amino acids 63-69;83-89;317-323

Prokaryotic membrane lipoprotein lipid attachment site:

Amino acids 260-271

13/392

FIGURE 13

TGCCGGGCTGCGGGGCGCCTTGACTCTCCCTCCACCCTGCCTCCTCGGGCTCCACTCGTC
TGCCCCCTGGACTCCCGTCTCCTCCTGTCTCCGGCTTCCCAGAGCTCCCTCCTTATGGCA
GCAGCTTCCCGCGTCTCCGGCGCAGCTTCTCAGCGGACGACCCTCTCGCTCCGGGGCTGA
GCCCAGTCCCTGGATGTTGCTGAAACTCTCGAGATCATGCGCGGGTTTGGCTGCTGCTTC
CCCGCCGGGTGCCACTGCCACCGCCCGCCGCTCTGCTGCCGCGGTCCGCGGGATGCTCAG
TAGCCCGCTGCCCGGCCCCCGCGATCCTGTGTTCTCGGAAGCCGTTTGCTGCTGCAGAG
TTGCACGAAC TAGTCATGGTGCTGTGGGAGTCCCCGCGGCAGTGCAGCAGCTGGACACTT
TGCGAGGGCTTTTGCTGGCTGCTGCTGCTGCCCGTCATGCTACTCATCGTAGCCCCCG
GTGAAGCTCGCTGCTTTCCCTACCTCCTTAAGTGACTGCCAAACGCCACCGGCTGGAAT
TGCTCTGGTTATGATGACAGAGAAAATGATCTCTTCTCTGTGACACCAACACCTGTAAA
TTTGATGGGGAATGTTTAAGAATTGGAGACACTGTGACTTGCGTCTGTCAAGTCAAGTGC
AACATGACTATGTGCCTGTGTGTGGCTCCAATGGGGAGAGCTACCAGAATGAGTGTTAC
CTGCGACAGGCTGCATGCAAACAGCAGAGTGAGATACTTGTGGTGTCAGAAGGATCATGT
GCCACAGATGCAGGATCAGGATCTGGAGATGGAGTCCATGAAGGCTCTGGAGAACTAGT
CAAAAGGAGACATCCACCTGTGATATTTGCCAGTTTGGTGCGAATGTGACGAAGATGCC
GAGGATGTCTGGTGTGTGTGTAATATTGACTGTTCTCAAACCAACTTCAATCCCCTCTGC
GCTTCTGATGGGAAATCTTATGATAATGCATGCCAAATCAAAGAAGCATCGTGTCAGAAA
CAGGAGAAAATTGAAGTCATGTCTTTGGGTCGATGTCAAGATAACACA ACTACA ACTACT
AAGTCTGAAGATGGGCATTATGCAAGAACAGATTATGCAGAGAATGCTAACAAATTAGAA
GAAAGTGCCAGAGAACACCACATACCTTGTCCGGAACATTACAATGGCTTCTGCATGCAT
GGGAAGTGTGAGCATTCTATCAATATGCAGGAGCCATCTTGCAGGTGTGATGCTGGTTAT
ACTGGACAACACTGTGAAAAAAGGACTACAGTGTTCTATACGTTGTTCCCGGTCCTGTA
CGATTTTCAGTATGTCTTAATCGCAGCTGTGATTGGAACAATTCAGATTGCTGTCTGT
GTGGTGGTCCTCTGCATCACAAGGAAATGCCCCAGAAGCAACAGAATTCACAGACAGAAG
CAAAATACAGGGCACTACAGTTCAGACAATACAACAAGAGCGTCCACGAGGTTAATCTAA
AGGGAGCATGTTTCACAGTGGCTGGACTACCGAGAGCTTGGACTACACAATACAGTATTA
TAGACAAAAGAATAAGACAAGAGATCTACACATGTTGCCTTGCATTTGTGGTAATCTACA
CCAATGAAAACATGTACTACAGCTATATTTGATTATGTATGGATATATTTGAAATAGTAT
ACATTGTCTTGATGTTTTTTCTGTAATGTAAATAAACTATTTATATCACACAATATAGTT
TTTTCTTCCCATGTATTTGTTATATATAATAAATACTCAGTGATGAG

14/392

FIGURE 14

MVLWESPRQCSSWTLCGFCWLLLLPVMLLIVARPVKLAAFPSTLSDCQTPTGWNC SGY
DDRENDLFLCDTNTCKFDGECLRIGDTVTCVCQFKCNNDYVPVCGSNGESYQNECYLRQ
AACKQQSEILVVSEGSCATDAGSGSGDGVHEGSGETSQKETSTCDICQFGAECDEDAED
VWCVCNIDCSQTNFNPLCASDGKSYDNACQIKEASCQKQEKIEVMSLGRCQDNTTTT TK
SEDGHYARTDYAENANKLEESAREHHIPCPEHYNGFCMHGKCEHSINMQEPSCRC DAGY
TGQHCEKKDYSVLYVVP GPVRFQYVLIAAVIGTIQIAVICVVVLCITRKCPRSNRIHRQ
KQNTGHYSSDNTTRASTRLI

15/392

FIGURE 15

GGAGCTCAGCCCAGTGGGCAGTCTGAAGATGGCCAATTACACGCTGGCACCAGAGGATGA
ATATGATGTCCTCATAGAAGGTGAACTGGAGAGCGATGAGGCAGAGCAATGTGACAAGTA
TGACGCCCAGGCACTCTCAGCCCAGCTGGTGCCATCACTCTGCTCTGCTGTGTTTGTGAT
CGGTGTCCTGGACAATCTCCTGGTTGTGCTTATCCTGGTAAAATATAAAGGACTCAAACG
CGTGAAAAATATCTATCTTCTAAACTTGGCAGTTTCTAACTTGTGTTTCTTGCTTACCCT
GCCCTTCTGGGCTCATGCTGGGGGCGATCCCATGTGTAAAATTCTCATTGGACTGTACTT
CGTGGGCCTGTACAGTGAGACATTTTTCAATTGCCTTCTGACTGTGCAAAGGTACCTAGT
GTTTTTGCACAAGGGCAACTTTTTCTCAGCCAGGAGGAGGGTGCCCTGTGGCATCATTAC
AAGTGTCTTGGCATGGGTAAACAGCCATTCTGGCCACTTTGCCTGAATACGTGGTTTTATAA
ACCTCAGATGGAAGACCAGAAATACAAGTGTGCATTTAGCAGAACTCCCTTCCTGCCAGC
TGATGAGACATTCTGGAAGCATTTTTCTGACTTTAAAAATGAACATTTCCGTTCTTGTCTCT
CCCCCTATTTATTTTTACATTTCTCTATGTGCAAATGAGAAAAACACTAAGGTTCAAGGA
GCAGAGGTATAGCCTTTTCAAGCTTGTTTTTGCCATAATGGTAGTCTTCCCTTCTGATGTG
GGCGCCCTACAATATTGCATTTTTCTGTCCACTTTCAAAGAACACTTCTCCCTGAGTGA
CTGCAAGAGCAGCTACAATCTGGACAAAAGTGTTTACATCACTAAACTCATCGCCACCAC
CCACTGCTGCATCAACCCTCTCCTGTATGCGTTTCTTGATGGGACATTTAGCAAATACCT
CTGCCGCTGTTTCCATCTGCGTAGTAACACCCCACTTCAACCCAGGGGGCAGTCTGCACA
AGGCACATCGAGGGAAGAACCTGACCATTCCACCGAAGTGTA~~AA~~ACTAGCATCCACCAAAT
GCAAGAAGAATAAACATGGATTTTCATCTTCTGCATTATTTTCATGTAAATTTTCTACAC
ATTTGTATACAAAATCGGATACAGGAAGAAAAGGGAGAGGTGAGCTAACATTTGCTAAGC
ACTGAATTTGTCTCAGGCACCGTGCAAGGCTCTTTACAAACGTGAGCTCCTTCGCCTCCT
ACCACTTGTCCATAGTGTGGATAGGACTAGTCTCATTTCTCTGAGAAGAAAATAAGGCG
CGGAAATTTGTCTAAGATCACATAACTAGGAAGTGGCAGAACTGATTCTCCAGCCCTGGT
AGCATTTGCTCAGAGCCTACGCTTGGTCCAGAACATCAAACCTCAAACCCTGGGGACAAA
CGACATGAAATAAATGTATTTTAAAACATCTAAAA

16/392

FIGURE 16

MANYTLAPEDEYDVLIEGELESDEAEQCDKYDAQALSAQLVPSLCSAVFVIGVLDNLLVV
LILVKYKGLKRVENIYLLNLAVSNLCFLLTLPFWAHAGGDEPMCKILIGLYFVGLYSETFF
NCLLTVQRYLVFLHKGNNFSARRRVPCGIITSVLAWVTAILATLPEYVVYKPQMEDQKYK
CAFSRTPFLPADETFWKHFLTLKMNISVLVLPLFIPTFLYVQMRKTLRFREQRYSLFKLV
FAIMVVFLLMWAPYNIAFFLSTFKEHFSLSDCKSSYNLDKSVHITKLIATTHCCINPLLY
AFLDGTFSKYLRCRCHLRNTPLOPRGQSAQGTSREEPDHSTEV

Signal sequence:

None

Transmembrane domain:

41-61, 76-96, 109-129, 147-167, 199-219, 237-257, 285-305

7 transmembrane receptor (rhodopsin family):

55-300

N-glycosylation site:

3-6, 205-208

Tyrosine kinase phosphorylation site:

70-76, 171-179, 228-234

N-myristoylation site:

52-57, 136-141, 148-153

G-protein coupled receptors:

55-85, 96-136, 209-220, 235-254, 292-308

17/392

FIGURE 17

CGGACGCGTGGGCGGACGCGTGGGCGGCCACGGCGCCCGCGGGCTGGGGCGGTGCGTTCTTCCTTCTCCGTGGCCTACGAGGGTCCCCAGCCTGGGTAAAGATGGCCCCATGGCCCCCGAAGGGCCTAGTCCCAGCTGTGCTCTGGGGCCTCAGCCTCTTCCTCAACCTCCCAGGACCTATCTGGCTCCAGCCCTCTCCACCTCCCCAGTCTTCTCCCCCGCCTCAGCCCCATCCGTGTCATACCTGCCGGGGACTGGTTGACAGCTTTAAACAAGGGCCTGGAGAGAACCATCCGGGACAACCTTGGAGGTGGAAACACTGCCTGGGAGGAAGAGAATTTGTCCAAATACAAAGACAGTGAGACCCGCTGGTAGAGGTGCTGGAGGGTGTGTGCAGCAAGTCAGACTTCGAGTGCCACCGCCTGCTGGAGCTGAGTGAGGAGCTGGTGGAGAGCTGGTGGTTTCACAAGCAGCAGGAGGCCCCGGACCTCTTCCAGTGGCTGTGCTCAGATTCCCTGAAGCTCTGCTGCCCCGCAGGCACCTTCGGGGCCCTCCTGCCCTTCCCTGTCTGGGGGAACAGAGAGGCCCTGCGGTGGCTACGGCAGTGTGAAGGAGAAGGGACACGAGGGGCAGCGGGCACTGTGACTGCCAAGCCGGCTACGGGGGTGAGGCCTGTGGCCAGTGTGGCCCTTGGCTACTTTGAGGCAGAACGCAACGCCAGCCATCTGGTATGTTTCGGCTTGTTTGGCCCCCTGTGCCCGATGCTCAGGACCTGAGGAATCAAACTGTTTGCAATGCAAGAAGGGCTGGGCCCTGCATCACCTCAAGTGTGTAGACATTGATGAGTGTGGCACAGAGGGAGCCAACCTGTGGAGCTGACCAATTCTGCGTGAACACTGAGGGCTCCTATGAGTGCCGAGACTGTGCCAAGGCCTGCCTAGGCTGCATGGGGGCAGGGCCAAGTTCGCTGTAAGAAGTGTAGCCCTGGCTATCAGCAGGTGGGCTCCAAGTGTCTCGATGTGATGAGTGTGAGACAGAGGTGTGTCCGGGAGAGAACAAGCAGTGTGAAAACACCGAGGGCGTTATCGCTGCATCTGTGCCGAGGGCTACAAGCAGATGGAAGGCATCTGTGTGAAGGAGCAGATCCCAGAGTCAGCAGGCTTCTTCTCAGAGATGACAGAAGACGAGTTGGTGGTGTGTCAGCAGATGTTCTTTGGCATCATCATCTGTGCACTGGCCACGCTGGCTGCTAAGGGCGACTTGGTGTTCACCGCCATCTTCATTGGGGCTGTGGCGGCCATGACTGGCTACTGGTTGTCAAGCGCAGTGACCGTGTGCTGGAGGGCTTCATCAAGGGCAGATAATCGCGGCCACCACCTGTAGGACCTCCTCCCACCCACGCTGCCCCCAGAGCTTGGGCTGCCCTCCTGCTGGACACTCAGGACAGCTTGGTTTATTTTTGAGAGTGGGGTAAGCACCCCTACCTGCCTTACAGAGCAGCCCCAGGTACCCAGGCCCCGGGCAGACAAGGCCCTGGGGTAAAAAGTAGCCCTGAAGGTGGATACCATGAGCTCTTCACCTGGCGGGGACTGGCAGGCTTCACAATGTGTGAATTTCAAAAGTTTTTCCTTAATGGTGGCTGCTAGAGCTTTGGCCCCCTGCTTAGGATTAGGTGGTCCTCACAGGGGTGGGGCCATCACAGCTCCCTCCTGCCAGCTGCATGCTGCCAGTTCCTGTCTGTGTTTACCACATCCCCACCCCCATTGCCACTTATTTATTCATCTCAGGAAATAAAGAAAGGTCTTGAAAGTTAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

18/392

FIGURE 18

MAPWPPKGLVPAVLWGLSLFLNLP GPIWLQPSPPPQSSPPPQPHPCHTCRGLVDSFNKGL
ERTIRDNFGGGNTAWEEENLSKYKDSETRLVEVLEGVCSKSDFECHRLELSEELVESWW
FHKQQEAPDLFQWLCSDSLKLCCPAGTFGPSCLPCPGGTERPCGGYGQCEGEGTRGGSGH
CDCQAGYGGEACGQCGLGYFEAERNASHLVCSACFGPCARCSGPPEESNCLQCKKGWALHH
LKCVDIDECEGTEGANCGADQFCVNTEGSYECRCAKACLGCMGAGPGRCKKCSPGYQQVG
SKCLDVDECETEVC PGENKQCENTEGGYRCICAEGYKQMEGICVKEQIPESAGFFSEMTE
DELVLQQMFFGIIICALATLAAKGDLVFTAIFIGAVAAMTGYWLSERSDRVLEGFIKGR

19/392

FIGURE 19

GCCCCGGGACTGGCGCAAGGTGCCCAAGCAAGGAAAGAAATAATGAAGAGACACATGTGTT
AGCTGCAGCCTTTTGAACACGCAAGAAGGAAATCAATAGTGTGGACAGGGCTGGAACCT
TTACCACGCTTGTTGGAGTAGATGAGGAATGGGCTCGTGATTATGCTGACATTCCAGCAT
GAATCTGGTAGACCTGTGGTTAACCCGTTCCCTCTCCATGTGTCTCCTCCTACAAAGTTT
TGTTCTTATGATACTGTGCTTTTCATTCTGCCAGTATGTGTCCCAAGGGCTGTCTTTGTTC
TTCCTCTGGGGGTTTAAATGTACCTGTAGCAATGCAAAATCTCAAGGAAATACCTAGAGA
TCTTCCTCCTGAAACAGTCTTACTGTATCTGGACTCCAATCAGATCACATCTATTCCCAA
TGAAATTTTTTAAGGACCTCCATCAACTGAGAGTTCTCAACCTGTCCAAAAATGGCATTGA
GTTTATCGATGAGCATGCCTTCAAAGGAGTAGCTGAAACCTTGCAGACTCTGGACTTGTC
CGACAATCGGATTCAAAGTGTGCACAAAAATGCCTTCAATAACCTGAAGGCCAGGGCCAG
AATTGCCAACAACCCCTGGCACTGCGACTGTACTCTACAGCAAGTTCTGAGGAGCATGGC
GTCCAATCATGAGACAGCCCACAACGTGATCTGTAAAACGTCCGTGTTGGATGAACATGC
TGGCAGACCATTCTCAATGCTGCCAACGACGCTGACCTTTGTAACCTCCCTAAAAAAC
TACCGATTATGCCATGCTGGTCACCATGTTTGGCTGGTTCACTATGGTGATCTCATATGT
GGTATATTATGTGAGGCAAAATCAGGAGGATGCCCGGAGACACCTCGAATACTTGAAATC
CCTGCCAAGCAGGCAGAAGAAAGCAGATGAACCTGATGATATTAGCACTGTGGTATAGTG
TCCAAACTGACTGTCAATTGAGAAAGAAAGAAAGTAGTTTGCGATTGCAGTAGAAATAAGT
GGTTTACTTCTCCCATCCATTGTAAACATTTGAAACTTTGTATTTTCAAGTTTTTTTGAAT
TATGCCACTGCTGAACTTTTAAACAAACACTACAACATAAATAATTTGAGTTTAGGTGATC
CACCCCTTAATTGTACCCCGATGGTATATTTCTGAGTAAGCTACTATCTGAACATTAGT
TAGATCCATCTCACTATTTAATAATGAAATTTATTTTTTTAATTTAAAAGCAAATAAAAG
CTTAACTTTGAACCATGGGAAAAAAAAAAAAAAAAAAAAAAAAACA

20/392

FIGURE 20

MNLVDLWLTRSLSMCLLLQSFVLMILCFHSASMC PKGCLCSSSGGLNVTCSNANLKEIPR
DLPPETVLLYLD SNQITSIPNEIFKDLHQLRVLNLSKNGIEFIDEHAFKGVAETLQTLDL
SDNRIQSVHKNAFNNLKARARIANNPWHCDCTLQQVLRSMASNHETAHN VICKTSVLDEH
AGRPFLNAANDADLCNLPKKT TDYAMLVTMFGWFTMVISYVVYYVRQNQEDARRHLEYLK
SLPSRQKKADEPDDISTVV

Signal sequence:
amino acids 1-33

Transmembrane domain:
amino acids 205-220

N-glycosylation site:
amino acids 47-51, 94-98

cAMP- and cGMP-dependent protein kinase phosphorylation site:
amino acids 199-203

Casein kinase II phosphorylation site:
amino acids 162-166, 175-179

N-myristoylation site:
amino acids 37-43, 45-51, 110-116

21/392

FIGURE 21

CGCCACCACTGCGGCCACCGCCAATGAAACGCCTCCCGCTCCTAGTGGTTTTTTCCACTT
TGTTGAATTGTTTCTTATACTCAAAATTGCACCAAGACACCTTGTCTCCCAAATGCAAAAT
GTGAAATACGCAATGGAATTGAAGCCTGCTATTGCAACATGGGATTTTCAGGAAATGGTG
TCACAATTTGTGAAGATGATAATGAATGTGGAAATTTAACTCAGTCCTGTGGCGAAAATG
CTAATTGCACTAACACAGAAGGAAGTTATTATTGTATGTGTGTACCTGGCTTCAGATCCA
GCAGTAACCAAGACAGGTTTATCACTAATGATGGAACCGTCTGTATAGAAAATGTGAATG
CAAACCTGCCATTTAGATAATGTCTGTATAGCTGCAAAATATTAATAAACTTTAACAAAA
TCAGATCCATAAAAGAACCTGTGGCTTTGCTACAAGAAGTCTATAGAAATTCTGTGACAG
ATCTTTTACCAACAGATATAATTACATATATAGAAATATTAGCTGAATCATCTTCATTAC
TAGGTTACAAGAACAACACTATCTCAGCCAAGGACACCCTTTCTAACTCAACTCTTACTG
AATTTGTAAAAACCGTGAATAAATTTGTTCAAAGGGATACATTTGTAGTTTGGGACAAGT
TATCTGTGAATCATAGGAGAACACATCTTACAAAACCTCATGCACACTGTTGAACAAGCTA
CTTTAAGGATATCCCAGAGCTTCCAAAAGACCACAGAGTTTGATACAAATTCACCGGATA
TAGCTCTCAAAGTTTTCTTTTTTGATTTCATATAACATGAAACATATTCATCCTCATATGA
ATATGGATGGAGACTACATAAATATATTTCCAAAAGAGAAAAGCTGCATATGATTCAAATG
GCAATGTTGCAGTTGCATTTTTATATTATAAGAGTATTGGTCCTTTGCTTTTCATCATCTG
ACAACCTTCTTATTGAAACCTCAAATTTATGATAATTCTGAAGAGGAGGAAAGAGTCATAT
CTTCAGTAATTTTCACTCTCAATGAGCTCAAACCCACCCACATTATATGAACTTGAAAAAA
TAACATTTACATTAAAGTCATCGAAAGGTCACAGATAGGTATAGGAGTCTATGTGCATTTT
GGAATTACTCACCTGATACCATGAATGGCAGCTGGTCTTCAGAGGGCTGTGAGCTGACAT
ACTCAAATGAGACCCACACCTCATGCCGCTGTAATCACCTGACACATTTTGCAATTTTGA
TGTCCTCTGGTCCCTTCATTGGTATTAAAGATTATAATATTCTTACAAGGATCACTCAAC
TAGGAATAATTATTTCACTGATTTGTCTTGCCATATGCATTTTTTACCTTCTGGTTCTTCA
GTGAAATTCAAAGCACCAGGACAACAATTCACAAAAATCTTTGCTGTAGCCTATTTCTTG
CTGAACTTGTTTTCTTGTGGGATCAATACAAATACTAATAAGCTCTTCTGTTCAATCA
TTGCCGGAAGTGTACACTACTTCTTTTTTAGCTGCTTTTGCATGGATGTGCATTGAAGGCA
TACATCTCTATCTCATTGTTGTGGGTGTCTATACAACAAGGGATTTTTGCACAAGAATT
TTTATATCTTTGGCTATCTAAGCCCAGCCGTGGTAGTTGGATTTTCGGCAGCACTAGGAT
ACAGATATTATGGCACAACCAAGTATGTTGGCTTAGCACCGAAAAACAATTTATTTGGA
GTTTTATAGGACCAGCATGCCTAATCATTCTTGTTAATCTCTTGGCTTTTGGAGTCATCA
TATACAAAGTTTTTCGTCACTGCAGGGTTGAAACCAGAAAGTTAGTTGCTTTGAGAAC
TAAGGTCTTGTGCAAGAGGAGCCCTCGCTCTTCTGTTCTTCTCGGCACCACCTGGATCT
TTGGGGTTCTCCATGTTGTGCACGCATCAGTGGTTACAGCTTACCTCTTCACAGTCAGCA
ATGCTTTCCAGGGGATGTTCAATTTTTTATTCCTGTGTGTTTTATCTAGAAAGATTCAAG
AAGAATATTACAGATTGTTCAAAAATGTCCCTGTTGTTTTGGATGTTTAAGGTAAACAT
AGAGAATGGTGGATAATTACAACCTGCACAAAAATAAAAAATCCAAGCTGTGGATGACCAA
TGTATAAAAAATGACTCATCAAATTATCCAATTATTAACCTACTAGACAAAAAGTATTTTAA
ATCAGTTTTTCTGTTTATGCTATAGGAACTGTAGATAATAAGGTAAAAATTATGTATCATA
TAGATATACTATGTTTTTCTATGTGAAATAGTTCTGTCAAAAATAGTATTGCAGATATTT
GGAAAGTAATTGGTTTTCTCAGGAGTGATATCACTGCACCCAAGGAAAGATTTCTTTCTA
ACACGAGAAGTATATGAATGTCTGAAGGAAACCACTGGCTTGATATTTCTGTGACTCGT
GTTGCCTTTGAACTAGTCCCCTACCACCTCGGTAATGAGCTCCATTACAGAAAGTGGA
CATAAGAGAATGAAGGGGCAGAAATATCAACAGTGAAAAGGGAATGATAAGATGTATTTT
GAATGAACTGTTTTTCTGTAGACTAGCTGAGAAATTGTTGACATAAAATAAAGAATTGA
AGAAACACATTTTACCATTTTGTGAATTGTTCTGAACTTAAATGTCCACTAAAACAACCTT
AGACTTCTGTTTGTCTAAATCTGTTTCTTTTTCTAATATTCTAAAAAAGGTTT
ACCTCCACAAATTGAAA

22/392

FIGURE 22

MKRLPLLVVVFSTLLNCSYTQNCTKTPCLPNAKCEIRNGIEACYCNMGFSGNGVTICEDDN
ECGNLTQSCGENANCTNTEGSYYCMCVPGFRSSSNQDRFITNDGTVCIENTVNANCHLDNV
CIAANINKTLTKIRSIKEPVALLQEVYRNSVTDLSPTDIITYIEILAESSSLGYKNNTI
SAKDTLSNSTLTFVKTVNNFVQRDTFVVWDKLSVNHRRTHLTKLMHTVEQATLRISQS
QKTTEFDTNSTDIALKVFFFDSDYNMKHIHPHMNMDGDYINIFPKRKAAYDSNGNVAVAF
YYKSIGPLLSSSDNFLLKPQNYDNSEEEERVISSVISVSMSSNPPTLYELEKITFTLSHR
KVTDRYRSLCAFWNYSPTMNGSWSSEGCETYSNETHTSCRCNHLTHFAILMSSGPSIG
IKDYNILTRITQLGIIISLICLAICIFTFWFFSEIQSTRTTIHKNLCCSLFLAELVFLVG
INTNTNKLFCSSIAGLLHYFFLAFAWMCIEGIHLYLIVVGVIYNKGFLHKNFYIFGYLS
PAVVVGFSAAALGYRYYGTTKVCWLSTENNFIWSFIGPACLIILVNLLAFGVIIYKVRHT
AGLKPEVSCFENIRSCARGALALLFLLGTTWIFGVLHVHASVVTAYLFTVSNAFQGMFI
FLFLCVLSRKIQEEYYRLFKNPCCFGCLR

23/392

FIGURE 23

CTCCTCTTAACATACTTGCAGCTAAAACTAAATATTGCTGCTTGGGGACCTCCTTCTAGC
CTTAAATTTTCAGCTCATCACCTTCACCTGCCTTGGTCATGGCTCTGCTATTCTCCTTGAT
CCTTGCCATTTGCACCAGACCTGGATTCCCTAGCGTCTCCATCTGGAGTGCGGCTGGTGGG
GGGCCTCCACCGCTGTGAAGGGCGGGTGGAGGTGGAACAGAAAGGCCAGTGGGGCACCGT
GTGTGATGACGGCTGGGACATTAAGGACGTGGCTGTGTTGTGCCGGGAGCTGGGCTGTGG
AGCTGCCAGCGGAACCCCTAGTGGTATTTTGTATGAGCCACCAGCAGAAAAAGAGCAAAA
GGTCCTCATCCAATCAGTCAGTTGCACAGGAACAGAAGATACATTGGCTCAGTGTGAGCA
AGAAGAAGTTTATGATTGTTTACATGATGAAGATGCTGGGGCATCGTGTGAGAAGCCAGA
GAGCTCTTTCTCCCCAGTCCCAGAGGGTGTGAGGCTGGCTGACGGCCCTGGGCATTGCAA
GGGACGCGTGGAAGTGAAGCACCAGAACCAGTGGTATACCGTGTGCCAGACAGGCTGGAG
CCTCCGGGCGCGAAAGGTGGTGTGCCGGCAGCTGGGATGTGGGAGGGCTGTACTGACTCA
AAAACGCTGCAACAAGCATGCCCTATGGCCGAAAACCCATCTGGCTGAGCCAGATGTCATG
CTCAGGACGAGAAGCAACCCTTCAGGATTGCCCTTCTGGGCCTTGGGGGAAGAACACCTG
CAACCATGATGAAGACACGTGGGTGGAATGTGAAGATCCCTTTGACTTGAGACTAGTAGG
AGGAGACAACCTCTGCTCTGGGCGACTGGAGGTGCTGCACAAGGGCGTATGGGGCTCTGT
CTGTGATGACAACCTGGGGAGAAAAGGAGGACCAGGTGGTATGCAAGCAACTGGGCTGTGG
GAAGTCCCTCTCTCCCTCCTTCAGAGACCGGAAATGCTATGGCCCTGGGGTTGGCCGCAT
CTGGCTGGATAATGTTTCGTTGCTCAGGGGAGGAGCAGTCCCTGGAGCAGTGCCAGCACAG
ATTTTGGGGGTTTTCACGACTGCACCCACCAGGAAGATGTGGCTGTCATCTGCTCAGTGTA
GGTGGGCATCATCTAATCTGTTGAGTGCCTGAATAGAAGAAAAACACAGAAGAAGGGAGC
ATTTACTGTCTACATGACTGCATGGGATGAACACTGATCTTCTTCTGCCCTTGGACTGGG
ACTTATACTTGGTGCCCTGATTCTCAGGCCTTCAGAGTTGGATCAGAACTTACAACATC
AGGTCTAGTTCTCAGGCCATCAGACATAGTTTGGAACTACATCACCACCTTTCCTATGTC
TCCACATTGCACACAGCAGATTCCAGCCTCCATAATTGTGTGTATCAACTACTTAAATA
CATTTCTCACACACACACACACACACACACACACACACACACATACACCATTGTCC
TGTTTCTCTGAAGAACTCTGACAAAATACAGATTTTGGTACTGAAAGAGATTCTAGAGGA
ACGGAATTTTAAGGATAAATTTTCTGAATTGGTTATGGGGTTTCTGAAATTGGCTCTATA
ATCTAATTAGATATAAAATTTCTGGTAACTTTATTTACAATAATAAAGATAGCACTATGTG
TTCAAA

24/392

FIGURE 24

MALLFSLILAICTRPGFLASPSGVRLVGGLHRCEGRVEVEQKGQWGTVCDDGWDIKDVAV
LCRELGCGAASGTPSGILYEPPAEKEQKVLIQSVSCTGTEDTLAQCEQEVEYDCSHDEDA
GASCENPESSFSVPPEGVRLADGPGHCKGRVEVKHQNQWYTVCTGWSLRAAKVVCRLG
CGRAVLTQKRCNKHAYGRKPIWLSQMSCSGREATLQDCPSGPWGKNTCNHDEDTWVECED
PFDLRLVGDNLCSGRLEVLHKGWGSVCDDNWGEKEDQVCKQLGCGKSLSPSFRDRKC
YGPVGRIWLDNVRCSGEEQSLEQCQHRFWGFHDCTHQEDVAVICSV

Signal sequence:
amino acids 1-15

Casein kinase II phosphorylation site:
amino acids 47-51, 97-101, 115-119, 209-213, 214-218, 234-238,
267-271, 294-298, 316-320, 336-340

N-myristoylation site:
amino acids 29-35, 43-49, 66-72, 68-74, 72-78, 98-104, 137-143,
180-186, 263-269, 286-292

Amidation site:
amino acids 196-200

Speract receptor repeated domain signature:
amino acids 29-67, 249-287

25/392

FIGURE 25

CGGACGCGTGGGCGTCCGGCGGTTCGACAGCCAGGAGGCGGAGGCGCGCGGGCCAGCCTG
GGCCCCAGCCACACCTTCACCAGGGCCAGGAGCCACCATGTGCGGATGTCCACTGGGG
CTACTGCTGTTGCTGCCGCTGGCTGGCCACTTGGCTCTGGGTGCCAGCAGGGTCGTGGG
CGCCGGGAGCTAGCACCGGGTCTGCACCTGCGGGGCATCCGGGACGCGGGAGGCCGGTAC
TGCCAGGAGCAGGACCTGTGCTGCCGCGGCCGTGCCGACGACTGTGCCCTGCCCTACCTG
GGCGCCATCTGTTACTGTGACCTCTTCTGCAACCGCACGGTCTCCGACTGCTGCCCTGAC
TTCTGGGACTTCTGCCTCGGCGTGCCACCCCTTTTCCCCGATCCAAGGATGTATGCAT
GGAGGTCGTATCTATCCAGTCTTGGAACGTACTGGGACAACGTGAACCGTTGCACCTGC
CAGGAGAACAGGCAGTGGCATGGTGGATCCAGACATGATCAAAGCCATCAACCAGGGCAA
TATGGCTGGCAGGCTGGGAACCACAGCGCCTTCTGGGGCATGACCCTGGATGAGGGCAT
TCGCTACCGCCTGGGCACCATCCGCCCATCTTCTCGGTCTGAACATGCATGAAATTTA
TACAGTGCTGAACCCAGGGGAGGTGCTTCCACAGCCTTCGAGGCCTCTGAGAAGTGGCC
CAACCTGATTCATGAGCCTCTTGACCAAGGCAACTGTGCAGGCTCCTGGGCCTTCTCCAC
AGCAGCTGTGGCATCCGATCGTGTCTCAATCCATTCTCTGGGACACATGACGCCTGTCTCT
GTGCCCCAGAACCTGCTGTCTTGTGACACCCACCAGCAGCAGGGCTGCCGCGGTGGGCG
TCTCGATGGTGCCTGGTGGTTCTGCGTCCGCGAGGGGTGGTGTCTGACCATGCTATACC
CTTCTCGGGCCGTGAACGAGACGAGGCTGGCCCTGCGCCCCCTGTATGATGCACAGCCG
AGCCATGGGTCCGGGCAAGCGCCAGGCCACTGCCCACTGCCCAACAGCTATGTTAATAA
CAATGACATCTACCAGGTCACTCCTGTCTACCGCCTCGGCTCCAACGACAAGGAGATCAT
GAAGGAGCTGATGGAGAATGGCCCTGTCCAAGCCCTCATGGAGGTGCATGAGGACTTCTT
CCTATACAAGGGAGGCATCTACAGCCACAGCCAGTGAGCCTTGGGAGGCCAGAGAGATA
CCGCCGGCATGGGACCCACTCAGTCAAGATCACAGGATGGGGAGAGGAGACGCTGCCAGA
TGGAAGGACGCTCAAATACTGGACTGCGGCCAACTCCTGGGGCCAGCCTGGGGCGAGAG
GGGCCACTTCCGCATCGTGC CGCGGTCAATGAGTGCGACATCGAGAGCTTCGTGCTGGG
CGTCTGGGGCCGCGTGGGCATGGAGGACATGGGTCACTGAGGCTGCGGGCACACGC
GGGGTCCGGCCTGGGATCCAGGCTAAGGGCCGGCGGAAGAGGCCCAATGGGGCGGTGAC
CCCAGCCTCGCCGACAGAGCCCGGGGCGCAGGCGGGCGCCAGGGCGCTAATCCCGGCGC
GGGTTCCGCTGACGCGAGCGCCCGCCTGGGAGCCGCGGGCAGGCGAGACTGGCGGAGCCC
CCAGACCTCCCAGTGGGGACGGGGCAGGGCCTGGCCTGGGAAGAGCACAGCTGCAGATCC
CAGGCCTCTGGCGCCCCCACTCAAGACTACCAAAGCCAGGACACCTCAAGTCTCCAGCCC
CAATACCCACCCCAATCCCGTATTCTTTTTTTTTTTTTTTTAGACAGGGTCTTGCTCCG
TTGCCCAGGTTGGAGTGCAGTGGCCCATCAGGGCTCACTGTAACCTCCGACTCCTGGGTT
CAAGTGACCTCCACCTCAGCCTCTCAAGTAGCTGGGACTACAGGTGCACCACCACACC
TGGCTAATTTTGTATTTTGTAAAGAGGGGGGTCTCACTGTGTTGCCAGGCTGGTTT
CGAACTCCTGGGCTCAAGCGGTCCACCTGCCTCCGCCTCCCAAAGTGCTGGGATTGCAGG
CATGAGCCACTGCACCCAGCCCTGTATTCTTATTCTTCAGATATTTATTTTCTTTTCAC
TGTTTTTAAATAAAACCAAAGTATTGATAAAAAAAA

26/392

FIGURE 26

MWRCPLGLLLLLPLAGHLALGAQQGRGRRELAPGLHLRGIRDAGGRYCQEQLCCRGRAD
DCALPYLGAICYCDLFCNRTVSDCCPDFWDFCLGVPPFPPIQGCMHGGRIYPVLGTYWD
NCNRCTCQENRQWHGGSRHDQSHQPGQLWLAGWEPQRLLGHDPG

N-glycosylation site:

amino acids 78-82, 161-165

Casein kinase II phosphorylation site:amino acids 80-84, 117-121, 126-130, 169-173, 205-209, 296-300,
411-415**N-myristoylation site:**amino acids 21-27, 39-45, 44-50, 104-110, 160-164, 224-230,
269-275, 378-384, 442-448**Amidation site:**

amino acids 26-30, 318-322

Eukaryotic thiol (cysteine) proteases histidine active site:

amino acids 398-409

27/392

FIGURE 27

CCCACGCGTCCGGCAGGTTTTTCTTCAAGCCAAGAAGGACACGGATTGGCTGAAGGAGAA
AGTGCAGAGCTTGCAGACACTGGCTGCCAACAACCTCTGCGTTGGCCAAAGCCAACAACGA
CACCCTGGAGGATATGAACAGCCAGCTCAACTCATTACAGGTCAGATGGAGAACATCAC
CACTATCTCTCAAGCCAACGAGCAGAACCCTGAAAGACCTGCAGGACTTACACAAAGATGC
AGAGAATAGAACAGCCATCAAGTTCAACCAACTGGAGGAACGCTTCCAGCTCTTTGAGAC
GGATATTGTGAACATCATTAGCAATATCAGTTACACAGCCCACCACCTGCGGACGCTGAC
CAGCAATCTAAATGAAGTCAGGACCACCTTGACAGATACCTTACCAAACACACAGATGAT
CTGACCTCCTTGAATAATACCCTGGCCAACATCCGTTTGGATTCTGTTTCTCTCAGGATG
CAACAAGATTTGATGAGGTCGAGGTAGACACTGAAGTAGCCAACTTATCAGTGATTATG
GAAGAAATGAAGCTAGTAGACTCCAAGCATGGTCAGCTCATCAAGAATTTTACAATACTA
CAAGGTCCACCGGGCCCCAGGGGTCCAAGAGGTGACAGAGGATCCCAGGGACCCCCTGGC
CCAAGTGGCAACAAGGGACAGAAAGGAGAGAAGGGGGAGCCTGGACCACCTGGCCCTGCG
GGTGAGAGAGGCCCAATTGGACCAGCTGGTCCCCCGGAGAGCGTGGCGGCAAAGGATCT
AAAGGCTCCCAGGGCCCCAAAGGCTCCCGTGGTTCCCTGGGAAGCCCGGCCCTCAGGGC
CCCAGTGGGGACCCAGGCCCCCCGGGGCCACCAGGCAAGAGGGGACTCCCCGGGCCCTCAG
GGCCCTCCTGGCTTCCAGGGACTTCAGGGCACCGTTGGGGAGCCTGGGGTGCCCTGGACCT
CGGGGACTGCCAGGCTTGCCTGGGGTACCAGGCATGCCAGGCCCAAGGGCCCCCCCCGGC
CCTCCTGGCCCATCAGGAGCGGTGGTGCCCCCTGGCCCTGCAGAATGAGCCAACCCCGGCA
CCGGAGGACAATAGCTGCCCGCCTCACTGGAAGAACTTCACAGACAAATGCTACTATTTT
TCAGTTGAGAAAGAAATTTTTGAGGATGCAAAGCTTTTCTGTGAAGACAAGTCTTCACAT
CTTGTTTTTCATAAACACTAGAGAGGAACAGCAATGGATAAAAAAACAGATGGTAGGGAGA
GAGAGCCACTGGATCGGCCTCACAGACTCAGAGCGTGAAAATGAATGGAAGTGGCTGGAT
GGGACATCTCCAGACTACAAAATTGGAAAGCTGGACAGCCGGATAACTGGGGTCATGGC
CATGGGCCAGGAGAAGACTGTGCTGGGTGATTTATGCTGGGCAGTGGAACGATTTCCAA
TGTGAAGACGTCAATAACTTCATTTGCGAAAAAGACAGGGAGACAGTACTGTCATCTGCA
TTATAACCGGACTGTGATGGGATCACATGAGCAAATTTTCAGCTCTCAAAGGCAAAGGACA
CTCCTTTCTAATTGCATCACCTTCTCATCAGATTGAAAAAAAAAAGCACTGAAAACCAA
TTACTGAAAAAAAAATTGACAGCTAGTGTTTTTTACCATCCGTCATTACCCAAAGACTTGG
GAACTAAATGTTCCCCAGGGTGATATGCTGATTTTCATTGTGCACATGGACTGAATCAC
ATAGATTCTCCTCCGTCAGTAACCGTGCGATTATACAAATTATGTCTTCCAAAGTATGGA
ACACTCCAATCAGAAAAAGGTTATCATTGGTCGTTGAGTTATGGGAAGAACTTAAGCATA
TACTGTGTAAACAGTGCCATACATTTCTAAAATCCCAAGTGTAGGAAAAATATGCAGACA
TACAGATATATAGGCCAACTATTAGTAATAATATGAAATATACTTAAAGAGCTTTTAAAA
CTTTGTATTTTTGTACAAAAAAA

28/392

FIGURE 28

MQQDLMSRLDTEVANLSVIMEEMKLVD SKHGQLIKNFTILQGPPGPRGPRGDRGSQGGP
GPTGNKGQKGEKGEPPGPAGERGPAGPPGERGGKGSKGSQGPKGSRGSPGKPGPQ
GPSGDPGPPGPPGKEGLPGPQGPFGQLQGTVGEPGVPGPRGLPGLPGVPGMPGPKGPP
GPPGPSGAVVPLALQNEPTPAPEDNSCPHWNFTDKCYYFSVEKEIFEDAKLFCEDKSS
HLVFINTREEQQWIKQMVGRESHWIGLTD SERENWKWLDGTSPDYKNWKAGQPDNWDH
GHGPGEDCAGLIYAGQWDFQCEDVNNFICEKDRETVLSSAL

Signal sequence:

None

Transmembrane domain:

None

N-glycosylation site:

16-19, 37-40, 213-216

Tyrosine kinase phosphorylation site:

212-220

N-myristoylation site:

97-102, 100-105, 148-153, 267-272, 293-298, 310-315

Cell attachment sequence:

51-53

C-type lectin domain signature:

308-330

Lectin C-type domain:

233-330

Collagen triple helix repeat:

43-102, 127-186

29/392

FIGURE 29

GGACTAATCTGTGGGAGCAGTTTATTCCAGTATCACCCAGGGTGCAGCCACACCAGGACT
GTGTTGAAGGGTGTTTTTTTCTTTTAAATGTAATACCTCCTCATCTTTTCTTCTTACAC
AGTGTCTGAGAACATTTACATTATAGATAAGTAGTACATGGTGGATAACTTCTACTTTTA
GGAGGACTACTCTCTTCTGACAGTCCTAGACTGGTCTTCTACACTAAGACACCATGAAGG
AGTATGTGCTCCTATTATTCTGGCTTTGTGCTCTGCCAAACCTTCTTTAGCCCTTCAC
ACATCGCACTGAAGAATATGATGCTGAAGGATATGGAAGACACAGATGATGATGATGATG
ATGATGATGATGATGATGATGAGGACAACTCTCTTTTCCAACAAGAGAGCCAAGAA
GCCATTTTTTTCCATTTGATCTGTTTTCCAATGTGTCCATTTGGATGTCAGTGCTATTTCAC
GAGTTGTACATTGCTCAGATTTAGGTTTGACCTCAGTCCCAACCAACATTCCATTTGATA
CTCGAATGCTTGATCTTCAAAACAATAAAATTAAGGAAATCAAAGAAAATGATTTTAAAG
GACTCACTTCACTTTATGGTCTGATCCTGAACAACAAGCTAACGAAGATTCAACCCAA
AAGCCTTTCTAACCACAAAAGAAGTTGCGAAGGCTGTATCTGTCCCACAATCAACTAAGTG
AAATACCACTTAATCTTCCCAAATCATTAGCAGAAGCTCAGAATTCATGAAAATAAAGTTA
AGAAAATACAAAAGGACACATTCAAAGGAATGAATGCTTTACACGTTTTGGAAATGAGTG
CAAACCTCTTGATAATAATGGGATAGAGCCAGGGGCATTTGAAGGGGTGACGGTGTTCC
ATATCAGAATTGCAGAAGCAAACTGACCTCAGTTCCTAAAGGCTTACCACCAACTTTAT
TGGAGCTTCACTTAGATTATAATAAAATTTCAACAGTGGAAGCTTGAGGATTTTAAACGAT
ACAAAGAACTACAAAGGCTGGGCCTAGGAAACAACAAAATCACAGATATCGAAAATGGGA
GTCTTGCTAACATAACCAGTGTGAGAGAAATACATTTGGAAAACAATAAACTAAAAAAA
TCCCTTCAGGATTACCAGAGTTGAAATACCTCCAGATAATCTTCCTTCATTCTAATTCAA
TTGCAAGAGTGGGAGTAAATGACTTCTGTCCAACAGTGCCAAAGATGAAGAAATCTTTAT
ACAGTGCAATAAGTTTTATTCAACAACCCGGTGAAATACTGGGAAATGCAACCTGCAACAT
TTCGTTGTGTTTTGAGCAGAATGAGTGTTTCAGCTTGGGAAGCTTTGGAATGTAATAATTAG
TAATTGGTAATGTCCATTTAATATAAGATTCAAAAATCCCTACATTTGGAATACTTGAAC
TCTATTAATAATGGTAGTATTATATATACAAGCAAATATCTATTCTCAAGTGGTAAGTCC
ACTGACTTATTTTATGACAAGAAATTTCAACGGAATTTTGCCAAACTATTGATACATAAG
GGGTGAGAGAAACAAGCATCTATTGCAGTTTCCTTTTTTGCGTACAAATGATCTTACATA
AATCTCATGCTTGACCATTCTTTCTTCATAACAAAAAGTAAGATATTCGGTATTTAAC
ACTTTGTTATCAAGCACATTTTAAAAGAACTGTACTGTAAATGGAATGCTTGACTTAGC
AAAATTTGTGCTCTTTCATTTGCTGTTAGAAAAACAGAATTAACAAAGACAGTAATGTGA
AGAGTGCAATTACACTATTCTTATTCTTTAGTAAGTTGGGTAGTACTGTAATATTTTAAAT
CATCTTAAAGTATGATTTGATATAATCTTATTGAAATTACCTTATCATGTCTTAGAGCCC
GTCTTTATGTTTTAAACTAATTTCTTAAAAATAAGCCTTCAGTAAATGTTCAATTACCAAC
TTGATAAATGCTACTCATAAGAGCTGGTTTTGGGGCTATAGCATATGCTTTTTTTTTTTTA
ATTATTACCTGATTTAAAAATCTCTGTAAAAACGTGTAGTGTTTCATAAAATCTGTAACT
CGCATTTTAAATGATCCGCTATTATAAGCTTTTAAATAGCATGAAAATTGTTAGGCTATATA
ACATTGCCACTTCAACTCTAAGGAATATTTTTGAGATATCCCTTTGGAAGACCTTGCTTG
GAAGAGCCTGGACACTAACAATTCTACACCAAATTGTCTCTTCAAATACGTATGGACTGG
ATAACTCTGAGAAACACATCTAGTATAACTGAATAAGCAGAGCATCAAATTAACAGACA
GAAACCGAAAGCTCTATATAAATGCTCAGAGTTCTTTATGTATTTCTTATTGGCATTCAA
CATATGTAAATCAGAAAACAGGGAAATTTTCATTAAAAATATTGGTTTGAAAT

30/392

FIGURE 30

MKEYVLLLFLALCSAKPFFSPSHIALKNMMLKDMEDTDDDDDDDDDDDDDEDNSLFPTRE
PRSHFFPFDLFPMCPFGCQCYSRVVHCSDLGLTSVPTNIPFDTRMLDLQNNKIKEIKEND
FKGLTSLYGLILNNKLTKIHPKAFLTTKKLRRLYLSHNQLSEIPLNLPKSLAELRIHEN
KVKKIQKDTFKGMNALHVLEMSANPLDNNGIEPGAEGVTVFHIRIAEAKLTSVPKGLPP
TLELHLDYNKISTVELEDFKRYKELQRLGLGNKKITDIENGLANIPRVREIHLNENKL
KKIPSGLPPELKYLQIIFLHSNSIARVGVNDFCPTVPKMKKSLYSAISLFNNPVKYWEMQP
ATFRCVLSRMSVQLGNFGM

Signal sequence:
amino acids 1-15

N-glycosylation site:
amino acids 281-285

N-myristoylation sites:
amino acids 129-135, 210-216, 214-220, 237-243, 270-276,
282-288

Leucine zipper pattern:
amino acids 154-176

31/392

FIGURE 31

AGCAGGGAAATCCGGATGTCTCGGTTATGAAAGTGGAGCAGTGAGTGTGAGCCTCAACATA
GTTCCAGAACTCTCCATCCGGACTAGTTATTGAGCATCTGCCTCTCATATCACCAGTGGC
CATCTGAGGTGTTTCCCTGGCTCTGAAGGGGTAGGCACGATGGCCAGGTGCTTCAGCCTG
GTGTTGCTTCTCACTTCCATCTGGACCACGAGGCTCCTGGTCCAAGGCTCTTTGCGTGCA
GAAGAGCTTTCCATCCAGGTGTCTGAGCAATTATGGGGATCACCCTTGTGAGCAAAAAG
GCGAACCAGCAGCTGAATTTACAGAAGCTAAGGAGGCCTGTAGGCTGCTGGGACTAAGT
TTGGCCGGCAAGGACCAAGTTGAAACAGCCTTGAAAGCTAGCTTTGAAACTTGCAGCTAT
GGCTGGGTGGGATGGATTTCGTGGTCTCTAGGATTAGCCCAAACCCCAAGTGTGGG
AAAAATGGGGTGGGTGTCCTGATTTGGAAGGTTCCAGTGAGCCGACAGTTTGCAGCCTAT
TGTTACAACCTCATCTGATACTTGGACTAACTCGTGCAATTCCAGAAATTATCACCACCAA
GATCCCATATTCAACACTCAAACTGCAACACAAACAACAGAATTTATTGTGAGTGACAGT
ACCTACTCGGTGGCATCCCCTTACTCTACAATACTGCCCCCTACTACTCCTCCTGCT
CCAGCTTCCACTTCTATTCCACGGAGAAAAAATTGATTTGTGTACAGAAGTTTTTATG
GAACTAGCACCATGTCTACAGAACTGAACCATTGTTGAAAATAAAGCAGCATTCAAG
AATGAAGCTGCTGGGTTTGGAGGTGTCCCCACGGCTCTGCTAGTGCTTGCTCTCCTCTTC
TTTGGTGCTGCAGCTGGTCTTGGATTTTGCTATGTCAAAAGGTATGTGAAGGCCTTCCCT
TTTACAAACAAGAATCAGCAGAAGGAAATGATCGAAACCAAAGTAGTAAAGGAGGAGAAG
GCCAATGATAGCAACCCTAATGAGGAATCAAAGAAAAGTGAATAAAACCCAGAAGAGTCC
AAGAGTCCAAGCAAACTACCGTGCGATGCCTGGAAGCTGAAGTTTAGATGAGACAGAAA
TGAGGAGACACACCTGAGGCTGGTTTTCTTTCATGCTCCTTACCCTGCCCCAGCTGGGGAA
ATCAAAGGGCCAAAGAACCAAGAAGAAAGTCCACCCTTGGTTCTTAAGTGAATCAGC
TCAGGACTGCCATTGGACTATGGAGTGCACCAAGAGAATGCCCTTCTCCTTATTGTAAC
CCTGTCTGGATCCTATCCTCCTACCTCCAAAGCTTCCCACGGCCTTTCTAGCCTGGCTAT
GTCCTAATAATATCCCACTGGGAGAAAGGAGTTTTGCAAAGTGCAAGGACCTAAACATC
TCATCAGTATCCAGTGGTAAAAAGGCCTCCTGGCTGTCTGAGGCTAGGTGGGTTGAAAGC
CAAGGAGTCACTGAGACCAAGGCTTTCTCTACTGATTCCGCAGCTCAGACCCTTTCTTCA
GCTCTGAAAGAGAAACACGTATCCCACCTGACATGTCCTTCTGAGCCCGGTAAGAGCAAA
AGAATGGCAGAAAAGTTTAGCCCCTGAAAGCCATGGAGATTCTCATAACTTGAGACCTAA
TCTCTGTAAAGCTAAAATAAAGAAATAGAACAAAGGCTGAGGATACGACAGTACACTGTCA
GCAGGGACTGTAAACACAGACAGGGTCAAAGTGTCTTCTCTGAACACATTGAGTTGGAAT
CACTGTTTAGAACACACACACTTACTTTTTCTGGTCTCTACCCTGCTGATATTTTCTCT
AGGAAATATACTTTTACAAGTAACAAAAATAAAAACTCTTATAAATTTCTATTTTTATCT
GAGTTACAGAAATGATTACTAAGGAAGATTACTCAGTAATTTGTTTAAAAAGTAATAAAA
TTCAACAAACATTTGCTGAATAGCTACTATATGTCAAGTGCTGTGCAAGGTATTACACTC
TGTAATTGAATATTATTCTCAAAAAATTGCACATAGTAGAACGCTATCTGGGAAGCTAT
TTTTTTCAGTTTTGATATTTCTAGCTTATCTACTTCCAACTAATTTTTATTTTTGCTGA
GACTAATCTTATTCATTTTCTCTAATATGGCAACCATTATAACCTTAATTTATTATTAAC
ATACCTAAGAAGTACATTGTTACCTCTATATACCAAGCACATTTTAAAAGTGCCATTAA
CAAATGTATCACTAGCCCTCCTTTTTTCCAACAAGAAGGGACTGAGAGATGCAGAAATATT
TGTGACAAAAAATTAAAGCATTTAGAAAACCT

32/392

FIGURE 32

MARCFSLVLLLLTSIWTTTRLLVQGSRLRAEELSIQVSCRIMGITLVSKKANQQQLNFTEAKEA
CRLGLSLAGKDQVETALKASFETCSYGWVGDFVVISRISPNPKCGKNGVGVLIWKVPV
SRQFAAYCYNSSDTWTNSCIPEIITTKDPIFNTQTATQTTEFIVSDSTYSVASPYSTIPA
PTTTPPAPASTSIPRRKKLICVTEVFMETSTMSTETEPFVENKAAFKNEAGFGGVPTAL
LVLALLFFGAAAGLGFCYVKRYVKAFFFTNKNQOKEMIETKVVKEEKANDSNPNEESKKT
DKNPEESKSPSKTTVRCLEAEV

Signal sequence:
amino acids 1-16

Transmembrane domain:
amino acids 235-254

N-glycosylation site:
amino acids 53-57, 130-134, 289-293

Casein kinase II phosphorylation site:
amino acids 145-149, 214-218

Tyrosine kinase phosphorylation site:
amino acids 79-88

N-myristoylation site:
amino acids 23-29, 65-71, 234-240, 235-239, 249-255, 253-259

FIGURE 33

[illegible]

34/392

TCTACTTATGTTGGACACTTGGCAGAAGGACCGTGCCCGGCGGCCTCATTTTGACCAGCT
GGTGGCTGCATTTGACAAGATGATCCGCAAGCCAGATACCCTGCAGGCTGGCGGGGACCC
AGGGGAAAGGCCTTCCCAGGCCCTTCTGACCCCTGTGGCCCTGGACTTTCCTTGTCTGGA
CTCACCCCAGGCCTGGCTTTTCAGCCATTGGACTGGAGTGCTACCAGGACAACCTTCTCCAA
GTTTGGCCTCTGTACCTTCAGTGATGTGGCTCAGCTCAGCCTAGAAGACCTGCCTGCCCT
GGGCATCACCCCTGGCTGGCCACCAGAAGAAGCTGCTGCACCACATCCAGCTCCTTCAGCA
ACACCTGAGGCAGCAGGGCTCAGTGGAGGTCTGAGAATGACGATACCCGTGACTCAGCCC
TGGACACTGGTCCGAGAAGGGACATGTGGGACGTGAGCCGGGCTCCAACAGCCTCTGTGA
GAGATGCCCCACACCAAACCAACCCTCCGATGGCTGCATTCCCTGGTCCCTCCGCTTTTC
CACCAGCCCCCTCCTCATTAAGGGAAAGAAGGGAATTTGCAAAAAAAAAAAAAAAAAAAAA
AAAAAAA

35/392

FIGURE 34

MATEGAAQLGNRVAGMVCSLWVLLLVSSVLALAEVLLD TTGETSEIGWLTYP PGGWDEVS
VLDDQRRRLTRTFEACHVAGAPPGTGQDNWLQTHFVERRGAQRAHIRLHFSVRACSSSLGVS
GGTCRETFTLYYRQAEPPDSPDSVSSWHLKRWTKVDTIAADESFPSSSSSSSSSSSSSSAAW
AVGPHGAGQRAGLQLNVKERSFGPLTQRGFYVAFQDTGACLALVAVRLFSYTCPAVLRSF
ASFPETQASGAGGASLVAAVGTCVAHAEEEDGVGGQAGGSPRLHNCN GEGKWMVAVGGC
RCQPGYQPARGDKACQACPRGLYKASAGNAPCSPCPARSHAPNPAAPVPCPCLEGFYRASS
DPPEAPCTGPPSAPQELWFEVQGSALMLHWRLPRELGGRGDLLFN VVCKECEGRQEPASG
GGGTCHRCRDEVHFDPRQRGLTESRVLVGGGLRAHVPIILEVQAVNGVSELSPDPPQAAAI
NVSTSHEVPSAVPVVHQVSRASNSITVSWPQPDQTNGNILDYQLRYDDQAEDESHSFTLT
SETNTATVTQLSPGHIYGFQVRARTAAGHPYGGKVYFQTL PQGELSSQLPERLSLVIGS
TLGALAFLLLLAAITVLAVVFQKRRTGYTEQLQQYSSPGLGVKYYIDPSTYEDPCQAIR
ELAREVDPAYIKIEEVIGTGSFGEVRQGRQLQPRGRREQTVAIQALWAGGAESLQMTFLGR
AAVLGQFQHPNILRLEGVVTKSRLPLVLTEFMELGPLDSFLRQREGQFSSQLVAMQRGV
AAAMQYLSSFAFVHRSLSAHSVLVNSHLVCKVARLGHSPQGPSCLLRWAAPEVIAHGKHT
HVGSDDELWRTALLGHE

signal sequence:
Amino acids 1-31

Transmembrane domains:
Amino acids 217-234;598-618

N-glycosylation site:
Amino acids 481-485

Glycosaminoglycan attachment sites:
Amino acids 249-253;419-423

cAMP- and cGMP-dependent protein kinase phosphorylation sites:
Amino acids 66-70;150-154;624-628

Tyrosine kinase phosphorylation sites:
Amino acids 644-673;664-671

N-myristoylation sites:
Amino acids 10-16;15-21;79-85;99-105;118-124;188-194;
192-198;218-224;250-256;261-267;275-281;276-282;298-304;321-
327;328-334;420-426;421-427;440-446;449-455;599-605;626-632;
708-714;766-772;779-785

Amidation site:
Amino acids 693-697

Cell attachment sequences:
Amino acids 310-313;399-402

36/392

FIGURE 35

GGGGTCTCCCTCAGGGCCGGGAGGCACAGCGGTCCCTGCTTGCTGAAGGGCTGGATGTAC
GCATCCGCAGGTTCCCGCGGACTTGGGGGCGCCCGCTGAGCCCCGGCGCCCGCAGAAGAC
TTGTGTTTGCCTCCTGCAGCCTCAACCCGGAGGGCAGCGAGGGCCTACCACCATGATCAC
TGGTGTGTTTCTGAGCATGCGCTTGTGGACCCAGTGGGCGTCCTGACCTCGCTGGCGTACTG
CCTGCACCAGCGGCGGGTGGCCCTGGCCGAGCTGCAGGAGGCCGATGGCCAGTGTCCGGT
CGACCGCAGCCTGCTGAAGTTGAAAATGGTGCAGGTCGTGTTTCGACACGGGGCTCGGAG
TCCTCTCAAGCCGCTCCCGCTGGAGGAGCAGGTAGAGTGGAACCCCGAGCTATTAGAGGT
CCCACCCCAAACCTCAGTTTGATTACACAGTCACCAATCTAGCTGGTGGTCCGAAACCATA
TTCTCCTTACGACTCTCAATACCATGAGACCACCCTGAAGGGGGGCATGTTTGCTGGGCA
GCTGACCAAGGTGGGCATGCAGCAAATGTTTGCCCTTGGGAGAGAGACTGAGGAAGAATA
TGTGGAAGACATTCCCTTTCTTTCACCAACCTTCAACCCACAGGAGGTCTTTATTTCGTTT
CACTAACATTTTTTCGGAATCTGGAGTCCACCCGTTGTTTGCTGGCTGGGCTTTTCCAGTG
TCAGAAAGAAGGACCCATCATCATCCACACTGATGAAGCAGATTCAGAAGTCTTGTATCC
CAACTACCAAAGCTGCTGGAGCCTGAGGCAGAGAACCAGAGGCCGGAGGCAGACTGCCTC
TTTACAGCCAGGAATCTCAGAGGATTTGAAAAAGGTGAAGGACAGGATGGGCATTGACAG
TAGTGATAAAGTGGACTTCTTCATCCTCCTGGACAACGTGGCTGCCGAGCAGGCACACAA
CCTCCCAAGCTGCCCCATGCTGAAGAGATTTGCACGGATGATCGAACAGAGAGCTGTGGA
CACATCCTTGACATACTGCCCAAGGAAGACAGGGAAAGTCTTCAGATGGCAGTAGGCCC
ATTCTCCACATCCTAGAGAGCAACCTGCTGAAAGCCATGGACTCTGCCACTGCCCCGA
CAAGATCAGAAAGCTGTATCTCTATGCGGCTCATGATGTGACCTTCATACCGCTCTTAAT
GACCCCTGGGGATTTTGTACCACAAATGGCCACCGTTTGCTGTTGACCTGACCATGGAAC
TTACCAGCACCTGGAATCTAAGGAGTGGTTTGTGCAGCTCTATTACCACGGGAAGGAGCA
GGTGCCGAGAGGTTGCCCTGATGGGCTCTGCCCGCTGGACATGTTCTTGAATGCCATGTC
AGTTTATACCTTAAGCCAGAAAAATACCATGCACTCTGCTCTCAAACCTCAGGTGATGGA
AGTTGGAAATGAAGAGTAACTGATTTATAAAAGCAGGATGTGTTGATTTTAAATAAAGT
GCCTTTATACAATG

37/392

FIGURE 36

MITGVFSMRLWTPVGVLTSLAYCLHQRRVALAELQEQADGQCPVDRSLLKLKVMQVVFRRHG
ARSPLKPLPLEEQVEWNPQLLEVPPQTQFDYTVTNLAGGPKPYSPYDSQYHETTLKGGMF
AGQLTKVGMQQMFA LGERLRKNYVEDIPFLSPTFNPQEVFIRSTNIFRNLESTRCLLAGL
FQCQKEGP IIIHTDEADSEVLYPNYQSCWSLRQRTGRRRQTASLQPGISED LKKVKDRMG
IDSSDKVDF FILLDNVAAEQAHNLPSCPMLKRFARMIEQRAVDTSLYILPKEDRESLQMA
VGPFLHILES NLLKAMDSATAPDKIRKLYLYAAHDVTFIPLMLT LGIFDHKWPPFAVDLT
MELYQHLESKEWFVQLYYHGKEQVPRGCPDGLCPLDMFLNAMS VYTLSP EKYHALCSQTQ
VMEVGNEE

Signal sequence:

amino acids 1-23

cAMP- and cGMP-dependent protein kinase phosphorylation site:

amino acids 218-222

Casein kinase II phosphorylation site:

amino acids 87-91, 104-108, 320-324

Tyrosine kinase phosphorylation site:

amino acids 280-288

N-myristoylation site:

amino acids 15-21, 117-123, 118-124, 179-185, 240-246, 387-393

Amidation site:

amino acids 216-220

Leucine zipper pattern:

amino acids 10-32

Histidine acid phosphatases phosphohistidine signature:

amino acids 50-65

38/392

FIGURE 37

ACTGCACTCGGTTCTATCGATTGAATTCCCCGGGGATCCTCTAGAGATCCCTCGACCTCG
ACCCACGCGTCCGCGGACGCGTGGGCGGACGCGTGGGCCGGCTACCAGGAAGAGTCTGCC
GAAGGTGAAGGCCATGGACTTCATCACCTCCACAGCCATCCTGCCCCCTGCTGTTCCGGCTG
CCTGGGCGTCTTCGGCCTCTTCCGGCTGCTGCAGTGGGTGCGCGGGAAGGCCTACCTGCG
GAATGCTGTGGTGGTGATCACAGGCGCCACCTCAGGGCTGGGCAAAGAATGTGCAAAAGT
CTTCTATGCTGCGGGTGCTAAACTGGTGCTCTGTGGCCGGAATGGTGGGGCCCTAGAAGA
GCTCATCAGAGAACTTACCGCTTCTCATGCCACCAAGGTGCAGACACACAAGCCTTACTT
GGTGACCTTCGACCTCACAGACTCTGGGGCCATAGTTGCAGCAGCAGCTGAGATCCTGCA
GTGCTTTGGCTATGTGACATACTTGTCAACAATGCTGGGATCAGCTACCGTGGTACCAT
CATGGACACCACAGTGGATGTGGACAAGAGGGTCATGGAGACAACTACTTTGGCCCACT
TGCTCTAACGAAAGCACTCCTGCCCTCCATGATCAAGAGGAGGCAAGGCCACATTGTCGC
CATCAGCAGCATCCAGGGCAAGATGAGCATTCCCTTTTCGATCAGCATATGCAGCCTCCAA
GCACGCAACCCAGGCTTTCTTTGACTGTCTGCGTGCCGAGATGGAACAGTATGAAATTGA
GGTGACCGTCATCAGCCCCGGCTACATCCACACCAACCTCTCTGTAAATGCCATCACCGC
GGATGGATCTAGGTATGGAGTTATGGACACCACCACAGCCCAGGGCCGAAGCCCTGTGGA
GGTGGCCCAGGATGTTCTTGCTGCTGTGGGGAAGAAGAAGAAAGATGTGATCCTGGCTGA
CTTACTGCCTTCCTTGGCTGTTTATCTTCGAACTCTGGCTCCTGGGCTCTTCTTCAGCCT
CATGGCCTCCAGGGCCAGAAAAGAGCGGAAATCCAAGAACTCCTAGTACTCTGACCAGCC
AGGGCCAGGGCAGAGAAGCAGCACTCTTAGGCTTGCTTACTCTACAAGGGACAGTTGCAT
TTGTTGAGACTTTAATGGAGATTTGTCTCACAAGTGGGAAAGACTGAAGAAACACATCTC
GTGCAGATCTGCTGGCAGAGGACAATCAAAAACGACAACAAGCTTCTTCCCAGGGTGAGG
GGAAACACTTAAGGAATAAATATGGAGCTGGGGTTTAACTAACTAACTAGAAATAAACA
TCTCAAACAGTAAAAAAAAAAAAAAAAAGGGCGGCCGCGACTCTAGAGTCGACCTGCAGAAG
CTTGGCCGCCATGGCCCACTTGTTTATTGCAGCTTATAATGGTTAC

39/392

FIGURE 38

MDFITSTAILPLLFGCLGVFGLFRLLQWVRGKAYLRNAVVVITGATSGLGKECAKVFYAA
GAKLVLCGRNGGALEELIRELTASHATKVQTHKPYLVTFDLTDSGAIVAAAAILQCFGY
VDILVNNAGISYRGTIMDTTVDVDKRVMETNYFGPVALTKALLPSMIKRRQGHIVAIS
QGKMSIPFRSAYAASKHATQAFFDCLRAEMEQYEIEVTVISPGYIHTNLSVNAITADGSR
YGVMDTTTAQGRSPVEVAQDVLA AVGKKKKDVILADLLPSLAVYLRTLAPGLFFSLMASR
ARKERKSKNS

Signal sequence:
amino acids 1-21

Transmembrane domain:
amino acids 104-120, 278-292

N-glycosylation site:
amino acids 228-232

Glycosaminoglycan attachment site:
amino acids 47-51

Casein kinase II phosphorylation site:
amino acids 135-139, 139-143, 253-257

Tyrosine kinase phosphorylation site:
amino acids 145-153, 146-153

N-myristoylation site:
amino acids 44-50, 105-111, 238-244, 242-248, 291-297

Amidation site:
amino acids 265-269

Prokaryotic membrane lipoprotein lipid attachment site:
amino acids 6-17

40/392

FIGURE 39

GCAAGCCAAGGCGCTGTTTGAGAAGGTGAAGAAGTTCCGGACCCATGTGGAGGAGGGGGACATTGT
GTACCGCCTCTACATGCGGCAGACCATCATCAAGGTGATCAAGTTCATCCTCATCATCTGCTACAC
CGTCTACTACGTGCACAACATCAAGTTCGACGTGGACTGCACCGTGGACATTGAGAGCCTGACGGG
CTACCGCACCTACCGCTGTGCCACCCCTGGCCACACTCTTCAAGATCCTGGCGTCTTCTACAT
CAGCCTAGTCATCTTCTACGGCCTCATCTGCATGTACACACTGTGGTGGATGCTACGGCGCTCCCT
CAAGAAGTACTCGTTTGAGTCGATCCGTGAGGAGAGCAGCTACAGCGACATCCCCGACGTCAAGAA
CGACTTCGCCTTCATGTGCACCTCATTGACCAATACGACCCGCTCTACTCCAAGCGCTTCGCCGT
CTTCCTGTGCGAGGTGAGTGAGAACAAGCTGCGGCAGCTGAACCTCAACAACGAGTGAGCGCTGGA
CAAGCTCCGGCAGCGGCTCACCAAGAACGCGCAGGACAAGCTGGAGCTGCACCTGTTTCATGCTCAG
TGGCATCCCTGACACTGTGTTTGACCTGGTGGAGCTGGAGGTCTCAAGCTGGAGCTGATCCCCGA
CGTGACCATCCCCGCCAGCATTGCCAGCTCACGGGCCTCAAGGAGCTGTGGCTCTACCACACAGC
GGCCAAGATTGAAGCGCCTGCGCTGGCCTTCCTGCGCGAGAACCTGCGGGCGCTGCACATCAAGTT
CACCGACATCAAGGAGATCCCGCTGTGGATCTATAGCCTGAAGACACTGGAGGAGCTGCACCTGAC
GGGCAACCTGAGCGCGGAGAACCAACCGCTACATCGTCATCGACGGGCTGCGGGAGCTCAAACGCCT
CAAGGTGCTGCGGCTCAAGAGCAACCTAAGCAAGCTGCCACAGGTGGTCAAGATGTGGGCGTGCA
CCTGCAGAAGCTGTCCATCAACAATGAGGGCACCAAGCTCATCGTCTCAACAGCCTCAAGAAGAT
GGCGAACCTGACTGAGCTGGAGCTGATCCGCTGCGACCTGGAGCGCATCCCCACTCCATCTTCAG
CCTCCACAACCTGCAGGAGATTGACCTCAAGGACAACAACCTCAAGACCATCGAGGAGATCATCAG
CTTCCAGCACCTGCACCGCCTCACCTGCCTTAAGCTGTGGTACAACCACATCGCCTACATCCCCAT
CCAGATCGGCAACCTCACCAACCTGGAGCGCCTCTACCTGAACCGCAACAAGATCGAGAAGATCCC
CAGGAGCTCTTCTACTGCCGCAAGCTGCGCTACCTGGACCTCAGCCACAACAACCTGACCTTCCT
CCCTGCCGACATCGGCTCCTGCAGAACCTCAGAACCTAGCCATCACGGCCAACCGGATCGAGAC
GCTCCCTCCGGAGCTCTTCCAGTGCCGGAAGCTGCGGGCCCTGCACCTGGGCAACAACGTGTGCA
GTCACTGCCCTCCAGGGTGGGCGAGCTGACCAACCTGACGCAGATCGAGCTGCGGGGCAACCGGCT
GGAGTGCTGCTGTGGAGCTGGGCGAGTGCCCACTGCTCAAGCGCAGCGGCTTGGTGGTGGAGGA
GGACCTGTTCAACACACTGCCACCCGAGGTGAAGGAGCGGCTGTGGAGGGCTGACAAGGAGCAGGC
CTGAGCGAGGCCGGCCAGCACAGCAAGCAGCAGGACCGCTGCCAGTCTCAGGCCCGGAGGGGC
AGGCCTAGCTTCTCCAGAACTCCCGGACAGCCAGGACAGCCTCGCGGCTGGGCAGGAGCCTGGGG
CCGCTTGTGAGTCAGGCCAGAGCGAGAGGACAGTATCTGTGGGGCTGGCCCCCTTTTCTCCCTCTGA
GACTCAGTCCCCAGGGCAAGTGCTTGTGGAGGAGAGCAAGTCTCAAGAGCGCAGTATTTGGATA
ATCAGGGTCTCCTCCCTGGAGGCCAGCTCTGCCCCAGGGGCTGAGCTGCCACCAGAGGTCTGGGA
CCCTCACTTTAGTTCTTGGTATTTATTTTTCTCCATCTCCACCTCCTTCATCCAGATAACTATA
CATTCCCAAGAAAGTTCAAGCCAGATGGAAGGTGTTCAAGGAAAGGTGGGCTGCCTTTTCCCTTG
TCCTTATTTAGCGATGCCGCCGGGCATTTAACACCCACCTGGACTTCAGCAGAGTGGTCCGGGGCG
AACCAGCCATGGGACGGTCACCCAGCAGTGCCGGGCTGGGCTCTGCGGTGCGGTCCACGGGAGAGC
AGGCCTCCAGCTGGAAGGCCAGGCCTGGAGCTTGCTCTTCAGTTTTTGTGGCAGTTTTAGTTTT
TTGTTTTTTTTTTTTTAAATCAAAAAACAATTTTTTTTTAAAAAAGCTTTGAAAATGGATGGTTTT
GGGTATTAAGAAAGAAAAAATTTAAAAAAGACACTAACGGCCAGTGAGTTGGAGTCTC
AGGGCAGGGTGGCAGTTTCCCTTGAGCAAAGCAGCCAGACGTTGAAGTGTGTTTCCCTTCCCTGGG
CGCAGGGTGCAGGGTGTCTTCCGGATCTGGTGTGACCTTGGTCCAGGAGTTCTATTTGTTCCCTGGG
GAGGGAGGTTTTTTTGTGTTTTTTGGGTTTTTTTGGTGTCTTGTTTTCTTCTCCTCCATGTGT
CTTGGCAGGCACTCATTTCTGTGGCTGTGCGCCAGAGGGAATGTTCTGGAGCTGCCAAGGAGGGAG
GAGACTCGGGTTGGCTAATCCCGGATGAACGGTGCTCCATTGCGACCTCCCTCCTCGTGCCCTGC
CCTGCCCTCTCCACGCACAGTGTTAAGGAGCCAAGAGGAGCCACTTCGCCCAGACTTTGTTTCCCCA
CCTCCTGCGGCATGGGTGTGTCCAGTGCCACCGCTGGCCTCCGCTGCTTCCATCAGCCCTGTGCGC
ACCTGGTCCCTCATGAAGAGCAGACACTTAGAGGCTGGTCCGGAATGGGGAGGTGCGCCCTGGGAG
GGCAGGGCTTGGTTCCAAGCCGGTTCCCGTCCCTGGCGCTGGAGTGACACAGCCAGTCCGGCAC
CTGTTGGCTGGAAGCCAACTGCTTTAGATCACTCGGGTCCCACTTAGAAGGGTCCCCGCTTA
GATCAATCACGTGGACACTAAGGCACGTTTTTAGAGTCTCTTGTCTTAATGATTATGTCATCCGT
TGTCGGTCCATTTGTGTTTTCTGCGTGTGTCTTGGATATAATCCTCAGAAATAATGCACACTAG
CCTCTGACAACCATGAAGCAAAAATCCGTACATGTGGGTCTGAACTTGTAGACTCGGTCCACAGTA
TCAAATAAATCTATAACAGAAAAA

41/392

FIGURE 40

MRQTIKVIKFILIIICYTVYYVHNIKFDVDCTVDIESLTGYRTYRCAHPLATLFKILASF
YISLVIFYGLICMYTLWWMLRRSLKKYSFESIREESSYSDIPDVKNDFAFMLHLIDQYDP
LYSKRFAVFLSEVSENKLRQLNLNNEWTLDKLRQLTKNAQDKLELHFLMLSGIPDTVFD
LVELEVCLKLELIPDVTIPPSIAQLTGLKELWLYHTAAKIEAPALAFLRENLRALHIKFTD
IKEIPLWIYSLKTLEELHLTGNLSAENNRYIVIDGLRELKRLKVLRLKSNLSKLPQVVTD
VGVHLQKLSINNEGTKLIVLNSLKMANLTELELIRCDLERIPHSIFSLHNLQEIDLKDN
NLKTIEEIIISFQHLHRLTCLKLWYNHAIYIPIQIGNLTNLERLYLNRNKIEKIPTQLFYC
RKLRYLDLSHNNLTFLPADIGLLQNLQNLAITANRIETLPPELFQCRKLRLHGLGNNVLQ
SLPSRVGELTNLTQIELRGNRLECLPVELGECPLLKRSGLVVEEDLFNTLPPEVKERLWR
ADKEQA

Transmembrane domain:

amino acids 51-75 (type II)

N-glycosylation site:amino acids 262-266, 290-294, 328-332, 396-400, 432-436,
491-495**cAMP- and cGMP-dependent protein kinase phosphorylation site:**

amino acids 85-89

Casein kinase II phosphorylation site:amino acids 91-95, 97-101, 177-181, 253-257, 330-334, 364-368,
398-402, 493-497**N-myristoylation site:**

amino acids 173-179, 261-267, 395-401, 441-447

42/392

FIGURE 41

GGGGGAGAAGGCGGCCGAGCCCCAGCTCTCCGAGCACCGGGTCGGAAGCCGCGACCCGAG
CCGCGCAGGAAGCTGGGACCGGAACCTCGGCGGACCCGGCCCCACCCAACTCACCTGCGC
AGGTCACCAGCACCCCTCGGAACCCAGAGGCCCGCGCTCTGAAGGTGACCCCCCTGGGGAG
GAAGGCGATGGCCCCCTGCGAGGACGATGGCCCCGCGCCCGCTCGCCCCGGCCGGCATCCC
TGCCGTGCGCCTTGTGGCTTCTGTGCACGCTCGGCCTCCAGGGCACCCAGGCCGGGCCACC
GCCCCGCCCCCTGGGCTGCCCCGCGGAGCCGACTGCCTGAACAGCTTTACCGCCGGGGT
GCCTGGCTTCGTGCTGGACACCAACGCCCTCGGTGAGCAACGGAGCTACCTTCCTGGAGTC
CCCCACCGTGCGCCGGGGCTGGGACTGCGTGCAGCGCTGCTGCACCACCCAGAACTGCAA
CTTGGCGCTAGTGGAGCTGCAGCCCGACCGCGGGGAGGACGCCATCGCCGCTGCTTCCT
CATCAACTGCCTCTACGAGCAGAACTTCGTGTGCAAGTTCGCGCCAGGAGGGCTTCAT
CAACTACCTCACGAGGGAAGTGTACCGCTCCTACCGCCAGCTGCGGACCCAGGGCTTTGG
AGGGTCTGGGATCCCCAAGGCCTGGGCAGGCATAGACTTGAAGGTACAACCCAGGAACC
CCTGGTGCTGAAGGATGTGAAAAACACAGATTGGCGCCTACTGCGGGGTGACACGGATGT
CAGGGTAGAGAGGAAAGACCCAAACCAGGTGGAACCTGTGGGGACTCAAGGAAGGCACCTA
CCTGTTCCAGCTGACAGTGACTAGCTCAGACCACCCAGAGGACACGGCCAACGTCACAGT
CACTGTGCTGTCCACCAAGCAGACAGAAGACTACTGCCTCGCATCCAACAAGGTGGGTGCG
CTGCCGGGGCTCTTTCCACGCTGGTACTATGACCCACGGAGCAGATCTGCAAGAGTTT
CGTTTATGGAGGCTGCTTGGGCAACAAGAACAACCTACCTTCGGGAAGAAGAGTGCAATTCT
AGCCTGTGCGGGGTGTGCAAGGTGGGCCTTTGAGAGGCAGCTCTGGGGCTCAGGCGACTTT
CCCCCAGGGCCCCCTCCATGGAAGGCGCCATCCAGTGTGCTCTGGCACCTGTGACCCAC
CCAGTTCGCGTGCAGCAATGGCTGCTGCATCGACAGTTTCCTGGAGTGTGACGACACCCC
CAACTGCCCCGACGCTCCGACGAGGCTGCCTGTGAAAAATACACGAGTGGCTTTGACGA
GCTCCAGCGCATCCATTTCCCCAGTGACAAAGGGCACTGCGTGGACCTGCCAGACACAGG
ACTCTGCAAGGAGAGCATCCCCGCGTGGTACTACAACCCCTTCAGCGAACACTGCGCCCCG
CTTTACCTATGGTGGTTGTTATGGCAACAAGAACAACCTTTGAGGAAGAGCAGCAGTGCCT
CGAGTCTTGTGCGGGCATCTCCAAGAAGGATGTGTTTGGCCTGAGGCGGGAAATCCCCAT
TCCCAGCACAGGCTCTGTGGAGATGGCTGTACAGTGTTCCTGGTCATCTGCATTGTGGT
GGTGGTAGCCATCTTGGGTACTGCTTCTTCAAGAACCAGAGAAAGGACTTCCACGGACA
CCACCACCACCCACCCACCCCTGCCAGCTCCACTGTCTCCACTACCGAGGACACGGGA
GCACCTGGTCTATAACCACACACCCCGCCCCCTCTGAGCCTGGGTCTCACCGGCTCTCAC
CTGGCCCTGCTTCCTGCTTGCCAAGGCAGAGGCCTGGGCTGGGAAAAACTTTGGAACCAG
ACTCTTGCCCTGTTTCCAGGCCCACTGTGCCTCAGAGACCAGGGCTCCAGCCCCTCTTGG
AGAAGTCTCAGCTAAGCTCACGTCTGAGAAAGCTCAAAGGTTTGAAGGAGCAGAAAAC
CCTTGGGCCAGAAAGTACCAGACTAGATGGACCTGCCTGCATAGGAGTTTGGAGGAAGTTG
GAGTTTTGTTTCTCTGTTCAAAGCTGCCTGTCCCTACCCCATGGTGCTAGGAAGAGGAG
TGGGGTGGTGTGACAGCCCTGGAGGCCCAACCCCTGTCTCCTCCGAGCTCCTCTTCATGCT
GTGCGCCACAGGCTGGGAGGAAGGACTTCCCTGTGTAGTTTGTGCTGTAAAGAGTTGCTT
TTTGTTTATTTAATGCTGTGGCATGGGTGAAGAGGAGGGGAAGAGGCCTGTTTGGCCTCT
CTGTCTCTCTTCTCTTCCCCCAAGATTGAGCTCTCTGCCCTTGATCAGCCCCACCCCTG
GCCTAGACCAGCAGACAGAGCCAGGAGAGGCTCAGCTGCATTCCGAGCCCCCACCCTCA
AGGTTCTCCAACATCACAGCCAGCCACCCACTGGGTAATAAAAGTGGTTTGTGGAAAA
AAAAAAAAAAAAAAAAAAAAA

43/392

FIGURE 42

MAPARTMARARLAPAGIPAVALWLLCTLGLQGTQAGPPPAPPGLPAGADCLNSFTAGVPG
 FVLDTNASVSNGATFLESPTVRRGWDCVRACCTTQNCNLALVELQPDREGDAIAACFLIN
 CLYEQNFVCKFAPREGFINYLTVREYRSYRQLRTQGFGGSGIPKAWAGIDLKVQPQEPLV
 LKDVENTDWRLLRGDTDVRVERKDPNQVELWGLKEGTYLFQLTVTSSDHPEDTANVTVTV
 LSTKQTEDYCLASNKVGRCRGSFPRWYYDPTEQICKSFVYGGCLGNKNNYLREEECILAC
 RGVQGGPLRGSSGAQATFPQGSPMERRHPVCSGTCQPTQFRCSNGCCIDSFLECDDTPNC
 PDASDEAAACEKYTSGFDELQRIHFPSDKGHCVDL PDTGLCKESI PRWYYPFSEHCARFT
 YGGCYGNKNNFEEEEQQCLESCRGISKDVFGLRREIPI PSTGSVEMAVTVFLVICIVVVV
 AILGYCFFKNQRKDFHGHHPPTPASSTVSTTEDTEHLVYNHTTRPL

signal sequence:

Amino acids 1-35

transmembrane domain:

Amino acids 466-483

N-glycosylation sites:

Amino acids 66-70;235-239;523-527

N-myristoylation sites:

A m i n o a c i d s
 29-35;43-49;161-167;212-218;281-287;282-288;285-291;
 310-316;313-319;422-428;423-429;426-432

Cell attachment sequence:

Amino acids 193-199

Pancreatic trypsin inhibitor (Kunitz) family signatures:

Amino acids 278-298;419-438

44/392

FIGURE 43

CCCACGCGTCCGCACCTCGGCCCCGGGCTCCGAAGCGGCTCGGGGGCGCCCTTTCGGTCA
ACATCGTAGTCCACCCCCCTCCCCATCCCCAGCCCCCGGGGATTCAGGCTCGCCAGCGCCC
AGCCAGGGAGCCGGCCGGGAAGCGCGATGGGGGGCCCCAGCCGCTCGCTCCTGCTCCTGC
TCCTGCTGTTTCGCCTGCTGCTGGGCGCCCCGGCGGGGCCAACCTCTCCCAGGACGACAGCC
AGCCCTGGACATCTGATGAAACAGTGGTGGCTGGTGGCACCGTGGTGGTCAAGTGCCAAAG
TGAAAGATCACGAGGACTCATCCCTGCAATGGTCTAACCTGCTCAGCAGACTCTCTACT
TTGGGGAGAAGAGAGCCCTTCGAGATAATCGAATTCAGCTGGTTACCTCTACGCCCCACG
AGCTCAGCATCAGCATCAGCAATGTGGCCCTGGCAGACGAGGGCGAGTACACCTGCTCAA
TCTTCACTATGCCTGTGCGAACTGCCAAGTCCCTCGTCACTGTGCTAGGAATTCACAGA
AGCCCATCATCAGTGGTTATAAATCTTCATTACGGGAAAAAGACACAGCCACCCTAAACT
GTCAGTCTTCTGGGAGCAAGCCTGCAGCCCGGCTCACCTGGAGAAAGGGTGACCAAGAAC
TCCACGGGAGAACCAACCCGCATACAGGAAGATCCCAATGGTAAACCTTCACTGTCTAGCA
GCTCGGTGACATTCCAGGTTACCCGGGAGGATGATGGGGCGAGCATCGTGTGCTCTGTGA
ACCATGAATCTCTAAAGGGAGCTGACAGATCCACCTCTCAACGCATTGAAGTTTTATACA
CACCAACTGCGATGATTAGGCCAGACCCTCCCCATCCTCGTGAGGGCCAGAAGCTGTTGC
TACACTGTGAGGGTCGCGGCAATCCAGTCCCCCAGCAGTACCTATGGGAGAAGGAGGGCA
GTGTGCCACCCCTGAAGATGACCCAGGAGAGTGCCCTGATCTTCCCTTTCCTCAACAAGA
GTGACAGTGGCACCTACGGCTGCACAGCCACCAGCAACATGGGCAGCTACAAGGCCTACT
ACACCCTCAATGTTAATGACCCAGTCCGGTGCCCTCCTCCTCCAGCACCTACCACGCCA
TCATCGGTGGGATCGTGGCTTTCATTGTCTTCTGCTGCTCATCATGCTCATCTTCCTTG
GCCACTACTTGATCCGGCACAAAGGAACCTACCTGACACATGAGGCAAAAGGCTCCGACG
ATGCTCCAGACGCGGACACGGCCATCATCAATGCAGAAGGCGGGCAGTCAGGAGGGGACG
ACAAGAAGGAATATTTATCTAGAGGGCGCTGCCCACTTCTGCGCCCCCAGGGGCCCT
GTGGGGACTGCTGGGGCCGTCACCAACCCGGACTTGACAGAGCAACCGCAGGGCCGCCCC
CTCCCGCTTGCTCCCCAGCCCACCCACCCCTGTACAGAATGTCTGCTTTGGGTGCGGT
TTTGTACTCGGTTTGGAATGGGGAGGGAGGAGGGCGGGGGGAGGGGAGGGTTGCCCTCAG
CCCTTTCGCTGGCTTCTCTGCATTTGGGTATTATTATTTTGTAAACATCCCAAATCAA
ATCTGTCTCCAGGCTGGAGAGGCAGGAGCCCTGGGGTGAGAAAAGCAAAAAACAAACAA
AAACA

45/392

FIGURE 44

MGAPAASLLLLLLLLFACCWAPGGANLSQDDSQPWTSDETVVAGGTVVVKCQVKDHEDSSL
QWSNPAQQTLYFGEKRALRDNRIQLVTSTPHELSSISNVALADEGEYTCSTFTMPVRTA
KSLVTVLGIPQKPIITGYKSSLREKDTATLNCQSSGSKPAARLTWRKGDQELHGEPTRIQ
EDPNGKTFTVSSSVTFQVTREDDGASIVCSVNHESLKGADRSTSQRIEVLYTPTAMIRPD
PPHPREGQKLLHCEGRGNPVPQQYLWEKEGSVPPLKMTQESALIFPFLNKSDSGTYGCT
ATSNMGSYKAYYTLNVNDPSPVPSSSSTYHAIIGGIVAFIVFLLLIIMLIFLGHYLIRHKG
TYLTHEAKGSDDAPDADTAIINAEGGQSGGDDKKEYFI

Signal sequence:
amino acids 1-20

Transmembrane domain:
amino acids 331-352

N-glycosylation site:
amino acids 25-29, 290-294

Casein kinase II phosphorylation site:
amino acids 27-31, 35-39, 89-93, 141-145, 199-203, 388-392

N-myristoylation site:
amino acids 2-8, 23-29, 156-162, 218-224, 295-301, 298-304,
306-310, 334-340, 360-364, 385-389, 386-390

Prokaryotic membrane lipoprotein lipid attachment site:
amino acids 7-18

46/392

FIGURE 45

ACTTGCCATCACCTGTTGCCAGTGTGGAAAAATTCTCCCTGTTGAATTTTTTGCACATGG
AGGACAGCAGCAAAGAGGGCAACACAGGCTGATAAGACCAGAGACAGCAGGGAGATTATT
TTACCATACGCCCTCAGGACGTTCCCTCTAGCTGGAGTTCTGGACTTCAACAGAACCCCA
TCCAGTCATTTTGATTTTGCTGTTTATTTTTTTTTTTCTTTTCTTTTCCCACCACATTG
TATTTTATTTCCGTACTTCAGAAATGGGCCTACAGACCACAAAGTGGCCCAGCCATGGGG
CTTTTTTCTGAAGTCTTGGCTTATCATTTCCTGGGGCTCTACTCACAGGTGTCCAAAC
TCCTGGCCTGCCCTAGTGTGTGCCGCTGCGACAGGAACTTTGTCTACTGTAATGAGCGAA
GCTTGACCTCAGTGCCTCTTGGGATCCCGGAGGCGTAACCGTACTCTACCTCCACAACA
ACCAAATTAATAATGCTGGATTTCCTGCAGAACTGCACAATGTACAGTCGGTGCACACGG
TCTACCTGTATGGCAACCAACTGGACGAATTCCTCATGAACCTTCCCAAGAATGTCAGAG
TTCTCCATTTGCAGGAAAACAATATTAGACCATTTTCACGGGCTGCTCTTGGCCAGCTCT
TGAAGCTTGAAGAGCTGCACCTGGATGACAACTCCATATCCACAGTGGGGGTGGAAGACG
GGGCCTTCCGGGAGGCTATTAGCCTCAAATTGTTGTTTTTGTCTAAGAATCACCTGAGCA
GTGTGCCTGTTGGGCTTCTGTGGACTTGCAAGAGCTGAGAGTGGATGAAAATCGAATTG
CTGTCATATCCGACATGGCCTTCCAGAATCTCACGAGCTTGGAGCGTCTTATTGTGGACG
GGAACCTCCTGACCAACAAGGGTATCGCCGAGGGCACCTTCAGCCATCTCACCAAGCTCA
AGGAATTTTCAATTGTACGTAATTCGCTGTCCACCCCTCCTCCCGATCTCCCAGGTACGC
ATCTGATCAGGCTCTATTTGCAGGACAACCAGATAAACCACATTCCTTTGACAGCCTTCT
CAAATCTGCGTAAGCTGGAACGGCTGGATATATCCAACAACCAACTGCGGATGCTGACTC
AAGGGGTTTTTGTATAATCTCTCCAACCTGAAGCAGCTCACTGCTCGGAATAACCCTTGGT
TTTGTGACTGCAGTATTAAATGGGTACAGAATGGCTCAAATATATCCCTTCATCTCTCA
ACGTGCGGGGTTTCATGTGCCAAGGTCCTGAACAAGTCCGGGGGATGGCCGTCAGGGAAT
TAAATATGAATCTTTTGTCTGTCCCACCACGACCCCCGGCCTGCCTCTCTTCACCCAG
CCCCAAGTACAGCTTCTCCGACCACTCAGCCTCCCACCCTCTCTATTCCAAACCCCTAGCA
GAAGCTACACGCCTCCAACCTCCTACCACATCGAACTTCCCACGATTCCTGACTGGGATG
GCAGAGAAAGAGTGACCCACCTATTTCTGAACGGATCCAGCTCTCTATCCATTTTGTGA
ATGATACTTCCATTCAAGTCAGCTGGCTCTCTCTCTTCACCGTGATGGCATACAAACCTCA
CATGGGTGAAAATGGGCCACAGTTTAGTAGGGGGCATCGTTCAGGAGCGCATAGTCAGCG
GTGAGAAGCAACACCTGAGCCTGGTTAACTTAGAGCCCCGATCCACCTATCGGATTTGTT
TAGTGCCACTGGATGCTTTTAACTACCGCGCGGTAGAAGACACCATTGTTTCAGAGGCCA
CCACCCATGCCTCCTATCTGAACAACGGCAGCAACACAGCGTCCAGCCATGAGCAGACGA
CGTCCCACAGCATGGGCTCCCCCTTCTGCTGGCGGGCTTGATCGGGGGCGCGGTGATAT
TTGTGCTGGTGGTCTTGCTCAGCGTCTTTTGTGTCATATGCACAAAAAGGGGCGCTACA
CCTCCCAGAAGTGGAATAACAACGGGGCGGGCGGAAAGATGATTATTGCGAGGCAGGCA
CCAAGAAGGACAACTCCATCCTGGAGATGACAGAAACCAGTTTTTCAGATCGTCTCCTTAA
ATAACGATCAACTCCTTAAAGGAGATTTAGACTGCAGCCCATTTACACCCCAAATGGGG
GCATTAATTACACAGACTGCCATATCCCCAACAACATGCGATACTGCAACAGCAGCGTGC
CAGACCTGGAGCACTGCCATACGTGACAGCCAGAGGCCAGCGTTATCAAGGCGGACAAT
TAGACTCTTGAGAACACACTCGTGTGTGCACATAAAGACACGCAGATTACATTTGATAAA
TGTTACACAGATGCATTTGTGCATTTGAATACTCTGTAATTTATACGGTGTACTATATAA
TGGGATTTAAAAAAGTGCTATCTTTTCTATTTCAAGTTAATTACAAACAGTTTTGTAAC
TCTTTGCTTTTTTAAATCTT

47/392

FIGURE 46

MGLQTTKWPSHGAFFLKSWLIISLGLYSQVSKLLACPSVCRCDRNFVYCNERSLTSVPLG
IPEGVTVLYLHNNQINNAGFPAELHNVQSVHTVYLYGNQLDEFPMNLPKNVRVLHLQENN
IQTISRALAQLLKLEELHLLDDNSISTVGVEDGAFREAI SLKLLFLSKNHLSSVPVGLPV
DLQELRV DENRIAVISDMAFQNLTSLERLIVDGNLLTNKGIAEGTFSHLTKLKEFSIVRN
SLSHPPPDLPGTHLIRLYLQDNQINHIPLTAFSNLRKLERLDISNNQLRMLTQGVFDNLS
NLKQLTARNNPWFCDCSI KWVTEWLKYIPSSLNVRGFMCGPEQVRGMAVRELNMNLLSC
PTTTPGLPLFTPAPSTASPTTQPPTLSIPNPSRSYTPPTPTTSKLPTIPDWDGRERVTPP
ISERIQLSIHVNDTSIQVSWLSLFTVMAYKLTWVKMGHSLVGGIVQERIVSGEKQHLSL
VNLEPRSTYRICLVPLDAFN YRAVEDTICSEATTHASYLNNGSNTASSHEQTTSHSMGSP
FLLAGLIGGAVIFVLVLLSVFCWHMHKKGRYTSQKWKNRGRRKDDYCEAGTKKDNSIL
EMTETSFQIVSLNNDQLLKGD FRLQPIYTPNGGINYTDCHI PNMRYCNSSVPDLEHCHT

Signal peptide:

amino acids 1-42

Transmembrane domain:

amino acids 542-561

N-glycosylation site:amino acids 202-206, 298-302, 433-437, 521-525, 635-639,
649-653**Casein kinase II phosphorylation site:**amino acids 204-208, 407-411, 527-531, 593-597, 598-602,
651-655**Tyrosine kinase phosphorylation site:**

amino acids 319-328

N-myristoylation site:amino acids 2-8, 60-66, 149-155, 213-219, 220-226, 294-300,
522-528, 545-551, 633-639**Amidation site:**

amino acids 581-585

Leucine zipper pattern:

amino acids 164-186

Phospholipase A2 aspartic acid active site:

amino acids 39-50

48/392

FIGURE 47

GCAGCGAGCGCCGGGTGCGGCCCTGCCGCCGACGGGATGTGACCTTCACCGTCGCTTAGC
CAGGATGACCGGAGCCCGTGTCTCGCGGCGTCCGCGCCTCGCTTCAGCCTCCCGGGTGCT
CTGACCGCACGCTCCCGGTGCTAGGCTCCCCGGCACC GGCCCTCGCCATGCCCGCCACCGC
CCGGGCCCGCCGCCCGCCCTGGGCACTGCGCTTCTGCTGCTCCTGCTGGCTTCCGAGTCTT
CTCACACTGTGCTGTTGCGGGCGCGTGAGGCGGCGCAGTTTCTGCGGCCCAGGCAGCGCC
GCGCCTACCAAGTCTTCGAGGAGGCCAAGCAGGGCCACCTGGAACGGGAGTGCGTGAGG
AGGTGTGCAGCAAAGAGGAGGCCAGAGAGGTGTTGAGAACGACCCCGAGACGGAGTATT
TCTATCCACGATATCAAGAGTGCATGAGAAAATATGGCAGGCCTGAAGAAAAAACCCAG
ATTTCCGCCAAATGTGTTTCAAACTTGCCCTGACCACTGCACCCCAAACCCCTTGATGAAGA
AGGGTACTCATATCTGCCAAGACCTCATGGGCAACTTCTTCTGCGTGTGCACAGATGGCT
GGGGAGGCCGGCTCTGTGACAAAGATGTCAATGAGTGTGTCCAGAAGAATGGGGCTGCA
GCCAGGTCTGCCACAACAAACCAGGAAGCTTCCAATGTGCCTGCCATAGTGGCTTCTCGC
TTGCATCAGACGGCCAGACCTGCCAAGATATCGATGAATGCACAGACTCAGACACCTGTG
GGGACGCGCGATGCAAGAACTTGCCAGGCTCCTACTCTTGCCTCTGCGATGAGGGATATA
CATACAGCTCCAAGGAGAAGACCTGCCAAGATGTGGACGAGTGCCAGCAGGATCGCTGTG
AGCAGACCTGTGTCAACTCCCCAGGCAGCTATACCTGCCACTGTGATGGGCGAGGGGGCC
TAAAACTATCCCCAGACATGGATACCTTGTGAGGACATCTTACCATGTGTGCCCTTCAGCA
TGGCCAAGAGCGTGAAGTCCTTGTACCTGGGCGCATGTTTACGCGGGACCCCGTGATTA
GACTACGCTTCAAGAGGCTTCAGCCTACCAGGCTGCTGGCTGAATTTGACTTCCGCACTT
TTGACCCCTGAAGGAGTCTCTTCTTCGCTGGAGGCCGTTTACAGACAGCACCTGGATTGTCC
TGGGCCTAAGAGCTGGGCGGCTTGAGCTGCAGCTTCCGTACAATGGCGTTGGGCGCATCA
CCAGCAGCGGGCCAACCATCAACCACGGCATGTGGCAAACTATCTCCGTGGAAGAGCTGG
AACGTAACCTTGTATCAAGGTCAACAAAGATGCTGTAATGAAGATCGCGGTAGCTGGGG
AGCTGTTTACAGCTGGAGAGGGGCTCTATCACCTGAATCTCACCGTGGGCGGCATTCCCT
TCAAGGAGAGTGTGCTCGTCCAGCCGATTAACCTCGCCTGGATGGGTGCATGAGGAGTT
GGAAGTGGCTGAACGGGGAAGACAGCGCCATCCAGGAGACAGTCAAGGCAAACAAAAA
TGCAGTGTCTTCTGTGACAGAAAGGGCTCCTTCTTCCCGGGGAATGGATTTGCTACCT
ACAGGCTCAACTACACCCGAACATCGCTGGATGTGCGCACGGAAACACCTGGGAAGTTA
AAGTTGTGGCTCGGATCCGCCCTGCCACGGACACGGGGGTGCTGCTGGCGCTGGTGGGGG
ACGACGATGTGCTCATCTCTGTGGCCCTAGTCGACTACCACTCTACAAAGAAGCTCAAGA
AGCAGTTGGTGGTCTTGGCAGTTGAGGATGTTGCCCTGGCACTGATGGAAATCAAGGTGT
GCGACAGCCAGGAACACACGGTCACTGTCTCCCTGCGGGAGGTTGAGGCCACCCTAGAAG
TGGATGGCACAAAGGGCCAGAGTGAAGTGAGCACTGCCCAGCTGCAGGAGCGACTGGACA
CACTTAAGACACATCTGCAAGGCTCTGTGCACACCTATGTTGGAGGCCTGCCAGAAGTAT
CGGTGATTTCTGCACCCGTCCTGCGTTCTACCGCGGATGCATGACTCTGGAGGTAAACG
GGAAAATCCTGGACCTGGATACGGCCTCGTACAAGCACAGTGACATCACCTCCCACTCCT
GCCCCCTGTGGAGCATGCCACCCCTTAGACCGAGCTGCAAGAGGGCTCCACACCTAAAG
ACAAAAATGAAGCAGGGTTTGGACACACAGCACTGGCTCCTCTCGCATGGTCTTGCAACA
CTGGAGCAGCGTGGACCGCCCTTGTGGTTTTTTTTTCTTGAGATCTTTCTTTTGCCTTG
TAACATATCTGTACATAATGGACGGGTGTGCGGTACCGGCTGCTCAGAGAGAGCCACGT
GACCTGGTGGGAGCTGGCTGGAAGGGGCTGGGCTAGAGGGGCTGGCAGTTTGCAGCAGAA
CGGATGTGAAGAAAATAATCTCTATTATTTTATTACTACATGCTTCTTTCTGACTCTA
AAATATGGAAAATAAAATATTTACAGAAACCTTTTAAAAA

49/392

FIGURE 48

MPPPPGPAAALGTALLLLLLLASESSHTVLLRAREAAQFLRPRQRRAYQVFEEAKQGHLE
ECVVEVCSKEEAREVFENDPETEYFYPRYQECMRKYGRPEEKNPDAKCVQNLDPDQCTPN
PCDKKGTHICQDLMGNFVCVCTDGGWGGRLCDKDVNECVQKNGGCSQVCHNKPGSFQCACH
SGFSLASDGQTCQDIDECTSDTCGDARCKNLPGSYSCLCDEGYTYSSKEKTCQDVDECQ
QDRCEQTCVNSPGSYTCHCDGRGGLKLSPDMDTCEDILPCVPFMAKSVKSLYLGRMFSG
TPVIRLRFKRLQPTRLLAEFDFTFDPEGVLFFAGGRSDSTWIVLGLRAGRLELQLRYNG
VGRITSSGPTINHGMWQTISVEELERNLVIKVNKDAVMKIAVAGELFQLERGLYHLNLT
GGIPFKESELVQPINPRLDGCMRSWNWLNGEDSAIQETVKANTKMQCFSVTERGSFFPGN
GFATYRLNYTRTSLDVGTETTWEVKVVARIRPATDTGVLLALVGDDDVVISVALVDYHST
KKLKKQLVVLAVEDVALALMEIKVCDSQEHTVTVSLREGEATLEVDGTKGQSEVSTAQLQ
ERLDTLKTHLQGSVHTYVGGLPEVSVISAPVTAFYRGCMTLEVNGKILDLDTASYKHSDI
TSHSCPPVEHATP

50/392

FIGURE 49

CGCCGCGCTCCCGCACCCGCGGCCCGCCACCGCGCCGCTCCCGCATCTGCACCCGCAGC
CCGGCGGCCTCCCGGCGGGAGCGAGCAGATCCAGTCCGGCCCGCAGCGCAACTCGGTCCA
GTCGGGCGGCGGGCTGCGGGCGCAGAGCGGAGATGCGAGCGGCTTGGGGCCACCCTGCTGT
GCCTGCTGCTGGCGGCGGCGGTCCCCACGGCCCCCGCGCCCGCTCCGACGGCGACCTCGG
CTCCAGTCAAGCCCGGCCCGGCTCTCAGTACCCGCAGGAGGAGGCCACCCTCAATGAGA
TGTTCCGCGAGGTTGAGGAACTGATGGAGGACACGCAGCACAAATTGCGCAGCGCGGTGG
AAGAGATGGAGGCAGAGAAGCTGCTGCTAAAGCATCATCAGAAAGTGAACCTGGCAAAC
TACCTCCCAGCTATCACAATGAGACCAACACAGACACGAAGGTTGGAAATAATACCATCC
ATGTGCACCGAGAAATTCACAAGATAACCAACAACCAGACTGGACAAATGGTCTTTTCAG
AGACAGTTATCACATCTGTGGGAGACGAAGAAGGCAGAAGGAGCCACGAGTGCATCATCG
ACGAGGACTGTGGGCCCAGCATGTACTGCCAGTTTGCCAGCTTCCAGTACACCTGCCAGC
CATGCCGGGGCCAGAGGATGCTCTGCACCCGGGACAGTGAGTGCTGTGGAGACCAGCTGT
GTGTCTGGGGTCACTGCACCAAATGGCCACCAGGGGCAGCAATGGGACCATCTGTGACA
ACCAGAGGGACTGCCAGCCGGGGCTGTGCTGTGCCCTTCCAGAGAGGCCTGCTGTTCCCTG
TGTGCACACCCCTGCCCGTGGAGGGCGAGCTTTGCCATGACCCCGCCAGCCGGCTTCTGG
ACCTCATACCTGGGAGCTAGAGCCTGATGGAGCCTTGGACCGATGCCCTTGTGCCAGTG
GCCTCCTCTGCCAGCCCCACAGCCACAGCCTGGTGTATGTGTGCAAGCCGACCTTCTGTG
GGAGCCGTGACCAAGATGGGGAGATCCTGCTGCCCAGAGAGGTCCCGATGAGTATGAAG
TTGGCAGCTTCATGGAGGAGGTGCCCCAGGAGCTGGAGGACCTGGAGAGGAGCCTGACTG
AAGAGATGGCGCTGGGGGAGCCTGCCGGCTGCCCGCGCTGCACTGCTGGGAGGGGAAGAGA
TTTAGATCTGGACCAGGCTGTGGGTAGATGTGCAATAGAAATAGCTAATTTATTTCCCCA
GGTGTGTGCTTTAGGCGTGGGCTGACCAGGCTTCTTCCCTACATCTTCTTCCCAGTAAGTT
TCCCCTCTGGCTTGACAGCATGAGGTGTTGTGCATTTGTTTCAGCTCCCCCAGGCTGTTCT
CCAGGCTTCACAGTCTGGTGTCTGGGAGAGTCAGGCAGGGTTAAACTGCAGGAGCAGTTT
GCCACCCCTGTCCAGATTATTGGCTGCTTTGCCTCTACCAGTTGGCAGACAGCCGTTTGT
TCTACATGGCTTTGATAATTGTTTGAGGGGAGGAGATGGAAACAATGTGGAGTCTCCCTC
TGATTGGTTTTTGGGAAATGTGGAGAAGAGTGCCCTGCTTTGCAAACATCAACCTGGCAA
AAATGCAACAAATGAATTTTCCACGCAGTCTTTTCCATGGGCATAGGTAAGCTGTGCCTT
CAGCTGTTGCAGATGAAATGTTCTGTTACCCCTGCATTACATGTGTTTATTCATCCAGCA
GTGTTGCTCAGCTCCTACCTCTGTGCCAGGGCAGCATTTTTCATATCCAAGATCAATTCCC
TCTCTCAGCACAGCCTGGGGAGGGGGTCATTGTTCTCCTCGTCCATCAGGGATCTCAGAG
GCTCAGAGACTGCAAGCTGCTTGCCCAAGTCACACAGCTAGTGAAGACCAGAGCAGTTTC
ATCTGGTTGTGACTCTAAGCTCAGTGCTCTCTCCACTACCCACACCAGCCTTGGTGCCA
CCAAAAGTGCTCCCCAAAAGGAAGGAGAATGGGATTTTCTTGGAGGCATGCACATCTGGA
ATTAAGGTCAAACATAATTCTCACATCCCTCTAAAAGTAACTACTGTTAGGAACAGCAGT
GTTCTCACAGTGTGGGGCAGCCGTCTTCTAATGAAGACAATGATATTGACACTGTCCCT
CTTTGGCAGTTGCATTAGTAACTTTGAAAGGTATATGACTGAGCGTAGCATACAGGTTAA
CCTGCAGAAACAGTACTTAGGTAATTGTAGGGCGAGGATTATAAATGAAATTTGCAAAAT
CACTTAGCAGCAACTGAAGACAATTATCAACCACGTGGAGAAAATCAAACCGAGCAGGGC
TGTGTGAAACATGGTTGTAATATGCGACTGCGAACACTGAACTCTACGCCACTCCACAAA
TGATGTTTTTCAGGTGTCATGGACTGTTGCCACCATGTATTTCATCCAGAGTTCTTAAAGTT
TAAAGTTGCACATGATTGTATAAGCATGCTTTCTTTGAGTTTTTAAATTATGTATAAACAT
AAGTTGCATTTAGAAATCAAGCATAAATCACTTCAACTGCAAAAAAAAAAAAAAAAAAAAA
AAAAAA

51/392

FIGURE 50

MQRLGATLLCLLLAAVPTAPAPAPTATSAPVKPGPALSYPQEEATLNEMFREVEELMED
TQHKLRSAVEEMEAEEAAKASSEVNLANLPPSYHNETNTDTKVGNNTIHVHREIHKITN
NQTGQMVFSETVITSVGDEEGRRSHECIIDEDCGPSMYCQFASFQYTCQPCRGQRM LCTR
DSECCGDQLCVWGHCTKMATRGSNGTICDNQRDCQPGLCCAFQRGLLFPVCTPLPVEGEL
CHDPASRLLDLITWELEPDGALDRCPCASGLLCQPHSHSLVYVCKPTFVGSRDQDGEILL
PREVPDEYEVGSFMEEVRQELEDLERSLTEEMALGEPAAAAAALLGGEI

Signal sequence:

amino acids 1-19

N-glycosylation site:

amino acids 96-100, 106-110, 121-125, 204-208

Casein kinase II phosphorylation site:amino acids 46-50, 67-71, 98-102, 135-139, 206-210, 312-316,
327-331**N-myristoylation site:**

amino acids 202-208, 217-223

Amidation site:

amino acids 140-144

52/392

FIGURE 51

GCCTGTTGCTGATGCTGCCGTGCGGTACTTGTCATGGAGCTGGCACTGCGGCGCTCTCCC
GTCCCGCGGTGGTTGCTGCTGCTGCCGCTGCTGCTGGGCCTGAACGCAGGAGCTGTCATT
GACTGGCCACAGAGGAGGGCAAGGAAGTATGGGATTATGTGACGGTCCGCAAGGATGCC
TACATGTTCTGGTGGCTCTATTATGCCACCAACTCCTGCAAGAACTTCTCAGAACTGCCC
CTGGTCATGTGGCTTCAGGGCGGTCCAGGCGGTTCTAGCACTGGATTTGGAACTTTGAG
GAAATTGGGCCCCCTTGACAGTGATCTCAAACCACGGAAAACCACCTGGCTCCAGGCTGCC
AGTCTCCTATTTGTGGATAATCCCGTGGGCACTGGGTTCA GTTATGTGAATGGTAGTGGT
GCCTATGCCAAGGACCTGGCTATGGTGGCTTCAGACATGATGGTTCCTCTGAAGACCTTC
TTCAGTTGCCACAAAGAATTCCAGACAGTTCATTCTACATTTTCTCAGAGTCTTATGGA
GGAAAAATGGCAGCTGGCATTGGTCTAGAGCTTTATAAGGCCATTACAGCGAGGGACCATC
AAGTGCAACTTTGCGGGGGTTGCCTTGGGTGATTCCCTGGATCTCCCCTGTTGATTCCGGTG
CTCTCCTGGGGACCTTACCTGTACAGCATGTCTCTTCTCGAAGACAAAGGTCTGGCAGAG
GTGTCTAAGGTTGCAGAGCAAGTACTGAATGCCGTAAATAAGGGGCTCTACAGAGAGGCC
ACAGAGCTGTGGGGGAAAGCAGAAATGATCATTGAACAGAACACAGATGGGGTGAACCTTC
TATAACATCTTAACTAAAAGCACTCCACGTCTACAATGGAGTCTGAGTCTAGAATTCACA
CAGAGCCACCTAGTTTGTCTTTGTGTCAGCGCCACGTGAGACACCTACAACGAGATGCCTTA
AGCCAGCTCATGAATGGCCCCATCAGAAAGAAGCTCAAAATTATTCTGAGGATCAATCC
TGGGGAGGCCAGGCTACCAACGTCTTTGTGAACATGGAGGAGGACTTCATGAAGCCAGTC
ATTAGCATTGTGGACGAGTTGCTGGAGGCAGGGATCAACGTGACGGTGTATAATGGACAG
CTGGATCTCATCGTAGATACCATGGGTGAGGAGGCCCTGGGTGCGGAACTGAAGTGGCCA
GAACTGCCTAAATTGAGTCAGCTGAAGTGAAGGCCCTGTACAGTGACCCTAAATCTTTG
GAAACATCTGCTTTTGTCAAGTCCTACAAGAACCTTGCTTTCTACTGGATTCTGAAAGCT
GGTCATATGGTTCTTCTGACCAAGGGGACATGGCTCTGAAGATGATGAGACTGGTGACT
CAGCAAGAATAGGATGGATGGGGCTGGAGATGAGCTGGTTTGGCCTTGGGGCACAGAGCT
GAGCTGAGGCGGCTGAAGCTGTAGGAAGCGCCATTCTTCCCTGTATCTAACTGGGGCTGT
GATCAAGAAGGTTCTGACCAGCTTCTGCAGAGGATAAAATCATTGTCTCTGGAGGCAATT
TGGAATTTATTTCTGCTTCTTAAAAAACCTAAGATTTTTTAAAAAATTGATTTGTTTTG
ATCAAAATAAAGGATGATAATAGATATTAA

53/392

FIGURE 52

MELALRRSPVPRWLLLLPLLLGLNAGAVIDWPTEEGKEVWDYVTVRKDAYMFWWLYYATN
SCKNFSELPLVMWLQGGPGGSSTGFGNFEEIGPLDSLKPRKTTWLQAASLLFVDNPVGT
GFSYVNGSGAYAKDLAMVASDMMVLLKTFFSCHKEFQTVPFYIFSESYGGKMAAGIGLEL
YKAIQRGTIKCNFAGVALGDSWISPVDSVLSWGPYLYSMSLLEDKGLAEVSKVAEQVLNA
VNKGLYREATELWGKAEMIEQNTDGVNFYNILTKSTPTSTMESSLEFTQSHLVCLCQRH
VRHLQRDALSQLMNGPIRKKLKIIPEDQSWGGQATNVFVNMEEDFMKPVISIVDELLEAG
INVTVYNGQLDLIVDTMGQEAWVRKLKWPELPKFSQLKWKALYSDPKSLETSAFVKSYKN
LAFYWILKAGHMVPSDQGDMAKMMRLVTQQE

Signal sequence:
amino acids 1-25

N-glycosylation site:
amino acids 64-68, 126-130, 362-366

cAMP- and cGMP-dependent protein kinase phosphorylation site:
amino acids 101-105

Casein kinase II phosphorylation site:
amino acids 204-208, 220-224, 280-284, 284-288, 351-355,
449-453

N-myristoylation site:
amino acids 22-28, 76-82, 79-85, 80-86, 119-125, 169-175,
187-193, 195-201, 331-337, 332-338, 360-366

54/392

FIGURE 53

GTCTGTTCCCAGGAGTCCTTCGGCGGCTGTTGTGTCAGTGGCCTGATCGCGATGGGGACA
AAGGCGCAAGTCGAGAGGAACTGTTGTGCCTCTTCATATTGGCGATCCTGTTGTGCTCC
CTGGCATTGGGCAGTGTTACAGTGCACCTCTTCTGAACCTGAAGTCAGAATTCCTGAGAAT
AATCCTGTGAAGTTGTCTGTGCCTACTCGGGCTTTTCTTCTCCCCGTGTGGAGTGGAAAG
TTTGACCAAGGAGACACCACCAGACTCGTTTGCTATAATAACAAGATCACAGCTTCCTAT
GAGGACCGGGTGACCTTCTTGCCAACTGGTATCACCTTCAAGTCCGTGACACGGGAAGAC
ACTGGGACATACACTTGTATGGTCTCTGAGGAAGGCGGCAACAGCTATGGGGAGGTCAAG
GTCAAGCTCATCGTGCTTGTGCCTCCATCCAAGCCTACAGTTAACATCCCCCTCCTCTGCC
ACCATGGGAACCGGGCAGTGCTGACATGCTCAGAACAAGATGGTTCCCCACCTTCTGAA
TACACCTGGTTCAAAGATGGGATAGTGATGCCTACGAATCCCAAAAGCACCCGTGCCTTC
AGCAACTCTTCCATGTCTCTGAATCCCACAACAGGAGAGCTGGTCTTTGATCCCCCTGTCA
GCCTCTGATACTGGAGAATACAGCTGTGAGGCACGGAATGGGTATGGGACACCCATGACT
TCAAATGCTGTGCGCATGGAAGCTGTGGAGCGGAATGTGGGGGTTCATCGTGGCAGCCGTC
CTTGTAACCCCTGATTCTCCTGGGAATCTTGGTTTTTGGCATCTGGTTTGCCTATAGCCGA
GGCCACTTTGACAGAACAAAGAAAGGGACTTCGAGTAAGAAGGTGATTTACAGCCAGCCT
AGTGCCCGAAGTGAAGGAGAATTCAAACAGACCTCGTCATTCTTGGTGTGAGCCTGGTCCG
GCTCACCGCCTATCATCTGCATTTGCCTTACTCAGGTGCTACCGGACTCTGGCCCCCTGAT
GTCTGTAGTTTTCACAGGATGCCTTATTTGTCTTCTACACCCACAGGGCCCCCTACTTCT
TCGGATGTGTTTTTAATAATGTCTAGCTATGTGCCCCATCCTCCTTCATGCCCTCCCTCCC
TTTCCTACCACTGCTGAGTGGCCTGGAACCTTGTTTAAAGTGTTTATTCCCCATTTCTTTG
AGGGATCAGGAAGGAATCCTGGGTATGCCATTGACTTCCCTTCTAAGTAGACAGCAAAAA
TGGCGGGGGTCCGAGGAATCTGCACTCAACTGCCCACCTGGCTGGCAGGGATCTTTGAAT
AGGTATCTTGAGCTTGGTTCTGGGCTCTTTCTTGTGTACTGACGACCAGGGCCAGCTGT
TCTAGAGCGGGAATTAGAGGCTAGAGCGGCTGAAATGGTTGTTTGGTGATGACACTGGGG
TCCTTCCATCTCTGGGGCCCACTCTCTTCTGTCTTCCCATGGGAAGTGCCACTGGGATCC
CTCTGCCCTGTCTCCTGAATACAAGCTGACTGACATTGACTGTGTCTGTGGAAAATGGG
AGCTCTTGTTGTGGAGAGCATAGTAAATTTTCAGAGAACTTGAAGCCAAAAGGATTTAAA
ACCGCTGCTCTAAAGAAAAGAAAACCTGGAGGCTGGGCGCAGTGGCTCACGCCTGTAATCC
CAGAGGCTGAGGCAGGCGGATCACCTGAGGTCGGGAGTTCGGGATCAGCCTGACCAACAT
GGAGAAACCCTACTGGAAATACAAAGTTAGCCAGGCATGGTGGTGCATGCCTGTAGTCCC
AGCTGCTCAGGAGCCTGGCAACAAGAGCAAACTCCAGCTCAAAAAAAAAAAAAAAAAA

55/392

FIGURE 54

MGTKAQVERKLLCLFILAILLCSLALGSVTVHSSEPEVRIPENNPVKLS CAYSGFSSPRV
EWKFDQGD TTRLVCYNNKITASYEDRVTF LPTGITFKSVTREDTGT YTCMVSEEGGNSYG
EVKVKLIVLVPPSKPTVNI PSSATIGNRAVLTCSEQDGSP PSEYTWFKDGI VMP TNP KST
RAFSNSSYVLNPTTGELVFDPLSASDTGEYSCEARNGYGT P MTSNAVRME AVERN VGVIV
AAVLVTLILLGILVFGIWFAYS RGHFDR TKKGTSSKKVIYSQPSARSEGEFKQTSSFLV

Signal sequence:
amino acids 1-27

Transmembrane domain:
amino acids 238-255

N-glycosylation site:
amino acids 185-189

cAMP- and cGMP-dependent protein kinase phosphorylation site:
amino acids 270-274

Casein kinase II phosphorylation site:
amino acids 34-38, 82-86, 100-104, 118-122, 152-156,
154-158, 193-197, 203-207, 287-291

N-myristoylation site:
amino acids 105-111, 116-122, 158-164, 219-225, 237-243,
256-262

56/392

FIGURE 55

GTTGTGTCCTTCAGCAAAACAGTGGATTTAAATCTCCTTGCACAAGCTTGAGAGCAACAC
AATCTATCAGGAAAGAAAGAAAGAAAAAACCGAACCTGACAAAAAGAAGAAAAAGAAG
AAGAAAAAAATCATGAAAACCATCCAGCCAAAAATGCACAATTCTATCTCTTGGGCAAT
CTTCACGGGGCTGGCTGCTCTGTGTCTCTTCCAAGGAGTGCCCGTGCGCAGCGGAGATGC
CACCTTCCCCAAAGCTATGGACAACGTGACGGTCCGGCAGGGGGAGAGCGCCACCCCTCAG
GTGCACTATTGACAACCGGGTCACCCGGGTGGCCTGGCTAAACCGCAGCACCATCCTCTA
TGCTGGGAATGACAAGTGGTGCCTGGATCCTCGCGTGGTCTTCTGAGCAACACCCAAAC
GCAGTACAGCATCGAGATCCAGAACGTGGATGTGTATGACGAGGGCCCTTACACCTGCTC
GGTGCAGACAGACAACACCCAAAGACCTCTAGGGTCCACCTCATTGTGCAAGTATCTCC
CAAAATTGTAGAGATTTCTTCAGATATCTCCATTAATGAAGGGAACAATATTAGCCTCAC
CTGCATAGCAACTGGTAGACCAGAGCCTACGGTTACTTGGAGACACATCTCTCCAAAGC
GGTTGGCTTTGTGAGTGAAGACGAATACTTGGAAATTTCAGGGCATCACCCGGGAGTGC
AGGGGACTACGAGTGCAGTGCCTCCAATGACGTGGCCGCGCCCGTGGTACGGAGAGTAAA
GGTCACCGTGAACCTATCCACCATACATTTCAGAAGCCAAGGGTACAGGTGTCCCCGTGGG
ACAAAAGGGGACACTGCAGTGTGAAGCCTCAGCAGTCCCCTCAGCAGAATTCCAGTGGTA
CAAGGATGACAAAAGACTGATTGAAGGAAAGAAAGGGGTGAAAGTGGAACAGACCTTT
CCTCTCAAACTCATCTTCTTCAATGTCTCTGAACATGACTATGGGAATACTTGCCT
GGCCTCCAACAAGCTGGGCCACACCAATGCCAGCATCATGCTATTTGGTCCAGGCGCCGT
CAGCGAGGTGAGCAACGGCACGTGAGGAGGGCAGGCTGCGTCTGGCTGCTGCCTCTTCT
GGTCTTGACCTGCTTCTCAAATTTTGATGTGAGTGCCACTTCCCCACCCGGGAAAGGCT
GCCGCCACCACCACCACCAACACAACAGCAATGGCAACACCGACAGCAACCAATCAGATA
TATACAAATGAAATTAGAAGAAACACAGCCTCATGGGACAGAAATTTGAGGGAGGGGAAC
AAAGAATACTTTGGGGGGAAAAGAGTTTTAAAAAAGAAATTGAAAATTGCCTTGAGATA
TTTAGGTACAATGGAGTTTTCTTTCCCAAACGGGAAGAACACAGCACACCCGGCTTGA
CCCACTGCAAGCTGCATCGTGCAACCTCTTTGGTGCAGTGTGGGCAAGGGCTCAGCCTC
TCTGCCCACAGAGTGCCCCACGTGGAACATTCTGGAGCTGGCCATCCCAAATTCATCA
GTCCATAGAGACGAACAGAATGAGACCTTCCGGCCCCAAGCGTGGCGCTGCGGGCACTTTG
GTAGACTGTGCCACCACGGCGTGTGTTGTGAAACGTGAAATAAAAAAGAGCAAAAAAAA

57/392

FIGURE 56

MKTIQPKMHNSISWAI FTGLAALCLFQGVPRSGDATFPKAMDNVTVRQGESATLRCTID
NRVTRVAWLNRSTILYAGNDKWCLDPRVVLLSNTQTQYSIEIQNVDVYDEGPYTCVQTD
NHPKTSRVHLIVQVSPKIVEISSDISINEGNNISLTCIATGRPEPTVTWRHISPKAVGFV
SEDEYLEIQGITREQSGDYECASNDVAAPVRRVKVTVNYPPYISEAKGTGVPVGQKGT
LQCEASAVPSAEFQWKDDKRLIEGKKGVKVENRPFLSKLIFFNVSEHDYGNVTCVASNK
LHTNASIMLFGPGAVSEVSNGTSRRAGCVWLLPLLVLHLLKLF

58/392

FIGURE 57

GCTGCGCCGGCTGCGGCTGCAGGGGAATCCGCTGTGGTGCGGCTGCCAGGCGCGGCCCT
ACTCGAGTGGCTGGCGCGGGCGCGCTGCGCTCGGACGGCGCGTGCCAGGGGCGCGGCG
CCTGCGGGGCGAGGCTCTGGACGCCCTGCGGCCCTGGGACCTGCGCTGCCCTGGGGACGC
GGCGCAGGAAGAGGAAGAGCTGGAAGAGCGGGCTGTGGCCGGGCCCCGCGCCCCCTCCGCG
CGGCCCTCCGCGCGGCCCCGGGGAGGAGCGGGCAGTCGCGCCTTGCCCTCGCGCCTGCGT
GTGCGTCCCCGAGTCCCGGCACAGCAGCTGCGAGGGCTGCGGCCTGCAGGCGGTGCCCCG
CGGCTTCCCCAGCGACACCCAGCTCCTGGACCTGAGGCGGAACCACTTCCCCCTCGGTGCC
CCGAGCGGCCTTCCCCGGNCTGGGCCACCTGGTGTGCTGCACCTGCAGCACTGCGGCAT
CGCGGAGCTGGAAGCGGGCGCCCTGGCCGGGCTGGGCCGCTGATCTACCTGTACCTCTC
CGACAACCAGCTCGCAGGCCTCAGCGCTGCTGCCCTTGAAGGGGCTCCCCGCCCTCGGCTA
CCTGTACCTAGAACGCAACCGTTTCTGAGGTGCCAGGGGCTGCCNTGCGCGCCCTGCC
CAGCCTCTTCTCCCTGCACCTGCAGGACAACGCTGTGGACCGCCTGGCACCTGGGGACCT
GGGGAGAACACGGGCCTTGCGCTGGGTCTACCTGAGTGGAACCGCATCACCGAAGTGTC
CCTTGGGGCGCTGGGCCAGCTCGGGAGCTGGAGAAGCTGCACCTGGACAGGAATCAGCT
GCGAGAGGTGCCCACTGGGGCCTTGGAGGGGCTGCCTGCCCTCCTGGAGCTGCAGCTCTC
GGGCAACCCACTCAGGGCCTTGCGTGACGGAGCCTTCCAGCCTGTGGGCAGGTGCTGCA
GCACCTCTTCTGAACAGCAGTGGCCTGGAGCAGATTTGTCTGGGGCCTTTTCAGGCCT
GGGGCCCGGGCTCCAGAGCCTGCACCTGCAGAAGAACAGCTTCGGGCCCTGCCTGCCCT
GCCCAGTCTCAGCCAGCTGGAGCTCATCGACCTCAGCAGCAATCCCTTCCCCTGTGACTG
CCAGCTGCTTCCGCTGCACAGGTGGCTTACTGGGCTGAACCTGCGGGTGGGGGCCACCTG
CGCCACCCCTCCCAATGCCCGTGGCCAGAGGTGAAGGCTGCAGCTGCTGTCTTTGAAGA
CTGCCCCGGGCTGGGCTGCCAGAAAGGCCAAGCGGACACCAGCCTCCAGGCCCACTGCCAG
GAGAACCCCCATCAAAGGAAGACAGTGTGGAGCAGATAAGAACATCCTCTTCCCCACATG
GTACCACACTGTGGAGCCCACCTCGCTGTCATAGGCCCTGCGGCTCTGAAGGATGGCTTTG
CCCGCTCCCGCTCTGCCCCCTCAAGTGGAACCCAAGCTGGGCTCAGAATCTGTAGAGTGAG
GCCCCACCAAGGGAAACGACACCCACGGCCTGAGAGCCAGGTGGAGTCCTGCCACTCAGC
TGCCTGCCTTTGCTCCACCCCTCTCCACCCCTCAAAGAGGTCTCGAGGGGACACTCTGAA
GGCACCTGGCTCAGAACCACTGCCATCCAAGGAGCGAGGAGTCCCAGGGCTGAGCAAATG
CAGCGGGGAGGTGGCAGTTCCCCTGCTTCCCGATCCTCATTTTCTGCTTCACTTGACTC
CTCCAGATAGGAGCTGCTCTCACTGCCCACACTGCTG

59/392

FIGURE 58

LRRLRLQGNPLWCGCQARPLLEWLLARARVRSDGACQGPRRLRGEALDALRPWDLRCPGDA
AQEEEELEERAVAGPRAPPRGPPRGPEERAVAPCPACVCPESRHSSCEGCGLQAVPR
GFPSDTQLLDLRRNHFPSPRAAFPGGLHVLVSLHLQHCGIAELEAGALAGLGRLIYLYLS
DNQLAGLSAAALEGAPRLGYLYLERNRFLQVPGAAXRALPSLFSHLQDNAVDR LAPGDL
GRTRALRWVYLSGNRITEVSLGALGPARELEKLHLDRNQLREVPTGALEGLPALLELQLS
GNPLRALRDGAFQPVGRSLQHLFLNSSGLEQICPGAFSGLGPGLSHLQKNQLRALPAL
PSLSQLELIDLSSNPFPCDCQLLPLHRWLTGLNLRVGATCATPPNARGQRVKAAA VFE
CPGWAARKAKRTPASRPSARRTPIKGRQCGADKNILFPTWYHTVEPTSLS

Signal sequence:

None

Transmembrane domain:

None

N-glycosylation site:

325-328

Glycosaminoglycan attachment site:

338-341

Protein kinase C phosphorylation site:

438-440

N-myristoylation site:

166-171, 186-191, 253-258, 286-291, 335-340, 339-344, 450-455

Leucine rich repeat N-terminal domain:

94-123

Leucine Rich Repeat:

125-148, 149-172, 173-196, 197-220, 221-244, 245-268, 269-292, 293-316, 318-341, 343-364, 365-386

Leucine rich repeat C-terminal domain:

374-422

60/392

FIGURE 59

CTCCACGGTGTCCAGCGCCAGAAATGCGGCTTCTGGTCCTGCTATGGGGTTGCCTGCTG
CTCCAGGTTATGAAGCCCTGGAGGGCCAGAGGAAATCAGCGGGTTCGAAGGGGACACT
GTGTCCCTGCAGTGCACCTACAGGGAAGAGCTGAGGGACCACCGGAAGTACTGGTGCAGG
AAGGGTGGGATCCTCTTCTCTCGCTGCTCTGGCACCATCTATGCAGAAGAAGAAGGCCAG
GAGACAATGAAGGGCAGGGTGTCCATCCGTGACAGCCGCCAGGAGCTCTCGCTCATTTGTG
ACCCTGTGGAACCTCACCTGCAAGACGCTGGGGAGTACTGGTGTGGGGTCGAAAAACGG
GGCCCCGATGAGTCTTTACTGATCTCTCTGTTCGTCTTTCCAGGACCCTGCTGTCTCTCCC
TCCCCTTCTCCACCTTCCAGCCTCTGGCTACAACACGCCTGCAGCCCAAGGCAAAAGCT
CAGCAAACCCAGCCCCCAGGATTGACTTCTCCTGGGCTCTACCCGGCAGCCACCACAGCC
AAGCAGGGGAAGACAGGGGCTGAGGCCCTCCATTGCCAGGGACTTCCCAGTACGGGCAC
GAAAGGACTTCTCAGTACACAGGAACCTCTCCTCACCCAGCGACCTCTCCTCCTGCAGGG
AGCTCCCGCCCCCCCCATGCAGCTGGACTCCACCTCAGCAGAGGACACCAGTCCAGCTCTC
AGCAGTGGCAGCTCTAAGCCCAGGGTGTCCATCCCGATGGTCCGCATACTGGCCCCAGTC
CTGGTGTCTGCTGAGCCTTCTGTGAGCCGAGGCTGATCGCCTTCTGCAGCCACCTGCTC
CTGTGGAGAAAGGAAGCTCAACAGGCCACGGAGACACAGAGGAACGAGAAGTTCTGGCTC
TCACGCTTGACTGCGGAGGAAAAGGAAGCCCCCTTCCAGGCCCTGAGGGGGACGTGATC
TCGATGCCTCCCCTCCACACATCTGAGGAGGAGCTGGGCTTCTCGAAGTTTGTCTCAGCG
TAGGGCAGGAGGCCCTCCTGGCCAGGCCAGCAGTGAAGCAGTATGGCTGGCTGGATCAGC
ACCGATTCCCAGAAAGCTTTCCACCTCAGCCTCAGAGTCCAGCTGCCCGGACTCCAGGGCT
CTCCCCACCCTCCCAGGCTCTCCTCTTGATGTTCCAGCCTGACCTAGAAGCGTTTGTCTC
AGCCCTGGAGCCCAGAGCGGTGGCCTTGCTCTTCCGGCTGGAGACTGGGACATCCCTGAT
AGGTTACATCCCTGGGCAGAGTACCAGGCTGCTGACCCTCAGCAGGGCCAGACAAGGCT
CAGTGGATCTGGTCTGAGTTTCAATCTGCCAGGAACCTCCTGGGCCCTCATGCCAGTGTCTG
GACCTTGCCTTCTCCTCCACTCCAGACCCACCTTGTCTTCCCTCCCTGGCGTCTCAGAC
TTAGTCCCACGGTCTCCTGCATCAGCTGGTGTATGAAGAGGAGCATGCTGGGGTGAGACTG
GGATTCTGGCTTCTCTTTGAACCACCTGCATCCAGCCCTTCAGGAAGCCTGTGAAAAACG
TGATTCTGGCCCCACCAAGACCCACCAAAACCATCTCTGGGCTTGGTGCAGGACTCTGA
ATTCTAACAATGCCAGTGAAGTGTGCACTTGAGTTTGAGGGCCAGTGGGCCTGATGAAC
GCTCACACCCCTTCACTTAGAGTGTGATTTGGGCTGTGACGTCTCCACCTGCCCCAAT
AGATCTGCTCTGTCTGCGACACCAGATCCACGTGGGGACTCCCCTGAGGCCTGCTAAGTC
CAGGCCTTGGTCAGGTGAGGTGCACATTGCAAGGATAAGCCCAGGACCGGCACAGAAGTGG
TTGCCCTTNCCTATTGCCCCCTCCCTGGNCCATGCCTTCTTGCCTTTGGAAAAAATGATGAA
GAAAACCTTGGCTCCTTCTGCTGGAAAGGGTTACTTGCCTATGGGTTCTGGTGGCTA
GAGAGAAAAGTAGAAAACAGAGTGCACGTAGGTGTCTAACACAGAGGAGAGTAGGAACA
GGGCGGATACCTGAAGGTGACTCCGAGTCCAGCCCCCTGGAGAAGGGGTGGGGGTGGTG
GTAAAGTAGCACAACCTACTATTTTTTTTCTTTTTTCCATTATTATTGTTTTTTAAGACAGA
ATCTCGTGCTGCTGCCAGGCTGGAGTGCAGTGGCACGATCTGCAAACTCCGCCTCCTGG
GTTCAAGTGATTCTTCTGCCTCAGCCTCCCGAGTAGCTGGGATTACAGGCACGCACCACC
ACACCTGGCTAATTTTTTGTACTTTTAGTAGAGATGGGGTTTACCATTGTTGGCCAGGCTG
GTCTTGAACCTCCTGACCTCAAATGAGCCTCCTGCTTCAGTCTCCCAAATTGCCGGGATTA
CAGGCATGAGCCACTGTGTCTGGCCCTATTTCTTTAAAAAGTGAAATTAAGAGTTGTTT
AGTATGCAAACTTGGAAAGATGGAGGAGAAAAAGAAAAGGAAGAAAAAATGTACCCCA
TAGTCTCACCAGAGACTATCATTATTTCTGTTTTGTTGTACTTCTTCCACTCTTTTCTTC
TTCACATAATTTGCCGGTGTCTTTTTTACAGAGCAATTATCTTGTATATACAACCTTTGTA
TCCTGCCTTTTTCCACCTTATCGTTCCATCACTTTATTCCAGCACTTCTCTGTGTTTTACA
GACCTTTTTATAAATAAAATGTTTCATCAGCTGCATAAAAAAAAAAAAAA

61/392

FIGURE 60

MRLLVLLWGCLLLPGYEALEGP EEISGFEGDTVSLQCTYREELRDHRKYWCRKGGILFSR
CSGTIYAE EEGQETMKGRVSI RDSRQELSLIVTLWNLT LQDAGEYWCGVEKRGPD ELLI
SLFVFP GPCCPPSPSPTFQPLATTRLQPKAKAQQTQPPGLTSPGLYPAATTAKQGKTGAE
APPLPGTSQYGHERTSQYTGTSPHPATSPPPAGSSRPPMQLDSTSAEDTSPALSSGSSKPR
VSIPMVRILAPVLVLLSLLSAAGLIAFCSHLLLWRKEAQQATETQRNEKFWLSRLTAEEK
EAPSQAPEGDVISM PPLHTSEELGFSKFVSA

Important features:**Signal peptide:**

amino acids 1-17

Transmembrane domain:

amino acids 248-269

N-glycosylation site:

amino acids 96-99

Fibrinogen beta and gamma chains C-terminal domain:

amino acids 104-113

Ig like V-type domain:

amino acids 13-128

62/392

FIGURE 61

CGGGCCAGCCTGGGGCGGCCGGCCAGGAACCACCCGTTAAGGTGTCTTCTCTTTAGGGAT
GGTGAGGTTGGAAAAAGACTCCTGTAAACCTCCTCCAGGATGAACCACCTGCCAGAAGAC
ATGGAGAACGCTCTCACCGGGAGCCAGAGCTCCCATGCTTCTCTGCGCAATATCCATTCC
ATCAACCCACACAACCTCATGGCCAGGATTGAGTCCTATGAAGGAAGGGAAAAGAAAGGC
ATATCTGATGTCAGGAGGACTTTCTGTTTGTTTGTACCTTTGACCTCTTATTCGTAACA
TTACTGTGGATAATAGAGTTAAATGTGAATGGAGGCATTGAGAACACATTAGAGAAGGAG
GTGATGCAGTATGACTACTATTCTTCATATTTTGATATATTTCTTCTGGCAGTTTTCGA
TTTAAAGTGTTAATACTTGACATATGCTGTGTGCAGACTGCGCCATTGGTGGGCAATAGCG
TTGACAACGGCAGTGACCAGTGCCTTTTTACTAGCAAAAGTGATCCTTTTGAAGCTTTTC
TCTCAAGGGGCTTTTGGCTATGTGCTGCCCATCATTTTCATTTCATCCTTGCCTGGATTGAG
ACGTGGTTCCTGGATTTCAAAGTGTTACCTCAAGAAGCAGAAGAAGAAAACAGACTCCTG
ATAGTTCAGGATGCTTCAGAGAGGGCAGCACTTATACCTGGTGGTCTTTCTGATGGTCAG
TTTTATTCCCTCCTGAATCCGAAGCAGGATCTGAAGAAGCTGAAGAAAAACAGGACAGT
GAGAAACCACTTTTAGAACTATGAGTACTACTTTTGTTAAATGTGAAAAACCCTCACAGA
AAGTCATCGAGGCAAAAAGAGGCAGGCAGTGGAGTCTCCCTGTGACAGTAAAGTTGAAA
TGGTGACGTCCACTGCTGGCTTTATTGAACAGCTAATAAAGATTTATTTATTGTAATACC
TCACAAACGTTGTACCATATCCATGCACATTTAGTTGCCTGCCTGTGGCTGGTAAGGTAA
TGTCATGATTCATCCTCTCTTCAGTGAGACTGAGCCTGATGTGTTAACAATAGGTGAAG
AAAGTCTTGTGCTGTATTCCTAATCAAAAGACTTAATATATTGAAGTAACACTTTTTTAG
TAAGCAAGATACCTTTTTTATTTCAATTCACAGAATGGAATTTTTTTTGTTCATGTCTCAG
ATTTATTTTGTATTTCTTTTTTAACTCTACATTTCCCTTGTTTTTTAACTCATGCACA
TGTGCTCTTTGTACAGTTTTTAAAAAGTGTAATAAAATCTGACATGTCAATGTGGCTAGTT
TTATTTTTCTTGTTTTGCATTATGTGTATGGCCTGAAGTGTTGGACTTGCAAAAGGGGAA
GAAAGGAATTGCGAATACATGTAAAATGTCACCAGACATTTGTATTATTTTTATCATGAA
ATCATGTTTTTCTCTGATTGTTCTGAAATGTTCTAAATACTCTTATTTTGAATGCACAAA
ATGACTTAAACCATTCATATCATGTTTCCTTTGCGTTCAGCCAATTTCAATTAAATGAA
CTAAATTAAAAA

63/392

FIGURE 62

MNHLPEDMENALTGSQSSHASLRNIHSINPTQLMARIESYEGREKKGISDVRRTFCLFVT
FDLLFVTLWLWIELNVNGGIENTLEKEVMQYDYSSYFDIFLLAVFRFKVLILAYAVCRL
RHWWAIALTTAVTSAFLLAKVILSKLFSQGAFGYVLPPIISFILAWIETWFLDFKVLPQEA
EEENRLLIVQDASERAALIPGGLSDGQFYSPPESEAGSEEAEKQDSEKPLLEL

Important features of the protein:**Signal peptide:**

amino acids 1-20

Transmembrane domains:

amino acids 54-72, 100-118, 130-144, 146-166

N-myristoylation sites:

amino acids 14-20, 78-84, 79-85, 202-208, 217-223

64/392

FIGURE 63

GCGCCGGGAGCCCATCTGCCCCCAGGGGACGGGGCGCGGGGCCGGCTCCCGCCCCGGCAC
ATGGCTGCAGCCACCTCGCGCGCACCCCCGAGGCGCCGCGCCAGCTCGCCCCGAGGTCCGT
CGGAGGCGCCCCGGCCGCCCCGAGCCAAGCAGCAACTGAGCGGGGAAGCGCCCGCTCCG
GGGATCGGGATGTCCTCTCTCTCTCTGCTAGTTTCTACTATGTTGGAACCTTG
GGGACTCACACTGAGATCAAGAGAGTGGCAGAGGAAAAGGTCACCTTGCCCTGCCACCAT
CAACTGGGGCTTCCAGAAAAAGACACTCTGGATATTGAATGGCTGCTCACCGATAATGAA
GGGAACCAAAAAGTGGTGATCACTTACTCCAGTCGTCTACAATAACTTGACTGAG
GAACAGAAGGGCCGAGTGCCCTTTGCTTCCAATTTCTGGCAGGAGATGCCTCCTTGACAG
ATTGAACCTCTGAAGCCCACTGATGAGGGCCGGTACACCTGTAAGGTTAAGAATTCAGGG
CGCTACGTGTGGAGCCATGTCATCTTAAAGTCTTAGTGAGACCATCCAAGCCCAAGTGT
GAGTTGGAAGGAGAGCTGACAGAAGGAAGTGACCTGACTTTGCAGTGTGAGTCATCCTCT
GGCACAGAGCCCATTTGTGTATTACTGGCAGCGAATCCGAGAGAAAAGAGGGAGAGGATGAA
CGTCTGCCTCCCAAATCTAGGATTGACTACAACCACCCTGGACGAGTTCTGCTGCAGAAT
CTTACCATGTCTACTCTGGACTGTACAGTGACAGCAGGCAACGAAGCTGGGAAGGAA
AGCTGTGTGGTGCGAGTAACTGTACAGTATGTACAAAGCATCGGCATGGTTGCAGGAGCA
GTGACAGGCATAGTGGCTGGAGCCCTGCTGATTTTCTCTTGGTGTGGCTGCTAATCCGA
AGGAAAGACAAAGAAAGATATGAGGAAGAAGAGAGACCTAATGAAATTCGAGAAGATGCT
GAAGCTCCAAAAGCCCGTCTTGTGAAACCCAGCTCCTCTTCTCAGGCTCTCGGAGCTCA
CGCTCTGGTTCTTCTCCACTCGCTCCACAGCAAATAGTGCCTCACGCAGCCAGCGGACA
CTGTCAACTGACGCAGCACCCAGCCAGGGCTGGCCACCCAGGCATACAGCCTAGTGGGG
CCAGAGGTGAGAGGTCTTGAACCAAAGAAAGTCCACCATGCTAATCTGACCAAAGCAGAA
ACCACACCCAGCATGATCCCCAGCCAGAGCAGAGCCTTCCAAACGGTCTGAATTACAATG
GACTTGACTCCCACGCTTTCTAGGAGTCAGGGTCTTTGGACTCTTCTCGTCATTGGAGC
TCAAGTCACCAGCCACACAACCAGATGAGAGGTCACTAAGTAGCAGTGAGCATTGCACG
GAACAGATTGAGATGAGCATTTTCTTATACAATAACCAAACAAGCAAAAGGATGTAAGCT
GATTCATCTGTAAAAAGGCATCTTATTGTGCCTTTAGACCAGAGTAAGGGAAAGCAGGAG
TCCAAATCTATTTGTTGACCAGGACCTGTGGTGAGAAGGTTGGGGAAAGGTGAGGTGAAT
ATACCTAAAACCTTTAATGTGGGATATTTGTATCAGTGCTTTGATTACAATTTTCAAG
AGGAAATGGGATGCTGTTTGTAATTTTCTATGCATTTCTGCAAACTTATTGGATTATTA
GTTATTCAGACAGTCAAGCAGAACCCACAGCCTTATTACACCTGTCTACACCATGTACTG
AGCTAACCACTTCTAAGAACTCCAAAAAGGAAACATGTGTCTTCTATTTCTGACTTAAC
TTCATTTGTCATAAGGTTTGGATATTAATTTCAAGGGGAGTTGAAATAGTGGGAGATGGA
GAAGAGTGAATGAGTTTCTCCCACTCTATACTAATCTCACTATTTGTATTGAGCCCCAAAA
TAACTATGAAAGGAGACAAAAATTTGTGACAAAGGATTGTGAAGAGCTTCCATCTTCAT
GATGTTATGAGGATTGTTGACAAACATTAGAAATATATAATGGAGCAATTGTGGATTTC
CCTCAAATCAGATGCCTCTAAGGACTTTCCTGCTAGATATTTCTGGAAGGAGAAAATACA
ACATGTCAATTTATCAACGTCCTTAGAAAGAATTCTTCTAGAGAAAAAGGGATCTAGGAAT
GCTGAAAGATTACCCAACATAACATTATAGTCTCTTCTTCTGAGAAAAATGTGAAACCAG
AATTGCAAGACTGGGTGGACTAGAAAGGGAGATTAGATCAGTTTTCTCTTAATATGTCAA
GGAAGGTAGCCGGGCATGGTGCCAGGCACCTGTAGGAAAAATCCAGCAGGTGGAGGTTGCA
GTGAGCCGAGATTATGCCATTGCACTCCAGCCTGGGTGACAGAGCGGGACTCCGTCTC

65/392

FIGURE 64

MSLLLLLLLLVSYYVGTLGTHTEIKRVAEEKVTLPCHHQLGLPEKDTLDIEWLLTDNEGNQ
KVVITYSSRHVYNNLTEEQGRVAFASNFLAGDASLQIEPLKPSDEGRYTCKVKNSGRYV
WSHVILKVLVRPSKPKCELEGELETEGSDTLQCESSSGTEPIVYYWQRIREKEGEDERLP
PKSRIDYNHPGRVLLQNLTMSYSGLYQCTAGNEAGKESCVVRVTVQYVQSIGMVAGAVTG
IVAGALLIFLLVWLLIRRDKERYEEEEERPNEIREDAEAPKARLVKPSSSSSSGSRSSRSG
SSSTRSTANSASRSQRTLSTDAAPQPGLATQAYSLVGPEVRGSEPKKVHHANLTKAETTP
SMIPSQSRAFQTV

Signal sequence:
amino acids 1-16

Transmembrane domain:
amino acids 232-251

66/392

FIGURE 65

GTCGGGGCTGCGCGACGGCGCAGGGGCTGCGGGGAGCGCCGCGCAGGCCGTGCAGTTCCT
AGCGAGGAGGCGCCGCGCCGCGCCATTGCCGCTCTCTCGGTGAGCGCAGCCCCGCTCTCCGGGC
CGGGCCTTCGCGGGCCACGGCGCCATGGGGCCAGTGCGGCATCACCTCCTCCAAGACCGT
GCTGGTCTTTCTCAACCTCATCTTCTGGGGGGCAGCTGGCATTATGCTATGTGGGAGC
CTATGTCTTCATCACTTATGATGACTATGACCACTTCTTTGAAGATGTGTACACGCTCAT
CCCTGCTGTAGTGATCATAGCTGTAGGAGCCCTGCTTTTCATCATTGGGCTAATTGGCTG
CTGTGCCACAATCCGGGAAAGTCGCTGTGGACTTGCCACGTTTGTATCATCCTGCTCTT
GGTTTTTGTGCAGAAAGTTGTTGTAGTGGTTTTTGGGATATGTTTACAGAGCAAAGGTGGA
AAATGAGGTTGATCGCAGCATTGAGAAAGTGTATAAGACCTACAATGGAACCAACCCCTGA
TGCTGCTAGCCGGGCTATTGATTATGTACAGAGACAGCTGCATTGTTGTGGAATTACAA
CTACTCAGACTGGGAAAATACAGATTGGTTCAAAGAAACCAAAACCAGAGTGTCCCTCT
TAGCTGCTGCAGAGAGACTGCCAGCAATTGTAATGGCAGCCTGGCCCCACCTTCCGACCT
CTATGCTGAGGGGTGTGAGGCTCTAGTAGTGAAGAAGCTACAAGAAATCATGATGCATGT
GATCTGGGCCGCACTGGCATTTCAGCTATTTCAGCTGCTGGGCATGCTGTGTGCTTGCAT
CGTGTGTGTGCAGAAGGAGTAGAGATCCTGCTTACGAGCTCCTCATCACTGGCGGAACCTA
TGCATAGTTGACAACTCAAGCCTGAGCTTTTTGGTCTTGTCTGATTGGAAGGTGAATT
GAGCAGGTCTGCTGCTGTTGGCCTCTGGAGTTCATTTAGTTAAAGCACATGTACACTGGT
GTTGGACAGAGCAGCTTGGCTTTTCATGTGCCACCTACTTACCTACTACCTGCGACTTT
CTTTTTCCTTGTCTAGCTGACTCTTCATGCCCCAAGATTTTAAGTACGATGGTGAACG
TTCTAATTTTCAAGCAATTTGCGAGTCATGTAGTGTGGTAGAATTAAAGGAGGACACGAG
CCTGCTTCTGTTACCTCCAAGTGGTAACAGGACTGATGCCGAAATGTCACCAGGTCCTTT
CAGTCTTCACAGTGGAGAACTCTTGGCCAAAGTTTTTTCGGGGAGGAGGAGGAAACCAG
CTTTCTGGTTAAGGTTAACACCAGATGGTGCCCTCATTTGGTGTCTTTTAAAAAATATT
TACTGTAGTCCAATAAGATAGCAGCTGTACAAAATGACTAAAATAGATTGTAGGATCATA
TGGCGTATATCTTGGTTCATCTTCAAATCAGAGACTGAGCTTTGAAACTAGTGGTTTTT
AATCAAAGTTGGCTTTATAGGAGGAGTATAATGTATGCACTACTGTTTTTAAAGAATTAG
TGTGAGTGTGTTTTTGTATGAATGAGCCCATTCATGGTAAGTCTTAAGCTTGTGGAAT
AATGTACCCATGTAGACTAGCAAAATAGTATGTAGATGTGATCTCAGTTGTAAATAGAAA
AATCTAATTCATAAACTCTGTATCAGCCCCCAAAAAAAAAAAAAAAAAA

67/392

FIGURE 66

MGQCGITSSKTVLVFLNLI FWGAAGILCYVGAYVFITYDDYDHHFFEDVYTLIPAVVIIAV
GALLFIIGLIGCCATIRESRCGLATFVI ILLLVFVTEV VVVVLGYVYRAKVENEVDRSIQ
KVYKTYNGTNPDAASRAIDYVQRQLHCCGIHNYSDWENTDWFKETKNQSVPLSCCRETAS
NCNGSLAHPSDLYAEGCEALVVKLQEIMMHVIWAALAFAAIQLLGMLCACIVLCRRSRD
PAYELLITGGTYA

Signal peptide:
none

Type II transmembrane domain:
11-38

Other transmembrane domains:
48-68, 87-107, 208-235

N-glycosylation site:
127-131, 152-156, 167-171, 183-187

Tyrosine kinase phosphorylation site:
236-244

N-myristoylation site:
5-11, 68-74, 71-77, 226-232

Prokaryotic membrane lipoprotein lipid attachment site:
62-73, 221-232

Transmembrane 4 family proteins:
7-35, 56-106

68/392

FIGURE 67

GCGGCACCTGGAAGATGCGCCCATTTGGCTGGTGGCCTGCTCAAGGTGGTGTTCGTGGTCT
TCGCCTCCTTGTGTGCCTGGTATTCGGGGTACCTGCTCGCAGAGCTCATTCCAGATGCAC
CCCTGTCCAGTGCTGCCATATAGCATCCGCAGCATCGGGGAGAGGCCTGTCCTCAAAGCTC
CAGTCCCCAAAAGGCAAAAATGTGACCACTGGACTCCCTGCCCATCTGACACCTATGCCT
ACAGGTTACTCAGCGGAGGTGGCAGAAGCAAGTACGCCAAAATCTGCTTTGAGGATAACC
TACTTATGGGAGAACAGCTGGGAAATGTTGCCAGAGGAATAAACATTGCCATTGTCAACT
ATGTAACCTGGGAATGTGACAGCAACACGATGTTTTGATATGTATGAAGGCGATAACTCTG
GACCGATGACAAAGTTTATTCAGAGTGCTGCTCCAAAATCCCTGCTCTTCATGGTGACCT
ATGACGACGGAAGCACAGACTGAATAACGATGCCAAGAATGCCATAGAAGCACTTGGAA
GTAAAGAAATCAGGAACATGAAATTCAGGTCTAGCTGGGTATTTATTGCAGCAAAAGGCT
TGGAACCTCCCTTCCGAAATTCAGAGAGAAAAGATCAACCACTCTGATGCTAAGAACAACA
GATATTCCTGGCTGGCCTGCAGAGATCCAGATAGAAGGCTGCATACCCAAAGAACGAAGCT
GACACTGCAGGGTCCTGAGTAAATGTGTTCTGTATAAAACAAATGCAGCTGGAATCGCTCA
AGAATCTTATTTTTCTAAATCCAACAGCCCATATTTGATGAGTATTTTGGGTTTGTGTA
AACCAATGAACATTTGCTAGTTGTATCAAATCTTGGTACGCAGTATTTTTATACCAGTAT
TTTATGTAGTGAAGATGTCAATTAGCAGGAACTAAAATGAATGGAAATTTTAAAAAAA
AAA

69/392

FIGURE 68

MRPLAGGLLKVVVFVVFASLCAWYSGYLLAELIPDAPLSSAAYSIRSIGERFVLKAPVPKR
QKCDHWTPCPSDTYAYRLLSGGGRSKYAKICFEDNLLMGEQLGNVARGINIAIVNYVTGN
VTATRCFDMYEGDNSGPMTKFIQSAAPKSLLFMVTYDDGSTRLNNDAKNAIEALGSKEIR
NMKFRSSWVFIAAKGLELPSEIQREKINHSDAKNNRYSGWPAEIQIEGCIPKERS

Signal sequence:

amino acids 1-20

N-glycosylation sites:

amino acids 120-124, 208-212

Glycosaminoglycan attachment site:

amino acids 80-84

N-myristoylation sites:

amino acids 81-87, 108-114, 119-125

70/392

FIGURE 69

ACACAACTTTACACCTGAATGAACGCCAAACCTCTATGGATATATAAAGGGAAGCTTGAG
GAGGAATTTACAGTTACAGTGCAGAAGCAGAAGCAAAGAATTAACCAGCTCTTCAGTC
AAGCAAATCCTCTACTCACCATGCTTCCTCCTGCCATTTCATTTCTATCTCCTTCCCCCTTG
CATGCATCCTAATGAAAAGCTGTTTGGCTTTTAAAAATGATGCCACAGAAATCCTTTATT
CACATGTGGTTAAACCTGTTCCAGCACACCCAGCAGCAACAGCACGTTGAATCAAGCCA
GAAATGGAGGCAGGCATTTCACTAACACTGGACTGGATCGGAACACTCGGGTTCAAGTGG
GTTGCCGGGAAGTGGCTTCCACCAAATACATCTCTGATGGCCAGTGCACCAGCATCAGCC
CTCTGAAGGAGCTGGTGTGTGCTGGCGAGTGCTTGCCCCCTGCCAGTGCTCCCTAACTGGA
TTGGAGGAGGCTATGGAACAAAGTACTGGAGCAGGAGGAGCTCCCAGGAGTGGCGGTGTG
TCAATGACAAAACCCGTACCCAGAGAATCCAGCTGCAGTGCCAAGATGGCAGCACACGCA
CCTACAAAATCACAGTAGTCACCTGCCTGCAAGTGCAAGAGGTACACCCGGCAGCACAAACG
AGTCCAGTCACAACCTTTGAGAGCATGTACCTGCCAAGCCAGTCCAGCATCACAGAGAGC
GGAAAAGAGCCAGCAAATCCAGCAAGCACAGCATGAGTTAGAACTCAGACTCCATAACT
AGACTTACTAGTAACCATCTGCTTTACAGATTTGATTGCTTGGAAGACTCAAGCCTGCCA
CTGCTGTTTTCTCACTTGAAAGTATATGCTTTCTGCTTTGATCAAACCCAGCAAGCTGTC
TTAAGTATCAGGACCTTCTTTGGGAATAGTTTTTCCTTTTAAAGTTTTTCAAGATGTAGG
TATATCCATGAATGCAATTTGCATTTAAATTCCACGTATCCCTGTAGTTTAAATTCCTCA
TTGGTCTTAAAAGACTGTTGATACTATAAACATCAGTGAATCAATTATATTTTAAACA
GAAAAGGGCTT

71/392

FIGURE 70

MLPPAIHFYLLPLACILMKSCLAFFKNDATILYSHVVKPVPAPHPSSNSTLNQARNNGGRHF
SNTGLDRNTRVQVGCRELRSTKYISDGQCTSIPLKELVCAGECLPLPVLPNWIGGGYGT
KYWSRRSSQEWRCVNDKTRTQRIQLQCQDGSTRITYKITVVTACKCKRYTRQHNESSHNFE
SMSPAKFPVQHHRRERKRASKSSKHSMS

Signal sequence:

1-23

Transmembrane domain:

None

N-glycosylation site:

47-50, 173-176

**cAMP- and cGMP-dependent protein kinase phosphorylation
site:**

125-128, 166-169, 195-198

N-myristoylation site:

64-69, 87-92, 115-120, 116-121, 150-155

72/392

FIGURE 71

CCCAGGCTCTAGTGCAGGAGGAGAAGGAGGAGGAGCAGGAGGTGGAGATTCCCAGTTAAA
AGGCTCCAGAATCGTGTACCAGGCAGAGAACTGAAGTACTGGGGCCTCCTCCACTGGGTC
CGAATCAGTAGGTGACCCCGCCCCTGGATTCTGGAAGACCTCACCATGGGACGCCCCGA
CCTCGTGCGGCCAAGACGTGGATGTTCCCTGCTCTTGCTGGGGGGAGCCTGGGCAGGACAC
TCCAGGGCACAGGAGGACAAGGTGCTGGGGGGTCA TGAGTGCCAACCCCATTCGCAGCCT
TGGCAGGCGGCCTTGTTCCAGGGCCAGCAACTACTCTGTGGCGGTGTCCTTG TAGGTGGC
AACTGGGTCCCTTACAGCTGCCCACTGTAAAAAACCGAAATACACAGTACGCCCTGGGAGAC
CACAGCCTACAGAAATAAAGATGGCCCAGAGCAAGAAATACCTGTGGTTCAGTCCATCCCA
CACCCCTGCTACAACAGCAGCGATGTGGAGGACCACAACCATGATCTGATGCTTCTTCAA
CTGCGTGACCAGGCATCCCTGGGGTCCAAAGTGAAGCCCATCAGCCTGGCAGATCATTGC
ACCCAGCCTGGCCAGAAGTGCACCGTCTCAGGCTGGGGCACTGTCACCAGTCCCCGAGAG
AATTTTCCTGACACTCTCAACTGTGCAGAAGTAAAAATCTTTCCCCAGAAGAAGTGTGAG
GATGCTTACCCGGGGCAGATCACAGATGGCATGGTCTGTGCAGGCAGCAGCAAAGGGGCT
GACACGTGCCAGGGCGATTCTGGAGGCCCCCTGGTGTGTGATGGTGCACTCCAGGGCATC
ACATCCTGGGGCTCAGACCCCTGTGGGAGGTCCGACAAACCTGGCGTCTATACCAACATC
TGCCGCTACCTGGACTGGATCAAGAAGATCATAGGCAGCAAGGGCTGGATTCTAGGATAAG
CACTAGATCTCCCTTAATAAACTCACAACCTCTCTGGTTC

73/392

FIGURE 72

MGRPRPRAAKTWMFLLLLGGAWAGHSRAQEDKVLGGHECQPHSQPWQAALFOGQQLLCGG
VLVGGNWVLTAAHCKKPKYTVRLGDHSLQNKDGPEQEIPVVQSIPHPCYNSSDVEDHNHD
LMLLQLRDQASLGSKVKPISLADHCTQPGQKCTVSGWGTVTSPRENFDTLNCAEVKIFP
QKKCEDAYPGQITDGMVCAGSSKGADTCQGDSSGGLVCDGALQGITSWGSDDPCGRSDKPG
VYTNICRYLDWIKKIIGSKG

Important Features:**Signal peptide:**

amino acids 1-23

Transmembrane domain:

amino acids 51-71

N-glycosylation site:

amino acids 110-113

Serine proteases, trypsin family, histidine active site:

amino acids 69-74 and 207-217

Tyrosine kinase phosphorylation site:

amino acids 182-188

Kringle domain proteins motif:

amino acids 205-217

74/392

FIGURE 73

CTCGGGCGCGCACAGGCAGCTCGGTTTGGCCCTGCGATTGAGCTGCGGGTTCGCGGCCGGCG
CCGGCCTCTCCAATGGCAAATGTGTGTGGCTGGAGGCGAGCGCGAGGCTTTCGGCAAAGG
CAGTCGAGTGTTTGCAGACCGGGGCGAGTCCTGTGAAAGCAGATAAAAGAAAACATTTAT
TAACGTGTCAATTACGAGGGGAGCGCCCGGCCGGGGCTGTGCGACTCCCCGCGGAACATTT
GGCTCCCTCCAGCTCCGAGAGAGGAGAAGAAGAAAGCGGAAAAGAGGCAGATTCACGTG
TTTCCAGCCAAGTGGACCTGATCGATGGCCCTCCTGAATTTATCACGATATTTGATTTAT
TAGCGATGCCCCCTGGTTTGTGTGTTACGCACACACACGTGCACACAAGGCTCTGGCTCG
CTTCCCTCCCTCGTTTCCAGCTCCTGGGCGAATCCACATCTGTTTCAACTCTCCGCCGA
GGGCGAGCAGGAGCGAGAGTGTGTCGAATCTGCGAGTGAAGAGGGACGAGGGAAAAGAAA
CAAAGCCACAGACGCAACTTGAGACTCCCGCATCCCAAAGAAGCACCAGATCAGCAAAA
AAAGAAGATGGGCCCCCGAGCCTCGTGCTGTGCTTGCTGTCCGCAACTGTGTTCTCCCT
GCTGGGTGGAAGCTCGGCCTTCTGTGCGACCACCGCTGAAAGGCAGGTTTCAGAGGGA
CCGCAGGAACATCCGCCCAACATCATCCTGGTGCTGACGGACGACCAGGATGTGGAGCT
GGGTTCATGCAGGTGATGAACAAGACCCGGCGCATCATGGAGCAGGGCGGGGCGCACTT
CATCAACGCCTTCGTGACCACACCCATGTGCTGCCCCTCACGCTCCTCCATCCTCACTGG
CAAGTACGTCCACAACCACAACACCTACACCAACAATGAGAACTGCTCCTCGCCCTCCTG
GCAGGCACAGCACGAGAGCCGCACCTTTGCCGTGTACCTCAATAGCACTGGCTACCGGAC
AGCTTCTTTCGGGAAGTATCTTAATGAATACAACGGCTCCTACGTGCCACCCGGCTGGAA
GGAGTGGGTTCGACTCCTTAAAACTCCCGCTTTTATAACTACACGCTGTGTGCGAACGG
GGTGAAGAGAAGCACGGCTCCGACTACTCCAAGGATTACCTCACAGACCTCATCACC
TGACAGCGTGAGCTTCTTCCGCACGTCCAAGAAGATGTACCCGCACAGGCCAGTCCTCAT
GGTCATCAGCCATGCAGCCCCCACGGCCCTGAGGATTCAGCCCCACAATATTCACGCCT
CTTCCCAAACGCATCTCAGCACATCACGCCGAGCTACAACCTACGCGCCCAACCCGGACAA
ACACTGGATCATGCGCTACACGGGGCCCATGAAGCCCATCCACATGGAATTCACCAACAT
GCTCCAGCGGAAGCGCTTGACAGCCCTCATGTGCGGTGGACGACTCCATGGAGACGATTTA
CAACATGCTGGTTGAGACGGGCGAGCTGGACAACACGTACATCGTATACACCGCCGACCA
CGGTTACCACATCGGCCAGTTTGGCCTGGTGAAAGGGAAAATCCATGCCATATGAGTTTGA
CATCAGGGTCCCGTTCTACGTGAGGGGGCCCCAACGTGGAAGCCGGCTGTCTGAATCCCCA
CATCGTCTCAACATTGACCTGGCCCCCACCATCCTGGACATTGCAGGCCTGGACATACC
TGCGGATATGGACGGGAAATCCATCCTCAAGCTGCTGGACACGGAGCGGCCGGTGAATCG
GTTTCACTTGAAAAAGAAGATGAGGGTCTGGCGGGACTCCTTCTTGGTGGAGAGAGGCAA
GCTGCTACACAAGAGAGACAATGACAAGGTGGACGCCAGGAGGAGAACTTTCTGCCCAA
GTACCAGCGTGTGAAGGACCTGTGTGAGCGTGTGAGTACCAGACGGCGTGTGAGCAGCT
GGGACAGAAGTGGCAGTGTGTGGAGGACGCCACGGGGAAGCTGAAGCTGCATAAGTGCAA
GGGCCCCATGCGGCTGGGCGGCAGCAGAGCCCTCTCCAACCTCGTGCCCAAGTACTACGG
GCAGGGCAGCGAGGCTGCACCTGTGACAGCGGGGACTACAAGCTCAGCCTGGCCGACG
CCGGAAAAAATCTTTCAAGAAGAAGTACAAGGCCAGCTATGTCCGCACTCGCTCCATCCG
CTCAGTGGCCATCGAGGTGGACGGCAGGGTGTACCACGTAGGCCTGGGTGATGCCGCCA
GCCCCGAAACCTCACCAAGCGGCACTGGCCAGGGGCCCTGAGGACCAAGATGACAAGGA
TGGTGGGGACTTCAGTGGCACTGGAGGCCTTCCCGACTACTCAGCCGCCAACCCCATTA
AGTGACACATCGGTGCTACATCCTAGAGAACGACACAGTCCAGTGTGACCTGGACCTGTA
CAAGTCCCTGCAGGCCTGGAAGACCACAAGCTGCACATCGACCACGAGATTGAAACCT
GCAGAACAAAATTAAGAACCTGAGGGAAGTCCGAGGTACCTGAAGAAAAAGCGGCCAGA
AGAATGTGACTGTCAAAAATCAGCTACCACACCCAGCACAAAGGCCGCCTCAAGCACAG
AGGCTCCAGTCTGCATCCTTTCAGGAAGGGCCTGCAAGAGAAGGACAAGGTGTGGCTGTT
GCGGGAGCAGAAGCGCAAGAAGAACTCCGCAAGCTGCTCAAGCGCCTGCAGAACACGA
CACGTGCAGCATGCCAGGCCTCACGTGCTTACCCACGACAACCAGCACTGGCAGACGGC

75/392

GCCTTTCTGGACACTGGGGCCTTTCTGTGCCTGCACCAGCGCCAACAATAACACGTACTG
GTGCATGAGGACCATCAATGAGACTCACAATTCCTCTTCTGTGAATTTGCAACTGGCTT
CCTAGAGTACTTTGATCTCAACACAGACCCCTACCAGCTGATGAATGCAGTGAACACACT
GGACAGGGATGTCCTCAACCAGCTACACGTACAGCTCATGGAGCTGAGGAGCTGCAAGGG
TTACAAGCAGTGTAACCCCCGGA~~CT~~CGAAACATGGACCTGGATGGAGGAAGCTATGAGCA
ATACAGGCAGTTTCAGCGTCGAAAGTGGCCAGAAATGAAGAGACCTTCTTCCAAATCACT
GGGACAACTGTGGGAAGGCTGGGAAGGTTAAGAAACAACAGAGGTGGACCTCCAAAAACA
TAGAGGCATCACCTGACTGCACAGGCAATGAAAAACCATGTGGGTGATTTCCAGCAGACC
TGTGCTATTGGCCAGGAGGCTGAGAAAGCAAGCACGCACTCTCAGTCAACATGACAGAT
TCTGGAGGATAACCAGCAGGAGCAGAGATAACTTCAGGAAGTCCATTTTGGCCCTGCTT
TTGCTTTGGATTATACCTCACCAGCTGCACAAAATGCATTTTTTCGTATCAAAAAGTCAC
CACTAACCCCTCCCCCAGAAGCTCACAAAGGAAAACGGAGAGAGCGAGCGAGAGAGATTTT
CTTGGAATTTCTCCCAAGGGCGAAAGTCATTGGAATTTTAAATCATAGGGGAAAAGCA
GTCCTGTTCTAAATCCTCTTATTCTTTTGGTTTGTCAAAAGAAGGAAC~~TA~~AGAAGCAGG
ACAGAGGCAACGTGGAGAGGCTGAAAACAGTGCAGAGACGTTTGACAATGAGTCAGTAGC
ACAAAAGAGATGACATTTACCTAGCACTATAAACCCCTGGTTGCCTCTGAAGAACTGCCT
TCATTGTATATATGTGACTATTTACATGTAATCAACATGGGAAC~~TTT~~TAGGGGAACCTAA
TAAGAAATCCCAATTTTCAGGAGTGGTGGTGTCAATAAACGCTCTGTGGCCAGTGTAAAA
GAAAAA

76/392

FIGURE 74

MGPPSLVLCLLSATVFSLGGSSAFLSHRLKGRFQDRRNIRPNIILVLTDDQDVELGS
MQVMNKTRRIMEQGGAHFINAFVTPMCCPSRSSILTGKYVHNHNTYTNNENCSSPSWQA
QHESRTFAVYLNSTGYRTAFFGKYLNEYNGSYVPPGWKEWVGLLKNSRFYNYTLCRNGVK
EKHGSDYSKDYLTDLITNDSVSFFRTSKKMPHRPVLVISHAAPHGPEDSAPQYSRLFP
NASQHITPSYNYAPNPDKHWIMRYTGPMKPIHMEFTNMLQRKRLQTLMSVDDSMETIYNM
LVETGELDNTYIVYTADHGYHIGQFGLVKGKSMPEFDIRVPFYVRGPNVEAGCLNPHIV
LNIDLAPTILDIAGLDIPADMKGKSLKLLDTERPVMRFLKKKMRVWRDSFLVERGKLL
HKRDNDKVDAQEENFLPKYQRVKDLQRAEYQTACEQLGQKWQCVEDATGKLKLHKCKGP
MRLGGSRALSNLVPKYQGQSEACTCDSDYKLSLAGRRKKLFKKKYKASYVRSRSIRSV
AIEVDGRVYHVLGDAQAQRNLTKRHWPGAPEDQDDKDGGDFSGTGGLPDYSAANPIKVT
HRCYILENDTVQCDLDLYKSLQAWKDHLHIDHEIETLQNKIKNLREVRGHLKKKRPEEC
DCHKISYHTQHKGRCLKHRGSSLHPFRKGLQEKDKVWLLREQKRKKKLRKLLKRLQNNDTC
SMPGLTCFTHDNQHWQTAPFWTLGPFCACTSANNNTYWCMTINETHNLFCEFATGFLE
YFDLNTDPYQLMNAVNTLDRDVLNQLHVQLMELRSCKGYKQCNPRTRNMDLDGGSYEQYR
QFQRRKWPEMKRPSSKSLGQLWEGWEG

Important features:**Signal peptide:**

amino acids 1-17

Sulfatases signature 1:

amino acids 86-99

Homologous region to sulfatase:

amino acids 87-106, 133-146, 216-229, 291-320, 365-375

N-glycosylation sites:amino acids 65-69, 112-116, 132-136, 149-153, 171-175,
198-202, 241-245, 561-565, 608-612, 717-721, 754-758,
764-768

77/392

FIGURE 75

CCCACGCGTCCGCCCACGCGTCCGGTGGACTATGGGCCAGTTTTTGTGCAAGAACCAGAT
GATATTATTTTTTCCAAGTCTGATGAAAAGAAGGTAGCATTGAATTGTGAAGTTCGT
GGCAATCCAGTTCCAGTTACAGATGGCTTCGAAATGGAACAGAAATAGATCTGGAAAGT
GATTATCGCTACAGTTTGATAGATGGCACCTTCATTATAAGCAATCCAAGTGAAGCAAAG
GATTCTGGTCATTATCAGTGTTTAGCAACCAACACTGTGGGGAGTATTCTTAGTAGAGAA
GCTACACTGCAGTTTGCTATCTGGGAAATTTTAGTGGCCGGACAAGAAGTGCAGTCTCT
GTGAGGGAAGGCCAGGGTGTCTGTTCTGATGTGCTCTCTCCGCCACATTCACCAGAGATC
ATCTATAGCTGGGTATTTAATGAGTTCCTTCTTTGTGGCGGAAGACAGCCGGCCGGTTC
ATCTCCCAGGAGACAGGCAACCTTTATATTTCTAAAGTCCAAACATCAGATGTTGGCAGC
TATATTTGTCTGGTGAAAAACACAGTGACGAATGCTAGAGTCCTTAGTCCTCCAACGCCA
CTCACTCTGCGTAATGATGGTGTGATGGGAGAATATGAGCCGAAAATTGAGGTCCATTTT
CCTTTCACGGTTACAGCTGCTAAAGGAACAACCTGTTAAGATGGAATGCTTTGCACTTGGC
AACCCCGTTCCAACAATCACATGGATGAAGGTTAATGGTTATATTCCTAGTAAGGCACGT
CTGCGGAAATCTCAGGCGGTGCTGGAAATACCGAATGTACAGCTGGATGATGCAGGCATT
TATGAGTGCAGAGCTGAAAACCTCACGTGGAAAAAATTCCTTTTCGTGGACAATTACAAGTA
TACACCTACCCACACTGGGTAGAAAACTGAATGATACTCAGTTAGACAGTGGGAGCCCT
CTCCGATGGGAATGTAAGGCTACTGGAAAACCCAGACCCACGTATCGTTGGCTGAAGAAT
GGAGTACCCCTCTCACCTCAGAGTAGGGTTGAGATGGTTAATGGAGTATTGATGATCCAC
AATGTGAATCAATCAGATGCTGGAATGTATCAGTGTTTGGCTGAAAATAAGTATGGAGCC
ATTTACGCTAGTGCTGAGCTGAAGATTCTAGCTTCAGCTCCCACTTTTGCACTGAATCAA
CTGAAGAAAACAATAATTGTTACCAAAGACCAAGAAGTTGTCATAGAGTGCAAACCCCAA
GGCTCTCCAAAACCAACCATCTCTTGGAAGAAAGGAGACAGAGCAGTTAGAGAAAACAAA
AGAATAGCTATTCTCCAGACGGGAGTCTACGGATCCTAAATGCTTCCAATCAGACGAG
GGAAAGTACGTTTGCCGAGGGGAAAACGTCTTTGGTTCTGCTGAAAT

78/392

FIGURE 76

MCSPPPHSPEIIYSWVFNEFPSFVAEDSRRFISQETGNLYISKVQTS DVGSIYICLVKNTV
TNARVLSPPPTPLTLRNDGVMGEYEPKIEVHFPPFTVTAAGTTVKMECFALGNPVPTITWM
KVNGYIPSKARLRKSQAVLEIPNVQLDDAGIYECRAENSRGKNSFRGQLQVYTYPHWVEK
LNDTQLDSGSPLRWECKATGKPRPTYRWLKNQVPLSPQSRVEMVNGVLMIHNVNQSDAGM
YQCLAENKYGAIYASAELKILASAPTFALNQLKKTIIIVTKDQEVVIECKPQGS PKPTISW
KKGDRVAVRENKRIAILPDGSLRILNASKSDEGKYVCRGENVFGSAE

Signal sequence:

None

Transmembrane domain:

None

N-glycosylation site:

182-185, 234-237, 325-328

Tyrosine kinase phosphorylation site:

328-334

N-myristoylation site:

50-55, 150-155, 239-244, 250-255

Immunoglobulin domain:

2-56, 100-156, 189-245, 281-338

79/392

FIGURE 77

GCTCCCAGCCAAGAACCTCGGGGCCGCTGCGCGGTGGGGAGGAGTTCCCCGAAACCCGGC
CGCTAAGCGAGGCCTCCTCCTCCCGCAGATCCGAACGGCCTGGGCGGGGTACCCCGGGT
GGGACAAGAAGCCCGCCGCTGCCTGCCCGGGCCCGGGAGGGGGCTGGGGCTGGGGCCGG
AGGCGGGGTGTGAGTGGGTGTGTGCGGGGGCGGAGGCTTGATGCAATCCCGATAAGAAA
TGCTCGGGTGTCTTGGGCACCTACCCGTGGGGCCCCGTAAGGCGCTACTATATAAGGCTGC
CGGCCCCGAGCCCGCGCCGTGAGAGCAGGAGCGCTGCGTCCAGGATCTAGGGCCACGA
CCATCCCAACCCGGCACTCACAGCCCCGAGCGCATCCCGGTGCGCGCCAGCCTCCCCG
ACCCCATCGCCGGAGCTGCGCCGAGAGCCCCAGGGAGGTGCCATGCGGAGCGGGTGTGT
GGTGGTCCACGTATGGATCCTGGCCGGCCTCTGGCTGGCCGTGGCCGGGCGCCCCCTCGC
CTTCTCGGACGCGGGGCCCCACGTGCATACGGCTGGGGCGACCCCATCCGCCTGCGGCA
CCTGTACACCTCCGGCCCCCACGGGCTCTCCAGCTGCTTCTGCGCATCCGTGCGGACGG
CGTGGTGGACTGCGCGCGGGGCCAGAGCGCGCACAGTTTGCTGGAGATCAAGGCAGTCGC
TCTGCGGACCGTGGCCATCAAGGGCGTGACAGCGTGCGGTACCTCTGCATGGGCGCCGA
CGGCAAGATGCAGGGGCTGCTTCAGTACTCGGAGGAAGACTGTGCTTTCGAGGAGGAGAT
CCGCCCAGATGGCTACAATGTGTACCGATCCGAGAAGCACCGCCTCCCGGTCTCCCTGAG
CAGTGCCAAACAGCGGCAGCTGTACAAGAACAGAGGCTTTCTTCCACTCTCTCATTTCT
GCCCATGCTGCCATGGTCCCAGAGGAGCCTGAGGACCTCAGGGGCCACTTGGAATCTGA
CATGTTCTCTTCGCCCCCTGGAGACCGACAGCATGGACCCATTTGGGCTTGTACCCGACT
GGAGGCCGTGAGGAGTCCCAGCTTTGAGAAGTAACTGAGACCATGCCCCGGCCTCTTCAC
TGCTGCCAGGGGCTGTGGTACCTGCAGCGTGGGGGACGTGCTTCTACAAGAACAGTCCTG
AGTCCACGTTCTGTTTAGCTTTAGGAAGAAACATCTAGAAGTTGTACATATTCAGAGTTT
TCCATTGGCAGTGCCAGTTTCTAGCCAATAGACTTGTCTGATCATAACATTGTAAGCCTG
TAGCTTGCCCAGCTGCTGCTGGGCCCCCATTCCTGCTCCCTCGAGGTTGCTGGACAAGCT
GCTGCACTGTCTCAGTTCTGCTTGAATACCTCCATCGATGGGGAACCTCACTTCCTTTGGA
AAAATTCTTATGTCAAGCTGAAATTCTTAATTTTTTCTCATCACTTCCCCAGGAGCAGC
CAGAAGACAGGCAGTAGTTTTAATTTAGGAACAGGTGATCCACTCTGTAAAACAGCAGG
TAAATTTCACTCAACCCCATGTGGGAATTGATCTATATCTCTACTTCCAGGGACCATTG
CCCTTCCCAAATCCCTCCAGGCCAGAACTGACTGGAGCAGGCATGGCCACCAGGCTTCA
GGAGTAGGGGAAGCCTGGAGCCCCACTCCAGCCCTGGGACAACTTGAGAATTCCCCCTGA
GGCCAGTTCTGTGATGGATGCTGTCTGAGAATAACTTGCTGTCCCGGTGTACCTGCTT
CCATCTCCAGCCCCACCAGCCCTCTGCCCACCTCACATGCCTCCCATGGATTGGGGCCT
CCCAGGCCCCCACCTTATGTCAACCTGCATTCTTGTTCAAAAATCAGGAAAAGAAAAG
ATTTGAAGACCCCAAGTCTTGTCAATAACTTGCTGTGTGGAAGCAGCGGGGAAGACCTA
GAACCCTTTCCCCAGCACTTGGTTTTCCAACATGATATTTATGAGTAATTTATTTTGATA
TGTACATCTCTTATTTTCTTACATTATTTATGCCCCCAAATTATATTTATGTATGTAAGT
GAGGTTTGTTTTGTATATTAAATGGAGTTTGTTTGT

80/392

FIGURE 78

MRSGCVVHVWILAGLWLA VAGRPLAFSDAGPHVHYGWGDP IRLRHLYTSGPHGLSSCFL
RIRADGVVDCARGQSAHSLLEIKAV ALRTVAIKGVH SVRYLCMGADGKM QGLLQYSEEDC
AFEEEIRPDGYNVYRSEK HRLPVSLSSAKQRQLYKNRGFLPLSHFLPMLPMVPEEPEDLR
GHLESDMFSSPLETDSMDPFGLVTGLEAVRSPSF EK

Signal peptide:
amino acids 1-22

Casein kinase II phosphorylation site:
amino acids 78-82, 116-120, 190-194, 204-208

N-myristoylation site:
amino acids 15-21, 54-60, 66-72, 201-207

Prokaryotic membrane lipoprotein lipid attachment site:
amino acids 48-59

81/392

FIGURE 79

CGGACGCGTGGGCGGACGCGTGGGCCTGGGCAAGGGCCGGGGCGCCGGGCGGAGCCACCTCTTCCC
CTCCCCCGCTTCCCTGTCEGCTCCGCTGGCTGGACGCGCTGGAGGAGTGGAGCAGCACCCGGCCG
GCCCTGGGGGCTGACAGTCCGCAAAAGTTTGGCCCCGAAGAGGAAGTGGTCTCAAACCCCGGCAGGTG
GCGACCAGGCCAGACCAGGGGCGCTCGCTGCCCTGCGGGCGGGCTGTAGGCGAGGGCGCGCCCCAGT
GCCGAGACCCGGGGCTTCAAGAGCCGGCCCCGGGAGAGAAGAGTGC GGCGGACGGAGAAAAACA
ACTCCAAAGTTGGCGAAAGGCACCGCCCCCTACTCCCGGGCTGCCGCCGCCCTCCCCGCCCCCAGCCC
TGGCATCCAGAGTACGGGTGAGCCCCGGGCCATGGAGCCCCCTGGGGAGGCGGCACCAGGGAGCC
TGGGCGCCCCGGGGCTCCGCCGCGACCCCATCGGGTAGACCACAGAAGCTCCGGGACCCCTTCCGGCA
CCTCTGGACAGCCCAGGATGCTGTTGGCCACCCTCCTCCTCCTCCTCCTTGGAGGCGCTCTGGCCC
ATCCAGACCCGATTATTTTCCAAATCATGCTTGTGAGGACCCCCCAGCAGTGTCTTAGAAGTGC
AGGGCACCTTACAGAGGCCCTGGTCCGGGACAGCCGCACCTCCCCCTGCCAACTGCACCTGGGTCA
TCCTGGGCAGCAAGGAACAGACTGTCAACATCAGGTTCCAGAAGCTACACCTGGCCTGTGGCTCAG
AGCGCTTAACCTACGCTCCCCCTCCAGCCATGATCTCCCTGTGTGAGGCACCTCCCAGCCCTC
TGCAGCTGCCCGGGGGCAACGTCAACATCACCTTACAGCTATGCTGGGGCCAGAGCACCCATGGGCC
AGGGCTTCTGTCTCCTACAGCCAAGATTGGCTGATGTGCCTGCAGGAAGAGTTTCAGTGCCTGA
ACCACCGCTGTGTATCTGCTGTCCAGCGCTGTGATGGGGTTGATGCCTGTGGCGATGGCTCTGATG
AAGCAGGTTGCAGCTCAGACCCCTTCCCTGGCTGACCCCAAGACCCGTCCCCCTCCTGCCTTGCA
ATGTCACCTTGGAGGACTTCTATGGGGTCTTCTCCTCCTCCTGGATATACACCTAGCCTCAGTCT
CCCACCCCCAGTCCCTGCCATTGGCTGTGGACCCCCATGATGGCCGGCGGTGGCCGTGCGCTTCA
CAGCCCTGGACTTGGGCTTTGGAGATGCAGTGCATGTGTATGACGGCCCTGGGCCCCCTGAGAGCT
CCCGACTACTGCGTAGTCTCACCCACTTCAAGCAATGGCAAGGCTGTCACTGTGGAGACACTGTCTG
GCCAGGCTGTGTCTCCTACCAACAGATTGCTTGGAGCAATGGTCTGTGGCTTCAATGCCACCTACC
ATGTGCGGGCTATTGCTTGCCTTGGGACAGACCTGTGGCTTAGGCTTGGCTTGGGAGCTGGCG
AAGGCCTAGGTGAGCGCTGCTACAGTGAGGCACAGCGCTGTGACGGCTCATGGGACTGTGCTGACG
GCACAGATGAGGAGGACTGCCCAGGCTGCCACCTGGACACTTCCCCCTGTGGGGCTGCTGGCACCT
CTGGTGCCACAGCCTGCTACCTGCTGCTGACCGCTGCAACTACCAGACTTCTGTGCTGATGGAG
CAGATGAGAGACGCTGTCCGCATTGCCAGCCTGGCAATTTCCGATGCCGGGACGAGAAGTGCCTGT
ATGAGACGTGGGTGTGCGATGGGCAGCCAGACTGTGCGGACGGCAGTGATGAGTGGGACTGCTCCT
ATGTTCTGCCCCGCAAGGTCATTACAGCTGCAGTCATTGGCAGCCTAGTGTGCGGCCTGCTCCTGG
TCATCGCCCTGGGCTGCACCTGCAAGCTCTATGCCATTTCGACCCAGGAGTACAGCATCTTTGCCC
CCCTCTCCCGGATGGAGGCTGAGATTGTGCAGCAGGCAGGCCACCCCTTCTACCGGCAGCTCATTG
CCCAGGGTGCCATCCACCTGTAGAAGACTTTCCTACAGAGAATCCTAATGATAACTCAGTGTGG
GCAACCTGCGTTCTCTGCTACAGATCTTACGCCAGGATATGACTCCAGGAGGTGGCCAGGTGCC
GCCGTGCTCAGCGGGGCGCTTGTGTCGACGCTGGTACGCGCTCTCCGCCGTGGGGCTTGCTCC
CTCGAACCAACACCCCGGCTCGGGCCTCTGAGGCCAGATCCAGGTCACACCTTCTGCTGCTCCCC
TTGAGGCCCTAGATGGTGGCACAGGTCCAGCCCGTGAGGGCGGGGCGAGTGGGTGGGCAAGATGGGG
AGCAGGCACCCCCACTGCCCATCAAGGCTCCCCCTCCCATCTGCTAGCACGTCTCCAGCCCCCACTA
CTGTCCCTGAAGCCCCAGGGCCACTGCCCTCACTGCCCTTAGAGCCATCACTATTGTCTGGAGTGG
TGCAGGCCCTGCGAGGCCGCTGTTGCCAGCCTGGGGCCCCCAGGACCAACCCGAGCCCCCTG
GACCCCAACAGCAGTCTTGGCCCTGGAAGATGAGGACGATGTGCTACTGGTGCCACTGGCTGAGC
CGGGGGTGTGGGTAGCTGAGGCAGAGGATGAGCCACTGCTTACCTGAGGGGACCTGGGGGCTCTAC
TGAGGCCTCTCCCCTGGGGGCTCTACTCATAGTGGCAACCTTTTAGAGGTGGGTGAGCCTCCCC
TCCACCACTTCCCTTCCCTGTCCCTGGATTTCAGGGACTTGGTGGGCCTCCCGTTGACCTATGTAG
CTGCTATAAAGTTAAGTGTCCCTCAGGCAGGGAGAGGGCTCACAGAGTCTCCTCTGTACGTGGCCA
TGGCCAGACACCCAGTCCCTTACCACCACCTGCTCCCCACGCCACCACCTTTGGGTGGCTGTT
TTTAAAAAGTAAAGTTCTTAGAGGATCATAGGTCTGGACACTCCATCCTTGCCAAACCTCTACCCA
AAAGTGGCCTTAAGCACCGGAATGCCAATTAACATAGAGACCTCCAGCCCCAAGGGGAGGATTGT
GGCAGAACCTGAGGTTTTGCCATCCACAATCCCTCCTACAGGCCTGGCTCACAAAAGAGTGCAA
CAAATGCTTCTATTCCATAGCTACGGCATTGCTCAGTAAGTTGAGGTCAAAAATAAAGGAATCATA
CATCTC

82/392

FIGURE 80

MLLATLLLLLLGGALAHPDRIIFPNHACEDPPAVLLEVQGTLQRPLVRDSRTSPANCTWL
ILGSKEQTVTIRFQKLHLACGSERLTLRSPLQPLISLCEAPPSPLQLPGGNVTITYSYAG
ARAPMGQGFLLSYSQDWLMCLQEETFQCLNHRCVSAVQRCDGVDACGDGSDEAGCSSDPFP
GLTPRPVPSLPCNVTLEDFYGVFSSPGYTHLASVSHPQSCHWLLDPHDGRRRLAVRFTALD
LGFGDAVHVYDGPPESSRLLRSLTHFSNGKAVTVETLSGQAVVSYHTVAWSNGRGFNA
TYHVRGYCLPWDRPCGLGSGLGAGEGLGERCYSEAQRCDGSWDCADGTDEEDCPGCPPGH
FPCGAAGTSGATACYLPADRCNYQTFCADGADERRCRHCQPGNFRRCRDEKCVYETWVCDG
QPD CADGSDEWDCSYVLPRKVITA AVIGSLVCGLLLVIALGCTCKLYAIRTQEYSIFAPL
SRMEAEIVQQAPP SYGQLIAQGAIPPVEDFPTENPNDNSVLGNLRSLLQILRQDMTPGG
GPGARRRQRGRMLMRRLVRRLLRWGLLPRTNTPARASEARSQVTPSAAPLEALDGGTGPAR
EGGAVGGQDGEQAPPLPIKAPLPSASTSPAPTTVPEAPGPLPSLPLEPSLLSGVVQALRG
RLLPSLGP PGPTRSPPGPHTAVLALEDEDDVLLVPLAEPGVWVAEAEDEPLLT

Important features:

Signal peptide:

amino acids 1-16

Transmembrane domain:

amino acids 442-462

LDL-receptor class A (LDLRA) domain proteins:

amino acids 411-431, 152-171, 331-350 and 374-393

83/392

FIGURE 81

CTTCTGTGCTGTTCCCTTCTTGCCTCTAACTTGTAACAAGACGTACTAGGACGATGCTAA
TGAAAGTCAAAACCGCTGGGTTTTTGAAGGATCCTTGGGACCTCATGCACATTTGTG
GAACTGGATGGAGAGATTTGGGGAAGCATGGACTCTTTAGCCAGCTTAGTTCTCTGTGG
AGTCAGCTTGCTCCTTTCTGGAAGTGTGGAAGGTGCCATGGACTTGATCTTGATCAATTC
CCTACCTCTTGTATCTGATGCTGAAACATCTCTCACCTGCATTGCCTCTGGGTGGCGCCC
CCATGAGCCCATCACCATAGGAAGGGACTTTGAAGCCTTAATGAACCAGCACCAGGATCC
GCTGGAAGTTACTCAAGATGTGACCAGAGAATGGGCTAAAAAGTTGTTTTGGAAGAGAGA
AAAGGCTAGTAAGATCAATGGTGCTTATTTCTGTGAAGGGCGAGTTCGAGGAGAGGCAAT
CAGGATACGAACCATGAAGATGCGTCAACAAGCTTCCTTCCCTACCAGCTACTTTAACTAT
GACTGTGGACAAGGGAGATAACGTGAACATATCTTTCAAAAAGGTATTGATTAAAGAAGA
AGATGCAGTGATTTACAAAAATGGTTCCTTCATCCATTAGTGCCCCGGCATGAAGTACC
TGATATTCTAGAAGTACACCTGCCTCATGCTCAGCCCCAGGATGCTGGAGTGTACTCGGC
CAGGTATATAGGAGGAAACCTCTTCACCTCGGCCTTCACCAGGCTGATAGTCCGGAGATG
TGAAGCCCAGAAGTGGGGACCTGAATGCAACCATCTCTGTACTGCTTGTATGAACAATGG
TGTCTGCCATGAAGATACTGGAGAATGCATTTGCCCTCCTGGGTTTATGGGAAGGACGTG
TGAGAAGGCTTGTGAAGTGCACACGTTTGGCAGAAGCTTGTAAGAAAGGTGCAGTGGACA
AGAGGGATGCAAGTCTTATGTGTTCTGTCTCCCTGACCCCTATGGGTGTTCTGTGCCAC
AGGCTGGAAGGGTCTGCAGTGCAATGAAGCATGCCACCCTGGTTTTTACGGGGCCAGATTG
TAAGCTTAGGTGCAGCTGCAACAATGGGGAGATGTGTGATCGCTTCCAAGGATGTCTCTG
CTCTCCAGGATGGCAGGGGCTCCAGTGTGAGAGAGAAGGCATACCGAGGATGACCCCAA
GATAGTGGATTTGCCAGATCATATAGAAGTAAACAGTGGTAAATTTAATCCCATTTGCAA
AGCTTCTGGCTGGCCGCTACCTACTAATGAAGAAATGACCCTGGTGAAGCCGGATGGGAC
AGTGCTCCATCCAAAAGACTTTAACCATACGGATCATTTCTCAGTAGCCATATTACCAT
CCACCGGATCCTCCCCCTGACTCAGGAGTTTGGGTCTGCAGTGTGAACACAGTGGCTGG
GATGGTGGAAAAGCCCTTCAACATTTCTGTTAAAGTTCTTCCAAAGCCCTGAATGCCCC
AAACGTGATTGACACTGGACATAACTTTGCTGTCAACATCAGCTCTGAGCCTTACTT
TGGGGATGGACCAATCAAATCCAAGAAGCTTCTATACAAACCCGTTAATCACTATGAGGC
TTGGCAACATATTTCAAGTGACAAATGAGATTGTTACACTCAACTATTTGGAACCTCGGAC
AGAATATGAACTCTGTGTGCAACTGGTCCGTCGTGGAGAGGGTGGGGAAGGGCATCCTGG
ACCTGTGAGACGCTTCACAACAGCTTCTATCGGACTCCCTCCTCCAAGAGGTCTAAATCT
CCTGCCTAAAAGTCAGACCACTCTAAATTTGACCTGGCAACCAATATTTCCAAGCTCGGA
AGATGACTTTTATGTTGAAGTGGAGAGAAGGTCTGTGCAAAAAGTGATCAGCAGAATAT
TAAAGTTCCAGGCAACTTGACTTCGGTGCTACTTAACAACTTACATCCCAGGGAGCAGTA
CGTGGTCCGAGCTAGAGTCAACACCAAGGCCAGGGGAATGGAGTGAAGATCTCACTGC
TTGGACCCTTAGTGACATTCTTCCTCCTCAACCAGAAAACATCAAGATTTCCAACATTAC
ACACTCCTCGGCTGTGATTTCTTGGACAATATTGGATGGCTATTCTATTCTTCTATTAC
TATCCGTTACAAGGTTCAAGGCAAGAATGAAGACCAGCACGTTGATGTGAAGATAAAGAA
TGCCACCATCATTCAGTATCAGCTCAAGGGCCTAGAGCCTGAAACAGCATACCAGGTGGA
CATTTTTTGAGAGAAACAACATAGGGTCAAGCAACCCAGCCTTTTCTCATGAACTGGTGAC
CCTCCAGAACTCTCAAGCACCAGCGGACCTCGGAGGGGGGAAGATGCTGCTTATAGCCAT
CCTTGGCTCTGCTGGAATGACCTGCCTGACTGTGCTGTTGGCCTTTCTGATCATATTGCA
ATTGAAGAGGGCAAATGTGCAAAGGAGAATGGCCCAAGCCTTCCAAAACGTGAGGGAAGA
ACCAGCTGTGCAGTTCAACTCAGGGACTCTGGCCCTAAACAGGAAGGTCAAAAACAACCC
AGATCCTACAATTTATCCAGTGCTTGACTGGAATGACATCAAATTTCAAGATGTGATTGG
GGAGGGCAATTTTGGCCAAGTTCTTAAGGCGCGCATCAAGAAGGATGGGTACGGATGGA
TGCTGCCATCAAAAGAATGAAAGAATATGCCTCCAAAGATGATCACAGGGACTTTGCAGG
AGAACTGGAAGTTCTTTGTAAACTTGGACACCATCCAAACATCATCAATCTCTTAGGAGC

84/392

ATGTGAACATCGAGGCTACTTGTACCTGGCCATTGAGTACGCGCCCCATGGAAACCTTCT
GGACTTCCTTCGCAAGAGCCGTGTGCTGGAGACGGACCCAGCATTTGCCATTGCCAATAG
CACCGCGTCCACACTGTCTCCACAGCAGCTCCTTCACTTCGCTGCCGACGTGGCCCGGGG
CATGGACTACTTGAGCCAAAAACAGTTTATCCACAGGGATCTGGCTGCCAGAAACATTTT
AGTTGGTGAAAACCTATGTGGCAAAAATAGCAGATTTTGGATTGTCCCGAGGTCAAGAGGT
GTACGTGAAAAAGACAATGGGAAGGCTCCCAGTGCGCTGGATGGCCATCGAGTCACTGAA
TTACAGTGTGTACACAACCAACAGTGATGTATGGTCCTATGGTGTGTTACTATGGGAGAT
TGTTAGCTTAGGAGGCACACCCCTACTGCGGGATGACTTGTGCAGAACTCTACGAGAAGCT
GCCCCAGGGCTACAGACTGGAGAAGCCCCCTGAACTGTGATGATGAGGTGTATGATCTAAT
GAGACAATGCTGGCGGGAGAAAGCCTTATGAGAGGCCATCATTTGCCAGATATTGGTGTC
CTTAAACAGAATGTTAGAGGAGCGAAAGACCTACGTGAATACCACGCTTTATGAGAAGTT
TACTTATGCAGGAATTGACTGTTCTGCTGAAGAAGCGGCCCTAGGACAGAACATCTGTATA
CCCTCTGTTTCCCTTTCACTGGCATGGGAGACCCCTTGACAACCTGCTGAGAAAACATGCCT
CTGCCAAAGGATGTGATATATAAGTGTACATATGTGCTGGAATTCTAACAAGTCATAGGT
TAATATTTAAGACACTGAAAAATCTAAGTGATATAAATCAGATTCTTCTCTCTCATTTTA
TCCCTCACCTGTAGCATGCCAGTCCCGTTTCATTTAGTCATGTGACCACTCTGTCTTG
TTTCCACAGCCTGCAAGTTCAGTCCAGGATGCTAACATCTAAAAATAGACTTAAATCTCA
TTGCTTACAAGCCTAAGAATCTTTAGAGAAGTATACATAAGTTTAGGATAAAATAATGGG
ATTTTCTTTTCTTTTCTCTGGTAATATTGACTTGTATATTTTAAGAAATAACAGAAAGCC
TGGGTGACATTTGGGAGACATGTGACATTTATATATTGAATTAATATCCCTACATGTATT
GCACATTGTAAAAAGTTTTAGTTTTGATGAGTTGTGAGTTTACCTTGTATACTGTAGGCA
CACTTTGCACTGATATATCATGAGTGAATAAATGTCTTGCCTACTCAAAAAAAAAAAAA

85/392

FIGURE 82

MDSLASLVLCGVSLLLSGTVEGAMDLILINSLPLVSDAETSLTCIASGWRPHEPITIGRD
FEALMNQHQDPLEVTQDVTREWAKKVWVKREKASKINGAYFCEGRVRGEAIRIRTMKMRQ
QASFLPATLTMTVDKGDVNISFKKVLIKEEDAVIYKNGSFIHSVPRHEVPDILEVHLPH
AQPQDAGVYSARYIGGNLFTSAFTRLIVRRCEAQKWGPECNHLCTACMNNGVCHEDTGEC
ICPPGFMGRTCEKACELHTFGRTCKERCSCGQEGCKSYVFCCLPDYPGCSCATGWKGLQCNE
ACHPGFYGPDCKLRSCNNGEMCDRFQGCCLSPGWQGLQCEREGIPRMTPKIVDLPDHIE
VNSGKFNPICKASGWPLPTNEEMTLVKPDGTVLHPKDFNHTDHFSAIFTIHRILPPDSG
VWVCSVNTVAGMVEKPFENISVKVLPKPLNAPNVIDTGHNFVINISSEPYFGDGPIKSKK
LLYKPVNHYEAWQHIQVTNEIVTLNYLEPRTEYELCVQLVRRGEGGEGHPGVRRTTAS
IGLPPPRGLNLLPKSQTTILNLTWQPIFPSSDDFYVEVERRSVQKSDQONIKVPGNLTSV
LLNNLHPREQYVVRARVNTKAQGEWSEDLTAWTLSDILPPQPENIKISNITHSSAVISWT
ILDGYSISSITIRYKVQGNEDQHVDVKIKNATIIQYQLKGLEPETAYQVDIFAENNIGS
SNPAFSHELVTLPESQAPADLGGGKMLLIAILGSAGMTCLTVLLAFLIILQLKRANVQRR
MAQAFQNVREEPAVQFNSGTLALNRKVKNPDPTIYPVLDWNDIKFQDVI GEGNFGQVLK
ARIKKDGLRMDAAIKRMKEYASKDDHRDFAGELEVLCKLGHPNIINLLGACEHRGYLYL
AIEYAPHGNLLDFLRKSRVLETDPAFAIANSTASTLSSQQLLHFAADVARGMDYLSQKQF
IHRDLAARNILVGENYVAKIADFGLSRGQEVYVKKTMGRLPVRWMAIESLNYSVYTTNSD
VWSYGVLLWEIVSLGGTPYCGMTCAELYEKLPGQYRLEKPLNCDDEVYDLMRQCWREKPY
ERPSFAQILVSLNRMLEERKTYVNTTLYEKFTYAGIDCSAEEAA

Signal sequence:

1-38

Transmembrane domain:

750-770

N-glycosylation site:140-143, 158-161, 399-402, 438-441, 464-467, 560-563, 596-
599, 649-652, 691-694, 930-933, 1011-1014, 1104-1107**cAMP- and cGMP-dependent protein kinase phosphorylation
site:**

534-537

Tyrosine kinase phosphorylation site:

149-156, 808-816, 1094-1102

N-myristoylation site:18-23, 98-103, 187-192, 196-201, 270-275, 286-291, 295-300,
420-425, 595-600, 984-989, 1036-1041, 1041-1046, 1115-1120**Prokaryotic membrane lipoprotein lipid attachment site:**

882-892

86/392

EGF-like domain cysteine pattern signature:

240-251, 287-298, 329-340

Tyrosine protein kinases specific active-site signature:

960-972

Protein kinase domain:

824-1092

Fibronectin type III domain:

444-529, 543-626, 639-724

EGF-like domain:

220-251, 268-298

laminin_EGF Laminin EGF-like (Domains III and V):

219-268

Immunoglobulin domain:

156-193

Zinc finger:

295-313

Receptor tyrosine kinase:

844-868, 869-898, 936-982, 986-1024, 1025-1052, 1052-1088

87/392

FIGURE 83

CGCGCCGGGCGCAGGGAGCTGAGTGGACGGCTCGAGACGGCGGCGCGTGCAGCAGCTCCA
GAAAGCAGCGAGTTGGCAGAGCAGGGCTGCATTTCCAGCAGGAGCTGCGAGCACAGTGCCT
GGCTCACAACAAGATGCTCAAGGTGTGAGCCGTACTGTGTGTGTGTGCAGCCGCTTGGTG
CAGTCAGTCTCTCGCAGCTGCCGCGGCGGTGGCTGCAGCCGGGGGGCGGTCCGACGGCGG
TAATTTTCTGGATGATAAACAATGGCTCACCACAATCTCTCAGTATGACAAGGAAGTCGG
ACAGTGGAAACAAATTCCGAGACGAAGTAGAGGATGATTATTTCCGCACTTGGAGTCCAGG
AAAACCCCTTCGATCAGGCTTTAGATCCAGCTAAGGATCCATGCTTAAAGATGAAATGTAG
TCGCCATAAAGTATGCATTGCTCAAGATTCTCAGACTGCAGTCTGCATTAGTCACCGGAG
GCTTACACACAGGATGAAAGAAGCAGGAGTAGACCATAGGCAGTGGAGGGGTCCCATATT
ATCCACCTGCAAGCAGTGCCCGAGTGGTCTATCCCAGCCCTGTTTGTGGTTTCAGATGGTCA
TACCTACTCTTTTCAGTGCAAACTAGAAATATCAGGCATGTGTCTTAGGAAAACAGATCTC
AGTCAAATGTGAAGGACATTGCCCATGTCTTCAGATAAGCCCACCAGTACAAGCAGAAA
TGTTAAGAGAGCATGCAGTGACCTGGAGTTCAGGGAAGTGGCAAACAGATTGCGGGACTG
GTTCAAGGCCCTTCATGAAAGTGGAAGTCAAAACAAGAAGACAAAAACATTGCTGAGGCC
TGAGAGAAGCAGATTCGATAACCAGCATCTTGCCAATTTGCAAGGACTCACTTGGCTGGAT
GTTTAACAGACTTGATACAACTATGACCTGCTATTGGACCAGTCAGAGCTCAGAAGCAT
TTACCTTGATAAGAATGAACAGTGTACCAAGGCATTCTTCAATTCTTGTGACACATACAA
GGACAGTTTAAATATCTAATAATGAGTGGTGTCTACTGCTTCCAGAGACAGCAAGACCCACC
TTGCCAGACTGAGCTCAGCAATATTCAGAAGCGGCAAGGGGTAAAGAAGCTCCTAGGACA
GTATATCCCCCTGTGTGATGAAGATGGTTACTACAAGCCAACACAATGTCATGGCAGTGT
TGGACAGTGTGGTGTGTTGACAGATATGGAAATGAAGTCATGGGATCCAGAATAAATGG
TGTTGCAGATTGTGCTATAGATTTTGAGATCTCCGGAGATTTTGCTAGTGGCGATTTTCA
TGAATGGACTGATGATGAGGATGATGAAGACGATATTATGAATGATGAAGATGAAATTGA
AGATGATGATGAAGATGAAGGGGATGATGATGATGGTGGTATGACCATGATGTATACAT
TTGATTGATGACAGTTGAAATCAATAAATTTCTACATTTCTAATATTTACAAAAATGATAG
CCTATTTAAAATTATCTTCTTCCCCAATAACAAAATGATTCTAAACCTCACATATATTTT
GTATAATTATTTGAAAAATTGCAGCTAAAGTTATAGAACTTTATGTTTAAATAAGAATCA
TTTGCTTTGAGTTTTTATATTCTTACACAAAAAGAAAATACATATGCAGTCTAGTCAGA
CAAAATAAAGTTTTGAAGTGCTACTATAATAAATTTTTACGAGAACAACTTTGTAAAT
CTTCATAAGCAAAATGACAGCTAGTGCTTGGGATCGTACATGTTAATTTTTTGAAAGAT
AATTCTAAGTGAATTTAAATAAATAAATTTTTAATGACCTGGGTCTTAAGGATTTAGG
AAAAATATGCATGCTTTAATTGCATTTCCAAAGTAGCATCTTGCTAGACCTAGATGAGTC
AGGATAACAGAGAGATACCACATGACTCCAAAAAAAAAAAAA

88/392

FIGURE 84

MLKVSAVLCVCAAAWCSQSLAAAAVAAAGGRSDGGNFLDDKQWLTTISQYDKEVGQWNK
FRDEVEDDYFRTWSPGKPFQALDPAKDPCLMKCSRHKVCIAQDSQTAVCISHRRLTHR
MKEAGVDHRQWRGPILSTCKQCPVVYPSPVCGSDGHTYSFQCKLEYQACVLGKQISVKCE
GHCPCPSPDKPTSTSRNVKRACSDLEFREVANRLRDWFKALHESGSQNKKTKTLLRPERSR
FDTSILPICKDSLGMFNRDLDTNYDLLLDQSELRSIYLDKNEQCTKAFFNSCDTYKDSL
SNNEWCYCFQRQQDPPCQTELSNIQKRQGVKLLGQYIPLCDEDGYKPTQCHGSGVQCW
CVDRYGNEVMGSRINGVADCAIDFEISGDFASGDFHEWTDDDEDDIMNDEDEIEDDDE
DEGDDDDGGDDHVDVYI

Important features:**Signal peptide:**

amino acids 1-16

Leucine zipper pattern:

amino acids 246-267

N-myristoylation sites:

amino acids 357-362, 371-376 and 376-381

Thyroglobulin type-1 repeat proteins:

amino acids 353-365 and 339-352

89/392

FIGURE 85

CCCACGCGTCCGGCACTGCAGTCTCCAGCCTGAGCCATGGGCGCCGAGCCCTCCTGCTC
CTGCTTCTGTCTTTTCTGGCGCCCTGGGCCACCATAGCCCTCCGGCCGGCCTTAAGGGCC
CTCGGCAGCCTACACTTGCCAACCAACCCACATCCCTCCCGGCTGTAGCCAAGAACTAT
TCGGTTCTCTACTTCCAACAGAAGGTTGATCATTGTTGGATTAACTGTGAAAACTTTT
AATCAGCGGTACCTAGTAGCTGATAAATACTGGAAGAAAAATGGTGGATCAATACTTTTC
TACACTGGTAATGAAGGGGACATTATCTGGTTTTGTAATAACACGGGGTTTCATGTGGGAT
GTGGCTGAGGAACTGAAAGCTATGTTGGTGTGTTGCTGAACATCGATACTATGGAGAGTCT
CTCCCCCTTGGTGACAACCTCATTCAAGGATTCAGACACTTGAATTTCTGACATCAGAA
CAAGCTCTGGCTGATTTTGCAGAGTTAATCAAACACTTGAAAAGAACAATCCCAGGAGCT
GAAAATCAACCTGTTCATTGCCATAGGAGGCTCCTATGGTGGCATGCTTGCCGCTGGTTT
AGGATGAAATATCCTCATATGGTAGTTGGAGCTCTTGCAGCTTCTGCCCCCTATCTGGCAG
TTTGAGGATTTAGTACCTTGTGGTGTATTTATGAAGATCGTAACCTACAGATTTTAGGAAA
AGCGGTCCACATTGTTTCAGAGAGCATCCACAGGTCCTGGGATGCCATTAATCGACTCTCA
AATACTGGCAGTGGTTTTGCAGTGGCTTACTGGAGCCCTTCACTTATGCAGCCCATTAACT
TCTCAGGACATCCAACATTTGAAAGACTGGATCTCTGAAACCTGGGTGAATCTGGCAATG
GTGGACTATCCTTATGCCTCTAACTTTTTACAGCCTTTGCCTGCTTGGCCTATCAAGGTA
GTGTGCCAGTATTTGAAAAATCCCAATGTATCTGATTCACTGCTGCTGCAGAATATTTTC
CAAGCTCTGAATGTATATTACAATTATTCGGGCCAGGTGAAATGCCTGAATATTTTCAGAG
ACAGCAACTAGCAGTCTGGGAACACTGGGTTGGAGCTATCAGGCCTGCACAGAAGTAGTC
ATGCCCTTTTGTACTAATGGTGTGATGACATGTTTGAACCTCACTCATGGAACTTAAAG
GAACCTTTCTGATGACTGTTTTCAACAGTGGGGTGTGAGACCAAGGCCCTCCTGGATCACT
ACTATGTATGGAGGCAAAAACATTAGTTTACACACAAACATTGTTTTTTCAGCAATGGTGAA
CTAGACCCCTGGTCAGGAGGTGGAGTAAGGATATCACAGACACTCTGGTTGCAGTC
ACCATCTCAGAGGGGGGCCACCCTTAGATCTCCGCACCAAGAATGCCTTGGATCCTATG
TCTGTGCTGTTAGCCCGCTCCTTGGAAGTTAGACATATGAAGAATTGGATCAGAGATTTTC
TATGACAGTGCGGGAAAGCAGCACTGAGAACTTTTGATTGTTTTCAATTTCTTCTTTTA
TGTTTACACCACCACATTCCCATTCACTTTGATTTTCTACATGTAATTACCTTCTTTTGT
TTATCATTAGATTTGATGGGGCCAAAGTTGAGATAGAATAGAGGGTGATGACGGTAAGAG
CAAGTGTCCCATGAATGTGATTTCTGGGTTCTCACTGTCTTTTGACCCACGTCTAGGAA
GAATCTTCTTGATAGCTCTCCACACCATCAGTGGCCCTCATAACTGGAGTAGAGTTCCCT
GGTTGCTTTTTCATAAGAGGGGAGAGTTACTTTTC

90/392

FIGURE 86

MGRRALLLLLLSFLAPWATIALRPALRALGSLHLPTNPTSLPAVAKNYSVLYFQQKVDHF
GFNTVKTFFNQRYLVADKYWKKNNGGSILFYTGNEGDIWFNNTGFMWDVAEELKAMLVFA
EHRYYGESLPFGDNSFKDSRHLNFLTSEQALADFAELIKHLKRTIPGAENQPVIAIGGSY
GGMLAAWFRMKYPHVVVGALAASAPIWQFEDLVPCGVFMKIVTTDFRKSGPHCSESIHRS
WDAINRLSNTGSGLQWLTGALHLCSPILTSQDIQHLKDWISETWVNLAMVDYPYASNFLQP
LPAWPIKVVQCQYLKNPNVSDSLLLQNIQALNVYNYSGQVKCLNISETATSSLGTLGWS
YQACTEVVMPFCTNGVDDMFEPHSWNLKELSDDCFQQWGVPRPSWITTMYGKKNISSHT
NIVFSNGELDPWSGGGVTKDITDTLVAVTISEGAHHLDLRTKNALDPMSVLLARSLEVRH
MKNWIRDFYDSAGKQH

Signal sequence:

1-18

Transmembrane domain:

None

N-glycosylation site:

47-50, 101-104, 317-320, 336-339, 345-348, 415-418

Glycosaminoglycan attachment site:

433-436

N-myristoylation site:

178-183, 181-186, 182-187, 198-203, 339-344, 434-439

Amidation site:

1-4

alpha/beta hydrolase fold:

115-372

91/392

FIGURE 87

GGCGGCGTCCGTGAGGGGCTCCTTTGGGCAGGGGTAGTGTGTTGGTGTCCCTGTCTTGCGT
GATATTGACAACTGAAGCTTTCCCTGCACCACTGGACTTAAGGAAGAGTGTACTCGTAGG
CGGACAGCTTTAGTGGCCGGCCGGCCGCTCTCATCCCCCGTAAGGAGCAGAGTCCTTTGT
ACTGACCAAGATGAGCAACATCTACATCCAGGAGCCTCCACGAATGGGAAGGTTTATT
GAAAACCTACAGCTGGAGATATTGACATAGAGTTGTGGTCCAAAAGAGCTCCTAAAGCTTG
CAGAAATTTTATCCAACCTTGTGTTGGAAGCTTATTATGACAATACCATTTTTCATAGAGT
TGTGCCTGGTTTTCATAGTCCAAGGCGGAGATCCTACTGGCACAGGGAGTGGTGGAGAGTC
TATCTATGGAGCGCCATTCAAAGATGAATTCATTACGGTTGCGTTTTAATCGGAGAGG
ACTGGTTGCCATGGCAAATGCTGGTTCTCATGATAATGGCAGCCAGTTTTTCTTCACACT
GGGTCGAGCAGATGAACCTTAACAATAAGCATACCATCTTTGGAAAGGTTACAGGGGATAC
AGTATATAACATGTTGCGACTGTCAGAAGTAGACATTGATGATGACGAAAGACCACATAA
TCCACACAAAATAAAAAGCTGTGAGGTTTTGTTAATCCTTTTGATGACATCATTCCAAG
GGAAATTAAAAGGCTGAAAAAAGAGAAACAGAGGAGGAAGTAAAGAAATTGAAACCCAA
AGGCACAAAAAATTTTAGTTTTACTTTTCATTTGGAGAGGAAGCTGAGGAAGAAGAGGAGGA
AGTAAATCGAGTTAGTCAGAGCATGAAGGGCAAAAGCAAAAGTAGTCATGACTTGCTTAA
GGATGATCCACATCTCAGTTCTGTTCCAGTTGTAGAAAGTGAAAAAGGTGATGCACCAGA
TTTAGTTGATGATGGAGAAGATGAAAGTGCAGAGCATGATGAATATATTGATGGTGATGA
AAAGAACCTGATGAGAGAAAGAATTGCCAAAAAATTA AAAAAGGACACAAGTGCGAATGT
TAAATCAGCTGGAGAAGGAGAAGTGGAGAAGAAATCAGTCAGCCGCAGTGAAGAGCTCAG
AAAAGAAGCAAGACAATTAAAACGGGAACCTCTTAGCAGCAAAACAAAAAAGTAGAAAA
TGCAGCAAAACAAGCAGAAAAAAGAAGTGAAGAGGAAGAAGCCCCCTCAGATGGTGCTGT
TGCCGAATACAGAAGAGAAAAAGCAAAAGTATGAAGCTTTGAGGAAGCAACAGTCAAAGAA
GGGAACCTTCCCGGGAAGATCAGACCCTTGCACTGCTGAACCAGTTTAAATCTAAACTCAC
TCAAGCAATTGCTGAAACACCTGAAAATGACATTCTGAAACAGAAGTAGAAGATGATGA
AGGATGGATGTCACATGTACTTCAGTTTGAGGATAAAAGCAGAAAAGTGAAAGATGCAAG
CATGCAAGACTCAGATACATTTGAAATCTATGATCCTCGGAATCCAGTGAATAAAAGAAG
GAGGGAAGAAAGCAAAAAGCTGATGAGAGAGAAAAAAGAAAGAAGATAAAATGAGAATAA
TGATAACCAGAACTTGCTGGAAATGTGCCTACAATGGCCTTGTAACAGCCATTGTTCCCA
ACAGCATCACTTAGGGGTGTGAAAAGAAGTATTTTTGAACCTGTTGTCTGGTTTTGAAAA
ACAATTATCTTGTGTTTGCAAATTGTGGAATGATGTAAGCAAATGCTTTTGGTTACTGGTA
CATGTGTTTTTCTAGCTGACCTTTTATATTGCTAAATCTGAAATAAAATAACTTTTCT
TCCACAAAAA

92/392

FIGURE 88

MSNIYIQEPPTNGKVLLKTTAGDIDIELWSKEAPKACRNFIQLCLEAYYDNTIFHRVVP
FIVQGGDPTGTGSGGESIYGAPFKDEFHSRLRFNRRGLVAMANAGSHDNGSQFFFTLGRA
DELNNKHTIFGKVTGDTVYNMLRLSEVDIDDDERPHNPHKIKSCEVLFPFDDIIPREIK
RLKKEKPEEEVKKLKPVGTKNFSLLSFGEEAEVEVNRVSQSMKGKSKSSHDLLKDDP
HLSSVPVVESEKGDAPDLVDDGEDESAEHDEYIDGDEKNLMRERIAKKLKKDTSANVKS
A
GEGEVEKKSVSRSEELRKEARQLKRELLAAKQKKVENAAKQAEKRSEEEEAPPDGAVAEY
RREKQKYEALRKQQSKKGTSRDQTLALLNQFKSKLTQAIAETPENDIPETEVEDDEGWM
SHVLQFEDKSRKVKDASMQSDTFEIIDPRNPFVNKRREESKKLMREKKERR

Important features:**Signal peptide:**

amino acids 1-21

N-glycosylation sites:

amino acids 109-112 and 201-204

**Cyclophilin-type peptidyl-prolyl cis-trans isomerase
signature:**

amino acids 49-66

**Homologous region to Cyclophilin-type peptidyl-prolyl cis-
trans isomerase:**

amino acids 96-140, 49-89 and 22-51

93/392

FIGURE 89

CCCGGCTCCGCTCCCTCTGCCCCCTCGGGGTCGCGCGCCACGATGCTGCAGGGCCCTGG
CTCGCTGCTGCTGCTCTTCCTCGCCTCGCACTGCTGCCTGGGCTCGGCGCGGGGCTCTT
CCTCTTTGGCCAGCCCGACTTCTCCTACAAGCGCAGCAATTGCAAGCCCATCCCGGTCAA
CCTGCAGCTGTGCCACGGCATCGAATACCAGAACATGCGGCTGCCAACCTGCTGGGCCA
CGAGACCATGAAGGAGGTGCTGGAGCAGGCCGGCGCTTGGAATCCCGCTGGTCATGAAGCA
GTGCCACCCGGACACCAAGAAGTTCCTGTGCTCGCTCTTCGCCCCCGTCTGCCTCGATGA
CCTAGACGAGACCATCCAGCCATGCCACTCGCTCTGCGTGCAAGGACCGCTGCGC
CCCGGTCATGTCCGCCTTCGGCTTCCCCCTGGCCCGACATGCTTGAGTGCGACCGTTTCCC
CCAGGACAACGACCTTTGCATCCCCCTCGCTAGCAGCGACCACCTCCTGCCAGCCACCGA
GGAAGCTCCAAAGGTATGTGAAGCCTGCAAAAATAAAAATGATGATGACAACGACATAAT
GGAAACGCTTTGTAAAAATGATTTTGCCTGAAAATAAAAGTGAAGGAGATAACCTACAT
CAACCGAGATAACAAAATCATCCTGGAGACCAAGAGCAAGACCATTTACAAGCTGAACGG
TGTGTCCGAAAGGGACCTGAAGAAATCGGTGCTGTGGCTCAAAGACAGCTTGCAGTGCAC
CTGTGAGGAGATGAACGACATCAACGCGCCCTATCTGGTCATGGGACAGAAACAGGGTGG
GGAGCTGGTGATCACCTCGGTGAAGCGGTGGCAGAAGGGGCAGAGAGAGTTCAAGCGCAT
CTCCCGCAGCATCCGCAAGCTGCAGTGCTAGTCCCGGCATCCTGATGGCTCCGACAGGCC
TGCTCCAGAGCACGGCTGACCATTCTGCTCCGGGATCTCAGCTCCCGTTCCCCAAGCAC
ACTCCTAGCTGCTCCAGTCTCAGCCTGGGCAGCTTCCCCCTGCCTTTTGCACGTTTGCAT
CCCCAGCATTTCTGAGTTATAAGGCCACAGGAGTGGATAGCTGTTTTACCTAAAGGAA
AAGCCCACCCGAATCTTGTAGAAATATTCAAATAATAAAATCATGAATATTTTAA

94/392

FIGURE 90

MLQGPGSLLLLFLASHCCLGSAAGLFLFGQPDFS YKRSNCKPIPVNLQLCHGIEYQNMRL
PNLLGHETMKEVLEQAGAWIPLVMKQCHPDTKKFLCSLFAPVCLDDLDDETIQPCHS LCVQ
VKDRCAPVMSAFGFPWPDMLECDRFPQDNDLCIPLASSDHLLPATEEAPKVCEACKNKND
DDNDIMETLCKNDFALKIKVKEITYINRDTKIILETKSKTIYKLNGVSEKDLKKS VLWLK
DSLQCTCEEMNDINAPYLVMGQKQGGELVITSVKRWQKGQREFKRISRSIRKLQC

Important features:

Signal peptide:

amino acids 1-20

Cysteine rich domain, homologous to frizzled N terminus:

amino acids 6-153

95/392

FIGURE 91

GGAAGGGGAGGAGCAGGCCACACAGGCACAGGCCGGTGAGGGACCTGCCCAGACCTGGAG
GGTCTCGCTCTGTACACAGGCTGGAGTGCAGTGGTGTGATCTTGGCTCATCGTAACCTC
CACCTCCCGGGTTCAAGTGATTCTCATGCCTCAGCCTCCCGAGTAGCTGGGATTACAGGT
GGTGACTTCCAAGAGTGACTCCGTCCGAGGAAAATGACTCCCCAGTCGCTGCTGCAGACG
ACACTGTTCCCTGCTGAGTCTGCTCTTCCCTGGTCCAAGGTGCCACGGCAGGGGCCACAGG
GAAGACTTTTCGCTTCTGCAGCCAGCGGAACCAGACACACAGGAGCAGCCTCCACTACAAA
CCCACACCAGACCTGCGCATCTCCATCGAGAACTCCGAAGAGGCCCTCACAGTCCATGCC
CCTTTCCCTGCAGCCCACCCTGCTTCCCGATCCTTCCCTGACCCCAGGGGCCCTTACCAC
TTCTGCCTCTACTGGAACCGACATGCTGGGAGATTACATCTTCTCTATGGCAAGCGTGAC
TTCTTGCTGAGTGACAAAGCCTCTAGCCTCCTCTGCTTCCAGCACCAGGAGGAGAGCCTG
GCTCAGGGCCCCCGCTGTAGCCACTTCTGTACCTCCTGGTGGAGCCCTCAGAACATC
AGCCTGCCAGTGCCGCCAGCTTACCTTCTCCTTCCACAGTCCTCCCCACACGGCCGCT
CACAATGCCTCGGTGGACATGTGCGAGCTCAAAGGGACCTCCAGCTGCTCAGCCAGTTC
CTGAAGCATCCCCAGAAGGCCTCAAGGAGGCCCTCGGCTGCCCCCGCCAGCCAGCAGTTG
CAGAGCCTGGAGTCGAAACTGACCTCTGTGAGATTATGAGGGGACATGGTGTCTTCGAG
GAGGACCGGATCAACGCCACGGTGTGGAAGCTCCAGCCCACAGCCGGCCCTCCAGGACCTG
CACATCCACTCCCGGCAGGAGGAGGAGCAGAGCGAGATCATGGAGTACTCGGTGCTGCTG
CCTCGAACACTCTTCCAGAGGACGAAAGGCCGGAGCGGGGAGGCTGAGAAGAGACTCCTC
CTGGTGGACTTCAGCAGCCAAGCCCTGTTCCAGGACAAGAATTCAGCCAAGTCTGGGT
GAGAAGGTCTTGGGGATTGTGGTACAGAACACCAAAGTAGCCAACCTCACGGAGCCCGTG
GTGCTCACTTTCAGCACCAGCTACAGCCGAAGAATGTGACTCTGCAATGTGTGTTCTGG
GTTGAAGACCCCCACATTGAGCAGCCCCGGGGCATTTGGAGCAGTGCTGGGTGTGAGACCGTC
AGGAGAGAAACCCAAACATCCTGCTTCTGCAACCACTTGACCTACTTTGCAGTGCTGATG
GTCTCCTCGGTGGAGGTGGACGCCGTGCACAAGCACTACCTGAGCCTCCTCTCTACGTG
GGCTGTGTCTCTGCCCCTGGCCTGCCTTGTACCATTTGCCGCTACCTCTGCTCCAGG
GTGCCCCCTGCCGTGCAGGAGGAAACCTCGGGACTACACCATCAAGGTGCACATGAACCTG
CTGCTGGCCGTCTTCCCTGCTGGACACGAGCTTCCCTGCTCAGCGAGCCGGTGGCCCTGACA
GGCTCTGAGGCTGGCTGCCGAGCCAGTGCCATCTTCCCTGCACTTCTCCCTGCTCACCTGC
CTTTCCTGGATGGGCCTCGAGGGGTACAACCTCTACCGACTCGTGGTGGAGGTCTTTGGC
ACCTATGTCCCTGGCTACCTACTCAAGCTGAGCGCCATGGGCTGGGGCTTCCCCATCTTT
CTGGTGACGCTGGTGGCCCTGGTGGATGTGGACAACTATGGCCCCATCATCTTGGCTGTG
CATAGGACTCCAGAGGGCGTCATCTACCTTCCATGTGCTGGATCCGGGACTCCCTGGTC
AGCTACATCACCAACCTGGGCCTCTTCAGCCTGGTGTTCCTGTTCAACATGGCCATGCTA
GCCACCATGGTGGTGCAGATCCTGCGGCTGCGCCCCACACCCAAAAGTGGTCACATGTG
CTGACACTGCTGGGCCTCAGCCTGGTCCCTTGGCCTGCCCTGGGCCTTGATCTTCTTCTCC
TTTGCTTCTGGCACCTTCCAGCTTGTCTGCTCCTTACCTTTTCAGCATCATCACCTCCTTC
CAAGGCTTCCCTCATCTTCATCTGGTACTGGTCCATGCGGCTGCAGGCCCCGGGTGGCCCC
TCCCCCTCTGAAGAGCAACTCAGACAGCGCCAGGCTCCCCATCAGCTCGGGCAGCACCTCG
TCCAGCCGCATCTAGGCCTCCAGCCCACCTGCCCATGTGATGAAGCAGAGATGCGGCCTC
GTCGCACACTGCCTGTGGCCCCCGAGCCAGGCCAGCCCCAGGCCAGTCAGCCGCAGACT
TTGGAAAGCCCAACGACCATGGAGAGATGGGCGGTTGCCATGGTGGACGGACTCCCGGGC
TGGGCTTTTGAATTGGCCTTGGGGACTACTCGGCTCTCACTCAGCTCCCACGGGACTCAG
AAGTGCGCCGCATGCTGCCTAGGGTACTGTCCCCACATCTGTCCCAACCCAGCTGGAGG
CCTGGTCTCTCCTTACAACCCCTGGGCCAGCCCTCATTGCTGGGGGCCAGGCCTTGGAT
CTTGAGGGTCTGGCACATCCTTAATCCTGTGCCCTGCCTGGGACAGAAATGTGGCTCCA
GTTGCTCTGTCTCTCGTGGTCACCCTGAGGGCACTCTGCATCCTCTGTCATTTTAACCTC
AGGTGGCACCCAGGGCGAATGGGGCCCAGGGCAGACCTTCAGGGCCAGAGCCCTGGCGGA

97/392

FIGURE 92

MTPQSLLQTTLFLLSLLFLVQGAHGRGHREDFRFCSQRNQTHRSSLHYKPTPDLRISIEN
SEEALTVHAPFPAAHPASRSFPDPRGLYHFCLYWNRHAGRLHLLYGKRDFFLLSDKASSLL
CFQHQEESLAQGPPLLATSVTSWWSPQNISLPSAASFSTFSFHSPHTAAHNASVDMCELK
RDLQLLSQFLKHPQKASRRPSAAPASQQLOSLESKLTSVRFMGDMVSFEDRINATVWKL
QPTAGLQDLHIHSRQEEEQSEIMEYSVLLPRTLTFQRTKGRSGEAEKRLLLVDFSSQALFQ
DKNSSQVLGEKVLGIVVQNTKVANLTEPVVLTFFQHLQPKNVTLCVFWVEDPTLSSPGH
WSSAGCETVRRETQTSCFCNHLTYFAVLMVSSVEVDAVHKHYLSLLSYVGCVVVSALACL
TIAAYLCSRVPPLPCRKRPRDYTIKVHMNLLLAFLLDTSFLLSEPVALTGSEAGCRASAI
FLHFSLLTCLSWMGLEGYNLYRLVVEVFGTYVPGYLLKLSAMGWGFPIFLVTLVALVDVD
NYGPIILAVHRTPEGVIYPSMCWIRDSLVSYITNLGLFSLVFLFNMAMLATMVVQILRLR
PHTQKWSHVLTLGLSLVLGLPWALIFFSFASGTFQLVVLVLYLSIITSFQGFIFIFIWYS
MRLQARGGPSPLKSNSDSARLPISSGSTSSRI

Important features:**Signal peptide:**

amino acids 1-25

Putative transmembrane domains:amino acids 382-398, 402-420, 445-468, 473-491, 519-537,
568-590 and 634-657**Microbodies C-terminal targeting signal:**

amino acids 691-693

cAMP- and cGMP-dependent protein kinase phosphorylation sites:

amino acids 198-201 and 370-373

N-glycosylation sites:amino acids 39-42, 148-151, 171-174, 234-237, 303-306, 324-
327 and 341-344**G-protein coupled receptors family 2 proteins:**

amino acids 475-504

98/392

FIGURE 93

CCCACGCGTCCGAAGGCAGACAAAGGTTCAATTTGTAAAGAAGCTCCTTCCAGCACCTCCT
CTCTTCTCCTTTTGCCCAAACACCCAGTGAGTGTGAGCATTTAAGAAGCATCCTCTGC
CAAGACCAAAGGAAAGAAGAAAAAGGGCCAAAGCCAAATGAAACTGATGGTACTTGT
TTTCACCATTTGGGCTAACTTTGCTGCTAGGAGTTCAAGCCATGCCTGCAAATCGCCTCTC
TTGCTACAGAAAGATACTAAAAGATCACAACTGTCACAACCTTCCGGAAGGAGTAGCTGA
CCTGACACAGATTGATGTCAATGTCCAGGATCATTTCTGGGATGGGAAGGGATGTGAGAT
GATCTGTTACTGCAACTTCAGCGAATTGCTCTGCTGCCCAAAGACGTTTTCTTTGGACC
AAAGATCTCTTTTCGTGATTCCTTGCAACAATCAATGAGAATCTTCATGTATTCTGGAGAA
CACCATTCCTGATTTCCACAAACTGCACTACATCAGTATAACTGCATTTCTAGTTTCTA
TATAGTGCAATAGAGCATAGATTCTATAAATTCTTACTTGTCTAAGACAAGTAAATCTGT
GTTAAACAAGTAGTAATAAAAGTTAATTCATCTAAAAAAAAAAAAA

99/392

FIGURE 94

MKLMVLVFTIGLTLLLGVQAMPANRLSCYRKILKDHNCHNLPEGVADLTQIDVNVQDHFW
DGKGCEMICYCNFSELLCCPKDVFFGPKISFVIPCNNQ

Important features:

Signal peptide:

amino acids 1-20

N-glycosylation site:

amino acids 72-76

Tyrosine kinase phosphorylation site:

amino acids 63-71

100/392

FIGURE 95

GAATTCCGGGCCCCAGGATGCCAACTTTGAATAGGATGAAGACTACAACCTTGTTCCCTTC
TCATCTGCATCTCCCTGCTCCAGCTGATGGTCCCAGTGAATACTGATGAGACCATAGAGA
TTATCGTGGAGAATAAGGTCAAGGAACTTCTTGCCAATCCAGCTAACTATCCCTCCACTG
TAACGAAGACTCTCTCTTGCACTAGTGTCAAGACTATGAACAGATGGGCCTCCTGCCCTG
CTGGGATGACTGCTACTGGGTGTGCTTGTGGCTTTGCCTGTGGATCTTGGGAGATCCAGA
GTGGAGATACTTGCAACTGCCTGTGCTTACTCGTTGACTGGACCACTGCCCCTGCTGCC
AACTGTCCTAAGAATGAAGAGGTGGAGAACCCAGCTTTGATATGATGAATCTAACAAAA
CTGCAGTCTCAATTTGGAAATCTGACTCATGTGCCTTTAAATGTGTTTCATATTGCCCAT
TACCCTGCTTCTTGAAATGCTTCTTGAAAAATAAAGACAAATTTGCATGTG

101/392

FIGURE 96

MKTTTCSLLICISLLQLMVPVNTDETIEIIVENKVKELLANPANYPSTVTKTLSCTSVK
TMNRWASCPAGMTATGCACGFACGSWEIQSGDTCNCLCLLDWTTARCCQLS

102/392

FIGURE 97

GAGGCAGAAAGGCAGAAAGGAGAAAATTCAAGGATAACTCTCCTGAGGGGTGAGCCAAGCC
CTGCCATGTAGTGCACGCAGGACATCAACAAACACAGATAACAGGAAATGATCCATTCCC
TGTGGTCACTTATTCTAAAGGCCCAACCTTCAAAGTTCAAGTAGTGATATGGATGACTC
CACAGAAAGGGAGCAGTCACGCCTTACTTCTTGCCCTTAAGAAAAGAGAAGAAATGAAACT
GAAGGAGTGTGTTTCCATCCTCCCACGGAAGGAAAGCCCCTCTGTCCGATCCTCCAAAGA
CGGAAAGCTGCTGGCTGCAACCTTGCTGCTGGCACTGCTGTCTTGCTGCCTCACGGTGGT
GTCTTTCTACCAGGTGGCCGCCCTGCAAGGGGACCTGGCCAGCCTCCGGGCAGAGCTGCA
GGGCCACCACGCGGAGAAGCTGCCAGCAGGAGCAGGAGCCCCAAGGCCGGCCTGGAGGA
AGCTCCAGCTGTCACCGCGGACTGAAAATCTTTGAACCACCAGCTCCAGGAGAAGGCAA
CTCCAGTCAGAACAGCAGAAAATAAGCGTGCCGTTCAAGGTCCAGAAGAAACAGTCACTCA
AGACTGCTTGCAACTGATTGCAGACAGTGAACACCAACTATACAAAAAGGATCTTACAC
ATTTGTTCCATGGCTTCTCAGCTTTAAAAGGGGAAGTGCCCTAGAAGAAAAAGAGAATAA
AATATTGGTCAAAGAAACTGGTTACTTTTTTATATATGGTCAGGTTTTATATACTGATAA
GACCTACGCCATGGGACATCTAATTCAGAGGAAGAAGGTCCATGTCTTTGGGGATGAATT
GAGTCTGGTGACTTTGTTTCGATGTATTCAAAATATGCCTGAAACACTACCCAATAATTC
CTGCTATTGAGCTGGCATTGCAAACTGGAAGAAGGAGATGAACTCCAACCTTGCAATACC
AAGAGAAAATGCACAAATATCACTGGATGGAGATGTCACATTTTTTGGTGCATTGAAACT
GCTGTGAACCTACTTACACCATGTCTGTAGCTATTTTCCTCCCTTTCTCTGTACCTCTAAG
AAGAAAGAATCTAACTGAAAATACCAAAAAAAAAAAAAAAAAA

103/392

FIGURE 98

MDDSTEREQSRLTSCCLKKREEMKLKECVSILPRKESPSVRSSKDGKLLAATLLLALLSCC
LTVVSFYQVAALQGD LASLRAELQGHHA EKL PAGAGAPKAGLEEAPAVTAGLKI FEPPAP
GEGNSSQNSRNKRAVQGPEETVTQDCLQLIADSETPTIQKGSYTFVPWLLSFKRGSAL EE
KENKILVKETGYFFIYGQVLYTDKTYAMGH LIQRKKVHVFGDELSLVTLFR CIQNMPETL
PNNSCYSAGIAKLEEGDELQLAIPRENAQISLDGDVTFFGALKLL

Transmembrane domain:

amino acids 47-72

N-glycosylation site:

amino acids 124-127, 242-245

cAMP- and cGMP-dependent protein kinase phosphorylation site:

amino acids 33-36, 173-176

N-myristoylation site:

amino acids 96-101

TNF family proteins:

amino acids 172-206

104/392

FIGURE 99

CCGAGGTTGGCGATCGCTGAGAGGCAGGAGGGCCGAGGCGGGCCTGGGAGGCGGCCGGAG
GTGGGGCGCCGCTGGGGCCGGCCCGCACGGGCTTCATCTGAGGGCGCACGGCCCGCGACC
GAGCGTGCGGACTGGCCTCCCAAGCGTGGGGCGACAAGCTGCCGGAGCTGCAATGGGGCCG
CGGCTGGGGATTCTTGTTTGGCCTCCTGGGCGCCGTGTGGCTGCTCAGCTCGGGCCACGG
AGAGGAGCAGCCCCCGGAGACAGCGGCACAGAGGTGCTTCTGCCAGGTTAGTGGTTACTT
GGATGATTGTACCTGTGATGTTGAAACCATTGATAGATTTAATAACTACAGGCTTTTCC
AAGACTACAAAACTTCTTGAAAGTGACTACTTTAGGTATTACAAGGTAAACCTGAAGAG
GCCGTGTCCTTTCTGGAATGACATCAGCCAGTGTGGAAGAAGGGACTGTGCTGTCAAACC
ATGTCAATCTGATGAAGTTCCTGATGGAATTAAATCTGCGAGCTACAAGTATTCTGAAGA
AGCCAATAATCTCATTGAAGAATGTGAACAAGCTGAACGACTTGGAGCAGTGGATGAATC
TCTGAGTGAGGAAACACAGAAGGCTGTTCTTCAGTGGACCAAGCATGATGATTCTTCAGA
TAACTTCTGTGAAGCTGATGACATTGAGTCCCCTGAAGCTGAATATGTAGATTTGCTTCT
TAATCCTGAGCGCTACACTGGTTACAAGGGACCAGATGCTTGGAAAATATGGAATGTGAT
CTACGAAGAAAACCTGTTTTAAGCCACAGACAATTAAAGACCTTTAAATCCTTTGGCTTC
TGGTCAAGGGACAAGTGAAGAGAACACTTTTTACAGTTGGCTAGAAGGTCTCTGTGTAGA
AAAAAGAGCATTCTACAGACTTATATCTGGCCTACATGCAAGCATTAAATGTGCATTTGAG
TGCAAGATATCTTTTACAAGAGACCTGGTTAGAAAAGAAATGGGGACACAACATTACAGA
ATTTCAACAGCGATTTGATGGAATTTTGACTGAAGGAGAAGGTCCAAGAAGGCTTAAGAA
CTTGTATTTTCTCTACTTAATAGAACTAAGGGCTTTATCCAAAGTGTTACCATTCTTCGA
GCGCCCGAGATTTTCAACTCTTTACTGGAATAAAATTCAGGATGAGGAAAACAAAATGTT
ACTTCTGGAATACTTTCATGAAATCAAGTCATTTCCCTTTGCATTTTGATGAGAATTCATT
TTTTGCTGGGGATAAAAAAGAAGCACACAACTAAAGGAGGACTTTGACTGCATTTTAG
AAATATTTCAAGAATTATGGATTGTGTTGGTTGTTTTAAATGTGCTCTGTGGGGAAAGCT
TCAGACTCAGGGTTTGGGCACTGCTCTGAAGATCTTATTTTCTGAGAAATTGATAGCAAA
TATGCCAGAAAGTGGACCTAGTTATGAATTCCATCTAACCAGACAAGAAATAGTATCATT
ATTCAACGCATTTGGAAGAATTTCTACAAGTGTGAAAGAATTAGAAAACTTCAGGAACCTT
GTTACAGAAATATTCATTAAAGAAAACAAGCTGATATGTGCCTGTTTCTGGACAATGGAGG
CGAAAGAGTGGAAATTTCAATCAAAGGCATAATAGCAATGACAGTCTTAAGCCAAACATTT
TATATAAAGTTGCTTTTGTAAAGGAGAATTTATATTGTTTTAAGTAAACACATTTTTAAAA
ATTGTGTTAAGTCTATGTATAATACTACTGTGAGTAAAGTAATACTTTAATAATGTGGT
ACAAATTTTTAAGTTTTAATATTGAATAAAAGGAGGATTATCAAATTAATAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAA

105/392

FIGURE 100

MGRGWGFLFGLLGAVWLLSSGHGEEQPPETAARCFQCQVSGYLDDCTCDVETIDRFNNYR
LFPRLOKLLESDFRYKYNLKRPCPFWNDISQCGRRDCAVKPCQSDEVPDGIKSASYKY
SEEANNLIEECEQAERLGAVDESLSEETQKAVLQWTKHDDSSDNFCEADDIQSPEAEYVD
LLNPERYTGYPDAWKIWNVIYEENCFKPQTIKRPLNPLASGQGTSEENTFYSWLEGL
CVEKRAFYRLISGLHASINVHLSARYLLQETWLEKKWGHNITEFQQRFDGILTEGEGPRR
LKNLYFLYLIELRALSKVLPFFERPDFQLFTGNKIQDEENKMLLLEILHEIKSFPLHFDE
NSFFAGDKKEAHKLKEDFRLHFRNISRIMDCVGCFCRLWGKLQTQGLGTALKILFSEKL
IANMPESGPSYEFHLTRQEIVSLFNAFGRISTSVKELENFRNLLQNIH

Important features:

Signal peptide:

amino acids 1-23

N-glycosylation site:

amino acids 280-283 and 384-387

Amidation site:

amino acids 94-97

Glycosaminoglycan attachment site:

amino acids 20-23 and 223-226

Aminotransferases class-V pyridoxal-phosphate:

amino acids 216-222

Interleukin-7 proteins:

amino acids 338-343

106/392

FIGURE 101

GCCTAGCCAGGCCAAGAATGCAATTGCCCCGGTGGTGGGAGCTGGGAGACCCCTGTGCTT
GGACGGGACAGGGTCGGGGGACACGCAGGATGAGCCCCGCGACCACTGGCACATTCTTGC
TGACAGTGACAGTATTTTCTCCAAGGTACACTCCGATCGGAATGTATACCCATCAGCAG
GTGTCCTCTTTGTTTCATGTTTTGGAAAGAGAATATTTTAAGGGGAATTTCCACCTTACC
CAAAACCTGGCGAGATTAGTAATGATCCCATAAACATTTAATACAAATTTAATGGGTTACC
CAGACCGACCTGGATGGCTTCGATATATCCAAAGGACACCATATAGTGATGGAGTCCTAT
ATGGGTCCCCAACAGCTGAAAATGTGGGGAAGCCAACAATCATTGAGATAACTGCCTACA
ACAGGCGCACCTTTGAGACTGCAAGGCATAATTTGATAATTAATATAATGTCTGCAGAAG
ACTTCCCGTTGCCATATCAAGCAGAATTCTTCATTAAGAATATGAATGTAGAAGAAATGT
TGGCCAGTGAGGTTCTTGGAGACTTTCTTGGCGCAGTGAAAAATGTGTGGCAGCCAGAGC
GCCTGAACGCCATAAACATCACATCGGCCCTAGACAGGGGTGGCAGGGTGCCACTTCCCA
TTAATGACCTGAAGGAGGGCGTTTATGTCATGGTTGGTGCAGATGTCCCGTTTTCTTCTT
GTTTACGAGAAGTTGAAAATCCACAGAATCAATTGAGATGTAGTCAAGAAATGGAGCCTG
TAATAACATGTGATAAAAAATTTCTGACTCAATTTTACATTGACTGGTGCAAAATTTTCAT
TGTTTGATAAAACAAAGCAAGTGTCCACCTATCAGGAAGTGATTCGTGGAGAGGGGATTT
TACCTGATGGTGGAGAATACAAACCCCTTCTGATTCTTTGAAAAGCAGAGACTATTACA
CGGATTTTCTAATTACACTGGCTGTGCCCTCGGCAGTGGCACTGGTCCTTTTTCTAATAC
TTGCTTATATCATGTGCTGCCGACGGGAAGGCGTGAAAAGAGAAACATGCAACACCAG
ACATCCAACCTGGTCCATCACAGTGCTATTTCAGAAATCTACCAAGGAGCTTCGAGACATGT
CCAAGAATAGAGAGATAGCATGGCCCCTGTCAACGCTTCCTGTGTTCCACCCTGTGACTG
GGGAAATCATACCTCCTTTACACACAGACAACTATGATAGCACAAACATGCCATTGATGC
AAACGCAGCAGAACTTGCCACATCAGACTCAGATTCCCCAACAGCAGACTACAGGTAAAT
GGTATCCCTGAAGAAAGAAAACCTGACTGAAGCAATGAATTTATAATCAGACAATATAGCA
GTTACATCACATTTCTTTCTCTTCCAATAATGCATGAGCTTTTCTGGCATATGTTATGC
ATGTTGGCAGTATTAAGTGTATACCAATAATACAACTAACTTTCAATTTTACTAATGTA
TTTTTTTGTAATAAGCATTTTGTACAATTTGTAAACATTGATGACTTTATATTGTT
ACAATAAAAGTTGATCTTTAAAATAAATATTATTAATGAAGCCTAAAAAAAAAAAA

107/392

FIGURE 102

MQLPRWWEELGDPCAWTGQGRGTRRMSPATTGTFLLTVYSIFSKVHSDRNVYPSAGVLFVH
VLEREYFKGEFPPYPKPGEISNDPITFNTNLMGYPDPRPGWLRYIQRTPYSDGVLYGSPTA
ENVGKPTIIIEITAYNRRTFETARHNLIINIMSAEDFPLPYQAEFFIKNMNVEEMLASEVL
GDFLGAVKNVWQPERLNAINITSALDRGGRVPLPINDLKEGVYVMVGADVFPSSCLREVE
NPQNQLRCSQEMEPVITCDKKFRTQFYIDWCKISLVDTKQVSTYQEVIRGEGILPDGGE
YKPPSDLKSRDYYTDFLITLAVPSAVALVFLILAYIMCCRREGVEKRNMQTPDIQLVH
HSAIQKSTKELRDMSKNREIAWPLSTLPVFHPVTGEIIPPLHTDNYDSTNMPLMQTQQNL
PHQTQIPQQOTTGKWYP

signal sequence:

Amino acids 1-46

transmembrane domain:

Amino acids 319-338

N-glycosylation site:

Amino acids 200-204

cAMP- and cGMP-dependent protein kinase phosphorylation site:

Amino acids 23-27

Tyrosine kinase phosphorylation site:

Amino acids 43-52

N-myristoylation sites:

Amino acids 17-23;112-118;116-122;185-191

108/392

FIGURE 103

CAGAAGAGGGGGCTAGCTAGCTGTCTCTGCGGACCAGGGAGACCCCCGCGCCCCCCCCGGT
GTGAGGCGGCCTCACAGGGCCGGGTGGGCTGGCGAGCCGACGCGGCGGCGGAGGAGGCTG
TGAGGAGTGTGTGGAACAGGACCCGGGACAGAGGAACCATGGGCTCCGCAGAACCTGAGCA
CCTTTTGCCTGTTGCTGCTATACCTCATCGGGGCGGTGATTGCCGGACGAGATTTCTATA
AGATCTTGGGGGTGCCTCGAAGTGCCTCTATAAAGGATATTAAAAAGGCCTATAGGAAAC
TAGCCCTGCAGCTTCATCCCGACCGGAACCCTGATGATCCACAAGCCCAGGAGAAATTCC
AGGATCTGGGTGCTGCTTATGAGGTTCTGTCTAGATAGTGAGAAACGGAAACAGTACGATA
CTTATGGTGAAGAAGGATTAAAAGATGGTCATCAGAGCTCCCATGGAGACATTTTTTTCAC
ACTTCTTTGGGGATTTTGGTTTCATGTTTGGAGGAACCCCTCGTCAGCAAGACAGAAATA
TTCCAAGAGGAAGTGATATTATTGTAGATCTAGAAGTCACTTTGGAAGAAGTATATGCA
GAAATTTTTGTGGAAGTAGTTAGAAACAAACCTGTGGCAAGGCAGGCTCCTGGCAACCGGA
AGTGCAATTGTTCGGCAAGAGATGCGGACCACCCAGCTGGGCCCCTGGGCGCTTCCAAATGA
CCCAGGAGGTGGTCTGCGACGAATGCCCTAATGTCAAACCTAGTGAATGAAGAACGAACGC
TGGAAGTAGAAATAGAGCCTGGGGTGAGAGACGGCATGGAGTACCCCTTTATTGGAGAAG
GTGAGCCTCACGTGGATGGGGAGCCTGGAGATTTACGGTTCCGAATCAAAGTTGTCAAGC
ACCCAATATTTGAAAGGAGAGGAGATGATTTGTACACAAATGTGACAATCTCATTAGTTG
AGTCACTGGTTGGCTTTGAGATGGATATTACTCACTTGGATGGTCACAAGGTACATATTT
CCCGGATAAGATCACCAGGCCAGGAGCGAAGCTATGGAAGAAAGGGGAAGGGCTCCCCA
ACTTTGACAACAACAATATCAAGGGCTCTTTGATAATCACTTTTGATGTGGATTTTCCAA
AAGAACAGTTAACAGAGGAAGCGAGAGAAGGTATCAAACAGCTACTGAAACAAGGGTCAG
TGCAGAAGGTATACAATGGACTGCAAGGATATTGAGAGTGAATAAAATTGGACTTTGTTT
AAAATAAGTGAATAAGCGATATTTATTATCTGCAAGGTTTTTTTTGTGTGTGTTTTTGT
TTATTTTCAATATGCAAGTTAGGCTTAATTTTTTTTATCTAATGATCATCATGAAATGAAT
AAGAGGGCTTAAGAATTTGTCCATTTGCATTTCGGAAAAGAATGACCAGCAAAAGGTTTAC
TAATACCTCTCCCTTTGGGGATTTAATGTCTGGTGCTGCCGCCTGAGTTTCAAGAATTAA
AGCTGCAAGAGGACTCCAGGAGCAAAAAGAAACACAATATAGAGGGTTGGAGTTGTTAGCA
ATTTTCATTCAAATGCCAACTGGAGAAGTCTGTTTTTAAATACATTTTGTGTTATTTTTTA

109/392

FIGURE 104

MAPQNLSTFCLLLLYLIGAVIAGRDFYKILGVPRSASIKDIKKAYRKLALQLHPDRNPDD
PQAQEKFQDLGAAYEVLSDSEKRKQYDITYGEEGLKDGHQSSHGDI FSHFFGDFGFMFGGT
PRQQDRNIPRGSDIIVDLEVTLEEYAGNFVEVVRNKPVARQAPGKRKCNCRQEMRTTQL
GPGRFQMTQEVVCDPCPNVKLVNEERTLEVEIEPGVRDGMETPFGEPEPHVDGEPGDLR
FRIKVVKHPIFERRGDDLYTNVTISLVESLVGFEMDITHLDGHKVVHISRDKITRPGAKLW
KKGEGLPNFDNNNIKGSLIITFDVDFPKEQLTEEAREGIKQLLKQGSVQKVYNGLQGY

Important features:

Signal peptide:

amino acids 1-22

Cell attachment sequence:

amino acids 254-257

Nt-dnaJ domain signature:

amino acids 67-87

Homologous region to Nt-dnaJ domain proteins:

amino acids 26-58

N-glycosylation site:

amino acids 5-9, 261-265

Tyrosine kinase phosphorylation site:

amino acids 253-260

N-myristoylation site:

amino acids 18-24, 31-37, 93-99, 215-221

Amidation site:

amino acids 164-168

110/392

FIGURE 105

GGCACGAGGCGGCGGGGCAGTCGCGGGGATGCGCCCCGGGAGCCACAGCCTGAGGCCCTCAG
GTCTCTGCAGGTGTCTGTGGAGGAACCTAGCACCTGCCATCCTCTTCCCCAATTTGCCACT
TCCAGCAGCTTTAGCCCATGAGGAGGATGTGACCGGGACTGAGTCAGGAGCCCTCTGGAA
GCATGGAGACTGTGGTGATTGTTGCCATAGGTGTGCTGGCCACCATCTTTCTGGCTTCGT
TTGCAGCCTTGGTGCTGGTFTTGACGGCAGCGCTACTGCCGCGCGGAGACCTGCTGCAGC
GCTATGATTCTAAGCCCATTTGTGGACCTCATTGGTGCCATGGAGACCCAGTCTGAGCCCT
CTGAGTTAGAACTGGACGATGTCGTTATCACCAACCCCCACATTGAGGCCATTCTGGAGA
ATGAAGACTGGATCGAAGATGCCTCGGGTCTCATGTCCCACTGCATTGCCATCTTGAAGA
TTTGTCACTCTGACAGAGAAGCTTGTTGCCATGACAATGGGCTCTGGGGCCAAGATGA
AGACTTCAGCCAGTGTGACGACATCATTGTGGTGGCCAAGCGGATCAGCCCCAGGGTGG
ATGATGTTGTGAAGTCGATGTACCCTCCGTTGGACCCCAAACCTCCTGGACGCACGGACGA
CTGCCCTGCTCCTGTCTGTGTCAGTCACCTGGTGCTGGTGACAAGGAATGCCTGCCATCTGA
CGGGAGGCCTGGACTGGATTGACCAGTCTCTGTCGGCTGCTGAGGAGCATTTGGAAGTCC
TTCGAGAAGCAGCCCTAGCTTCTGAGCCAGATAAAGGCCTCCAGGCCCTGAAGGCTTCC
TGCAGGAGCAGTCTGCAATTTAGTGCCTACAGGCCAGCAGCTAGCCATGAAGGCCCTGC
CGCCATCCCTGGATGGCTCAGCTTAGCCTTCTACTTTTTCTATAGAGTTAGTTGTTCTC
CACGGCTGGAGAGTTTCACTGTGTGTGCATAGTAAAGCAGGAGATCCCCGTCACTTTATG
CCTCTTTTGCAGTTGCAAACTGTGGCTGGTGAGTGGCAGTCTAATACTACAGTTAGGGGA
GATGCCATTCACTCTCTGCAAGAGGAGTATTGAAAACCTGGTGGACTGTCAGCTTTATTTA
GCTCACCTAGTGTTTTCAAGAAAATTGAGCCACCGTCTAAGAAATCAAGAGGTTTCACAT
TAAAATTAGAATTTCTGGCCTCTCTCGATCGGTGAGAATGTGTGGCAATTCTGATCTGCA
TTTTTCAGAAGAGGACAATCAATTGAACTAAGTAGGGGTTTCTTCTTTTGGCAAGACTTG
TACTCTCTCACCTGGCCTGTTTCATTTATTTGTATTATCTGCCTGGTCCCTGAGGCGTCT
GGGTCTCTCCTCTCCCTTGCAAGTTTGGGTTTGAAGCTGAGGAACTACAAAGTTGATGAT
TTCTTTTTTATCTTTATGCCTGCAATTTTACCTAGCTACCACTAGGTGGATAGTAAATTT
ATACTTATGTTTCCCTCAAAAAAAAAAAAAA

111/392

FIGURE 106

METVVIVAIGVLATIFLASFAALVLVCRQRYCRPRDLLQRYDSKPIVDLIGAMETQSEPS
ELELDDVVITNPHIEAILENEDWIEDASGLMSHCIAILKICHTLTEKLVAMTMGSGAKMK
TSASVSDIIVVAKRISPRVDDVVKSMYPPLDPKLLDARTTALLSVSHLVLVTRNACHLT
GGLDWIDQSLSAABEHLEVLREAALASEPDKGLPGPEGFLQEQSAI

112/392

FIGURE 107

GCTTCATTTCTCCCGACTCAGCTTCCACCCCTGGGCTTTCCGAGGTGCTTTCGCCGCTGT
CCCCACCACTGCAGCCATGATCTCCTTAACGGACACGCAGAAAATTGGAATGGGATTAAC
AGGATTTGGAGTGTTTTTCCTGTTCTTTGGAATGATTCTCTTTTTTGACAAAGCACTACT
GGCTATTGGAAATGTTTTATTTGTAGCCGGCTTGGCTTTTGTAAATTGGTTTAGAAAGAAC
ATTCAGATTCTTCTTCCAAAAACATAAAATGAAAGCTACAGGTTTTTTCTGGGTGGTGT
ATTTGTAGTCCTTATTGGTTGGCCTTTGATAGGCATGATCTTCGAAATTTATGGATTTTT
TCTCTTGTTCAGGGGCTTCTTTCCTGTCGTTGTTGGCTTTATTAGAAGAGTGCCAGTCCT
TGGATCCCTCCTAAATTTACCTGGAATTAGATCATTGTAGATAAAGTTGGAGAAAGCAA
CAATATGGTATAAACAACAAGTGAATTTGAAGACTCATTTAAAATATTGTGTTATTTATAA
AGTCATTTGAAGAATATTCAGCACAAAATTAAATTACATGAAATAGCTTGTAATGTTCTT
TACAGGAGTTTAAAACGTATAGCCTACAAAGTACCAGCAGCAAATTAGCAAAGAAGCAGT
GAAAACAGGCTTCTACTCAAGTGAACCTAAGAAGAAGTCAGCAAGCAAACCTGAGAGAGGTG
AAATCCATGTTAATGATGCTTAAGAAACTCTTGAAGGCTATTGTGTTGTTTTCCACAA
TGTGCGAAACTCAGCCATCCTTAGAGAACTGTGGTGCCTGTTCTTTCTTTTTATTTTG
AAGGCTCAGGAGCATCCATAGGCATTTGCTTTTTTAGAAGTGTCCTGCAATGGCAAAAA
TATTTCCAGTTGCACTGTATCTCTGGAAGTGATGCATGAATTCGATTGGATTGTGTCATT
TTAAAGTATTAAAACCAAGGAAACCCCAATTTTGATGTATGGATTACTTTTTTTTGNGCN
CAGGGCC

113/392

FIGURE 108

MISLTDTQKIGMGLTGFGVFFLFFGMILFFDKALLAIGNVLFVAGLAFVIGLERTFRFFF
QKHKMKATGFFLGGVFVVLIGWPLIGMIFEIYGFFLLFRGFFPVVVGFIIRVPVLGSLN
LPGIRSFVDKVGESNNMV

Important features:

Transmembrane domains:

amino acids 12-30 (typeII), 33-52, 69-89 and 93-109

N-myristoylation sites:

amino acids 11-16, 51-56 and 116-121

Aminoacyl-transfer RNA synthetases class-II protein:

amino acids 49-59

114/392

FIGURE 109

CCAGTCTGTCGCCACCTCACTTGGTGTCTGCTGTCCCCGCCAGGCAAGCCTGGGGTGAGA
GCACAGAGGAGTGGGCGGGGACCATGCGGGGGACGCGGCTGGCGCTCCTGGCGCTGGTGC
TGGCTGCCTGCGGAGAGCTGGCGCGGGCCCTGCGCTGCTACGTCTGTCCGGAGCCCACAG
GAGTGTGCGACTGTGTCACCATCGCCACCTGCACCACCAACGAAACCATGTGCAAGACCA
CACTCTACTCCCGGGAGATAGTGTACCCCTTCCAGGGGGACTCCACGGTGACCAAGTCCT
GTGCCAGCAAGTGTAAGCCCTCGGATGTGGATGGCATCGGCCAGACCCTGCCCCGTGTCCT
GCTGCAATACTGAGCTGTGCAATGTAGACGGGGCGCCCGCTCTGAACAGCCTCCACTGCG
GGGCCCTCACGCTCCTCCCACTCTTGAGCCTCCGACTGTAGAGTCCCCGCCCACCCCAT
GGCCCTATGCGGCCAGCCCCGAATGCCTTGAAGAAGTGCCCCCTGCACCAGGAAAAAA
AAAAA

115/392

FIGURE 110

MRGTRLALLALVLAACGELAPALRCYVCPEPTGVSDCVTIATCTTNETMCKTTLYSREIV
YPFQGDSTVTKSCASKCKPSDVGIGQTLFVSCCNTELCNVDGAPALNSLHCGALTLLPL
LSLRL

Important features:

Signal peptide:

amino acids 1-17

N-glycosylation site:

amino acids 46-49

116/392

FIGURE 111

GCGCCGCCAGGCGTAGGCGGGGTGGCCCTTGCGTCTCCCGCTTCCTTGAAAAACCCGGCG
GGCGAGCGAGGCTGCGGGCCGGCCGCTGCCCTTCCCCACACTCCCCGCCGAGAAGCCTCG
CTCGGCGCCCAACATGGCGGGTGGGCGCTGCGGCCCCGAGCTAACGGCGCTCCTGGCCGC
CTGGATCGCGGCTGTGGCGGCGACGGCAGGCCCCCGAGGAGGCCGCGCTGCCGCCGGAGCA
GAGCCGGGTCCAGCCCATGACCGCCTCCAACCTGGACGCTGGTGATGGAGGGCGAGTGGAT
GCTGAAATTTTACGCCCCATGGTGTCCATCCTGCCAGCAGACTGATTCAGAATGGGAGGC
TTTTGCAAAGAATGGTGAAATACTTCAGATCAGTGTGGGAAGGTAGATGTCATTCAAGA
ACCAGGTTTGAGTGGCCGCTTCTTTGTCAACACTCTCCAGCATTTTTTCATGCAAAGGA
TGGGATATTCGCGCGTTATCGTGGCCCAGGAATCTTCGAAGACCTGCAGAATTATATCTT
AGAGAAGAAATGGCAATCAGTCGAGCCTCTGACTGGCTGGAAATCCCCAGCTTCTCTAAC
GATGTCTGGAATGGCTGGTCTTTTAGCATCTCTGGCAAGATATGGCATCTTCACAACTA
TTTCACAGTGACTCTTGGAATTCCTGCTTGGTGTTCTTATGTGTTTTTCGTCATAGCCAC
CTTGGTTTTTGGCCTTTTTATGGGTCTGGTCTTGGTGTTAATATCAGAATGTTTTCTATGT
GCCACTTCCAAGGCATTTATCTGAGCGTTCTGAGCAGAATCGGAGATCAGAGGAGGCTCA
TAGAGCTGAACAGTTGCAGGATGCGGAGGAGGAAAAAGATGATTCAAATGAAGAAGAAAA
CAAAGACAGCCTTGTAGATGATGAAGAAGAGAAAGAAAGATCTTGGCGATGAGGATGAAGC
AGAGGAAGAAGAGGAGGAGGACAACCTTGGCTGCTGGTGTGGATGAGGAGAGAAGTGAGGC
CAATGATCAGGGGCCCCCAGGAGAGGACGGTGTGACCCGGGAGGAAGTAGAGCCTGAGGA
GGCTGAAGAAGGCATCTCTGAGCAACCCTGCCAGCTGACACAGAGGTGGTGGAAAGACTC
CTTGAGGCAGCGTAAAGTCAGCATGCTGACAAGGGACTGTAGATTTAATGATGCGTTTT
CAAGAATACACACCAAACAATATGTCAGCTTCCCTTTGGCCTGCAGTTTGTACCAATC
CTTAATTTTTCTGAATGAGCAAGCTTCTCTTAAAGATGCTCTCTAGTCATTTGGTCTC
ATGGCAGTAAGCCTCATGTATACTAAGGAGAGTCTTCCAGGTGTGACAATCAGGATATAG
AAAAACAAACGTAGTGTTGGGATCTGTTTGGGAGACTGGGATGGGAACAAGTTCATTTACT
TAGGGGTCAGAGAGTCTCGACCAGAGGAGGCCATTCCCAGTCCTAATCAGCACCTTCCAG
AGACAAGGCTGCAGGCCCTGTGAAATGAAAGCCAAGCAGGAGCCTTGGCTCCTGAGCATC
CCCAAAGTGTAACGTAGAAGCCTTGCACTCTTTTTCTTGTGTAAAGTATTTATTTTTGTCA
AATTGCAGGAAACATCAGGCACCACAGTGCATGAAAAATCTTTCACAGCTAGAAATTGAA
AGGGCCTTGGGTATAGAGAGCAGCTCAGAAGTCATCCAGCCCTCTGAATCTCCTGTGCT
ATGTTTTATTTCTTACCTTTAATTTTTCCAGCATTTCCACCATGGGCATTCAGGCTCTCC
ACACTCTTCACTATTATCTCTTGGTCAGAGGACTCCAATAACAGCCAGGTTTACATGAAC
TGTGTTTGTTCATTCTGACCTAAGGGGTTTAGATAATCAGTAACCATAACCCCTGAAGCT
GTGACTGCCAAACATCTCAAATGAAATGTTGTGGCCATCAGAGACTCAAAGGAAGTAAG
GATTTTACAAGACAGATTAAAAAAAATGTTTTGTCCAAATATAGTTGTTGTTGATTT
TTTTTTAAGTTTTCTAAGCAATATTTTTCAAGCCAGAAGTCCTCTAAGTCTTGCCAGTAC
AAGGTAGTCTTGTGAAGAAAAGTTGAATACTGTTTTGTTTTCATCTCAAGGGTTCCCTG
GGTCTTGAACCTACTTTAATAATAACTAAAAAACCACTTCTGATTTTCCTTCAGTGATGTG
CTTTTGGTGAAAGAAATTAATGAACTCCAGTACCTGAAAGTGAAAGATTTGATTTTGTTC
CATCTTCTGTAATCTTCCAAGAATTATATCTTTGTAAATCTCTCAATACTCAATCTACT
GTAAGTACCCAGGGAGGCTAATTTCTTT

117/392

FIGURE 112

MAGGRCGPQLTALLAAWIAAVAATAGPEEAALPPEQSRVQPMTASNWTLVMEGEWMLKFY
APWCPSCQOTDSEWEAFKNGEILQISVGKVDVIQEPGLSGRFFVTTLPAFFHAKDGIFR
RYRGP G IFEDLQNYILEKKWQSVEPLTGWKSPASLTMSGMAGLFSISGKIWHLHNYFTVT
LGIPAWCSYVFFVIATLVFGLFMGLVLVVISECFYVPLPRHLSERSEQNRRSEEAHRAEQ
LQDAEEEEKDDSNEEENKDSLVDDEEEKEDLGDEDEAEEDNLAAGVDEERSEANDQG
PPGEDGVTRREEVEPEEAEGISEQPCPADTEVVEDSLRQRKSQHADKGL

Important features:

Signal peptide:

amino acids 1-22

Transmembrane domain:

amino acids 191-211

N-glycosylation site:

amino acids 46-49

Thioredoxin family proteins: (homologous region to disulfide isomerase)

amino acids 56-72

Flavodoxin proteins:

amino acids 173-187

118/392

FIGURE 113

GAGGAACCTACCGGTACCGGCCGCGCGCTGGTAGTCGCCGGTGTGGCTGCACCTCACCAA
TCCCGTGCGCCGCGGCTGGGCCGTGCGAGAGTGCCTGTGCTTCTCTCCTGCACGCGGTGC
TTGGGCTCGGCCAGGCGGGGTCCGCCGCCAGGGTTTGAGGATGGGGGAGTAGCTACAGGA
AGCGACCCCGCGATGGCAAGGTATATTTTTGTGGAATGAAAAGGAAGTATTAGAAATGAG
CTGAAGACCATTACAGATTAATATTTTTGGGGACAGATTTGTGATGCTTGATTACCCCT
TGAAGTAATGTAGACAGAAGTTCTCAAATTTGCATATTACATCAACTGGAACCAGCAGTG
AATCTTAATGTTCACTTAAATCAGAACTTGCATAAGAAAAGAGAATGGGAGTCTGGTTAAA
TAAAGATGACTATATCAGAGACTTGAAAAGGATCATTCTCTGTTTTCTGATAGTGTATAT
GGCCATTTTAGTGGGCACAGATCAGGATTTTACAGTTTACTTGGAGTGTCCAAAACCTGC
AAGCAGTAGAGAAATAAGACAAGCTTTCAAGAAATTGGCATTGAAGTTACATCCTGATAA
AAACCCGAATAACCCAAATGCACATGGCGATTTTTTAAAAATAAATAGAGCATATGAAGT
ACTCAAAGATGAAGATCTACGGAAAAAGTATGACAAATATGGAGAAAAGGGACTTGAGGA
TAATCAAGGTGGCCAGTATGAAAGCTGGAACATTATCGTTATGATTTTGGTATTTATGA
TGATGATCCTGAAATCATAACATTGGAAGAAGAGAATTTGATGCTGCTGTTAATTCTGG
AGAACTGTGGTTTGTAAATTTTTACTCCCCAGGCTGTTCCACTGCCATGATTTAGCTCC
CACATGGAGAGACTTTGCTAAAGAAGTGGATGGGTACTTCGAATTGGAGCTGTTAACTG
TGGTGATGATAGAATGCTTTGCCGAATGAAAGGAGTCAACAGCTATCCCAGTCTCTTCAT
TTTTCGGTCTGGAATGGCCCCAGTGAAATATCATGGAGACAGATCAAAGGAGAGTTTAGT
GAGTTTGCATGCAGCATGTTAGAAGTACAGTGACAGAACTTTGGACAGGAAATTTTGT
CAACTCCATACAACTGCTTTTGCTGCTGGTATTGGCTGGCTGATCACTTTTGTTCAAA
AGGAGGAGATTGTTGACTTCACAGACAGCACTCAGGCTTAGTGGCATGTTGTTTCTCAA
CTCTTGGATGCTAAAGAAATATATTGGGAAGTAATACATAATCTTCCAGATTTTGAAC
ACTTTTCGGCAACACACTAGAGGATCGTTTGGCTCATCATCGGTGGCTGTTATTTTTTCA
TTTTTGAAAAAATGAAAATTCAAATGATCCTGAGCTGAAAAAACTAAAACTCTACTTAA
AAATGATCATATTCAAGTTGGCAGGTTTGACTGTTTCTCTGCACCAGACATCTGTAGTAA
TCTGTATGTTTTTTCAGCCGTCTCTAGCAGTATTTAAAGGACAAGGAACCAAAGAATATGA
AATTCATCATGGAAGAAGATTCTATATGATATACTTGCCTTTGCCAAAGAAGTGTGAA
TTCTCATGTTACCACGCTTGGACCTCAAATTTTCTGCCAATGACAAAGAACCATGGCT
TGTTGATTTCTTTGCCCCCTGGTGTCCACCATGTGAGCTTTACTACCAGAGTTACGAAG
AGCATCAAATCTTCTTATGGTCAGCTTAAGTTTGGTACACTAGATTGTACAGTTTCATGA
GGGACTCTGTAACATGTATAACATTAGGCTTATCCAACAACAGTGGTATTCAACCAGTC
CAACATTCATGAGTATGAAGGACATCACTCTGCTGAACAAATCTTGGAGTTCATAGAGGA
TCTTATGAATCCTTCAGTGGTCTCCCTTACACCCACCACCTTCAACGAACTAGTTACACA
AAGAAAACACAACGAAGTCTGGATGGTTGATTTCTATTCTCCGTGGTGTCTCCTTGCCA
AGTCTTAATGCCAGAATGGAAAAGAATGGCCCGGACATTAACCTGGACTGATCAACGTGGG
CAGTATAGATTGCCAACAGTATCATTCTTTTTGTGCCCAGGAAAACGTTCAAAGATACCC
TGAGATAAGATTTTTTCCCCCAAATCAAATAAAGCTTATCAGTATCACAGTTACAATGG
TTGGAATAGGGATGCTTATTCCTGAGAATCTGGGGTCTAGGATTTTTACCTCAAGTATC
CACAGATCTAACACCTCAGACTTTAGTGAAAAAGTTCTACAAGGGAAAAATCATTGGGT
GATTGATTTCTATGCTCCTTGGTGTGGACCTTGCCAGAATTTTGCTCCAGAATTTGAGCT
CTTGGCTAGGATGATTAAAGGAAAAGTGAAAGCTGGAAAAGTAGACTGTGAGGCTTATGC
TCAGACATGCCAGAAAGCTGGGATCAGGGCCTATCCAAGTGTAAAGTTTATTTCTACGA
AAGAGCAAAGAGAAATTTTCAAGAAGAGCAGATAAATAACCAGAGATGCAAAAGCAATCGC
TGCTTAAATAAGTGAAAAATTTGGAACTCTCCGAAATCAAGGCAAGAGGAATAAGGATGA
ACTTTGATAATGTTGAAGATGAAGAAAAAGTTTAAAGAAATTTCTGACAGATGACATCAG
AAGACACCTATTTAGAATGTTACATTTATGATGGGAATGAATGAACATTATCTTAGACTT

119/392

GCAGTTGTACTGCCAGAATTATCTACAGCACTGGTGTAAAAGAAGGGTCTGCAAACCTTTT
TCTGTAAAGGGCCGGTTTATAAATATTTTAGACTTTGCAGGCTATAATATATGGTTCACA
CATGAGAACAAGAATAGAGTCATCATGTATTCCTTGTTATTTGCTTTTAAACAACCTTTAA
AAAATATTTAAACGATTCTTAGCTCAGAGCCATACAAAAGTAGGCTGGATTCCAGTCCATG
GACCATAGATTGCTGTCCCCCTCGACGGACTTATAATGTTTCAGGTGGCTGGCTTGAACA
TGAGTCTGCTGTGCTATCTACATAAATGTCTAAGTTGTATAAAGTCCACTTCCCTTCAC
GTTTTTTGGCTGACCTGAAAAGAGGTAACCTTAGTTTTTTGGTCACTTGTTCTCCTAAAAAT
GCTATCCCTAACCATATATTTATATTTTCGTTTTAAAAACACCCATGATGTGGCACAGTAA
ACAAACCCTGTTATGCTGTATTATTATGAGGAGATTCTTCATTGTTTTCTTTCCTTCTCA
AAGGTTGAAAAAATGCTTTTAATTTTTTACAGCCGAGAAACAGTGCAGCAGTATATGTGC
ACACAGTAAGTACACAAATTTGAGCAACAGTAAGTGCACAAATTCGTAGTTTGCTGTAT
CATCCAGGAAAACCTGAGGGAAAAAATTATAGCAATTAAGTGGGCATTGTAGAGTATCC
TAAATATGTTATCAAGTATTTAGAGTTCTATATTTTAAAGATATATGTGTTTCATGTATTT
TCTGAAATTGCTTTCATAGAAATTTTCCCACTGATAGTTGATTTTTGAGGCATCTAATAT
TTACATATTTGCCTTCTGAACCTTGTTTTGACCTGTATCCTTTATTTACATTGGGTTTTT
CTTTCATAGTTTTGGTTTTTCACTCCTGTCCAGTCTATTTATTATTCAAATAGGAAAAAT
TACTTTACAGGTTGTTTTACTGTAGCTTATAATGATACTGTAGTTATTCCAGTTACTAGT
TTACTGTCAGAGGGCTGCCTTTTTTCAGATAAATATTGACATAATAACTGAAGTTATTTTT
ATAAGAAAATCAAGTATATAAATCTAGGAAAGGGATCTTCTAGTTTCTGTGTTGTTTAGA
CTCAAAGAATCACAAATTTGTCAGTAACATGTAGTTGTTTAGTTATAAATTCAGAGTGTAC
AGAATGGTAAAAATTCCAATCAGTCAAAGAGGTCAATGAATTAAAGGCTTGCAACTTT
TTCAAAAAAAAAAAAAAAAAA

120/392

FIGURE 114

MGVWLNKDDYIRD LKRIILCFLIVYMAILVGT DQDFYSLLGVSKTASSREIRQAFKKLAL
KLHPDKNPNNPNAHGDFLKINRAYEVLKDEDLRKKYDKYGEKGL EDNQGGQYESWNYRY
DFGIYDDDPEIITLERREFDAAVNSGELW FVNFYSPGCSHCHDLAPTWRDFAKEVDGLLR
IGAVNCGDDRMLCRMKG VNSYPSLFIFRSGMAPVKYHGDRSKESLVSFAMQHVRSTVTEL
WTGNFVNSIQTAFAAGIGWLITFC SKGGDCLTSQTRLRLSGMLFLNSLDAKEIYLEVIHN
LPDFELLSANTLEDRLAHRWLLFFHFGKNENSNDPELKKLKTLLKNDHIQVGRFDCSSA
PDICSNLYVFQPSLAVFKGQGTKEYEIH HGKKILYDILAFAKESVNSHVTTLGPNFPAN
DKEPWLVDFFAPWCPPCRALLPELRRASNLLYGQLKFGTLDCTVHEGLCNMYNIQAYPTT
VVFNQSNIEHEYEGHSAEQILEFIEDLMNPSVVS LPTTTFNELVTQRKHNEVWMVDFYSP
WCHPCQVLMPEWKRMARTLTGLINVSIDCQQYHSFCAQENVQRYPEIRFFPPKSNKAYQ
YHSYNGWNRDAYSLRIWGLGFLPQVSTD LTPQTFSEKVLQGKNHWVIDFYAPWCGPCQNF
APEFELLARMIKGKV KAGKVD CQAYAQTCQKAGIRAYPTVKFYFYERAKRNFQEEQINTR
DAKAIAALISEKLETLRNQGKR NKDEL

Important features:

Endoplasmic reticulum targeting sequence:

amino acids 744-747

Cytochrome c family heme-binding site signature:

amino acids 158-163

Nt-dnaJ domain signature:

amino acids 77-96

N-glycosylation site:

amino acids 484-487

121/392

FIGURE 115

GCGGGCTGTTGACGGCGCTGCGATGGCTGCCTGCGAGGGCAGGAGAAGCGGAGCTCTCGG
TTCCTCTCAGTCGGACTTCCTGACGCCGCCAGTGGGCGGGGCCCCCTTGGGCGCTCGCCAC
CACTGTAGTCATGTACCCACCGCCGCCGCCGCCCTCATCGGGACTTCATCTCGGTGAC
GCTGAGCTTTGGCGAGAGCTATGACAACAGCAAGAGTTGGCGGCGGCGCTCGTGCTGGAG
GAAATGGAAGCAACTGTCGAGATTGCAGCGGAATATGATTCTCTTCTCCTTGCCTTTCT
GCTTTTCTGTGGACTCCTCTTCTACATCAACTTGGCTGACCATTTGGAAGCTCTGGCTTT
CAGGCTAGAGGAAGAGCAGAAGATGAGGCCAGAAATTGCTGGGTAAACCAGCAAATCC
ACCCGCTCTTACCAGCTCCTCAGAAGGCGGACCCGACCCTGAGAACTTACCTGAGATTTT
GTCACAGAAGACACAAAGACACATCCAGCGGGGACCACCTCACCTGCAGATTAGACCCCC
AAGCCAAGACCTGAAGGATGGGACCCAGGAGGAGGCCACAAAAGGCAAGAAGCCCCCTGT
GGATCCCCCGCCGGAAGGAGATCCGCAGAGGACAGTCATCAGCTGGAGGGGAGCGGTGAT
CGAGCCTGAGCAGGGCACCGAGCTCCCTTCAAGAAGAGCAGAAGTGCCACCAAGCCTCC
CCTGCCACCGGCCAGGACACAGGGCACACCAGTGTCATCTGAACTATCGCCAGAAGGGCGT
GATTGACGTCTTCTGTCATGCATGGAAGGATACCGCAAGTTTGCATGGGGCCATGACGA
GCTGAAGCCTGTGTCCAGGTCCTTCAGTGAGTGGTTTTGGCCTCGGTCTCACACTGATCGA
CGCGCTGGACACCATGTGGATCTTGGGTCTGAGGAAAGAATTTGAGGAAGCCAGGAAGTG
GGTGTGAAGAAGTTACACTTTGAAAAGGACGTGGACGTCAACCTGTTTGAGAGCACGAT
CCGCATCCTGGGGGGGCTCCTGAGTGCCTACCACCTGTCTGGGGACAGCCTCTTCTGAG
GAAAGCTGAGGATTTTGGAAATCGGCTAATGCCTGCCTTCAGAACACCATCCAAGATTCC
TACTCGGATGTGAACATCGGTACTGGAGTTGCCACCCGCCACGGTGGACCTCCGACAG
CACTGTGGCCGAGGTGACCAGCATTGAGCTGGAGTTCCGGGAGCTCTCCCGTCTCACAGG
GGATAAGAAGTTTCAGGAGGCAGTGGAGAAGGTGACACAGCACATCCACGGCCTGTCTGG
GAAGAAGGATGGGCTGGTGCCCATGTTTCATCAATACCCACAGTGGCCTCTTACCCACCT
GGGCGTATTACGCTGGGCGCCAGGGCCGACAGCTACTATGAGTACCTGCTGAAGCAGTG
GATCCAGGGCGGGAAGCAGGAGACACAGCTGCTGGAAGACTACGTGGAAGCCATCGAGGG
TGTCAGAAGCACCTGCTGCGGCACTCCGAGCCCAGTAAGCTCACCTTTGTGGGGGAGCT
TGCCCCAGGCGGCTTCAGTGCCAAGATGGACCACCTGGTGTGCTTCTGCCAGGGACGCT
GGCTCTGGGCGTCTACCACGGCCTGCCCGCCAGCCACATGGAGCTGGCCCAGGAGCTCAT
GGAGACTTGTACAGATGAACCGGCAGATGGAGACGGGGCTGAGTCCCGAGATCGTGCA
CTTCAACCTTTACCCCCAGCCGGGCCGTCGGGACGTGGAGGTCAAGCCAGCAGACAGGCA
CAACCTGCTGCGGCCAGAGACCGTGGAGAGCCTGTTCTACCTGTACCGCTCACAGGGGA
CCGCAAATACCAGGACTGGGGCTGGGAGATTCTGCAGAGCTTCAGCCGATTACACGGGT
CCCCCTCGGGTGGCTATTCTTCCATCAACAATGTCCAGGATCCTCAGAAGCCCCAGCCTAG
GGACAAGATGGAGAGCTTCTTCTGGGGGAGACGCTCAAGTATCTGTTCTTGCTCTTCTC
CGATGACCCAAACCTGCTCAGCCTGGACGCCTACGTGTTCAACACCGAAGCCACCTCT
GCCTATCTGGACCCCTGCCTAGGGTGGATGGCTGCTGGTGTGGGGACTTCGGGTGGGCAG
AGGCACCTTGCTGGGTCTGTGGCATTTCCTCAAGGGGCCACGTAGCACCGGCAACCGCCAA
GTGGCCCAGGCTCTGAACTGGCTCTGGGCTCCTCCTCGTCTCTGCTTTAATCAGGACACC
GTGAGGACAAGTGAGGCCGTGAGTCTTGGTGTGATGCGGGGTGGGCTGGGCCGCTGGAGC
CTCCGCTGCTTCTCCTCAGAAGACACGAATCATGACTCACGATTGCTGAAGCCTGAGCAG
GTCTCTGTGGGCCGACCAGAGGGGGGCTTCGAGGTGGTCCCTGGTACTGGGGTGACCGAG
TGGACAGCCCAGGGTGACGCTCTGCCCCGGGCTCGTGAAGCCTCAGATGTCCCAATCCAA
GGGTCTGGAGGGGCTGCCGTGACTCCAGAGGCCTGAGGCTCCAGGGCTGGCTCTGGTGT
TACAAGCTGGACTCAGGGATCCTCCTGGCCGCCCGCAGGGGGCTTGGAGGGCTGGACGG
CAAGTCCGTCTAGCTCACGGGCCCCCTCCAGTGGAATGGGTCTTTTCGGTGGAGATAAAAG
TTGATTTGCTCTAACCGCAA

122/392

FIGURE 116

MAACEGRRSGALGSSQSDFLTPPVGGAPWAVATTVMYPPPPPPPHRDFISVTLSFGESY
DNSKSWRRRSCWRKWKQLSRLQRMILFLLAFLLFCGLLFYINLADHWKALAFRLEEEQK
MRPEIAGLKPANPPVLPAPQKADTDPENLPEISSQKTQRHIQRGPPHLQIRPPSQDLKDG
TQEEATKRQEAPVDPRPEGDPQRTVISWRGAVIEPEQGTELPSSRAEVPTKPPLPPARTQ
GTPVHLNRYQKGVIDVFLHAWKGYRKFAWGHDELKPVSRSEWFGGLGLTLIDALDTMWI
LGLRKEFEEARKWVSKLHFEKDVDVNLFEISTIRILGGLLSAYHLSGDSLFLRKAEDFGN
RLMPAFRTPSKIPYSDVNIGTGVAHPPRWTSdstVAEVTsiQLEFRELSRLTGDKKFQEA
VEKVTQHIHGLSGKKDGLVPMFINTHSGLFTHLGVFTLGARADSYEYLLKQWIQGGKQE
TQLLEDYVEAIEGVRTHLLRHSEPSKLTfVGELAHGRFSakMDHLVCFLPGTLALGVYHG
LPASHMELAQELMETCYQMNRMETGLSPEIVHFNLYPQGRRDVEVKPADRHNLRLRPET
VESLFYLYRVTGDRKYQDWGWEILQSFSRfTRVPSGGYSSINNVDpQKPEPRDKMESFF
LGETLKYLFLLFSDDPNLLSLDAYVFNTeAHPLPIWTPA

Important features of the protein:

Transmembrane domain:

amino acids 21-40 and 84-105 (type II)

123/392

FIGURE 117

GTGGGATTTATTTGAGTGCAAGATCGTTTTCTCAGTGGTGGTGGGAAGTTGCCTCATCGCA
GGCAGATGTTGGGGCTTTGTCCGAACAGCTCCCCCTCTGCCAGCTTCTGTAGATAAGGGTT
AAAACTAATATTTATATGACAGAAGAAAAAGATGTCATTCCGTAAAGTAAACATCATCA
TCTTGGTCCTGGCTGTTGCTCTCTTCTTACTGGTTTTGTCACCATAACTTCCTCAGCTTGA
GCAGTTTGTTAAGGAATGAGGTTACAGATTCAGGAATTGTAGGGCCTCAACCTATAGACT
TTGTCCCAAATGCTCTCCGACATGCAGTAGATGGGAGACAAGAGGAGATTCTGTGGTCA
TCGCTGCATCTGAAGACAGGCTTGGGGGGGCCATTGCAGCTATAAACAGCATTTCAGCACA
ACACTCGCTCCAATGTGATTTTCTACATTGTTACTCTCAACAATACAGCAGACCATCTCC
GGTCTGGCTCAACAGTGATTCCCTGAAAAGCATCAGATACAAAATTGTCAATTTTGACC
CTAACTTTTGAAGGAAAAGTAAAGGAGGATCCTGACCAGGGGAATCCATGAAACCTT
TAACCTTTGCAAGGTTCTACTTGCCAATTCTGGTTCCCAGCGCAAAGAAGGCCATATACA
TGGATGATGATGTAATTGTGCAAGGTGATATTCTTGCCCTTTACAATACAGCACTGAAGC
CAGGACATGCAGCTGCATTTTTCAGAAGATTGTGATTTCAGCCTCTACTAAAGTTGTCATCC
GTGGAGCAGGAAACCAGTACAATTACATTGGCTATCTTGACTATAAAAAGGAAAGAATTC
GTAAGCTTTCCATGAAAGCCAGCACTTGCTCATTTAATCCTGGAGTTTTTGTGCAAACC
TGACGGAATGGAAACGACAGAATATACTAACCAACTGGAAAAATGGATGAAACTCAATG
TAGAAGAGGGACTGTATAGCAGAACCCTGGCTGGTAGCATCACAACACCTCCTCTGCTTA
TCGTATTTTATCAACAGCACTCTACCATCGATCCTATGTGGAATGTCGGCCACCTTGCTT
CCAGTGCTGGAAAACGATATTCACCTCAGTTTGTAAAGGCTGCCAAGTTACTCCATTGGA
ATGGACATTTGAAGCCATGGGGAAGGACTGCTTCATATACTGATGTTTGGGAAAAATGGT
ATATTCCAGACCCAACAGGCAAATTC AACCTAATCCGAAGATATACCGAGATCTCAAACA
TAAAGTGAAACAGAATTTGAACTGTAAGCAAGCATTTCTCAGGAAGTCCTGGAAGATAGC
ATGCATGGGAAGTAACAGTTGCTAGGCTTCAATGCCTATCGGTAGCAAGCCATGGAAAAA
GATGTGTCAGCTAGGTAAAGATGACAACTGCCCTGTCTGGCAGTCAGCTTCCCAGACAG
ACTATAGACTATAAATATGTCTCCATCTGCCTTACCAAGTGTCTTACTACAATGCTG
AATGACTGGAAAGAAGAACTGATATGGCTAGTTTCAGCTAGCTGGTACAGATAATTCAAAA
CTGCTGTTGGTTTTAATTTTGTAACTGTGGCCTGATCTGTAAATAAACTTACATTTTTC

124/392

FIGURE 118

MSFRKVNIIILVLAVALFLLVLHHNFLSLSSLLRNEVTDSGIVGPQPIDFVPNALRHAVD
GRQEEIPVVIAASEDRLGGAIAAINSIOHNTRSNVIFYIVTLNNTADHLRSWLNSDSLKS
IRYKIVNFDPKLLEGKVKEPDQGESMKPLTFARFYLPILVPSAKKAIYMDDDVIVQGDI
LALYNTALKPGHAAAFSEDCDSASTKVVIRGAGNQYNYIGYLDYKKERIRKLSMKASTCS
FNPGVFVANLTEWKRQNTITNQLEKWMKLNVEEGLYSRTLGSITTPPLLIVFYQQHSTID
PMWNVRLGSSAGKRYSPQFVKAACKLLHWNGHLKPWGRTASYTDVWEKWYIPDPTGKFNL
IRRYTEISNIK

125/392

FIGURE 119

CCATCCCTGAGATCTTTTTATAAAAAACCCAGTCTTTGCTGACCAGACAAAGCATACCAG
ATCTCACCAGAGAGTCGCAGACACTATGCTGCCTCCCATGGCCCTGCCAGTGTGTCTTG
GATGCTGCTTTCCTGCCTCATTCTCCTGTGTGTCAGGTTCAAGGTGAAGAAACCCAGAAGGA
ACTGCCCTCTCCACGGATCAGCTGTCCCAAAGGCTCCAAGGCCTATGGCTCCCCCTGCTA
TGCCTTGTTTTTGTACCAAAATCCTGGATGGATGCAGATCTGGCTTGCCAGAAGCGGCC
CTCTGGAAACTGGTGTCTGTGCTCAGTGGGGCTGAGGGATCCTTCGTGTCTCCCTGGT
GAGGAGCATTAGTAACAGCTACTCATACTCTGGATTGGGCTCCATGACCCACACAGGG
CTCTGAGCCTGATGGAGATGGATGGGAGTGGAGTAGCACTGATGTGATGAATTACTTTGC
ATGGGAGAAAAATCCCTCCACCATCTTAAACCCTGGCCACTGTGGGAGCCTGTCAAGAAG
CACAGGATTTCTGAAGTGGAAGATTATACTGTGATGCAAAGTTACCCTATGTCTGCAA
GTTCAAGGACTAGGGCAGGTGGGAAGTCAGCAGCCTCAGCTTGGCGTGCAGCTCATCATG
GACATGAGACCAGTGTGAAGACTCACCTGGAAGAGAATATTCTCCCCAACTGCCCTAC
CTGACTACCTTGTATGATCCTCCTTCTTTTTCCTTTTCTTCACCTTCATTTCAGGCTT
TTCTCTGTCTTCCATGTCTTGAGATCTCAGAGAATAATAATAAAAATGTTACTTTATAAA
AAAAAAAAAAAAAAAAAAAA

126/392

FIGURE 120

MLPPMALPSVSWMLLSCLILLCQVQGEETQKELPSPRISCPKGSKAYGSPCYALFLSPKS
WMDADLACQKRPSGKLVSVLSGAEGSFVSSLVRSISNSYSYIWIGLHDPTQGSEPDGDGW
EWSSTDVMNYFAWEKNPSTILNPGHCGSLSRSTGFLKWKDYNCDAKLPYVCKFKD

Important features:

Signal peptide:

amino acids 1-26

C-type lectin domain signature:

amino acids 146-171

127/392

FIGURE 121

AAAGTTACATTTTCTCTGGAACCTCTCCTAGGCCACTCCCTGCTGATGCAACATCTGGGTT
TGGGCAGAAAGGAGGGTGCTTCGGAGCCCGCCCTTTCTGAGCTTCCTGGGCCGCTCTAG
AACAAATTCAGGCTTCGCTGCGACTCAGACCTCAGCTCCAACATATGCATTCTGAAGAAAG
ATGGCTGAGATGGACAGAATGCTTTATTTTGGAAAGAAACAATGTTCTAGGTCAAACCTGA
GTCTACCAAATGCAGACTTTCACAATGGTTCTAGAAGAAATCTGGACAAGTCTTTTCATG
TGGTTTTTCTACGCATTGATTCCATGTTTGCTCACAGATGAAGTGGCCATTCTGCCTGCC
CCTCAGAACCTCTCTGTACTCTCAACCAACATGAAGCATCTCTTGATGTGGAGCCCAGTG
ATCGCGCTTGAGAAACAGTGTACTATTCTGTGGAATACCAGGGGGAGTACGAGAGCCTG
TACACGAGCCACATCTGGATCCCCAGCAGCTGGTGCTCACTCACTGAAGGTCTTGAGTGT
GATGTCACTGATGACATCACGGCCACTGTGCCATACAACCTTCGTGTCAGGGCCACATTG
GGCTCACAGACCTCAGCCTGGAGCATCCTGAAGCATCCCTTTAATAGAAACTCAACCATC
CTTACCCGACCTGGGATGGAGATCACCAGAGATGGCTTCCACCTGGTTATTGAGCTGGAG
GACCTGGGGCCCCAGTTTGAGTTCCTTGTTGGCCTACTGGAGGAGGGAGCCTGGTGCCGAG
GAACATGTCAAAATGGTGAGGAGTGGGGGTATTCCAGTGCACCTAGAAACCATGGAGCCA
GGGGCTGCATACTGTGTGAAGGCCAGACATTCTGTAAGGCCATTGGGAGGTACAGCGCC
TTCAGCCAGACAGAATGTGTGGAGGTGCAAGGAGAGGCCATTCCCTGGTACTGGCCCTG
TTTGCCTTTGTTGGCTTCATGCTGATCCTTGTTGGTCTGCTGCACTGTTCTGCTGAAAATG
GGCCGGCTGCTCCAGTACTCCTGTTGCCCCGTGGTGGTCTCCAGACACCTTGAAAATA
ACCAATTCACCCAGAGTTAATCAGCTGCAGAGGGAGGAGGTGGATGCCTGTGCCACG
GCTGTGATGTCTCCTGAGGAACTCCTCAGGGCCTGGATCTCATAAGTTTGCAGGAGGGCC
CAGGTGAAGCCGAGAACCTGGTCTGCATGACATGGAAACCATGAGGGGACAAGTTGTGTT
TCTGTTTTCCGCCACGGACAAGGGATGAGAGAAGTAGGAAGAGCCTGTTGTCTACAAGTC
TAGAAGCAACCATCAGAGGCAGGGTGGTTTGTCTAACAGAACTGACTGAGGCTTAGGG
GATGTGACCTCTAGACTGGGGCTGCCACTTGCTGGCTGAGCAACCCTGGGAAAAGTGAC
TTCATCCCTTCGGTCTTAAGTTTTCTCATCTGTAATGGGGGAATTACCTACACACCTGCT
AAACACACACACAGAGTCTCTCTATATATACACACGTACACATAAATACACCCAGC
ACTTGCAAGGCTAGAGGGAACTGGTGACACTCTACAGTCTGACTGATTAGTGTCTCTG
GAGAGCAGGACATAAATGTATGATGAGAATGATCAAGGACTCTACACACTGGGTGGCTTG
GAGAGCCCACTTTCCAGAAATAATCCTTGAGAGAAAAGGAATCATGGGAGCAATGGTGTT
GAGTTCACTTCAAGCCCAATGCCGTGCAGAGGGGAATGGCTTAGCGAGCTCTACAGTAG
GTGACCTGGAGGAAGGTCACAGCCACACTGAAAATGGGATGTGCATGAACACGGAGGATC
CATGAACTACTGTAAAGTGTGACAGTGTGTGCACACTGCAGACAGCAGGTGAAATGTAT
GTGTGCAATGCGACGAGAATGCAGAAGTCAGTAACATGTGCATGTTTGTGTGCTCCTTT
TTTCTGTTGGTAAAGTACAGAATTCAGCAAATAAAAAGGGCCACCCTGGCCAAAAGCGGT
AAAAAAAAAAAAAAAA

128/392

FIGURE 122

MQTFTMVLEEIIWTSLFMWFFYALIPCLLTDEVAILPAPQNLSVLSTNMKHLMLWSPVIAP
GETVYYSVEYQGEYESLYTSHIWIPSSWCSLTEGPECVTDITATVPYNLRVRATLGSQ
TSAWSILKHPFNRNSTILTRPGMEITKDGFLVIELEDLGPQFEFLVAYWRREPGAEHV
KMVRSGGIPVHLETMEPGAAYCVKAQTFVKAIGRYSAFSQTECVEVQGEAIPVLALFAF
VGFMLILVVVPLFVWKMGRLLQYSCCPVVVLPDTLKITNSPQKLISCRREEVDACATAVM
SPEELLRAWIS

Important features:

Signal peptide:

amino acids 1-29

Transmembrane domain:

amino acids 230-255

N-glycosylation sites:

amino acids 40-43 and 134-137

Tissue factor proteins homology:

amino acids 92-119

Integrins alpha chain protein homology:

amino acids 232-262

129/392

FIGURE 123

CGGACGCGTGGGCCGCCACCTCCGGAACAAGCCATGGTGGCGGCGACGGTGGCAGCGGCG
TGGCTGCTCCTGTGGGCTGCGGCCTGCGCGCAGCAGGAGCAGGACTTCTACGACTTCAAG
GCGGTCAACATCCGGGGCAAACCTGGTGTGCTGAGAAAGTACCGCGGATCGGTGTCCCTG
GTGGTGAATGTGGCCAGCGAGTGGGCTTACAGACCAGCACTACCGAGCCCTGCAGCAG
CTGCAGCGAGACCTGGGCCCCCACCCTTTAACGTGCTCGCCTTCCCCTGCAACCAGTTT
GGCCAACAGGAGCCTGACAGCAACAAGGAGATTGAGAGCTTTGCCCGCCGCACCTACAGT
GTCTCATTTCCCATGTTTAGCAAGATTGCAGTCACCGGTACTGGTGGCCATCCTGCCTTC
AAGTACCTGGCCCAGACTTCTGGGAAGGAGCCCACCTGGAACCTTCTGGAAGTACCTAGTA
GCCCCAGATGGAAAGGTGGTAGGGGCTTGGGACCCAACTGTGTGAGTGGAGGAGGTCAGA
CCCCAGATCACAGCGCTCGTGAGGAAGCTCATCCTACTGAAGCGAGAAGACTTATAACCA
CCGCGTCTCCTCCTCCACCACCTCATCCCGCCCACCTGTGTGGGGCTGACCAATGCAAAC
TCAAATGGTGCTTCAAAGGGAGAGACCCACTGACTCTCCTTCTTTACTCTTATGCCATT
GGTCCCATCATTTCTTGTGGGGGAAAAATTCTAGTATTTTGATTATTTGAATCTTACAGCA
ACAAATAGGAACTCCTGGCCAATGAGAGCTCTTGACCAGTGAATCACCAGCCGATACGAA
CGTCTTGCCAACAAAAATGTGTGGCAAATAGAAGTATATCAAGCAATAATCTCCCACCCA
AGGCTTCTGTAAACTGGGACCAATGATTACCTCATAGGGCTGTTGTGAGGATTAGGATGA
AATACCTGTGAAAGTGCCTAGGCAGTGCCAGCCAAATAGGAGGCATTCAATGAACATTTT
TTGCATATAAACCAAAAAATAACTTGTTATCAATAAAAACTTGCATCCAACATGAATTT
CAGCCGATGATAATCCAGGCCAAAGGTTTAGTTGTTGTTATTTCTCTGTATTATTTCT
TCATTACAAAAGAAATGCAAGTTCATTGTAACAATCCAAACAATACCTCACGATATAAAA
TAAAAATGAAAGTATCCTCCTCAAAA

130/392

FIGURE 124

MVAATVAAAWLLLWAAACAQQEQDFYDFKAVNIRGKLVSLEKYRGSVSLVVNVASECGFT
DQHYRALQQLQORDLGPHHFNVLAFFPCNQFGQQEPDSNKEIESFARRTYSVSFPMFSKIAV
TGTGAHPAFKYLAQTSGKEPTWNFWKYLVPDGTKVVGAWDPTVSVEEVRPQITALVRKLI
LLKREDL

131/392

FIGURE 125

CGGACGCGTGGGCGGACGCGTGGGCGGACGCGTGGGTGGGAGGGGGCAGGATGGGAGGG
AAAGTGAAGAAAACAGAAAAGGAGAGGGACAGAGGCCAGAGGACTTCTCATACTGGACAG
AAACCGATCAGGCATGGAACTCCCCCTTCGTCACTCACCTGTTCTTGCCCCCTGGTGTTCCT
GACAGGTCTCTGCTCCCCCTTTAACCTGGATGAACATCACCCACGCCTATTCCCAGGGCC
ACCAGAAGCTGAATTTGGATACAGTGTCTTACAACATGTTGGGGGTGGACAGCGATGGAT
GCTGGTGGGCGCCCCCTGGGATGGGCCTTCAGGCGACCGAGGGGGGACGTTTATCGCTG
CCCTGTAGGGGGGGGCCACAATGCCCCATGTGCCAAGGGCCACTTAGGTGACTACCAACT
GGGAAATTCATCTCATCCTGCTGTGAATATGCACCTGGGGATGTCTCTGTTAGAGACAGA
TGGTGATGGGGGATTCATGGTGAGCTAAGGAGAGGGTGGTGGCAGTGTCTCTGAAGGTCC
ATAAAAGAAAAAGAGAAGTGTGGTAAGGGAAAATGGTCTGTGTGGAGGGGTCAAGGAGT
TAAAAACCCTAGAAAGCAAAAGGTAGGTAATGTCAGGGAGTAGTCTTCATGCCTCCTTCA
ACTGGGAGCATGTTCTGAGGGTGCCCTCCCAAGCCTGGGAGTAACTATTTCCCCCATCCC
CAGGCCTGTGCCCCCTCTCTGGTCTCGTGCTTGTGGCAGCTCTGTCTTCAGTTCTGGGATA
TGTGCCCCGTGTGGATGCTTCATTCCAGCCTCAGGGAAGCCTGGCACCCACTGCCCCAAGCT
GAGCCAGAGGAAGGCTGAGTACTTGGTTCACAGAAGGAGATACTGGGTGGGAAAAAGATG
GGGCAAAGCGGTATGATGCCTGGCAAAGGGCCTGCATGGCTATCCTCATTGCTACCTAAT
GTGCTTGCAAAGCTCCATGTTTCCTAACAGATTGAGACTCCTGGCCAGGTGTGGTGGCC
CACACCTGTAATTCTAGCACTTTGGGAGGCCAAGGTGGGCAGATCACTTGAGGTCAGGAG
TTCAAGACCAGCCTGGCCAACATGGTGAACTCCATCTCTACTAAAAAATAAATAACA
AAAATTAGCTGGGTGCGCTAGTGCTGCCTGTAATCTCATCTACTCGGGAGGCTAAGACA
GGAGACTCTCACTTCAACCCAGGAGGTGGAGGTTGCGGTGAGCCAAGATTGTGCCTCTGC
ACTCTAGCGTGGGTGACAGAGTAAGCGAGACTCCATCTCAAAAATAAATAAATAAAT
TCAGACTCCTTATCAGGAGTCCATGATCTGGCCTGGCACAGTAACTCATGCCTGTAATCC
CAACATTTTGGGAGGCCAACGCAGGAGGATTGCTTGAGGTCTGGAGGTTTGAGACCAGCC
TGGGCAACATAGAAAGACCCCATCTCTAAATAAATGTTTTAAAAAT

132/392

FIGURE 126

MELPFVTHLFLPLVFLTGLCSPPFNLDHHPRLPFPGPPEAEFGYSVLQHVGGGQRWMLVGA
PWDGPGSGDRRGDVYRCPVGGAHNAPCAKGHLGDYQLGNSSHPAVNMHLGMSLLETGDGCG
FMVS

Important features:

Signal peptide:

amino acids 1-22

Cell attachment sequence:

amino acids 70-73

N-glycosylation site:

amino acids 98-101

Integrins alpha chain proteins:

amino acids 67-81

133/392

FIGURE 127

GAGAGGACGAGGTGCCGCTGCCTGGAGAATCCTCCGCTGCCGTCCGGCTCCCGGAGCCCAG
CCCTTTCCCTAACCCAACCCAACCTAGCCCAGTCCCAGCCGCCAGCGCCTGTCCCTGTCAC
GGACCCAGCGTTACCATGTCATCCTGCCGTCTTCCTATCCTTACCCGACCTCAGATGCTC
CCTTCTGCTCCTGGTAACTTGGGTTTTTACTCCTGTAACAACTGAAATAACAAGTCTTGC
TACAGAGAATATAGATGAAATTTTAAACAATGCTGATGTTGCTTTAGTAAATTTTATGC
TGACTGGTGTGCTTTCAGTCAGATGTTGCATCCAATTTTGGAGGAAGCTTCCGATGTCAT
TAAGGAAGAATTTCCAAATGAAAATCAAGTAGTGTGTTGCCAGAGTTGATTGTGATCAGCA
CTCTGACATAGCCCAGAGATACAGGATAAGCAAATACCCAACCCCTCAAATTTGTTTCGTAA
TGGGATGATGATGAAGAGAGAATACAGGGGTCAGCGATCAGTGAAAGCATTGGCAGATTA
CATCAGGCAACAAAAAGTGACCCCATTTCAAGAAATTCGGGACTTAGCAGAAATCACCAC
TCTTGATCGCAGCAAAAGAAATATCATTGGATATTTTGAGCAAAAGGACTCGGACAACTA
TAGAGTTTTTGAACGAGTAGCGAATATTTTGCATGATGACTGTGCCTTTCTTCTGCATT
TGGGGATGTTTTCAAAACCGGAAAGATATAGTGGCGACAACATAATCTACAAACCACCAGG
GCATTCTGCTCCGGATATGGTGTACTTGGGAGCTATGACAAATTTTGATGTGACTTACAA
TTGGATTCAAGATAAATGTGTTCTTGTCCGAGAAATAACATTTGAAAATGGAGAGGA
ATTGACAGAAGAAGGACTGCCTTTTCTCATACTCTTTCACATGAAAGAAGATACAGAAAG
TTTAGAAATATTCCAGAAATGAAGTAGCTCGGCAATTAATAAGTGAAAAAGGTACAATAAA
CTTTTACATGCCGATTGTGACAAATTTAGACATCCTCTTCTGCACATACAGAAACTCC
AGCAGATTGTCCTGTAATCGCTATTGACAGCTTTAGGCATATGTATGTGTTTGGAGACTT
CAAAGATGTATTAATTCCTGGAAACTCAAGCAATTCGTATTTGACTTACATTCTGGAAA
ACTGCACAGAGAATTCCATCATGGACCTGACCCAACCTGATACAGCCCCAGGAGAGCAAGC
CCAAGATGTAGCAAGCAGTCCACCTGAGAGCTCCTTCCAGAACTAGCACCCAGTGAATA
TAGGTATACTCTATTGAGGGATCGAGATGAGCTTTTAAAAACTTGAAAAACAGTTTGTAAG
CCTTTCAACAGCAGCATCAACCTACGTGGTGGAAATAGTAAACCTATATTTTCATAATTC
TATGTGTATTTTTATTTTGAATAAACAGAAAGAAATTTAAAAA
AAAAAAAAAAAAAAAAAAAAA

134/392

FIGURE 128

MHPAVFLSLPDLRCSLLLLVTWVFTPVTTEITSLATENIDEILNNADVALVNFYADWCRF
SQMLHPIFEEASDVIKEEFPNENQVVFARVDCDQHS DIAQRYRISKYPTLKLFRNGMMMK
REYRGQRSVKALADYIRQOKSDPIQEIRDLAEITTLDRSKRNIIGYFEQKDSDNRYRVFER
VANILHDDCAFLSAFGDVSKPERYSGDNIIYKPPGHSAPDMVYLGAMTNFDVTYNWIQDK
CVPLVREITFENGEEELTEEGLPFLILFHMKEDTESLEIFQNEVARQLISEKGTINFLHAD
CDKFRHPLLHIQKTPADCPVIAIDSFRHMYVFGDFKDVLIPGKLKQFVFDLHSGKLHREF
HHGPDPTDTAPGEQAQDVASSPPESSFQKLAPSEYRYTLLRDRDEL

Important features:

Signal peptide:

amino acids 1-29

Endoplasmic reticulum targeting sequence:

amino acids 403-406

Tyrosine kinase phosphorylation site:

amino acids 203-211

Thioredoxin family proteins:

amino acids 50-66

135/392

FIGURE 129

GAGCAGGACGGAGCCATGGACCCCGCCAGGAAAGCAGGTGCCCAGGCCATGATCTGGACT
GCAGGCTGGCTGCTGCTGCTGCTGCTTCGCGGAGGAGCGCAGGCCCTGGAGTGCTACAGC
TGCGTGACAGAAAGCAGATGACGGATGCTCCCCGAACAAGATGAAGACAGTGAAGTGCGCG
CCGGGCGTGAGCGTCTGCACCGAGGCCGTGGGGGCGGTGGAGACCATCCACGGACAATT
TCGCTGGCAGTGCGGGGTTGCGGTTTCGGGACTCCCCGGCAAGAATGACCGCGCCTGGAT
CTTCACGGGCTTCTGGCGTTTCATCCAGCTGCAGCAATGCGCTCAGGATCGCTGCAACGCC
AAGCTCAACCTCACCTCGCGGGCGCTCGACCCGGCAGGTAATGAGAGTGACATACCCGCC
AACGGCGTGAGTGCTACAGCTGTGTGGGCTGAGCCGGGAGGCGTGCCAGGGTACATCG
CCGCCGGTCGTGAGCTGCTACAACGCCAGCGATCATGTCTACAAGGGCTGCTTCGACGGC
AACGTACCTTGACGGCAGCTAATGTGACTGTGTTCCTTGCTGTCGGGGCTGTGTCCAG
GATGAATTCTGCACTCGGGATGGAGTAACAGGCCCAGGGTTCACGCTCAGTGGCTCCTGT
TGCCAGGGGTCCCGCTGTAACCTGACCTCCGCAACAAGACCTACTTCTCCCTCGAATC
CCACCCCTTGTCGGGCTGCCCCCTCCAGAGCCACGACTGTGGCCTCAACCACATCTGTC
ACCACTTCTACCTCGGCCCCAGTGAGACCCACATCCACCACCAACCCATGCCAGCGCCA
ACCAGTCAGACTCCGAGACAGGGAGTAGAACACGAGGCCTCCCGGGATGAGGAGCCAGG
TTGACTGGAGGCGCGCTGGCCACCAGGACCGCAGCAATTGAGGGCAGTATCCTGCAAAA
GGGGGGCCCCAGCAGCCCCATAATAAAGGCTGTGTGGCTCCACAGCTGGATTGGCAGCC
CTTCTGTTGGCCGTGGCTGCTGGTGTCTTACTGTGAGCTTCTCCACCTGGAAATTTCCCT
CTCACCTACTTCTCTGGCCCTGGGTACCCCTCTTCTCATCACTTCCTGTTCCCACTG
GACTGGGCTGGCCAGCCCCTGTTTTTCCAACATTCCCCAGTATCCCCAGCTTCTGCTGC
GCTGGTTTTGCGGCTTTGGGAAATAAAATACCGTTGTATATATTCTGCCAGGGGTGTTCTA
GCTTTTTGAGGACAGCTCCTGTATCCTTCTCATCCTTGTCTCTCCGCTTGTCTCTTGTG
ATGTTAGGACAGAGTGAGAGAAGTCAGCTGTACGGGGAAGGTGAGAGAGAGGATGCTAA
GCTTCCTACTCACTTTCTCCTAGCCAGCCTGGACTTTGGAGCGTGGGTGGGTGGGACAA
TGGCTCCCCACTCTAAGCACTGCCTCCCCTACTCCCCGCATCTTTGGGGAATCGGTTCCC
CATATGTCTTCTTACTAGACTGTGAGCTCCTCGAGGGGGGGCCCGGTACCCAATTCGCC
CTATAGTGAGTCGTA

136/392

FIGURE 130

MDPARKAGAQAMIWTAGWLLLLLLRGGAQALECYSCVQKADDGCS PNKMKT VKCAPGVDV
CTEAVGAVETIHGQFSLAVRGCGSGLPGKNDRGLDLHGLLAFIQLQQCAQDRCN AKLNLT
SRALDPAGNESAYPPNGVECYSCVGLSREACQGTSPPVVSCYNASDHVYKGC FDNVTLT
AANVTVSLPVRGCVQDEFCTRDGVTGPGFTLSGCCQGSRCNSDLRNKTYFSPRI PPLVR
LPPPEPTTVASTTSVTTST SAPVRPTSTTKPMPAPTSQTPRQGEHEASRDEEPRLTGGA
AGHQDRSNSGQYPAKGGPQQPHNKGCVAPTAGLAALLLAVAAGVLL

137/392

FIGURE 131

AAACTTGACGCCATGAAGATCCCGGTCTTCCTGCCGTGGTGCTCCTCTCCCTCCTGGTG
CTCCACTCTGCCCAGGGAGCCACCCTGGGTGGTCCTGAGGAAGAAAGCACCATTGAGAAT
TATGCGTCACGACCCGAGGCCTTTAACACCCCGTTCCTGAACATCGACAAATTGCGATCT
GCGTTTAAGGCTGATGAGTTCCTGAACTGGCACGCCCTCTTTGAGTCTATCAAAAGGAAA
CTTCCTTTCCTCAACTGGGATGCCTTTCCTAAGCTGAAAGGACTGAGGAGCGCAACTCCT
GATGCCCAGTGAACCATGACCTCCACTGGAAGAGGGGGCTAGCGTGAGCGCTGATTCTCAA
CCTACCATAACTCTTTCCTGCCTCAGGAACCTCCAATAAAACATTTTCCATCCAAA

138/392

FIGURE 132

MKIPVLPVVLVLLSLLVLHSAQGATLGGPEEESTIENYASRPEAFNTPFLNIDKLRSAFKA
DEFLNWHALFESIKRKLPFLNWDAPFKLKGLRSATPDAQ

139/392

FIGURE 133

CAGTTCTGAAATCAATGGAGTTAATTTAGGGAATACAAACCAGCCATGGGGGTGGAGATT
GCCTTTGCCTCAGTGATTCTCACCTGCCTCTCCCTTCTGGCAGCAGGAGTCTCCAGGTT
GTTCTTCTCCAGCCAGTTCCAACCTCAGGAGACAGGTCCCAAGGCCATGGGAGATCTCTCC
TGTGGCTTTGCCGGCCACTCATGAGAGTGTTTTTGTGTAAAGTATTTTTTTAGAATACTGT
TGA~~CTT~~CTTCATGATTTAATAACCATCCTTTGCGAAGTTTTATGAGGCTTTAGGGGAATG
TCAACCCTCAAATTTTTGTTATACTAGATGGCTTCCATTTACCCACCACTATTTTAAGGT
CCCTTTATTTTTAGGTTCAAGGTTCA~~TTT~~GACTTGAGAAAGTGCCCTTCTGCAGCTTCAT
TGATTTTGTTTATCTTCACTATTAATTGTAACGATTAAAAAAGAATAAGAGCACGCAGAC
CTCTAGGAGAATATTTTATCCCTGGGTGCCCCTGACACATTTATGTAGTGATCCACAAA
TGTGATTGTTAATTTAAATGTTATTCTAATATTAGTACATTCAGTTGTGATGTAATATGA
ATAACCAGAATCTATTTCTTAAAAGTTTTGAGTATATTTTTTCACTAGATATTTGTATAG
AAAGACTGAATAGTGATG

140/392

FIGURE 134

MGVEIAFASVILTCLSLAAGVSQVVLLQPVPQTQETGPKAMGDLSCGFAGHS

141/392

FIGURE 135

GGGGAATCTGCAGTAGGTCTGCCGGCGATGGAGTGGTGGGCTAGCTCGCCGCTTCGGCTC
TGGCTGCTGTTGTTCCCTCCTGCCCTCAGCGCAGGGCCGCCAGAAGGAGTCAGGTTCAAAA
TGGAAAGTATTTATTGACCAAATTAACAGGTCTTTGGAGAATTACGAACCATGTTCAAGT
CAAACTGCAGCTGCTACCATGGTGTCATAGAAGAGGATCTAACTCCTTTCCGAGGAGGC
ATCTCCAGGAAGATGATGGCAGAGGTAGTCAGACGGAAGCTAGGGACCCACTATCAGATC
ACTAAGAACAGACTGTACCGGGAAAATGACTGCATGTTCCCCTCAAGGTGTAGTGGTGTT
GAGCACTTTATTTTGGAAAGTGATCGGGCGTCTCCCTGACATGGAGATGGTGATCAATGTA
CGAGATTATCCTCAGGTTCTAAATGGATGGAGCCTGCCATCCCAGTCTTCTCCTTCAGT
AAGACATCAGAGTACCATGATATCATGTATCCTGCTTGGACATTTTGGGAAGGGGGACCT
GCTGTTTTGGCCAATTTATCCTACAGGTCTTGGACGGTGGGACCTCTTCAGAGAAGATCTG
GTAAGGTGAGCAGCACAGTGGCCATGGAAAAAGAAAACTCTACAGCATATTTCCGAGGA
TCAAGGACAAGTCCAGAACGAGATCCTCTCATTCTTCTGTCTCGGAAAAACCCAAAACT
GTTGATGCAGAATACACCAAAAACCGGCCTGGAAATCTATGAAAGATACCTTAGGAAAG
CCAGCTGCTAAGGATGTCCATCTTGTGGATCACTGCAAATACAAGTATCTGTTTAATTTT
CGAGGCGTAGCTGCAAGTTTCCGGTTTAAACACCTCTTCCCTGTGTGGCTCACTTGTTTTT
CATGTTGGTGATGAGTGGCTAGAATTCTTCTATCCACAGCTGAAGCCATGGGTTCACTAT
ATCCCAGTCAAAACAGATCTCTCCAATGTCCAAGAGCTGTTACAATTTGTAAAAGCAAAT
GATGATGTAGCTCAAGAGATTGCTGAAAGGGGAAGCCAGTTTATTAGGAACCATTTGCAG
ATGGATGACATCACCTGTTACTGGGAGAACCTCTTGAGTGAATACTCTAAATTCCTGTCT
TATAATGTAACGAGAAGGAAAGGTTATGATCAAATTATCCCAAAATGTTGAAAAC TGAA
CTATAGTAGTCATCATAGGACCATAGTCCTCTTTGTGGCAACAGATCTCAGATATCCTAC
GGTGAGAAGCTTACCATAAGCTTGGCTCCTATACCTTGAATATCTGCTATCAAGCCAAAT
ACCTGGTTTTTCCTTATCATGCTGCACCCAGAGCAACTCTTGAGAAAGATTTAAAATGTGT
CTAATACACTGATATGAAGCAGTTCAACTTTTTGGATGAATAAGGACCAGAAATCGTGAG
ATGTGGATTTTGAACCCAACCTCTACCTTTTCAATTTCTTAAGACCAATCACAGCTTGTGCC
TCAGATCATCCACCTGTGTGAGTCCATCACTGTGAAATTGACTGTGTCCATGTGATGATG
CCCTTTGTCCCATTTATTTGGAGCAGAAAATTCGTCAATTTGGAAGTAGTACAACCTCATTGC
TGGAATTGTGAAATTATTCAAGGCGTGATCTCTGTCACTTTATTTTAATGTAGGAAACCC
TATGGGGTTTATGAAAAATACTTGGGGATCATTCTCTGAATGGTCTAAGGAAGCGGTAGC
CATGCCATGCAATGATGTAGGAGTTCTCTTTTGTAACCATAAACTCTGTTACTCAGGA
GGTTTCTATAATGCCACATAGAAAGAGGCCAATTGCATGAGTAATTATTGCAATTGGATT
TCAGGTTCCCTTTTTTGTGCCTTCATGCCCTACTTCTTAATGCCTCTCTAAAGCCAAA

142/392

FIGURE 136

MEWWASSPLRLWLLLFLPSAQGRQKESGSKWKV FIDQINRSLENYEPCSSQNCSCYHGV
IEEDLTPFRGGISRKMMAEVRRKLGTHYQITKNRLYREND CMFPSRCSGVEHFILEVIG
RLPDMEMVINVRDYPQVPKWMEPAIPVFSFSKTSEYHDIMYPAWTFWEGGPAVWPIYPTG
LGRWDLFREDLVRSAQWPWKKKNSTAYFRGSRTSPERDPLILLSRKNPKLVDAEYTKNQ
AWKSMKDTLGKPAAKDVHLVDHCKYKYLNFNRGVAASFRFKHLFLCGSLVFHVGDEWLEF
FYPQLKPWVHYIPVKTDLSNVQELLQFVKANDDVAQEIAERGSQFIRNHLQMDDITCYWE
NLLSEYSKFLSYNVTRRKGYDQIIPKMLKTEL

143/392

FIGURE 137

ATTCTCCTAGAGCATCTTTGGAAGCATGAGGCCACGATGCTGCATCTTGGCTCTTGTCTG
CTGGATAACAGTCTTCCTCCTCCAGTGTTCAAAGGAACTACAGACGCTCCTGTTGGCTC
AGGACTGTGGCTGTGCCAGCCGACACCCAGGTGTGGGAACAAGATCTACAACCCCTTCAGA
GCAGTGCTGTTATGATGATGCCATCTTATCCTTAAAGGAGACCCGCCGCTGTGGCTCCAC
CTGCACCTTCTGGCCCTGCTTTGAGCTCTGCTGTCCCGAGTCTTTTGGCCCCCAGCAGAA
GTTTCTTGTGAAGTTGAGGGTTCTGGGTATGAAGTCTCAGTGTCACTTATCTCCCATCTC
CCGGAGCTGTACCAGGAACAGGAGGCACGTCCTGTACCCATAAAAACCCAGGCTCCACT
GGCAGACGGCAGACAAGGGGAGAAGAGACGAAGCAGCTGGACATCGGAGACTACAGTTGA
ACTTCGGAGAGAAGCAACTTGACTTCAGAGGGATGGCTCAATGACATAGCTTTGGAGAGG
AGCCCAGCTGGGGATGGCCAGACTTCAGGGGAAGAATGCCTTCCTGCTTCATCCCCTTTC
CAGCTCCCCTTCCCGCTGAGAGCCACTTTCATCGGCAATAAAATCCCCACATTTACCATCT

144/392

FIGURE 138

MRPRCCILALVCWITVFLQCSKGTTDAPVGSGLWLCQPTPRCGNKIYNPSEQCCYDDAI
LSLKETRRCGSTCTFWPCFELCCPESFGPQQKFLVKLRVLGMKSQCHLSPISRSCTRNR
HVLYP

Important features:

Signal sequence:

amino acids 1-21

N-myristoylation sites:

amino acids 33-39, 70-76

145/392

FIGURE 139

CCTCTGTCCACTGCTTTTCGTGAAGACAAGATGAAGTTCACAATTGTCTTTGCTGGACTTC
TTGGAGTCTTCTAGCTCCTGCCCTAGCTAACTATAATATCAACGTCAATGATGACAACA
ACAATGCTGGAAGTGGGCAGCAGTCAGTGAGTGTCAACAATGAACACAATGTGGCCAATG
TTGACAATAACAACGGATGGGACTCCTGGAATTCATCTGGGATTATGGAAATGGCTTTG
CTGCAACCAGACTCTTTCAAAAGAAGACATGCATTGTGCACAAAATGAACAAGGAAGTCA
TGCCCTCCATTCAATCCCTTGATGCACTGGTCAAGGAAAAGAAGCTTCAGGGTAAGGGAC
CAGGAGGACCACCTCCCAAGGGCCTGATGTACTCAGTCAACCCAAACAAAGTCGATGACC
TGAGCAAGTTCGGAAAAAACATTGCAAACATGTGTCTGGGATTCCAACATACATGGCTG
AGGAGATGCAAGAGGCAAGCCTGTTTTTTTACTCAGGAACGTGCTACACGACCAGTGTAC
TATGGATTGTGGACATTTCTTCTGTGGAGACACGGTGGAGAACTAAACAATTTTTTAAA
GCCACTATGGATTTAGTCATCTGAATATGCTGTGCAGAAAAATATGGGCTCCAGTGGTT
TTTACCATGTCATTCTGAAATTTTCTCTACTAGTTATGTTTGATTCTTTAAGTTTCAA
TAAAATCATTTAGCATTGAAAAAA

146/392

FIGURE 140

MKFTIVFAGLLGVFLAPALANYNINVNDDNNAAGSGQQSVSVNNEHNVANVDNNGWDSW
NSIWYDYGNGFAATRLFQKKTCIVHKMNKEVMPSIQSLDALVKEKKLQGKGPGGPPPKGLM
YSVNPKNKVDLDSKFGKNIANMCRGIPTYMAEEMQEASLFFYSGTCYTTSVLWIVDISFCG
DTVEN

Signal Peptide:
amino acids 1-20

N-myristoylation Sites:
amino acids 67-72, 118-123, 163-168

Flavodoxin protein homology:
amino acids 156-174

147/392

FIGURE 141

GGTCCTTAATGGCAGCAGCCGCCGCTACCAAGATCCTTCTGTGCCTCCCGCTTCTGCTCC
TGCTGTCCGGCTGGTCCCGGGCTGGGCGAGCCGACCCCTCACTCTCTTTGCTATGACATCA
CCGTCATCCCTAAGTTCAGACCTGGACCACGGTGGTGTGCGGTTCAAGGCCAGGTGGATG
AAAAGACTTTTCTTCACTATGACTGTGGCAACAAGACAGTCACACCTGTCAGTCCCCCTGG
GGAAGAACTAAATGTCACAACGGCCTGGAAAGCACAGAACCCAGTACTGAGAGAGGTGG
TGGACATACTTACAGAGCAACTGCGTGACATTAGCTGGAGAATTACACACCCAAGGAAC
CCCTCACCTGTCAGGCAAGGATGTCTGTGAGCAGAAAGCTGAAGGACACAGCAGTGGAT
CTTGGCAGTTCAGTTTCGATGGGCAGATCTTCCTCCTCTTTGACTCAGAGAAGAGAATGT
GGACAACGGTTCATCCTGGAGCCAGAAAGATGAAAGAAAAGTGGGAGAATGACAAGGTTG
TGGCCATGTCCTTCCATTACTTCTCAATGGGAGACTGTATAGGATGGCTTGAGGACTTCT
TGATGGGCATGGACAGCACCCCTGGAGCCAAGTGCAGGAGCACCCTCGCCATGTCCTCAG
GCACAACCCAACTCAGGGCCACAGCCACCACCCTCATCCTTTGCTGCCTCCTCATCATCC
TCCCTTGCTTCATCCTCCCTGGCATCTGAGGAGAGTCCTTTAGAGTGACAGGTTAAAGCT
GATACCAAAGGCTCCTGTGAGCACGGTCTTGATCAAACCTCGCCCTTCTGTCTGGCCAGC
TGCCCACGACCTACGGTGTATGTCCAGTGGCCTCCAGCAGATCATGATGACATCATGGAC
CCAATAGCTCATTCACTGCCTTGATTCTTTTGCCAACAATTTTACCAGCAGTTATACCT
AACATATTATGCAATTTTCTCTTGGTGCTACCTGATGGAATTCCTGCACTTAAAGTTCTG
GCTGACTAAACAAGATATATCATTTTCTTCTCTCTTTTGTGTTGGAAAATCAAGTACT
TCTTTGAATGATGATCTCTTTCTTGCAAATGATATTGTCAGTAAAATAATCACGTTAGAC
TTCAGACCTCTGGGGATTCTTTCCGTGTCCTGAAAGAGAATTTTAAATTATTTAATAAG
AAAAAATTTATATTAATGATTGTTTCCTTTAGTAATTTATTGTTCTGTACTGATATTTAA
ATAAGAGTTCTATTTCCCAAAAAAAAAAAAAAAAAAAAA

148/392

FIGURE 142

MAAAAATKILLCLPLLLLLLSGWSRAGRADPHSLCYDITVIPKFRPGPRWCAVQGQVDEKT
FLHYDCGNKTVTPVSPLGKKLNVTTAWKAQNPVLREVVDILTEQLRDIQLENYTPKEPLT
LQARMSCEQKAEGHSSGSWQFSFDGQIFLLFDSEKRMWTTVHPGARKMKEKWENDKVVAM
SFHYFSMGDCIGWLEDFLMGMDSTLEPSAGAPLAMSSGTTQLRATATTLLILCCLLIILPC
FILPGI

Important features:**Signal peptide:**

amino acids 1-25

Transmembrane domain:

amino acids 224-246

N-glycosylation site:

amino acids 68-72, 82-86

N-myristoylation site:

amino acids 200-206, 210-216

Amidation site:

amino acids 77-81

149/392

FIGURE 143

AATGTGAGAGGGGCTGATGGAAGCTGATAGGCAGGACTGGAGTGTTAGCACCCAGTACTGG
ATGTGACAGCAGGCAGAGGAGCACTTAGCAGCTTATTCAGTGTCCGATTCTGATTCCGGC
AAGGATCCAAGCATGGGAATGCTGCCGTCGGGCAACTCCTGGCACACTGCTCCTCTTCTG
GCTTTCCTGCTCCTGAGTTCAGGACCGCACGCTCCGAGGAGGACCGGGACGGCCTATGG
GATGCCTGGGGCCCATGGAGTGAATGCTCACGCACCTGCCGGGGAGGGGCCTCCTACTCT
CTGAGGCGCTGCCTGAGCAGCAAGAGCTGTGAAGGAAGAAATATCCGATACAGAACATGC
AGTAATGTGGACTGCCCACCAGAAGCAGGTGATTTCCGAGCTCAGCAATGCTCAGCTCAT
AATGATGTCAAGCACCATGGCCAGTTTTATGAATGGCTTCCTGTGTCTAATGACCCTGAC
AACCCATGTTCACTCAAGTGCCAAGCCAAAGGAACAACCCCTGGTTGTTGAAGTAGCACCT
AAGGTCTTAGATGGTACGCGTTGCTATACAGAATCTTTGGATATGTGCATCAGTGGTTTA
TGCCAAATTGTTGGCTGCGATCACCAGCTGGGAAGCACCGTCAAGGAAGATAAATCCCAGCT
GTCTGCAACGGAGATGGGTCCACCTGCCGGCTGGTCCGAGGGCAGTATAAATCCCAGCT
TCCGCAACCAAATCGGATGATACTGTGGTTGCACTTCCCTATGGAAGTAGACATATTCGC
CTTGTCTTAAAGGTCTGATCACTTATATCTGGAAACCAAAACCCCTCCAGGGGACTAAA
GGTGAACACAGTCTCAGCTCCACAGGAACCTTCTTGTGGACAATTCTAGTGTGGACTTC
CAGAAATTTCCAGACAAAGAGATACTGAGAATGGCTGGACCACTCACAGCAGATTTTATT
GTCAAGATTCGTAACCTCGGGCTCCGCTGACAGTACAGTCCAGTTCATCTTCTATCAACCC
ATCATCCACCGATGGAGGGAGACGGATTTCTTTCCTTGCTCAGCAACCTGTGGAGGAGGT
TATCAGCTGACATCGGCTGAGTGCTACGATCTGAGGAGCAACCGTGTGGTTGCTGACCAA
TACTGTCACTATTACCCAGAGAACATCAAACCCAAACCCAAGCTTCAGGAGTGCAACTTG
GATCCTTGTCCAGCCAGTGACGGATACAAGCAGATCATGCCTTATGACCTCTACCATCCC
CTTCTCGGTGGGAGGCCACCCCATGGACCGCGTGCTCCTCCTCGTGTGGGGGGGGCATC
CAGAGCCGGGCAGTTTCTTGTGTGGAGGAGGACATCCAGGGGCATGTCACTTCAGTGGAA
GAGTGGAAATGCATGTACACCCCTAAGATGCCCATCGCGCAGCCCTGCAACATTTTTGAC
TGCCCTAAATGGCTGGCACAGGAGTGGTCTCCGTGCACAGTGACATGTGGCCAGGGCCTC
AGATAACCGTGTGGTCTCTGCATCGACCATCGAGGAATGCACACAGGAGGCTGTAGCCCA
AAAACAAAGCCCCACATAAAAGAGGAATGCATCGTACCCACTCCCTGCTATAAACCCAAA
GAGAACTTCCAGTCGAGGCCAAGTTGCCATGGTTCAAACAAGCTCAAGAGCTAGAAGAA
GGAGCTGCTGTGTGTCAGAGGAGCCCTCGTTAAGTTGTAAAAGCACAGACTGTTCTATATTG
AAACTGTTTTGTTTAAAGAAAGCAGTGTCTCACTGGTTGTAGCTTTCATGGGTTCTGAAC
TAAGTGTAATCATCTCACCAAAGCTTTTTGGCTCTCAAATTAAAGATTGATTAGTTTCAA
AAAAAAA

150/392

FIGURE 144

MECCRRATPGTLLLFLAFLLLSSRTARSEEDRDGLWDAWGPWSECSRTC GGGGASYSLRRC
LSSKSCEGRNIRYRTCSNVDCPPEAGDFRAQQCSAHNDVKHHGQFYEWLPVSNDPDNPCS
LKCQAKGTTLVVELAPKVLDTGTRCYTESLDMCISGLCQIVGCDHQLGSTVKEDNCGVCNG
DGSTCRLVRGQYKSQLSATKSDDTVVALPYGSRHRLVLKGPDLHLYLETKTLQGTKGENS
LSSTGTFLVDNSSVDFQKFPDKEILRMAGPLTADFIVKIRNSGSADSTVQFIFYQPIIHR
WRETDFFPCSATCGGGYQLTSAECYDLRSNRVVADQYCHYYPENIKPKPKLQECNLDPCP
ASDGYKQIMPYDLYHPLPRWEATPWTACSSSCGGGIQSRVSCVEEDIQGHVTSVEEWKC
MYTPKMPIAQPCNIFDCPKWLAQEWSPCTVTGQGLRYRVVLCIDHRGMHTGGCSPKTKP
HIKEECIVPTPCYKPKEKLPVEAKLPWFKQAQELEEGA AVSEEPS

Important features:

Signal peptide:

amino acids 1-25

N-glycosylation site:

amino acids 251-254

Thrombospondin 1:

amino acids 385-399

von Willebrand factor type C domain proteins:

amino acids 385-399, 445-459 and 42-56

151/392

FIGURE 145

GGAGGAGGGAGGGCGGGCAGGCGCCAGCCCAGAGCAGCCCCGGGCACCAGCACGGACTCT
CTCTTCCAGCCCAGGTGCCCCCACTCTCGCTCCATTTCGGCGGGAGCACCCAGTCCTGTA
CGCCAAGGAACTGGTCTCTGGGGGCACCATGGTTTCGGCGGCAGCCCCCAGCCTCCTCATC
CTTCTGTTGCTGCTGCTGGGGTCTGTGCCTGCTACCGACGCCCCGCTCTGTGCCCCCTGAAG
GCCACGTTCTTGAGGATGTGGCGGGTAGTGGGGAGGCCGAGGGCTCGTCGGCCTCCTCC
CCGAGCCTCCCGCCACCCTGGACCCCCGGCCCTCAGCCCCACATCGATGGGGCCCCAGCCC
ACAACCCTGGGGGGCCCATCACCCCCACCAACTTCCTGGATGGGATAGTGGACTTCTTC
CGCCAGTACGTGATGCTGATTGCTGTGGTGGGCTCCCTGGCCTTTCTGCTGATGTTTCATC
GTCTGTGCCGCGGTATCACCCGGCAGAAGCAGAAGGCCTCGGCCTATTACCCATCGTCC
TTCCCCAAGAAGAAGTACGTGGACCAGAGTGACCGGGCCGGGGGCCCCCGGGCCTTCAGT
GAGGTCCCCGACAGAGCCCCCGACAGCAGGCCCGAGGAAGCCCTGGATTCTCCCGGCAG
CTCAGGCCGACATCTTGCCCGCCACCAGAACCTCAAGTCCCCCACCAGGGCTGCACTG
GGCGGTGGGGACCGAGCCAGGATGGTGGAGGGCAGGGGCGCAGAGGAAGAGGAGAAGGGC
AGCCAGGAGGGGGACAGGAAGTCCAGGGACATGGGGTCCAGTGGAGACACCAGAGGCG
CAGGAGGAGCCGTGCTCAGGGGTCTTTGAGGGGGCTGTGGTGGCCGGTGAGGGCCAAGGG
GAGCTGGAAGGGTCTCTCTTGTAGCCCAGGAAGCCCAGGGACCAGTGGGTCCCCCGAA
AGCCCCCTGTGCTTGCAGCAGTGTCCACCCCAGTGTCTAAACAGTCTCTCCCGGGCTGCCAGC
CCTGACTGTGCGGGCCCCAAGTGGTCACCTCCCCGTGTATGAAAAGGCCTTCAGCCCTGA
CTGCTTCCTGACACTCCCTCCTTGGCCTCCCTGTGGTGCCAATCCCAGCATGTGCTGATT
CTACAGCAGGCAGAAATGCTGGTCCCCGGTGCCCCGGAGGAATCTTACCAAGTGCCATCA
TCCTTCACCTCAGCAGCCCCAAAGGGCTACATCCTACAGCACAGTCCCCCTGACAAAGTG
AGGGAGGGCACGTGTCCCTGTGACAGCCAGGATAAAACATCCCCCAAAGTGCTGGGATTA
CAGGCGTGAGCCACCGTGCCCCGGCCCAAACCTACTTTTTTAAACAGCTACAGGGTAAATC
CTGCAGCACCCACTCTGGAAAATACTGCTCTTAATTTTCTGAAGGTGGCCCCCTGTTTC
TAGTTGGTCCAGGATTAGGGATGTGGGGTATAGGGCATTTAAATCCTCTCAAGCGCTCTC
CAAGCACCCCCGGCCTGGGGGTGAGTTTCTCATCCCGCTACTGCTGCTGGGATCAGGTTG
AATGAATGGAATCTTCTGTCTGGCCTCCAAAGCAGCCTAGAAGCTGAGGGGCTGTGTT
TGAGGGGACCTCCACCCTGGGGAAAGTCCGAGGGGCTGGGGAAAGGTTTCTGACGCCCAGC
CTGGAGCAGGGGGGCCCTGGCCACCCCTGTTGCTCACACATTGTCTGGCAGCCTGTGTC
CACAAATATTGCTCAGTCCTCGACAGGGAGCCTGGGCTCCGTCTGCTTTAGGGAGGCTCT
GGCAGGAGGTCTCTCCCCATCCCTCCATCTGGGGCTCCCCAACCTCTGCACAGCTCT
CCAGGTGCTGAGATATAATGCACCAGCACATAAACCTTTATTCCGGCCTGAAAAAAAAA
AAGA

152/392

FIGURE 146

MVSAAAPSLILLILLLLGSPATDARSVPLKATFLEDVAGSGEAGSSASSPSLPPPWTP
ALSPTSMGPQPTTLGGPSPPTNFLDGIVDFFRQYVMLIAVVGSLAFLLMFIVCAAVITRQ
KQKASAYYPSSFPPKKKYVDQSDRAGGPRAFSEVPDRAPDSRPEEALDSSRQLQADILAAAT
QNLKSPTRAALGGGDGARMVEGRGAEEEEKGSQEGDQEVQGHGVPVETPEAQEEPCSGVL
EGAVVAGEGQGELEGSLLLAQEAQGPVGPPEPCACSSVHPSV

Signal peptide:

amino acids 1-25

Transmembrane domain:

amino acids 94-118

N-myristoylation site:amino acids 18-24, 40-46, 46-52, 145-151, 192-198, 193-199,
211-217, 238-244, 242-248

153/392

FIGURE 147

GAAAGACGTGGTCCTGACAGACAGACAATCCTATTCCCTACCAAAATGAAGATGCTGCTG
CTGCTGTGTTTGGGACTGACCCTAGTCTGTGTCCATGCAGAAGAAGCTAGTTCTACGGGA
AGGAACTTTAATGTAGAAAAGATTAATGGGGAATGGCATACTATTATCCTGGCCTCTGAC
AAAAGAGAAAAGATAGAAGAACATGGCACTTTTAGACTTTTTTCTGGAGCAAATCCATGTC
TTGGAGAATTCCTTAGTTCTTAAAGTCCATACTGTAAGAGATGAAGAGTGCTCCGAATTA
TCTATGGTTGCTGACAAAACAGAAAAGGCTGGTGAATATTCTGTGACGTATGATGGATT
AATACATTTACTATACCTAAGACAGACTATGATAACTTTCTTATGGCTCACCTCATTAA
GAAAAGGATGGGGAAACCTTCCAGCTGATGGGGCTCTATGGCCGAGAACCAGATTTGAGT
TCAGACATCAAGGAAAGGTTTGCACTATGTGAGGAGCATGGAATCCTTAGAGAAAAT
ATCATTGACCTATCCAATGCCAATCGCTGCCTCCAGGCCCGAGAATGAAGAATGGCCTGA
GCCTCCAGTGTTGAGTGGACACTTCTCACCAGGACTCCACCATCATCCCTTCTATCCAT
ACAGCATCCCCAGTATAAATTCTGTGATCTGCATTCCATCCTGTCTCACTGAGAAGTCCA
ATTCCAGTCTATCAACATGTTACCTAGGATACCTCATCAAGAATCAAAGACTTCTTTAAA
TTTCTCTTTGATACACCTTGACAATTTTTTCATGAAATTATTCCTCTTCCTGTTCAATAA
ATGATTACCCTTGCACTTAA

154/392

FIGURE 148

MKMLLLLCLGLTLVCVHAEASSTGRNFNVEKINGEWHTIILASDKREKIEEHGNFRLFL
EQIHVLENSLVLKVHTVRDEECSELSMVADKTEKAGEYSVTYDGFNTFTIPKTDYDNFLM
AHLINEKDGETFQLMGLYGREPDLSSEIKERFAQLCEEHGILRENIIDLSNANRCLQARE

155/392

FIGURE 149

GTGGACTCTGAGAAGCCCAGGCAGTTGAGGACAGGAGAGAGAAGGCTGCAGACCCAGAGG
GAGGGAGGACAGGGAGTCGGAAGGAGGAGGACAGAGGAGGGCACAGAGACGCAGAGCAAG
GGCGGCAAGGAGGAGACCCTGGTGGGAGGAAGACACTCTGGAGAGAGAGGGGGCTGGGCA
GAGATGAAGTTCCAGGGGCCCCCTGGCCTGCCTCCTGCTGGCCCTCTGCCTGGGCAGTGGG
GAGGCTGGCCCCCTGCAGAGCGGAGAGGAAAAGCACTGGGACAAATATTGGGGAGGCCCTT
GGACATGGCCTGGGAGACGCCCTGAGCGAAGGGGTGGGAAAGGCCATTGGCAAAGAGGCC
GGAGGGGCAGCTGGCTCTAAAGTCAGTGAGGCCCTTGGCCAAGGGACCAGAGAAGCAGTT
GGCACTGGAGTCAGGCAGGTTCCAGGCTTTGGCGCAGCAGATGCTTTGGGCAACAGGGTC
GGGGAAGCAGCCCATGCTCTGGGAAACACTGGGCACGAGATTGGCAGACAGGCAGAAGAT
GTCATTTCGACACGGAGCAGATGCTGTCCGCGGCTCCTGGCAGGGGGTGCCTGGCCACAGT
GGTGCTTGGGAACTTCTGGAGGCCATGGCATCTTTGGCTCTCAAGGTGGCCTTGGAGGC
CAGGGCCAGGGCAATCCTGGAGGTCTGGGGACTCCGTGGGTCCACGGATACCCCGGAAAC
TCAGCAGGCAGCTTTGGAATGAATCCTCAGGGAGCTCCCTGGGGTCAAGGAGGCAATGGA
GGGCCACCAAACCTTTGGGACCAACACTCAGGGAGCTGTGGCCCAGCCTGGCTATGGTTCA
GTGAGAGCCAGCAACCAGAATGAAGGGTGCACGAATCCCCACCATCTGGCTCAGGTGGA
GGCTCCAGCAACTCTGGGGGAGGCAGCGGCTCACAGTCGGGCAGCAGTGGCAGTGGCAGC
AATGGTGACAACAACAATGGCAGCAGCAGTGGTGGCAGCAGCAGTGGCAGCAGCAGTGGC
AGCAGCAGTGGCGGCAGCAGTGGCGGCAGCAGTGGTGGCAGCAGTGGCAACAGTGGTGGC
AGCAGAGGTGACAGCGGCAGTGAGTCCTCCTGGGGATCCAGCACCGGCTCCTCCTCCGGC
AACCACGGTGGGAGCGGCGGAGGAAATGGACATAAACCCGGGTGTGAAAAGCCAGGGAAT
GAAGCCCGCGGGAGCGGGGAATCTGGGATTTCAGGGCTTCAGAGGACAGGGAGTTTCCAGC
AACATGAGGGAAATAAGCAAGAGGGCAATCGCCTCCTTGGAGGCTCTCGAGACGATATAT

156/392

FIGURE 150

MKFQGPLACLLIALCLGSGEAGPLQSGEESTGTNIGEALGHGLGDALSEGVGKAIGKEAG
GAAGSKVSEALGQGTREAVGTGVRQVPFGAADALGNRVGEAAHALGNTGHEIGRQAEDV
IRHGADAVRGSWQGVPGHSGAWETSGGHGIFGSQGGGLGGQGNPGLGTPWVHGYPGNS
AGSFGMNPQGAPWGQGGNNGPPNFGTNTQGAVAQPGYGSVRASNQNEGCTNPPPSGSGGG
SSNSGGSGSGSQSGSSGSGSNNGDNNNGSSSGSSSGSSSGSSSGSSSGSSSGSSSGNSGGS
RGDSGSESSWGSSTGSSSGNHGSGGGNGHKPGCEKPGNEARGSGESGIQGFRGQGVSSN
MREISKEGNRLGSGSDNYRGQSSWGSGGDAVGGVNTVNSETSPGMFNFDTFWKNFKS
KLGFINWDAINKDQRSSRIP

Signal peptide:
amino acids 1-21

N-glycosylation site:
amino acids 265-269

Glycosaminoglycan attachment site:
amino acids 235-239, 237-241, 244-248, 255-259, 324-328,
388-392

Casein kinase II phosphorylation site:
amino acids 26-30, 109-113, 259-263, 300-304, 304-308

N-myristoylation site:
amino acids 17-23, 32-38, 42-48, 50-56, 60-66, 61-67, 64-70,
74-80, 90-96, 96-102, 130-136, 140-146, 149-155, 152-158,
155-161, 159-165, 163-169, 178-184, 190-196, 194-200,
199-205, 218-224, 236-242, 238-244, 239-245, 240-246,
245-251, 246-252, 249-252, 253-259, 256-262, 266-272,
270-276, 271-277, 275-281, 279-285, 283-289, 284-290,
287-293, 288-294, 291-297, 292-298, 295-301, 298-304,
305-311, 311-317, 315-321, 319-325, 322-328, 323-329,
325-331, 343-349, 354-360, 356-362, 374-380, 381-387,
383-389, 387-393, 389-395, 395-401

Cell attachment sequence:
amino acids 301-304

157/392

FIGURE 151

CGGCCACAGCTGGCATGCTCTGCCTGATCGCCATCCTGCTGTATGTCTCGTCCAGTACC
TCGTGAACCCCGGGGTGCTCCGCACGGACCCCAGATGTCAAGAATATGAACACGTGGCTG
CTGTTCCCTCCCCCTGTTCCCGGTGCAGGTGCAGACCCTGATAGTCGTGATCATCGGGATG
CTCGTGCTCCTGCTGGACTTTCTTGGCTTGGTGCACCTGGGCCAGCTGCTCATCTTCCAC
ATCTACCTGAGTATGTCCCCCACCTAAGCCCCCGATCCCCCAAGGCTGGGTGGTCCAGA
GCTGCTCATCTTACACCTCTACTTGAGTATGTCCCTAACCCTGAGCCCCCACGCCTGGG
GCCAGAGTCTTTGTCCCCGTGTGCGCATGTGTTCCAGGTGAGCCTCTCCAGAAGTGAG
ATCATGGACAAAAAGGGCAAATCACAGGAAGAAATTAAATCCATGAGGACCCAGCAGGCC
CAGCAAGAAGCTGAACTCACGCCGAGACCTGCAGGAGTGGTGCCAGGTGCTTGAAGTAAC
AAGTTTAAATGTTTCAGAGACAATGGAATGGAATCTATTAGGCAAGAACAGGACATTATG
AAATAAGGACAGGTGGACTTCCAAAAACACAAGTAGAAATTCTAACAAATGAAATATATTA
CAGGCAGGTCACCCATAACCAACAACCTGAAGCGAGAGCTGTGGTCTTGCTTGGTCTCA
CAGTGGGCACAGCGGTAGGCGGTGAGTCATGTTGCTGAACGACGGAGGGTAAACTCCCCA
GCCCCAAGAAAACCTGTGTTGGAAGTAACAACAACCTCCCTGCTCCTGGCACCAGCCGTT
TTGGTCATGGTGGGCCAGCTGCAAAGCGTCTTCCATTCTCTGGGCAGTGGTGGCCCCGAG
GCTGTGGCCTCTCAGGGGGTTTCTGTGGACACGGGCAGCAGAGTGTGTCCAGGCCAGCCC
CCAAGAATGCCCTGCTCCTGACAGCTTGGCCAACCCCTGGTCAGGGCAGAGGGAGTTGGG
TGGGTGAGGCTCTGGGCTCACCTCCATCTCCAGAGCATCCCCTGCCTGCAGTTGTGGCAA
GAACGCCCAGCTCAGAATGAACACACCCACCAAGAGCCTCCTTGTTTATAACACAGGT
TACCCTACAAACCACTGTCCCCACACAACCTGGGGATGTTTTAAACACACACCTCTAA
CGCATATCTTACAGTCACTGTTGTCTTGCTGAGGGTTGAATTTTTTTTAAATGAAAGTGC
AATGAAAATCACTGGATTAAATCCTACGGACACAGAGCTGAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAA

158/392

FIGURE 152

MNTWLLFLPLFPVQVQTLIVVIIGMLVLLLDLGLVHLGQLLIFHIYLSMSPTLSPRSPQ
GWVVRAAHLTPLEYPNPEPPTPGARVFVPRVRMCSGSASPRSEIMDKKGKSQEEIKSM
RTQQAQQEALTPRPAGVVPGA

159/392

FIGURE 153

AACTGGAAGGAAAGAAAGAAAGGTCAGCTTTGGCCCAGATGTGGTTACCCCTTGGTCTCC
TGTCTTTATGTCTTTCTCCTCTTCCTATTCTGTTCATCTCCCTCACTTAAGTCTCAGGCCT
GTCAGCAGCTCCTGTGGACATTGCCATCCCCCTCTGGTAGCCTTCAGAGCAAACAGGACAA
CCTATGTTATGGATGTTTCCACCAACCAGGGTAGTGGCATGGAGCACCGTAACCATCTGT
GCTTCTGTGATCTCTATGACAGAGCCACTTCTCCACCTCTGAAATGTTCCCTGCTCTGAA
ATCTGGCATGAGATGGCACAGGTGACCACGCAGAAGCCACCAGAATCTTGCCTGCCCTAT
TCCTCCTCCCAAGTCTGTTCTCTTATTGTCAACCTCAGCACAAACAGGCTGGCGCCAATGG
CATTACAGAGAAAGCAATCTGTGTGGCTAGTGGGCAGATTACCATGCAAGCCCCAGGAGA
AATGGAGGAGCTTTGTAGCCACCTCCCTGTCAGCCAGTATTAACATGTCCCTTCCCCCT
GCCCCGCCGTAGATTACAGGACATTGCCCCCTGTGTGCCACCAACCAGGACTTCCCCCT
GGCTTGGCATCCCTGGCTCTCTCCTGGTACCCAGCAAGACGTCTGTTCCAGGGCAGTGTA
GCATCTTTCAAGCTCCGTTACTATGGCGATGGCCATGATGTTACAATCCCACTTGCCTGA
ATAATCAAGTGGGAAGGGGAAGCAGAGGGAAATGGGGCCATGTGAATGCAGCTGCTCTGT
TCTCCCTACCTGAGGAAAAACCAAGGGGAAGCAACAGGAACCTTCTGCAACTGGTTTTTA
TCGGAAGATCATCCTGCCTGCAGATGCTGTTGAAGGGGCACAAGAAATGTAGCTGGAGA
AGATTGATGAAAGTGCAGGTGTGTAAGGAAATAGAACAGTCTGCTGGGAGTCAGACCTGG
AATTCTGATTCCAACTCTTTATTACTTTGGGAAGTCACTCAGCCTCCCCGTAGCCATCT
CCAGGTGACGGAACCCAGTGTATTACCTGCTGGAACCAAGGAACTAACAATGTAGGTT
ACTAGTGAATACCCCAATGGTTTCTCCAATTATGCCCATGCCACCAAACAATAAAACAA
AATTCTCTAACACTGAAA

160/392

FIGURE 154

MWLPLGLLSLCLSPILSSPSLKSQACQQLWTLPSPLVAFRANRTTYVMDVSTNQSG
MEHRNHLCFCDLYDRATSPPLKCSLL

161/392

FIGURE 155

GTAGCGCGCTCTTGGGTCTCCCGGCTGCCGCTGCTGCCGCCGCCGCCCTCGGGTCGTGGAGC
CAGGAGCGACGTACCCGCCATGGCAGGCATCAAAGCTTTGATTAGTTTGTCTTTGGAGG
AGCAATCGGACTGATGTTTTTGTATGCTTGGATGTGCCCTTCCAATATACAACAAATACTG
GCCCCCTCTTTGTTCTATTTTTTTTACATCCTTTCACCTATTCCATACTGCATAGCAAGAAG
ATTAGTGGATGATACAGATGCTATGAGTAACGCTTGTAAGGAACCTTGCCATCTTTCTTAC
AACGGGCATTGTGTCGTGTCAGCTTTTGGACTCCCTATTGTATTGGCCAGAGCACATCTGAT
TGAGTGGGGAGCTTGTGCACCTGTTCTCACAGGAAACACAGTCATCTTTGCAACTATACT
AGGCTTTTTCTTGGTCTTTGGAAGCAATGACGACTTCAGCTGGCAGCAGTGGTGA¹AAAAGA
AATTACTGAACTATTGTCAAATGGACTTCCTGTCA²TTTGTGGCCATTACGCACACAGG
AGATGGGGCAGTTAATGCTGAATGGTATAGCAAGCCTCTTGGGGGTATTTTAGGTGCTCC
CTTCTCACTTTTATTGTAAGCATACTATTTTTCACAGAGACTTGCTGAAGGATTAAAAGGA
TTTTCTCTTTTGGAAAAGCTTGACTGATTTACACTTATCTATAGTATGCTTTTTTGTGGT
GTCCTGCTGAATTTAAATATTTATGTGTTTTTCTGTTAGGTTGATTTTTTTTTTGGAAATCA
ATATGCAATGTTAAACACTTTTTTAATGTAATCATTTGCATTGGTTAGGAATTCAGAATT
CCGCCGGCTCTATTACTGGTCAAGTACATCTTTTCTCTTAAAATTATTTAGCCTCCATTA
TTACAAAAAATTATAAAAAATAAGTTTTTCAGTCAGTCAGGATGACATCACTCCCAATGTTA
TGCAGACATACAGACGGTTGGCATACTGTTATAGACTGTATACTCAGTGCAAATATAGCTG
CATTTTATACCTCAGAGGGGGCCAAGTGTTAATGCCATGCCCTCCGTTAAGGGTTGTGGT
TTTACTGGTAGACAGATGTTTGTGGATTGAA³AATTATTTTATGGAATTGCTACGAGGA
GTGCTTTTCTTCTCAATTGTTTAGAAGAATTATGTTAAACTTTAAGGTAAAGGTGTAAAA
ACATTTTTTGAGATAAGGTTTTTATTTATGTTTATTTATTGTTAGAGTGAGTTGCAATGTGG
GAAGAAATGACATTGAAATTCCAGTTTTTGAATCCTGTTTCTATTTATAAGTGAAATTTG
TGATCTCCTATCAACCTTTTCATGTTTTACCCTGTTAAAATGGACATACATGGAACCACTA
CTGATGAGGGACAGTTGTATGTTTGCATCATATATGCCAGAAAACCTTCCTCTGCTTCCT
CCTTTTGA⁴CTTATTTGGTATGTTGTATATATTACATAAAATAACTTTTCAAATATAGTTT
AATAACACTTAGAAGTGTTTACTTACCTGGAAAATAATTGCTATGCCGTACATTACAGAT
GCCCCCTCCCCTGCAAGGCCTTGCCATGATTAA⁵CAAGTAACTTGTTAGTCTTACAGATAA
TTCATGCATTAAACAGTTTAA⁶GATTTAGACCATGGTAATAGTAGTTCCTTATTCTCTAAGGT
TATATCATATGTAATTTAAAGTATTTTTTAAGACAAGTTTCCTGTATACCTCTGAAC⁷TGT
TTTGATTTTTGAGTTCATCATGATAGATCTGCTGTTTCCTTATAAAAGGCATTTGTTGTGT
GAGTTAATGCAAAGTAGCCAAGTCCAGCTATATAGCAGCTTCAGAAACATACCTGACCAA
AAAAATTCCCAGTAACCAGGCATGATCAATTTATAGTGGTTCGTTTACATCTAATAATTATC
AGGACTTTTTTTCAGGAGTGGGTTATAAAAAACATTCAAGTTGGTCTGACAGTATTTTGTTA
AGGATATTTGTTTGTATGTTTATTCAGTATACTTACATAAAAAATTATTTCGCCATCAGCC
AAAAC⁸TAGTAATCATGACAGCTGTCTGTTGTTTTATGAAGTTTATTTCTCAAGAAAATG
GGAATAAATTG⁹GGATTGTTTCAGCTTTTTTACTAAAGATGCCTAAAGCCACAGGTTTTA
TTGCC¹⁰TAACTTAAGCCATGACTTTTAGATATGAGATGACGGGAAGCAGGACGAAATATCG
CGCTGTGGCTGGAGCCTTCCCCTGGAGGCTGAAAGTGGCTTGTTGGTATTATAATGTTCA
GATTTCAAGAGGAAGGTTCAGGTGACAGTACATGAGTTAGAGAGCTGGTGAGACAGTTGGGAAC
TCTTTGTGCTTGTGATCTACTGGACTTTTTTTTTTTCAGGAAGTGCATCTCTGGTCTCTC
CCTATTTTTCTGTTCTGGATGTCA¹¹GTGCAGTGCAGTGCATGCTACTGTTTTATCCACTTGGCCAC
AGACTTTTTTCTAACAGCTGCGTATTATTTCTATATACTAATTGCATTGGCAGCATTGTGT
CTTTGACCTTGTATACTAGCTTGACATAGTGTCTGTCTCTGATTTCTAGGCTAGTTACTTG
AGATATGAATTTTCCATAGAATATGCAC¹²TGATACAACATTACCATTCTTCTATGGAAAGA
AAACTTTTTGATGATGAAACAATAAAGATTTTAAATATCTATTTTAAAAA¹³AAAA

162/392

FIGURE 156

MAGIKALISLSFGGAIGLMFLMLGCALPIYNKYWPLFVLFFYILSPIPYCIARRLVDDTD
AMSNACKELAIFLTGTGIVVSAFGLPIVFARAHLEWGACALVLTGNTVIFATILGFFLVF
GSNDDFSQQW

163/392

FIGURE 157

GTTTCTCATAGTTGGCGTCTTCTAAAGGAAAAACACTAAAATGAGGAACTCAGCGGACCG
GGAGCGACGCAGCTTGAGGGAAGCATCCCTAGCTGTTGGCGCAGAGGGGCGAGGCTGAAG
CCGAGTGGCCCGAGGTGTCTGAGGGGCTGGGGCAAAGGTGAAAGAGTTTCAACAACAGCT
TCCTGGAACCCATGACCCATGAAGTCTTGTCGACATTTATACCGTCTGAGGGTAGCAGCT
CGAAACTAGAAGAAGTGGAGTGTGTCAGGGACGGCAGTATCTCTTTGTGTGACCCTGGC
GGCCTATGGGACGTTGGCTTCAGACCTTTGTGATACACCATGCTGCGTGGGACGATGACG
GCGTGGAGAGGAATGAGGCTGAGGTACACTGGCTTGCTCCTCCTAGCCACAGCAGGC
TGCTTTGCTGACTTGAACGAGGTCCCTCAGGTCACCGTCCAGCCTGCGTCCACCGTCCAG
AAGCCCGGAGGCACTGTGATCTTGGGCTGCGTGGTGGAACTCCAAGGATGAATGTAACC
TGGCGCCTGAATGGAAGGAGCTGAATGGCTCGGATGATGCTCTGGGTGTCTCATCACC
CACGGGACCCTCGTCATCACTGCCCTTAACAACCACTGTGGGACGGTACCAGTGTGTG
GCCCCGATGCTGCGGGGGCTGTGGCCAGCGTGCCAGCCACTGTGACACTAGCCAATCTC
CAGGACTTCAGATTAGATGTGCAGCAGTGATTGAAGTGGATGAGGGAAACACAGCAGTC
ATTGCCCTGCCACCTGCCTGAGAGCCACCCCAAAGCCAGGTCCGGTACAGCGTCAAACAA
GAGTGGCTGGAGGCCTCCAGAGGTAACCTGATCATGCCCTCAGGGAACCTCCAGATT
GTGAATGCCAGCCAGGAGGACGAGGGCATGTACAAGTGTGCAGCCTACAACCCAGTGACC
CAGGAAGTGAAAACCTCCGGCTCCAGCGACAGGCTACGTGTGCGCCGCTCCACCGCTGAG
GCTGCCCGCATCATCTACCCCCCAGAGGCCCAAACCATCATCGTCACCAAAGGCCAGAGT
CTCATTTCTGGAGTGTGTGGCCAGTGGAATCCCACCCCCACGGGTACCTGGGCCAAGGAT
GGGTCCAGTGTCAACGGCTACAACAAGGCTTGGCAGGATGCTGAGCAACCTCCTCATCGAC
ACCACCAGCGAGGAGGACTCAGGCACCTACCGCTGCATGGCCGACAATGGGGTTGGGCAG
CCCCGGGCGAGCGGTTCATCTCTACAATGTCCAGGTGTTTGAACCCCCCTGAGGTACCATG
GAGCTATCCCAGCTGGTTCATCCCCCTGGGGCCAGAGTGCCAAGCTTACCTGTGAGGTGCGT
GGGAACCCCCCGCCCTCCGTGCTGTGGCTGAGGAATGCTGTGCCCTCATCTCCAGCCAG
CGCCTCCGGCTCTCCCGCAGGGCCCTGCGCGTGCTCAGCATGGGGCCTGAGGACGAAGGC
GTCTACCAAGTGCATGGCCGAGAACGAGGTTGGGAGCGCCCATGCCGTAGTCCAGCTGCGG
ACCTCCAGGCCAAGCATAACCCCCAAGGCTTGTGTCAGGATGCTGAGCTGGCTACTGGCACA
CCTCCTGTATCACCTTCAAACCTCGGCAACCTTGAGCAGATGCTGAGGGGGCAACCGGCG
CTCCCCAGACCCCCAACGTCACTGGGGCCTGCTTCCCCGAAGTGTCCAGGAGAGAAGGGG
CAGGGGGCTCCCCCGGAGGCTCCCATCATCTCAGCTCGCCCCGCACCTCCAAGACAGAC
TCATATGAACTGGTGTGGCGGCCTCGGCATGAGGGCAGTGGCCGGGCGCCAATCCTCTAC
TATGTGGTGAAACACCGCAAGCAGGTCAAAATTCCTCTGACGATTGGACCATCTCTGGC
ATTCCAGCCAAACAGCACCGCCTGACCTTACCAGACTTGACCCCGGAGCTTGTATGAA
GTGGAGATGGCAGCTTACAACCTGTGCGGGAGAGGGCCAGACAGCCATGGTCACCTTCCGA
ACTGGACGGCGGCCCAAACCCGAGATCATGGCCAGCAAAGAGCAGCAGATCCAGAGAGAC
GACCCTGGAGCCAGTCCCCAGAGCAGCAGCCAGCCAGACCACGGCCGCTCTCCCCCCCCA
GAAGCTCCCGACAGGCCACCATCTCCACGGCCTCCGAGACCTCAGTGACCTGACCTGG
ATTCCCCGTGGGAATGGTGGGTTCCCAATCCAGTCCCTCCGTGTGGAGTACAAGAAGCTA
AAGAAAGTGGGAGACTGGATTCTGGCCACCAGCGCCATCCCCCATCGCGGCTGTCCGTG
GAGATCACGGGCCCTAGAGAAAGGCACCTCCTACAAGTTTCGAGTCCGGGCTCTGAACATG
CTGGGGGAGAGCGAGCCAGCGCCCCCTCTCGGCCCTACGTGGTGTGCGGGCTACAGCGGT
CGCGTGTACGAGAGGCCCCGTGGCAGGTCTTATATCACCTTACGGATGCGGGTCAATGAG
ACCACCATCATGCTCAAGTGGATGTACATCCAGCAAGTAACAACAACACCCCAATCCAT
GGCTTTTATATCTATTATCGACCCACAGACAGTGACAATGATAGTGAATAACAAGGAT
ATGGTGGAAAGGGGACAAGTACTGGCACTCCATCAGCCACCTGCAGCCAGAGACCTCCTAC
GACATTAAGATGCAGTGCTTCAATGAAGGAGGGGAGAGCGAGTTTCAACAACGTGATGATC
TGTGAGACCAAAGCTCGGAAGTCTTCTGGCCAGCCTGGTTCGACTGCCACCCCACTCTG
GCCCCACCACAGCGCGCCCTTCTGAAACCATAGAGCGCGCGGTGGGCATGGGCCATG
GTGGCTCGCTCCAGCGACCTGCCCTATCTGATTGTGCGGGTCTGCTGGGCTCCATCGTT
CTCATCATCGTCACCTTCATCCCCCTTCTGCTTGTGGAGGGCCTGGTCTAAGCAAAAACAT
ACAACAGACCTGGGTTTTCTCGAAGTGCCCTTCCACCTCCTGCCCCGTATACTATGGTG
CCATTGGGAGGACTCCCAGGCCACCAGGCCAGTGACAGCCCTACCTCAGTGGCATCAGT

164/392

GGACGGGCCTGTGCTAATGGGATCCACATGAATAGGGGCTGCCCCCTCGGCTGCAGTGGGC
TACCCGGGCATGAAGCCCCAGCAGCACTGCCCAGGCGAGCTTCAGCAGCAGAGTGACACC
AGCAGCCTGCTGAGGCAGACCCATCTTGGCAATGGATATGACCCCCAAAGTCACCAGATC
ACGAGGGGTCCCAAGTCTAGCCCGGACGAGGGCTCTTTCTTATACACACTGCCCCGACGAC
TCCACTCACCAGCTGCTGCAGCCCCATCACGACTGCTGCCAACGCCAGGAGCAGCCTGCT
GCTGTGGGCCAGTCAGGGGTGAGGAGAGCCCCGACAGTCCTGTCCTGGAAGCAGTGTGG
GACCCCTCCATTTCACTCAGGGCCCCCATGCTGCTTGGGCCTTGTGCCAGTTGAAGAGGTG
GACAGTCCTGACTCCTGCCAAGTGAGTGGAGGAGACTGGTGTCCCCAGCACCCCGTAGGG
GCCTACGTAGGACAGGAACCTGGAATGCAGCTCTCCCCGGGGCCACTGGTGCGTGTGTCT
TTTGAAACACCACCTCTCACAATTTAGGCAGAAGCTGATATCCAGAAAGACTATATATT
GTTTTTTTTTTAAAAAAAAGAGAGAAAAAGAGACAGAGAAAATTGGTATTTATTTTC
TATTATAGCCATATTTATATATTTATGCACTTGTAATAAATGTATATGTTTTATAATTC
TGGAGAGACATAAGGAGTCCTACCCGTTGAGGTTGGAGAGGGAAAATAAAGAAGCTGCCA
CCTAACAGGAGTCACCCAGGAAAGCACCGCACAGGCTGGCGCGGGACAGACTCCTAACCT
GGGGCCTCTGCAGTGGCAGGCGAGGCTGCAGGAGGCCACAGATAAGCTGGCAAGAGGAA
GGATCCCAGGCACATGGTTCATCACGAGCATGAGGGAACAGCAAGGGGCACGGTATCACA
GCCTGGAGACACCCACACAGATGGCTGGATCCGGTGCTACGGGAAACATTTTCCTAAGAT
GCCCATGAGAACAGACCAAGATGTGTACAGCACTATGAGCATTAAAAACCTTCCAGAAT
CAATAATCCGTGGCAACATATCTCTGTAAAAACAACACTGTAACCTCTAAATAAATGTT
TAGTCTTCCCTGTAAAA

165/392

FIGURE 158

MLRGTMATAWRGMRPEVTLACLALLATAGCFADLNEVPQVTVQPASTVQKPGGTIVILGCVVE
PPRMNVTWRLNGKELNGSDDALGVLITHGTLVITALNNHTVGRYQCVARMPAGAVASVPA
TVTLANLQDFKLDVQHVIEVDEGNTAVIACHLPESHKPAQVRYSVKQEWLEASRGNYLIM
PSGNLQIVNASQEDEGMYKCAAYNPVTQEVKTSGSSDRLRVRRSTAEAAARI IYPPEAQTI
IVTKGQSLILECVASGI PPPRVTWAKDGSSVTGYNKTRFLLSNLLIDTTSEEDSGTYRCM
ADNGVGQPGAAVILYNVQVFEPPEVTMELSOLVIPWGQSAKLTCEVRGNPPPSVLWLRNA
VPLISSQRLRLSRRALRVLSMGPEDEGVYQCMANEVGSAAHAVVQLRTSRPSITPRLWQD
AELATGTPPVSPSKLGNPEQMLRGQPALPRPPTS VGPASPKCPGEGKGQGAPEAPIILSS
PRTSKTDSYELVWRPRHEGSGRAPILYYVVKHRKQVTNSSDDWTISGIPANQHRLTLTRL
DPGSLYEVEMAAYNCAGEGQTAMVTFRTGRRPKPEIMASKEQQIQRDDPGASPOSSSQPD
HGRSLSPPEAPDRPTISTASETSVYVTWIPRGNGGFPIQSFRVEYKKLKKVGDWILATSAI
PPSRLSVEITGLEKGT SYKFRVRALNMLGESEPSAPSRPYVVS GYSGRVYERPVAGPYIT
FTDAVNETTIMLKWMIYIPASNNNTPIHGFIYYRPTSDNDSDYKKDMVEGDKYWHSISH
LQPETSYDIKMQCFNEGGESEFSNVMICETKARKSSGQPGRLPPPTLAPPQPLPETIER
PVG TGAMVARSSDLPYLIVGVVLGSIVLIIVTFIPFCLWRAW SKQKHTTDLGFP RSALPP
SCPYTMVPLGGLPGHQASGQPYLSGISGRACANGIHMNRGCPSAAVGYPGMKPQQHCPGE
LQQQSDTSSLLRQTHLGNGYDPQSHQITRGP KSSPDEGSFLYTL PDDSTHQLLQPHHDC
QRQEQPAAVGQSGVRRAPDSPVLEAVWDPPFHS GPPCCLGLVPVEEVDSPDSCQVSGGDW
CPQHVPV GAYVGQEPGMQLSPGPLVRVVSFETPPLTI

Signal peptide:

amino acids 1-30

Transmembrane domain:

amino acids 16-30 (type II), 854-879

166/392

FIGURE 159

CCCACGCGTCCGCCCACGCGTCCGCCCACGCGTCCGCCCACGCGTCCGCCCACGCGTCCG
CCCACGCGTCCGCCCACGCGTCCGGTGCAAGCTCGCGCCGCACACTGCCTGGTGAGGGGA
AGGAGCCCCGGGCGCCTCTCGCCGCTCCCCGCGCCGCGTCCGCACCTCCCCACCGCCCGC
CGCCCGCCGCCCCGCGCCCGCAAAGCATGAGTGAGCCCGCTCTCTGCAGCTGCCCGGGGC
GCGAATGGCAGGCTGTTTCCGCGGAGTAAAAGGTGGCGCCGGTCAGTGGTCGTTTCCAAT
GACGGACATTAACCAGACTGTCAGATCCTGGGGAGTCGCGAGCCCCGAGTTTGGAGTTTT
TTCCCCCACAACGTCACAGTCCGAACCTGCAGAGGGAAAGGAAGGCGGCAGGAAGGCGAA
GCTCGGGCTCCGGCACGTAGTTGGGAAACTTGCGGGTCCTAGAAGTCGCCTCCCCGCCTT
GCCGCGCCGCTTGACAGCCCCGAGCCGAGCAGCAAAGTGAGACATTGTGCGCCTGCCAGA
TCCGCGCGCCGCGGACCGGGGCTGCCTCGGAAACACAGAGGGGTCTTCTCTCGCCCTGCA
TATAATTAGCCTGCACACAAAGGGAGCAGCTGAATGGAGGTTGTCACTCTCTGGAAAAGG
ATTTCTGACCGAGCGCTTCCAATGGACATTCTCCAGTCTCTCTGGAAAGATTCTCGCTAA
TGGATTTCTGCTGCTCGGTCTCTGTCTATACTGGCTGCTGAGGAGGCCCTCGGGGGTGG
TCTTGTGTCTGCTGGGGGCTGCTTTCAGATGCTGCCCGCCGCCCCCAGCGGGTGCCCGC
AGCTGTGCCGGTGCGAGGGGCGGCTGCTGTACTGCGAGGCGCTCAACCTCACCGAGGCGC
CCCACAACCTGTCCGGCTGCTGGGCTTGTCCCTGCGCTACAACAGCCTCTCGGAGCTGC
GCGCCGGCCAGTTCACGGGGTTAATGCAGCTCACGTGGCTCTATCTGGATCACAATCACA
TCTGCTCCGTGCAGGGGGACGCCTTTCAGAACTGCGCCGAGTTAAGGAACTCACGCTGA
GTTCCAACAGATCACCCAACCTGCCCAACACCACTTCCGGCCCCATGCCCAACCTGCGCA
GCGTGGACCTCTCGTACAACAAGCTGCGAGGCGCTCGCGCCCGACCTCTTCCACGGGCTGC
GGAAGCTCACCAAGCTGCATATGCGGGCCAACGCCATCCAGTTTGTGCCCCGTGCGCATCT
TCCAGGACTGCCGACGCTCAAGTTTCTCGACATCGGATACAATCAGCTCAAGAGTCTGG
CGCGCAACTCTTTCGCGGGCTTGTTTAAGCTCACCGAGCTGCACCTCGAGCACAACGACT
TGGTCAAGGTGAACTTCGCCCCACTTCCCGCGCCTCATCTCCCTGCACTCGCTCTGCCTGC
GGAGGAACAAGGTGGCCATTGTGGTCAGCTCGCTGGACTGGGTTTGGAACTGGAGAAAA
TGGACTTGTGCGGCAACGAGATCGAGTACATGGAGCCCCATGTGTTGAGACCGTGCCGC
ACCTGCAGTCCCTGCAGCTGGACTCCAACCGCCTCACCTACATCGAGCCCCGGATCCTCA
ACTCTTGGAAGTCCCTGACAAGCATCACCTGGCCGGGAACCTGTGGGATTGCGGGCGCA
ACGTGTGTGCCCTAGCCTCGTGGCTCAGCAACTTCCAGGGGCGCTACGATGGCAACTTGC
AGTGCGCCAGCCCGGAGTACGCACAGGGCGAGGACGTCCTGGACGCCGTGTACGCCCTTCC
ACCTGTGCGAGGATGGGGCCGAGCCACCAGCGGCCACCTGCTCTCGGCCGTACCAACC
GCAGTGATCTGGGGCCCCCTGCCAGCTCGGCCACCACGCTCGCGGACGGCGGGGAGGGGC
AGCACGACGGCACATTCGAGCCTGCCACCGTGGCTCTTCCAGGCGGCGAGCACGCCGAGA
ACGCCGTGCAGATCCACAAGGTGGTCACGGGCACCATGGCCCTCATCTTCTCCTTCTCTCA
TCGTGGTCTTGGTGCTCTACGTGTCTTGGAAAGTGTTCCTCAGCCAGCCTCAGGCAGCTCA
GACAGTGCTTTGTACGCAGCGCAGGAAGCAAAAGCAGAAACAGACCATGCATCAGATGG
CTGCCATGTCTGCCAGGAATACTACGTTGATTACAAACCGAACCACATTGAGGGAGCCC
TGGTGATCATCAACGAGTATGGCTCGTGACCTGCCACCAGCAGCCCGCGAGGGAATGCG
AGGTGTGATTGTCCAGTGGCTCTCAACCATGCGCTACCAAATACGCCTGGGCAGCCGG
GACGGGCCGGCGGGCACCAGGCTGGGGTCTCCTTGTCTGTGCTCTGATATGCTCCTTGAC
TGAACCTTTAAGGGGATCTCTCCAGAGACTTGACATTTTAGCTTTATTGTGTCTTAAAC
ACAAAAGCGAATTAAAAACACAACAAAAACCCACCCACACCTTCAGGACAGTCTATC
TTAAATTTTCATATGAGAACTCCTTCTCCTTTTGAAGATCTGTCCATATTAGGAATCTG
AGAGTGTAATAAAGGTGGCCATAAGACAGAGAGAGAATAATCGTGCTTTTGTTTTATGCTA
CTCCTCCACCCCTGCCCATGATTAAACATCATGTATGTAGAAAGATCTTAAGTCCATACGC
ATTTTCATGAAGAACCATTGGAAAGAGGAATCTGCAATCTGGGAGCTTAAGAGCAAATGAT

167/392

GACCATAGAAAGCTATGTTCTTACTTTGTGTGTGTGTCTGTATGTTTCTGCGTTGTGTGT
CTTTGTAGGCAAGCAAACGTTGTCTACACAAACGGGAATTTAGCTCACATCATTTTCATGC
CCCTGTGCCTCTAGCTCTGGAGATTGGTGGGGGGAGGTGGGGGGAAACGGCAGGAATAAG
GGAAAGTGGTAGTTTTAACTAAGGTTTTGTAACTTGAATCTTTTCTTCTCAAATTA
ATTATCTTTAAGCTTCAAGAACTTGCTCTGACCCCTCTAAGCAAATACTAAGCATTTA
AAAGAGAATCTAATTTTTAAAGGTGTAGCACCTTTTTTTTTTATTCTTCCCACAGAGGGTG
CTAATCTCATTATGCTGTGCTATCTGAAAAGAACTTAAGGCCACAATTCACGTCTCGTCC
TGGGCATTGTGATGGATTGACCCTCCATTTGCAGTACCTTCCCAGCTGATTAAAGTTCAG
CAGTGGTATTGAGGTTTTTTCGAATATTTATATAGAAAAAAAGTCTTTTTCACATGACAAAT
GACACTCTCACACCAGTCTTAGCCCTAGTAGTTTTTTTAGGTTGGACCAGAGGAAGCAGGT
TAAATGAGACCTGTCCTCTGCTGCACTCAGAAAAAATAGGCAGTCCCTGATGCTCAGATC
TTAGCCTTGATATTAATAGTTGAGACCACCTACCCACAATGCAGCCTATACTCCCAAGAC
TACAAAGTTACCATCGCAAAGGAAAGGTTATTCCAGTAAAAGGAAATAGTTTTCTCAACC
ATTTAAAAATATTCTTCTGAACTCATCAAAGTAGAAGAGCCCCAACCTTTTCTCTCTGC
CTTCAAGAAGGCAGACATTTGGTATGATTTAGCATCAACAACACATTTATGAGTATATGT
AAGTAATCAGAGGGGCAAATGCCACTTGTTATTCCTCCCAAGTTTCCAAGCAAGTACAC
ACAGATCTCTGGTAGGATTAGGGGCCACTTGTGTTTCCGGCTTATTTTAGTCGACTTGTC
AGCAAGTTTGATGCCTAGTCTATCTGACATGGCCAGTAGAACAGGGCATTGATGGATCA
CATGAGATGGTAGAAGGAACATCATCACATACCCCTCTCACAGAGAAAATTATCAAAGAA
CCAGAAATTATATCTGTTTTGGAGCAAGAGTGTCATAATGTTTCAGGGTAGTCAAATAA
ACATAAATTATCTCCTCTAGATGAGTGGCGATGTTGGCTGATTTGGGTCTGCCATTGACA
GAATGTCAAATAAAAAGGAATTAGCTAGAATATGACCATTAAATGTGCTTCTGAAATATA
TTTTGAGATAGGTTTAGAATGTCA

168/392

FIGURE 160

MDFLLLGLCLYWLLRRPSGVVLCLLGACFQMLPAAPSGCPQLCRCEGRLLYCEALNLTEA
PHNLSGLLGLSLRYNSLSELRAGQFTGLMQLTWLYLDHNNHICSVQGDAFQKLRRVKELTL
SSNQITQLPNTTFRPMPNLRSDLSYNKLQALAPDLFHGLRKLTTLHMRANAIQFVPVRI
FQDCRSKFLDIGYNQLKSLARNSFAGLFKLTELHLEHNDLVKVNFAHFPRILISLHSLCL
RRNKVAIVVSSLDWVWNLEKMDLSGNEIEYMEPHVFETVPHLQSLQLDSNRLTYIEPRIL
NSWKSLSITLAGNLWDCGRNVCALASWLSNFQGRYDGNLQCASPEYAQGEDVLDAVYAF
HLCEDGAEPTSGHLLSAVTNRSDLGPPASSATTADGGEGQHDGTFEPATVALPGGEHAE
NAVQIHKVVTGTMALIFSFLIVVLVLYVSWKCFPASLRQLRQCFVTQRRKQKQKQTMHQM
AAMSAQEYYVDYKPNHIEGALVIINEYGSCTCHQQPARECEV

170/392

FIGURE 162

MAPQSLPSSRMAPLGMLLGLLMAACFTFCLSHQNLKEFALTNPESSTKETERKETKAEE
ELDAEVLEVFPHTHEWQALQPGQAVPAGSHVRLNLQTGEREAKLOYEDKFRNNLKGKRLD
INTNTYTSQDLKSALAKFKEGAEMESSKEDKARQAEVKRLFRPIEELKKDFDELNVVIET
DMQIMVRLINKFNSSSSSLEEKIAALFDLEYVYVHQMDNAQDLLSFGLQVINGLNSTEP
LVKEYAAFVLGAAFSSNPKVQVEAIEGGALQKLLVILATEQPLTAKKKVLFALCSLLRHF
PYAQRQFLKLGGLQVLRRTLVOEKGTEVLAVRVVTTLLYDLVTEKMFEEEEAELTQEMSPEK
LQQYRQVHLLPGLWEQGWCEITAHLLALPEHDAREKVLQTLGVLLTTCRDRYRQDPQLGR
TLASLQAEYQVLASLELQDGEDEGYFQELLGSVNSLLKELR

Important features:**Signal peptide:**

amino acids 1-29

Hypothetical YJL126w/YLR351c/yhcX family protein:

amino acids 364-373

N-glycosylation site:

amino acids 193-197, 236-240

N-myristoylation site:

amino acids 15-21, 19-25, 234-240, 251-257, 402-408, 451-457

Homologous region SLS1 protein:

amino acids 68-340

171/392

FIGURE 163

CAGAGAGGAGGCTTTGGGAATTGTCCAGCAGAAACAGAGAAGTCTGAGGTGGTGTCAAGA
CAAAAGATGCTTCAGCTTTGGAACTTGTTCTCCTGTGCGGCGTGCTCACTGGGACCTCA
GAGTCTCTTCTTGACAATCTTGGAATGACCTAAGCAATGTCTGGGATAAGCTGGAACCT
GTTCTTCACGAGGGACTTGAGACAGTTGACAATACTCTTAAAGGCATCCTTGAGAACTG
AAGGTCGACCTAGGAGTGCTTCAGAAATCCAGTGCTTGGCAACTGGCCAAGCAGAAGGCC
CAGGAAGCTGAGAAATTGCTGAACAATGTCATTTCTAAGCTGCTTCCAATAACACGGAC
ATTTTTGGGTGAAAATCAGCAACTCCCTCATCCTGGATGTCAAAGCTGAACCGATCGAT
GATGGCAAAGGCCTTAACCTGAGCTTCCCTGTCACCGCGAATGTCACTGTGGCCGGGCCC
ATCATTGGCCAGATTATCAACCTGAAAGCCTCCTTGGACCTCCTGACCGCAGTCACAATT
GAAACTGATCCCCAGACACACCAGCCTGTTGCCGTCTTGGGAGAATGCGCCAGTGACCCA
ACCAGCATCTCACTTTCCTTGCTGGACAAACACAGCCAAATCATCAACAAGTTCGTGAAT
AGCGTGATCAACACGCTGAAAAGCACTGTATCCTCCCTGCTGCAGAAGGAGATATGTCCA
CTGATCCGCATCTTCATCCACTCCCTGGATGTGAATGTCATTGAGCAGGTCGTCGATAAT
CCTCAGCACAAAACCCAGCTGCAAACCCTCATCTGAAGAGGACGAATGAGGAGGACCACT
GTGGTGCATGCTGATTGGTTCCAGTGGCTTGCCCCACCCCCTTATAGCATCTCCCTCCA
GGAAGCTGCTGCCACCACCTAACCAGCGTGAAAGCCTGAGTCCCACCAGAAGGACCTTCC
CAGATACCCCTTCTCCTCACAGTCAGAACAGCAGCCTCTACACATGTTGTCTGCCCCTG
GCAATAAAGGCCCATTTCTGCACCCTTAA

172/392

FIGURE 164

MLQLWKLVLLCGVLTGTSESLLDNLGNDLSNVVDKLEPVLHEGLETVDNLTGKILEKLKV
DLGVLQKSSAWQLAKQKAQEA EKLLNNVISKLLPTNTDIFGLKISNSLILDVKAEPIDDG
KGLNLSFPVTANVTVAGPIIGQIINLKASLDLLTAVTIETDPQTHQPVAVLGECASDPTS
ISLSLLDKHSQIINKFVNSVINTLKSTVSSLLQKEICPLIRIFIHSLDVNVIQQVVDNPQ
HKTQLQTLI

Important features:

Signal peptide:

1-15

Transmembrane domain:

none

N-glycosylation site:

124-128, 132-136

N-myristoylation site:

12-18, 16-22, 26-32, 101-107, 122-128, 141-147

Leucine zipper pattern:

44-66

173/392

FIGURE 165

GCAGTCAGAGACTTCCCCTGCCCCCTCGCTGGGAAAGAACATTAGGAATGCCTTTTAGTGCT
CTTGCTTCCTGAACTAGCTCACAGTAGCCCCGGCGCCAGGGCAATCCGACCACATTTCA
CTCTCACCGCTGTAGGAATCCAGATGCAGGCCAAGTACAGCAGCACGAGGGACATGCTGG
ATGATGATGGGGACACCACCATGAGCCTGCATTCTCAAGCCTCTGCCACAACCTCGGCATC
CAGAGCCCCGGCGCACAGAGCACAGGGCTCCCTCTTCAACGTGGCGACCAGTGGCCCTGA
CCCTGCTGACTTTGTGCTTGGTGCTGCTGATAGGGCTGGCAGCCCTGGGGCTTTTGTTTT
TTCAGTACTACCAGCTCTCCAATACTGGTCAAGACACCATTTCTCAAATGGAAGAAAGAT
TAGGAAATACGTCCCAAGAGTTGCAATCTCTTCAAGTCCAGAATATAAAGCTTGCAGGAA
GTCTGCAGCATGTGGCTGAAAACTCTGTGCTGAGCTGTATAACAAAGCTGGAGCACACA
GGTGCAGCCCTTGTACAGAACAATGGAAATGGCATGGAGACAATTGCTACCAGTTCTATA
AAGACAGCAAAAAGTTGGGAGGACTGTAAATATTTCTGCCTTAGTGAAAACCTACCATGC
TGAAGATAAAACAAGAAGACCTGGAATTTGCCGCGTCTCAGAGCTACTCTGAGTTTT
TCTACTCTTATTGGACAGGGCTTTTGCGCCCTGACAGTGGCAAGGCCTGGCTGTGGATGG
ATGGAACCCCTTTCACCTCTGAACGTGTCCATATTATAATAGATGTCACCAGCCCAAGAA
GCAGAGACTGTGTGGCCATCCTCAATGGGATGATCTTCTCAAAGGACTGCAAAGAATTGA
AGCGTTGTGTCTGTGAGAGAAGGGCAGGAATGGTGAAGCCAGAGAGCCTCCATGTCCCCC
CTGAAACATTAGGCGAAGGTGACTGATTCGCCCTCTGCAACTACAAATAGCAGAGTGAGC
CAGGCGGTGCCAAAGCAAGGGCTAGTTGAGACATTGGGAAATGGAAATAATCAGGAAAG
ACTATCTCTCTGACTAGTACAAAATGGGTTCTCGTGTTTCCTGTTCCAGGATCACCAGCAT
TTCTGAGCTTGGGTTTATGCACGTATTTAACAGTCACAAGAAGTCTTATTTACATGCCAC
CAACCAACCTCAGAAACCCATAATGTCATCTGCCTTCTTGGCTTAGAGATAACTTTTAGC
TCTCTTTCTTCTCAATGTCTAATATCACCTCCCTGTTTTCATGTCTTCCTTACACTGGT
GGAATAAGAACTTTTTGAAGTAGAGGAAATACATTGAGGTAACATCCTTTTCTCTGACA
GTCAAGTAGTCCATCAGAAATTGGCAGTCACTTCCAGATTGTACCAGCAAATACACAAG
GAATTCTTTTTGTGTTGTTTCAGTTCATACTAGTCCCTTCCCAATCCATCAGTAAAGACCC
CATCTGCCTTGTCCATGCCGTTTCCCAACAGGGATGTCACCTTGATATGAGAATCTCAAAT
CTCAATGCCTTATAAGCATTCCCTTCTGTGTCCATTAAAGACTCTGATAATTGTCTCCCT
CCATAGGAATTTCTCCAGGAAAGAAATATATCCCCATCTCCGTTTCATATCAGAACTAC
CGTCCCCGATATTCCCTTCCAGAGAGATTAAAGACCAGAAAAAAGTGAGCCTCTTCATCTG
CACCTGTAATAGTTTCAGTTCCTATTTTCTTCCATTGACCCATATTTATACCTTTCAGGT
ACTGAAGATTTAATAATAATAAATGTAAATACTGTGAAAAA

174/392

FIGURE 166

MQAKYSSTRDMLDDDGDTTMSLHSQASATTRHPEPRRTEHRAPSSTWRPVALTLLTLCLV
LLIGLAALGLLFFQYYQLSNTGQDTISQMEERLGNTSQELQSLQVQNIKLAGSLQHVAEK
LCRELYNKAGAHRCSPCTEQWKWHGDNCYQFYKDSKSWEDCKYFCLSENSTMLKINKQED
LEFAASQSYSEFFYSYWTGLLRPDSGKAWLWMDGTPFTSELFHIIIDVTSPRSRDCVAIL
NGMIFSKDCKELKRCVCERRAGMVKPESLHVPPETLGEGD

175/392

FIGURE 167

GC GACGGG CAGGACGCCCCGTT CGCCTAGCGCGTGCTCAGGAGTTGGTGTCTGCCTGCG
CTCAGGATGAGGGGGAATCTGGCCCTGGTGGGCGTTCTAATCAGCCTGGCCTTCCTGTCA
CTGCTGCCATCTGGACATCCTCAGCCGGCTGGCGATGACGCCTGCTCTGTGCAGATCCTC
GTCCCTGGCCTCAAAGGGGATGCGGGAGAGAAGGGAGACAAAGGCGCCCCCGGACGGCCT
GGAAGAGTCGGCCCCACGGGAGAAAAAGGAGACATGGGGGACAAAGGACAGAAAGGCAGT
GTGGGTTCGT CATGGAAAAATTGGTCCCATTGGCTCTAAAGGTGAGAAAAGGAGATTCCGGT
GACATAGGACCCCCCTGGTCCTAATGGAGAACCAGGCCTCCCATGTGAGTGCAGCCAGCTG
CGCAAGGCCATCGGGGAGATGGACAACCAGGTCTCTCAGCTGACCAGCGAGCTCAAGTTC
ATCAAGAATGCTGTGCGCGGTGTGCGCGAGACGGAGAGCAAGATCTACCTGCTGGTGAAG
GAGGAGAAGCGCTACGCGGACGCCCAGCTGTCTGCCAGGGCCGCGGGGGCACGCTGAGC
ATGCCCAAGGACGAGGCTGCCAATGGCCTGATGGCCGCATACCTGGCGCAAGCCGGCCTG
GCCCCGTGTCTTCATCGGCATCAACGACCTGGAGAAGGAGGGCGCCTTCGTGTACTCTGAC
CACTCCCCCATGCGGACCTTCAACAAGTGGCGCAGCGGTGAGCCCAACAATGCCTACGAC
GAGGAGGACTGCGTGGAGATGGTGGCCTCGGGCGGCTGGAACGACGTGGCCTGCCACACC
ACCATGTACTTTCATGTGTGAGTTTGACAAGGAGAACATGTGAGCCTCAGGCTGGGGCTGC
CCATTGGGGGCCCCACATGTCCCTGCAGGGTTGGCAGGGACAGAGCCCAGACCATGGTGC
CAGCCAGGGAGCTGTCCCTCTGTGAAGGGTGGAGGCTCACTGAGTAGAGGGCTGTTGTCTC
AAACTGAGAAAATGGCCTATGCTTAAGAGGAAAATGAAAGTGTTCTGGGGTGCTGTCTC
TGAAGAAGCAGAGTTTCATTACCTGTATTGTAGCCCCAATGTCATTATGTAATTATTACC
CAGAATTGCTCTTCCATAAAGCTTGTGCCTTTGTCCAAGCTATACAATAAAATCTTTAAG
TAGTGCAGTAGTTAAGTCCAAAAAAAAAAAAAAAAAAAAA

176/392

FIGURE 168

MRGNLALVGVLI SLAFLSLLPSGHPQPAGDDACSVQILVPGLKGDAGEKGDKGAPGRPGR
VGPTGEKGDMDKGQKGSVGRHGKIGPIGSKGEKGDSDIGPPGPNGEPGLPCECSQLRK
AIGEMDNQVSQLTSELKFIKNAVAGVRETESKIYLLVKEEKRYADAQLSCQGRGGTLSMP
KDEAANGLMAAYLAQAGLARVFIGINDLEKEGAFVYSDHSPMRTFNKWRSGEPNNAYDEE
DCVEMVASGGWNDVACHTTMYFMCEFDKENM

177/392

FIGURE 169

AGTGA CTGCAGCCTTCCTAGATCCCCTCCACTCGGTTTCTCTCTTTGCAGGAGCACCGGC
AGCACCA GTGTGTGAGGGGAGCAGGCAGCGGTCC TAGCCAGTTCCTTGATCCTGCCAGAC
CACCCAGCCCCCGGCACAGAGCTGCTCCACAGGCACCA ATGAGGATCATGCTGCTATTAC
AGCCATCCTGGCCTTCAGCCTAGCTCAGAGCTTTGGGGCTGTCTGTAAAGGAGCCACAGGA
GGAGGTGGTTCTTGGCGGGGGCCGCAGCAAGAGGGATCCAGATCTCTACCAGCTGCTCCA
GAGACTCTTCAAAGCCACTCATCTCTGGAGGGATTGCTCAAAGCCCTGAGCCAGGCTAG
CACAGATCCTAAGGAATCAACATCTCCCGAGAAACGTGACATGCATGACTTCTTTGTGGG
ACTTATGGGCAAGAGGAGCGTCCAGCCAGAGGGAAAGACAGGACCTTTCTTACCTTCAGT
GAGGGTTCCTCGGCCCCCTTCATCCCAATCAGCTTGGATCCACAGGAAAGTCTTCCCTGGG
AACAGAGGAGCAGAGACCTTTA TAAGACTCTCCTACGGATGTGAATCAAGAGAACGTCCC
CAGCTTTGGCATCCTCAAGTATCCCCGAGAGCAGAATAGGTACTCCACTTCCGGACTCC
TGGACTGCATTAGGAAGACCTCTTTCCCTGTCCCAATCCCCAGGTGCGCACGCTCCTGTT
ACCCTTTCTCTTCCCTGTTCTTGTAACATTCTTG TGCTTTGACTCCTTCTCCATCTTTTC
TACCTGACCCTGGTGTGGAACTGCATAGTGAATATCCCCAACCCCAATGGGCATTGACT
GTAGAATACCCTAGAGTTCCTGTAGTGTCTACATTAAAAATATAATGTCTCTCTCTATT
CCTCAACAATAAAGGATTTTTGCATATGAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAA

178/392

FIGURE 170

MRIMLLFTAILAFSLAQSFQAVCKEPQEEVVPGGGRSKRDPDLYQLLQRLFKSHSSLEGL
LKALSQASTDPKESTSPKRDMDHFFVGLMGKRSVQPEGKTGPFLPSVRVPRPLHPNQLG
STGKSSSLGTEEQRPL

Important features:

Signal peptide:

amino acids 1-18

Tyrosine kinase phosphorylation site:

amino acids 36-45

N-myristoylation site:

amino acids 33-39, 59-65

Amidation site:

amino acids 90-94

Leucine zipper pattern:

amino acids 43-65

Tachykinin family signature:

amino acids 86-92

179/392

FIGURE 171

TGGCCTCCCCAGCTTGCCAGGCACAAGGCTGAGCGGGAGGAAGCGAGAGGCATCTAAGCA
GGCAGTGTTTTGCCTTCACCCCAAGTGACCATGAGAGGTGCCACGCGAGTCTCAATCATG
CTCCTCCTAGTAACCTGTGTCTGACTGTGCTGTGATCACAGGGGCCTGTGAGCGGGATGTC
CAGTGTGGGGCAGGCACCTGCTGTGCCATCAGCCTGTGGCTTCGAGGGCTGCGGATGTGC
ACCCCGCTGGGGCGGGAAGGCGAGGAGTGCCACCCCGGCAGCCACAAGGTCCCCTTCTTC
AGGAAACGCAAGCACACACCTGTCTTGTCTTGCCCAACCTGCTGTGCTCCAGGTTCCTCG
GACGGCAGGTACCGCTGCTCCATGGACTTGAAGAACATCAATTTTTTAGGCGCTTGCCCTGG
TCTCAGGATACCCACCATCCTTTCTCTGAGCACAGCCTGGATTTTTATTTCTGCCATGAA
ACCCAGCTCCCATGACTCTCCAGTCCCTACACTGACTACCTTGATCTCTCTTGTCTAGT
ACGCACATATGCACACAGGCAGACATACTCCCATCATGACATGGTCCCCAGGCTGGCCT
GAGGATGTCACAGCTTGAGGCTGTGGTGTGAAAGGTGGCCAGCCTGGTTCTCTTCCCTGCT
TCAGGCTGCCAGAGAGGTGGTAAATGGCAGAAAGGACATTCCCCCTCCCCCTCCCCAGGTG
ACCTGCTCTCTTTCCCTGGGCCCTGCCCCCTCTCCCCACATGTATCCCTCGGTCTGAATTAG
ACATTCCTGGGCACAGGCTCTTGGGTGCATTGCTCAGAGTCCCAGGTCCTGGCCTGACCC
TCAGGCCCTTCACGTGAGGTCTGTGAGGACCAATTTGTGGGTAGTTTCATCTTCCCTCGAT
TGGTTAACTCCTTAGTTTCAGACCACAGACTCAAGATTGGCTCTTCCCAGAGGGCAGCAG
ACAGTCACCCCAAGGCAGGTGTAGGGAGCCAGGGAGGCCAATCAGCCCCCTGAAGACTC
TGGTCCCAGTCAGCCTGTGGCTTGTGGCCTGTGACCTGTGACCTTCTGCCAGAATTGTCA
TGCCTCTGAGGCCCCCTCTTACCACACTTTACCAGTTAACCCTGAAGCCCCCAATTCCC
ACAGCTTTTCCATTAAAATGCAAATGGTGGTGGTTCAATCTAATCTGATATTGACATATT
AGAAGGCAATTAGGGTGTTTCTTAAACAACCTCTTTCCAAGGATCAGCCCTGAGAGCAG
GTTGGTGACTTTGAGGAGGGCAGTCCTCTGTCCAGATTGGGGTGGGAGCAAGGGACAGGG
AGCAGGGCAGGGGCTGAAAGGGGCACTGATTGAGACCAGGGAGGCAACTACACACCAACA
TGCTGGCTTTAGAATAAAAGCACCAACTGAAAAAA

180/392

FIGURE 172

MRGATRVSIMLLLLVTVSDCAVITGACERDVQCGAGTCCAI~~SLWLRGLRMCTPLGREGE~~EC
HPGSHKVPFFRKRKHHTCPCLPNLLCSRFPDGRYRCSMDLKNINF

Signal peptide:
amino acids 1-19

Tyrosine kinase phosphorylation site:
amino acids 88-95

N-myristoylation sites:
amino acids 33-39, 35-41, 46-52

181/392

FIGURE 173

AGCGCCCGGGCGTTCGGGGCGGTAAAAGGCCGGCAGAAAGGGAGGCACTTGAGAAATGCTCTT
TCCTCCAGGACCCAAGTTTCTTCACCATGGGGATGTGGTCCATTGGTGCAGGAGCCCTGG
GGGCTGCTGCCTTGGCATTGCTGCTTGCCAACACAGACGTGTTTCTGTCCAAGCCCCAGA
AAGCGGCCCCTGGAGTACCTGGAGGATATAGACCTGAAAACACTGGAGAAGGAACCAAGGA
CTTTCAAAGCAAAGGAGCTATGGGAAAAAATGGAGCTGTGATTATGGCCGTGCGGAGGC
CAGGCTGTTTCCTCTGTCTGAGAGGAAGCTGCGGATCTGTCTCCCTGAAAAGCATGTTGG
ACCAGCTGGGCGTCCCCCTCTATGCAGTGGTAAAGGAGCACATCAGGACTGAAGTGAAGG
ATTTCCAGCCTTATTTCAAAGGAGAAATCTTCCTGGATGAAAAGAAAAAGTTCTATGGTC
CACAAAGGCGGAAGATGATGTTTATGGGATTTATCCGTCTGGGAGTGTGGTACAACCTTCT
TCCGAGCCTGGAACGGAGGCTTCTCTGGAACCTGGAAGGAGAAGGCTTCATCCTTGGGG
GAGTTTTTCGTGGTGGGATCAGGAAAGCAGGGCATTCTTCTTGAGCACCGAGAAAAAGAAT
TTGGAGACAAAGTAAACCTACTTTCTGTTCTGGAAGCTGCTAAGATGATCAAACACAGA
CTTTGGCCTCAGAGAAAAAATGATTGTGTGAAACTGCCCAGCTCAGGGATAACCAGGGAC
ATTCACCTGTGTTTCATGGGATGTATTGTTTCCACTCGTGTCCCTAAGGAGTGAGAAACCC
ATTTATACTCTACTCTCAGTATGGATTATTAATGTATTTTAATATTCTGTTTAGGCCCAC
TAAGGCAAAATAGCCCCAAAACAAGACTGACAAAAATCTGAAAAACTAATGAGGATTATT
AAGCTAAAACCTGGGAAATAGGAGGCTTAAAAATTGACTGCCAGGCTGGGTGCAGTGGCTC
ACACCTGTAATCCCAGCACTTTGGGAGGCCAAGGTGAGCAAGTCACTTGAGGTCGGGAGT
TCGAGACCAGCCTGAGCAACATGGCGAAACCCCGTCTCTACTAAAAATACAAAAATCACC
CGGGTGTGGTGGCAGGCACCTGTAGTCCCAGCTACCCGGGAGGCTGAGGCAGGAGAATCA
CTTGAACCTGGGAGGTGGAGGTTGCGGTGAGCTGAGATCACACCCTGTATTCCAGCCTG
GGTGA CTGAGACTCTAACTAA

182/392

FIGURE 174

MSFLQDPSFFTMGMWSIGAGALGAAALALLANTDVFLSKPQKALEYLEDIDLKTLEKE
PRTFKAKELWEKNGAVIMAVRRPGCFLCREEAADLSSLKSMLDQLGVPLYAVVKEHIRTE
VKDFQPYFKGEIFLDEKKKFYGPQRRKMMFMGFIRLGWYNFFRAWNGGFSGNLEGEGLI
LGGVFVVGSGKQGILLEHREKEFGDKVNLLSVLEAAKMIKPQTLASEKK

183/392

FIGURE 175

GACAGTGGAGGGCAGTGGAGAGGACCGCGCTGTCCTGCTGTCACCAAGAGCTGGAGACAC
CATCTCCCACCGAGAGTCATGGCCCCATTGGCCCTGCACCTCCTCGTCCTCGTCCCCATC
CTCCTCAGCCTGGTGGCCTCCCAGGACTGGAAGGCTGAACGCAGCCAAGACCCCTTCGAG
AAATGCATGCAGGATCCTGACTATGAGCAGCTGCTCAAGGTGGTGACCTGGGGGGCTCAAT
CGGACCCTGAAGCCCCAGAGGGTGATTGTGGTTGGCGCTGGTGTGGCCGGGCTGGTGGCC
GCCAAGGTGCTCAGCGATGCTGGACACAAGGTCACCATCCTGGAGGCAGATAACAGGATC
GGGGGCCGCATCTTCACCTACCGGGACCAGAACACGGGCTGGATTGGGGAGCTGGGAGCC
ATGCGCATGCCCAGCTCTCACAGGATCCTCCACAAGCTCTGCCAGGGCCTGGGGCTCAAC
CTGACCAAGTTACCCAGTACGACAAGAACACGTGGACGGAGGTGCACGAAGTGAAGCTG
CGCAACTATGTGGTGGAGAAGGTGCCCCGAGAAGCTGGGCTACGCCCTGCGTCCCCAGGAA
AAGGGCCACTCGCCCCGAAGACATCTACCAGATGGCTCTCAACCAGGCCCTCAAAGACCTC
AAGGCACTGGGCTGCAGAAAGGCGATGAAGAAGTTTGAAAGGCACACGCTCTTGGAATAT
CTTCTCGGGGAGGGGAACCTGAGCCGGCCGGCGTGACGCTTCTGGGAGACGTGATGTCC
GAGGATGGCTTCTTCTATCTCAGCTTCGCCGAGGCCCTCCGGGCCACAGCTGCCTCAGC
GACAGACTCCAGTACAGCCGCATCGTGGGTGGCTGGGACCTGCTGCCGCGCGCTGCTG
AGCTCGCTGTCCGGGCTTGTGCTGTTGAACGCGCCCCGTGGTGGCGATGACCCAGGGACCG
CACGATGTGCACGTGCAGATCGAGACCTCTCCCCGGCGCGGAATCTGAAGGTGCTGAAG
GCCGACGTGGTGCTGCTGACGGCGAGCGGACCGGCGGTGAAGCGCATCACCTTCTCGCCG
CCGCTGCCCCGCCACATGCAGGAGGCGCTGCGGAGGCTGCACTACGTGCCGGCCACCAAG
GTGTTCTTAAGCTTCCGCAGGCCCTTCTGGCGCGAGGAGCACATTGAAGGCGGCCACTCA
AACACCGATCGCCCGTCGCGCATGATTTTCTACCCGCCCGCGCGAGGGCGCGCTGCTG
CTGGCCTCGTACACGTGGTCCGACGCGCGGCGAGCGTTCGCCGGCTTGAGCCGGGAAGAG
GCGTTGCGCTTGGCGCTCGACGACGTGGCGGCATTGCACGGGCCTGTCGTGCGCCAGCTC
TGGGACGGCACCCGGCGTCGTCAAGCGTTGGGCGGAGGACCAGCACAGCCAGGGTGGCTTT
GTGGTACAGCCGCCGCGCTCTGGCAAACCGAAAAGGATGACTGGACCGTCCCTTATGGC
CGCATCTACTTTGCCGGCGAGCACACCGCCTACCCGCACGGCTGGGTGGAGACGGCGGTC
AAGTCGGCGCTGCGCGCCGCCATCAAGATCAACAGCCGGAAGGGGCTGCATCGGACACG
GCCAGCCCCGAGGGGCACGCATCTGACATGGAGGGGCAGGGGCATGTGCATGGGGTGGCC
AGCAGCCCCCTCGCATGACCTGGCAAAGGAAGAAGGCAGCCACCCTCCAGTCCAAGGCCAG
TTATCTCTCAAAACACGACCCACACGAGGACCTCGCATTAAAGTATTTTCGGAAAAAAA
AAA

184/392

FIGURE 176

MAPLALHLLVLPILLSIVASQDWKAERSQDPFEKCMQDPDYEQLLKVVWGLNRTLKPQ
RVIVVGAGVAGLVAAKVLSDAGHKVTILEADNRIGGRIFTYRDQNTGWIGELGAMRMPSS
HRILHKLCQGLGLNLTKFTQYDKNTWTEVHEVKLRNYVVEKVPEKLGALRPQEKGHSPE
DIYQMALNQALKDLKALGCRKAMKKFERHTLLEYLLGEGNLSRPAVQLLGDVMSDGGFFY
LSFAEALRAHSCLSDRLQYSRIVGGWDLPRALLSSLSGLVLLNAPVVAMTQGPHDVHVQ
IETSPPARNLKVLKADVLLTASGPAVKRITFSPPLPRHMQEALRRLHYVPATKVFLSFR
RPFWREEHIEGGHSNTDRPSRMIFYPPPREGALLASYTWSDAAAFAGLSREEALRLAL
DDVAALHGPVVRQLWDGTGVVKRWAEDQHSQGGFVVQPPALWQTEKDDWTVPYGRIYFAG
EHTAYPHGWVETAVKSALRAAIKINSRKGPASDTASPEGHASDMEGQGHVHGVASSPSHD
LAKEEGSHPPVQGQLSLQNTTHTRTSH

Signal peptide:
amino acids 1-21

185/392

FIGURE 177

CCGGGGAGGGGAGGGCCCCGTCCCGCCCCCTCCCCGTCTCTCCCCGCCCCCTCCCCGTCCCTC
CCGCCGAAGCTCCGTCCCGCCCCGCGGGCCGGCTCCGCCCTCACCTCCCGGCCGCGGCTGC
CCTCTGCCCGGGTTGTCCAAGATGGAGGGCGCTCCACCGGGGTCGCTCGCCCCCTCCGGCTC
CTGCTGTTTCGTGGCGCTACCCGCCTCCGGCTGGCTGACGACGGGCGCCCCCGAGCCGCCG
CCGCTGTCCGGAGCCCCACAGGACGGCATCAGAATTAATGTAACCTACACTGAAAGATGAT
GGGGACATATCTAAACAGCAGGTTGTTCTTAACATAACCTATGAGAGTGGACAGGTGTAT
GTAAATGACTTACCTGTAAATAGTGGTGTAACCCGAATAAGCTGTCAGACTTTGATAGTG
AAGAATGAAAATCTTGAAAATTTGGAGGAAAAAGAATATTTTGGAATTGTCAGTGTAAGG
ATTTTAGTTCATGAGTGGCCTATGACATCTGGTTCCAGTTTGCAACTAATTGTCATTCAA
GAAGAGGTAGTAGAGATTGATGGAAAACAAGTTCAGCAAAAGGATGTCACTGAAATTGAT
ATTTTAGTTAAGAACCGGGGAGTACTCAGACATTCAAACCTATACCCTCCCTTTGGAAGAA
AGCATGCTCTACTCTATTTCTCGAGACAGTGACATTTTATTTACCCTTCCTAACCTCTCC
AAAAAAGAAAGTGTTAGTTCACTGCAAACCACTAGCCAGTATCTTATCAGGAATGTGGAA
ACCACTGTAGATGAAGATGTTTTACCTGGCAAGTTACCTGAAACTCCTCTCAGAGCAGAG
CCGCCATCTTCATATAAGGTAATGTGTGTCAGTGGATGGAAAAGTTTAGAAAAGATCTGTGT
AGGTTCTGGAGCAACGTTTTCCAGTATTCTTTTCAGTTTTTGAACATCATGGTGGTTGGA
ATTACAGGAGCAGCTGTGGTAATAACCATCTTAAAGGTGTTTTTCCAGTTTCTGAATAC
AAAGGAATTCTTCAGTTGGATAAAGTGGACGTCATACCTGTGACAGCTATCAACTTATAT
CCAGATGGTCCAGAGAAAAGAGCTGAAAACCTTGAAGATAAAACATGTATTTAAACGCC
ATCTCATATCATGGACTCCGAAGTAGCCTGTTGCCTCCAAATTTGCCACTTGAATATAAT
TTTCTTTAAATCGTT

186/392

FIGURE 178

MEGAPPGSLALRLLLFVALPASGWLTTGAPEPPPLSGAPQDGIRINVTTLKDDGDISKQQ
VVLNITYESGQVYVNDLPVNSGVTRISCQTLIVKNENLENLEEKEYFGIVSVRILVHEWP
MTSGSSLQLIVIQEEVVEIDGKQVQKDVTEIDILVKNRGVLRHSNYTLPLEESMLYSIS
RDS DILFTLPNL SKKESVSSLQTTSQYLIRNVETTVD EDVLP GKLPETPLRAEPPSSYKV
MCQWMEKFRKDL CRFWSNVFPVFFQFLNIMVVGITGA AVVITILKVFFPVSEYKGILQLD
KVDVIPVTAINLYPDGPEKRAENLEDKTCI

Signal peptide:

1-23

Transmembrane domain:

266-284

Leucine zipper pattern:

155-177

N-glycosylation site:

46-50, 64-68, 166-170, 191-195

Motif name: N-myristoylation site:

3-9, 42-48, 273-279

187/392

FIGURE 179

CTCCTTAGGTGGAAACCTGGGAGTAGAGTACTGACAGCAAAGACCGGGAAAGACCATAC
GTCCCCGGGCAGGGGTGACAACAGGTGTCTATCTTTTGATCTCGTGTGTGGCTGCCTTCC
TATTTCAAGGAAAGACGCCAAGGTAATTTTGACCCAGAGGAGCAATGATGTAGCCACCTC
CTAACCTTCCCTTCTTGAACCCCCAGTTATGCCAGGATTTACTAGAGAGTGTCAACTCAA
CCAGCAAGCGGCTCCTTCGGCTTAACCTTGTGGTTGGAGGAGAGAACCCTTTGTGGGGCTGC
GTTCTCTTAGCAGTGCTCAGAAAGTGACTTGCCCTGAGGGTGGACCAGAAGAAAGGAAAGGT
CCCCTCTTGCTGTTGGCTGCACATCAGGAAGGCTGTGATGGGAATGAAGGTGAAAACCTTG
GAGATTTCACTTCAGTCATTGCTTCTGCCTGCAAGATCATCCTTTAAAAGTAGAGAAGCT
GCTCTGTGTGGTGGTTAACTCCAAGAGGCAGAACTCGTTCTAGAAGGAAATGGATGCAAG
CAGCTCCGGGGGGCCCCAAACGCATGCTTCCTGTGGTCTAGCCCAGGGAAGCCCTTCCGTG
GGGGCCCCGGCTTTGAGGGATGCCACCGGTTCTGGACGCATGGCTGATTCTGAATGATG
ATGGTTCGCGGGGGGCTGCTTGCCTGGATTTCCCGGGTGGTGGTTTGTGGTGTCTCCTC
TGCTGTGCTATCTCTGTCTGTACATGTTGGCCTGCACCCCCAAAAGGTGACGAGGAGCAG
CTGGCACTGCCCCAGGGCCAACAGCCCCACGGGGAAGGAGGGGTACCAGGCCGTCTTCAG
GAGTGGGAGGAGCAGCACCGCAACTACGTGAGCAGCCTGAAGCGGCAGATCGCACAGCTC
AAGGAGGAGCTGCAGGAGAGGAGTGAGCAGCTCAGGAATGGGCAGTACCAAGCCAGCGAT
GCTGCTGGCCTGGGTCTGGACAGGAGCCCCCAGAGAAAACCCAGGCCGACCTCCTGGCC
TTCTTGCACTCGCAGGTGGACAAGGCAGAGGTGAATGCTGGCGTCAAGCTGGCCACAGAG
TATGCAGCAGTGCCTTTTCGATAGCTTTACTCTACAGAAGGTGTACCAGCTGGAGACTGGC
CTTACCCGCCACCCCGAGGAGAAGCCTGTGAGGAAGGACAAGCGGGATGAGTTGGTGGAA
GCCATTGAATCAGCCTTGGAGACCTGAACAATCCTGCAGAGAACAGCCCCAATCACCGT
CCTTACACGGCCTCTGATTTCATAGAAGGGATCTACCGAACAGAAAGGGACAAAGGGACA
TTGTATGAGCTCACCTTCAAAGGGGACCACAAACACGAATTCAAACGGCTCATCTTATTT
CGACCATTGAGCCCCATCATGAAAGTGAAAAATGAAAAGCTCAACATGGCCAACACGCTT
ATCAATGTTATCGTGCCTCTAGCAAAAAGGGTGGACAAGTTCGGGCAGTTCATGCAGAAT
TTCAGGGAGATGTGCATTGAGCAGGATGGGAGAGTCCATCTCACTGTTGTTTACTTTGGG
AAAGAAGAAATAAATGAAGTCAAAGGAATACTTGAAAACACTTCCAAAGCTGCCAACTTC
AGGAACTTTACCTTCATCCAGCTGAATGGAGAATTTTCTCGGGGAAAGGGACTTGATGTT
GGAGCCCGCTTCTGGAAGGGAAGCAACGTCCTTCTCTTTTCTGTGATGTGGACATCTAC
TTCACATCTGAATTCCTCAATACGTGTAGGCTGAATACACAGCCAGGGAAGAAGGTATTT
TATCCAGTTCTTTTCAGTCAGTACAATCCTGGCATAATATACGGCCACCATGATGCAGTC
CCTCCCTTGGAACAGCAGCTGGTCATAAAGAAGGAACTGGATTTTGGAGAGACTTTGGA
TTTGGGATGACGTGTCAGTATCGGTGAGACTTCATCAATATAGGTGGGTTTGATCTGGAC
ATCAAAGGCTGGGGCGGAGAGGATGTGCACCTTTATCGCAAGTATCTCCACAGCAACCTC
ATAGTGGTACGGACGCCTGTGCGAGGACTTCCACCTCTGGCATGAGAAGCGCTGCATG
GACGAGCTGACCCCCGAGCAGTACAAGATGTGCATGCAGTCCAAGGCCATGAACGAGGCA
TCCCACGGCCAGCTGGGCATGCTGGTGTTCAGGCACGAGATAGAGGCTCACCTTCGCAAA
CAGAAACAGAAGACAAGTAGCAAAAAAACATGAAGTCCAGAGAAGGATTGTGGGAGACA
CTTTTTCTTTCTTTTGCAATTACTGAAAGTGGCTGCAACAGAGAAAAGACTTCCATAAA
GGACGACAAAAGAATTGGACTGATGGGTGAGAGATGAGAAAAGCCTCCGATTTCTCTCTGT
TGGGCTTTTTTACAACAGAAATCAAATCTCCGCTTTGCCTGCAAAAGTAACCCAGTTGCA
CCCTGTGAAGTGTCTGACAAAGGCAGAATGCTTGTGAGATTATAAGCCTAATGGTGTGGA
GGTTTTGATGGTGTTTACAATACACTGAGACCTGTTGTTTTGTGTGCTCATTGAAATATT
CATGATTTAAGAGCAGTTTGTAAAAAATTCATTAGCATGAAAGGCAAGCATATTTCTCC
TCATATGAATGAGCCTATCAGCAGGGCTCTAGTTTCTAGGAATGCTAAAATATCAGAAGG
CAGGAGAGGAGATAGGCTTATTATGATACTAGTGAGTACATTAAGTAAAATAAATGGAC
CAGAAAAGAAAAGAAACCATAAATATCGTGTCTATTTTTCCCAAGATTAACCAAAAATA

188/392

ATCTGCTTATCTTTTTGGTTGTCCTTTTAACTGTCCTCCGTTTTTTTCTTTTATTTAAAAA
TGCACTTTTTTTCCCTTGTGAGTTATAGTCTGCTTATTTAATTACCACTTTGCAAGCCTT
ACAAGAGAGCACAAAGTTGGCCTACATTTTTATATTTTTTAAGAAGATACTTTGAGATGCA
TTATGAGAACTTTTCAAGCATCAAATTGATGCCATATCCAAGGACATGCCAAATG
CTGATTCTGTGAGGCACTGAATGTCAGGCATTGAGACATAGGGAAGGAATGGTTTGTACT
AATACAGACGTACAGATACTTTCTCTGAAGAGTATTTTCGAAGAGGAGCAACTGAACACT
GGAGGAAAAGAAAATGACACTTTCTGCTTTACAGAAAAGGAACTCATTGAGACTGGTGA
TATCGTGATGTACCTAAAAGTCAGAAACCACATTTTCTCCTCAGAAGTAGGGACCGCTTT
CTTACCTGTTTAAATAAACCAAAGTATACCGTGTGAACCAAACAATCTCTTTTCAAAACA
GGGTGCTCCTCCTGGCTTCTGGCTTCCATAAGAAGAAATGGAGAAAAATATATATATATA
TATATATATTGTGAAAGATCAATCCATCTGCCAGAATCTAGTGGGATGGAAGTTTTTGCT
ACATGTTATCCACCCAGGCCAGGTGGAAGTAACTGAATTATTTTTTAAATTAAGCAGTT
CTACTCAATCACCAAGATGCTTCTGAAAATTGCATTTTATTACCATTTCAACTATTTTT
TAAAAATAAATACAGTTAACATAGAGTGGTTTCTTCATTCATGTGAAAATTATTAGCCAG
CACCAGATGCATGAGCTAATTATCTCTTTGAGTCCTTGCTTCTGTTTGCTCACAGTAAAC
TCATTGTTTAAAAGCTTCAAGAACATTCAAGCTGTTGGTGTGTTAAAAAATGCATTGTAT
TGATTTGTACTGGTAGTTTATGAAATTTAATTAAACACAGGCCATGAATGGAAGGTGGT
ATTGCACAGCTAATAAAATATGATTTGTGGATATGAA

189/392

FIGURE 180

MMVRRGLLAWISRVVLLVLLCCAISVLYMLACTPKGDEEQLALPRANSPTGKEGYQAV
LQEWEEQHRNYVSSLKRQIAQLKEELQERSEQLRNGQYQASDAAGLGGLDRSPPEKTQADL
LAFLHSQVDKAEVNAGVKLATEYAAVPFDSFTLQKVYQLETGLTRHPPEKPVKDKRDEL
VEAIESALETLNPAENSPNHRPYTASDFIEGIYRTERDKGTLTYELTFKGDHKHEFKRLI
LFRPFSPIMKVKNEKLNMANLINVIVPLAKRVDKFRQFMQNFREMCIEQDGRVHLTVVY
FGKEEINEVKGILENTSKAANFRNFTFIQLNGEFSRGKGLDVGARFWKGSNVLLFFCDVD
IYFTSEFLNTCRLNTQPGKKVFYPVLFSQYNPGIIYGHDAVPPLEQQLVIKKETGFWRD
FGFGMTCQYRSDFINIGGFDLDIKGWGGEDVHLYRKYLHSNLIVVRTPVRLFHLWHEKR
CMDELTPEQYKMCMQSKAMNEASHGQLGMLVFRHEIEAHLRKQKQKTSSKKT

190/392

FIGURE 181

CGTCTCTGCGTTCCGCCATGCGTCCCGGGGCGCCAGGGCCACTCTGGCCTCTGCCCTGGGG
GGCCCTGGCTTGGGCGGTGGGCTTCGTGAGCTCCATGGGCTCGGGGAACCCCGCGCCCGG
TGGTGTGTTGCTGGCTCCAGCAGGGCCAGGAGGCCACCTGCAGCCTGGTGCTCCAGACTGA
TGTCACCCGGGCGAGTGCTGTGCCTCCGGCAACATTGACACCGCCTGGTCCAACCTCAC
CCACCCGGGGAACAAGATCAACCTCCTCGGCTTCTTGGGCCTTGCTCCACTGCCTTCCCTG
CAAAGATTGCTGCGACGGCGTGAGTGCGGCCCCGGGCAAGGCGTGCCGCATGCTGGGGGG
CCGCCCCGCGCTGCGAGTGCGCGCCCCGACTGCTCGGGGCTCCCGGCGCGGCTGCAGGTCTG
CGGCTCAGACGGCGCCACCTACCGCGACGAGTGCGGAGCTGCGCGCCGCGCGCTGCCGCGG
CCACCCGGACCTGAGCGTCATGTACCGGGGCGGCTGCCGCAAGTCCTGTGAGCACGTGGT
GTGCCCGCGGGCCACAGTCGTGCGTCGTGGACCAGACGGGCAGCGCCCACTGCGTGGTGTG
TCGAGCGGCGCCCTGCCCTGTGCCCTCCAGCCCCGGCCAGGAGCTTTCGGCAACAACAA
CGTCACCTACATCTCCTCGTGCCACATGCGCCAGGCCACCTGCTTCTGGGCGCTCCAT
CGGCGTGCGCCACGCGGGCAGCTGCGCAGGCACCCCTGAGGAGCCGCCAGGTGGTGAGTC
TGCAGAAGAGGAAGAGAACTTCGTGTGAGCCTGCAGGACAGGCCTGGGCCTGGTGCCCGA
GGCCCCCATCATCCCTGTTATTTATTGCCACAGCAGAGTCTAATTTATATGCCACGGA
CACTCCTTAGAGCCCGGATTTCGGACCCTTGGGGATCCCAGAACCTCCCTGACGATATCC
TGGAAGGACTGAGGAAGGGAGGCCTGGGGGCGGCTGGTGGGTGGGATAGACCTGCGTTC
CGGACACTGAGCGCCTGATTTAGGGCCCTTCTCTAGGATGCCCCAGCCCTACCCTAAGA
CCTATTGCCGGGGAGGATTCCACACTTCCGCTCCTTTGGGGATAAACCTATTAATTATTG
CTACTATCAAGAGGGGCTGGGCATTCTCTGCTGGTAATTCCTGAAGAGGCATGACTGCTTT
TCTCAGCCCCAAGCCTCTAGTCTGGGTGTGTACGGAGGGTCTAGCCTGGGTGTGTACGGA
GGGTCTAGCCTGGGTGAGTACGGAGGGTCTAGCCTGGGTGAGTACGGAGGGTCTAGCCTG
GGTGAGTACGGAGGGTCTAGCCTGGGTGTGTATGGAGGATCTAGCCTGGGTGAGTATGGA
GGGTCTAGCCTGGGTGAGTATGGAGGGTCTAGCCTGGGTGTGTATGGAGGGTCTAGCCTG
GGTGAGTATGGAGGGTCTAGCCTGGGTGTGTATGGAGGGTCTAGCCTGGGTGAGTATGGA
GGGTCTAGCCTGGGTGTGTACGGAGGGTCTAGTCTGAGTGCGTGTGGGGACCTCAGAACA
CTGTGACCTTAGCCAGCAAGCCAGGCCCTTCATGAAGGCCAAGAAGGCTGCCACCATTTC
CCTGCCAGCCCAAGAACTCCAGCTTCCCCACTGCCTCTGTGTGCCCCCTTTCGCTCCTGTG
AAGGCCATTGAGAAATGCCAGTGTGCCCCCTGGGAAAGGGCACGGCCTGTGCTCCTGAC
ACGGGCTGTGCTTGGCCACAGAACCACCCAGCGTCTCCCCTGCTGCTGTCCACGTAGTT
CATGAGGCAACGTGCGGTGCTCAGACGTGGAGCAGCCAGCGGCAGCTCAGAGCAGGGC
ACTGTGTCCGGCGGAGCCAAGTCCACTCTGGGGGAGCTCTGGCGGGGACCACGGGCCACT
GCTCACCCACTGGCCCCGAGGGGGGTGTAGACGCCAAGACTCACGCATGTGTGACATCCG
GAGTCCTGGAGCCGGGTGTCCCAGTGGCACCCTAGGTGCCTGCTGCCTCCACAGTGGGG
TTCACACCCAGGGCTCCTTGGTCCCCCACAACCTGCCCCGGCCAGGCCTGCAGACCCAGA
CTCCAGCCAGACCTGCCTCACCCACCAATGCAGCCGGGGCTGGCGACACCAGCCAGGTGC
TGGTCTTGGGCCAGTTCTCCACGACGGCTCACCCCTCCCCTCCATCTGCGTTGATGCTCA
GAATCGCCTACCTGTGCCTGCGTGTAACCACAGCCTCAGACCAGCTATGGGGAGAGGAC
AACACGGAGGATATCCAGCTTCCCCGGTCTGGGGTGAGGAATGTGGGGAGCTTGGGCATC
CTCCTCCAGCCTCCTCCAGCCCCCAGGCAGTGCCTTACCTGTGGTGCCAGAAAAGTGCC
CCTAGGTTGGTGGGTCTACAGGAGCCTCAGCCAGGCAGCCACCCACCCCTGGGGCCCTG
CCTCACCAAGGAAATAAAGACTCAAGCCATAAAAAAA

191/392

FIGURE 182

MRPGAPGPLWPLPWGALAWVGFVSSMGSGNPAPGGVCWLQQGQEATCSLVLQTDVTRAE
CCASGNIDTAWSNLTHPGNKINLLGFLGLVHCLPCKDSCDGVECGPGKACRMLGGRPRCE
CAPDCSGLPARLQVCGSDGATYRDECELRAARCRGHPDLSVMYRGRCRKSCHEVVCPRPQ
SCVVDQTGSAHCVVCRAAPCFVPSSPGQELCGNNNVITYISSCHMRQATCFLGRSIGVRHA
GSCAGTPEEPPGGESAEEEENFV

Important features:

Signal peptide:

amino acids 1-20

N-glycosylation sites:

amino acids 73-77, 215-219

Osteonectin domain proteins:

amino acids 97-130, 169-202

192/392

FIGURE 183

CACTCATTCAATTCCAAAGGGTCTCTCAAGGCAATGGTAATGTGCAAGGAGGTGATACCTA
AATGAATGACCAAAAGAACATGCTTCTGCTTTTGTGTGTCTCCTACATTTTAGACATTTG
TTTGTTCCTCTTGGTAGCCTTTAAATTCCTTGAAGCCCAGGACCATGTCTCACTTACCTT
TGTGTTTCCACTAACTAGTCTACCTCCTGGAATTGGCAGATACTCAGTGAAAAGCCTGTGA
AATAAGTGATGTCTATTTCTAGCATATTATTCTGAGATTTAATGATAGATTTAGTGATTG
AATGAGATTTCCATTTTCAAATACAGCAAAAGCATAACTATTTTCATTCCATTCAATTCA
TTCAACTTCATTCTCAAATTAGGTCTGAGTTAACTAATAATTACCTTTGAAATGTGTG
GGTTATTTGAGGCAATCAGGTGGTGACATTGAGCTCTCAGCCAGAGTTTGTTCCTGGAAT
TGATTCAGTTCCATTGCATTGATTTTTGTTCTCAGAAGCCAAGGTTTCCCATGAAAAATC
ATTCCCACCTTGAATTGGGCTGTGATTCTTGCTGCGTTTAAAGTAAAGGAAGCCTCTTGGTT
CTAGTTCCTGCAAACTTACACACTGAACCTGGGACAAGTTTTTGTTCAGTAATGGCTGGG
AAAAGAGGAACCTTTCATTTTATTCAGAAGTCAAAAACAAAGGCCTCCAGCCACTTGGGA
GATGTTTTGTTCGAGACACCAGCCTGGCTCTGTCTTTATGCCTAACAATTGAGCATCCAG
TCTTCTTTGTGCTGGGACCATTGCTCAGCTCTGCAAGGGGAAAAGAGGGAGAAAGCCAGA
GCTGCCAGGCTTCTTGCACTGGGGCCGGGGAGGGTTCTTGGGAAGCAGGTGCTCTCTGG
CTTCTTGGTACGTGAGGCTCTCGGAGCTGCCTCTCCTCTGACCCTCAGGTCTCACCAG
TTTGCTCCAGGAGTATATTGAAAACATACCCAGTGCTCTCTCAAGCACCCACTGCTTAGA
GGGCCCAGATTTCTTTTCTTCTTTCCCTTGCAAGAGCTGGAGACTGCATCGGGCATCTGG
TGTTTAAACTAAACAGGAAAAGTGAATAAGGTCCACAGTGCTCATTGTGTAGACTAGCT
GCCCTCCGATGGGTGCTCTGATTATCAGTGCTTCCAGTGCCAGGGCCTGTCACTAAACAGG
CCTCACTTCCTCCTTGGGGGCTTTCCCATGGGAGGTGTGGCTTTTTTACTCTACATGGAAA
TGACTCTCTGCAGCCACAGAACACAGTCATTTTCTGAATTATCCAGTCTCTCATGCGCC
CTGGATTCTCCAGATGCCTTATATCTCTTGTGCAAAGTTGTCTAAAATTTGGTTCCAG
CTTCCAAGCCTTGCCTTTTGGCCTTCCCTGGAAGTATTTTTGTGTGATGAGTCGTCTGTCAT
TATTCTCTAAAATGATTTGCTTTTTGTTCCTTTTCATTCTCTATTTCCACCCACATATACA
CACATGCTTCTTAACTTAGGGGATTACATGCCAATAAATCTATTGTTGAAAATGCACTAA
TACTATCGCAAAGACGAAAATTACAGGCTGAACCGTTGTAAGTCCATATGCTCCTCAAC
TTACATGTGTGATGGAGTTATGCCCAAATAAGTCCATCGTCAAGTTGAAAATCAAAATC
AAGCCATCTTAGGTTGAGGACCATTTGTTTGTACCTCCAAAGATGTCATATCTTTAAACA
TACTCCCTAGCTTTTCTTTTACTTTTATTTTGAAGTAATTATAGAATCACAGAAAGTT
GCAAAAAA

193/392

FIGURE 184

MGALIISGSSAGPVTQASLPPWGLSHGRCGFLLYMENTLCSHRTQSFSELSQSLMRPGF
LQMPYISCAKLSKIWFPAKPCLLAFLEVFLLMSRLSLFSKMICFLFLSFLFPPHIYTHAS

Important features of the protein:

Signal peptide:

amino acids 1-41

Transmembrane domain:

amino acids 88-107

Casein kinase II phosphorylation site:

amino acids 47-50

N-myristoylation site:

amino acids 24-29

194/392

FIGURE 185

AACTCAAACCTCCTCTCTCTGGGAAAACGCGGTGCTTGCTCCTCCCGGAGTGGCCTTGGCA
GGGTGTTGGAGCCCTCGGTCTGCCCCGTCCGGTCTCTGGGGCCAAGGCTGGGTTTCCCTC
ATGTATGGCAAGAGCTCTACTCGTGCGGTGCTTCTTCTCCTTGGCATAACAGCTCACAGCT
CTTTGGCCTATAGCAGCTGTGGAAATTTATACCTCCCGGGTGCTGGAGGCTGTTAATGGG
ACAGATGCTCGGTAAAATGCACTTTCTCCAGCTTTGCCCCCTGTGGGTGATGCTCTAACA
GTGACCTGGAATTTTCGTCTCTAGACGGGGGACCTGAGCAGTTTGTATTCTACTACCAC
ATAGATCCCTTCCAACCCATGAGTGGGCGGTTTAAGGACCGGGTGCTTGGGATGGGAAT
CCTGAGCGGTACGATGCCTCCATCCTTCTCTGGAACTGCAGTTCGACGACAATGGGACA
TACACCTGCCAGGTGAAGAACCACCTGATGTTGATGGGGTGATAGGGGAGATCCGGCTC
AGCGTCGTGCACACTGTACGCTTCTCTGAGATCCACTTCCTGGCTCTGGCCATTGGCTCT
GCCTGTGCACTGATGATCATAATAGTAATTGTAGTGGTCCTCTTCCAGCATTACCGGAAA
AAGCGATGGGCCGAAAGAGCTCATAAAGTGGTGAGATAAAATCAAAAGAAGAGGAAAGG
CTCAACCAAGAGAAAAAGGTCTCTGTTTATTTAGAAAGACACAGACTTACAATTTTAGATG
GAAGCTGAGATGATTTCCAAGAACAAAGAACCCTAGTATTTCTTGAAGTTAATGGAACTT
TTCTTTGGCTTTTCCAGTTGTGACCCGTTTTCCAACCAGTTCTGCAGCATATTAGATTCT
AGACAAGCAACACCCCTCTGGAGCCAGCACAGTGCTCCTCCATATCACCAGTCATACACA
GCCTCATTATTAAGGTCTTATTTAATTTTCAGAGTGTAATTTTTTCAAGTGCTCATTAGG
TTTTATAAACAAGAAGCTACATTTTGGCCCTTAAGACACTACTTACAGTGTTATGACTTG
TATACACATATATTGGTATCAAAGGGGATAAAAGCCAATTTGTCTGTTACATTTCTTTC
ACGTATTTCTTTTAGCAGCACTTCTGCTACTAAAGTTAATGTGTTTACTCTCTTTCCTTC
CCACATTCTCAATTAAAAGGTGAGCTAAGCCTCCTCGGTGTTTCTGATTAACAGTAAATC
CTAAATTCAAACGTAAATGACATTTTATTTTTATGTCTCTCCTTAACATATGAGACAC
ATCTTGTTTTACTGAATTTCTTTCAATATTCAGGTGATAGATTTTTTGTCTG

195/392

FIGURE 186

MYGKSSTRAVLLLLGIQLTALWPAAVEIYTSRVLEAVNGTDARLKCTFSSSFAPVGDALT
VTWNFRPLDGGPEQFVFYYHIDPFQPMSEGRFKDRVSWDGNPERYDASILLWKLQFDDNGT
YTCQVKNPPDVGVIQEIRLSVVHTVRFSEIHFLALAIGSACALMIIIVIVVVVLFQHYRK
KRWAEARAHKVVEIKSKEEERLNQEKKVSVYLEDTD

196/392

FIGURE 187

GCATTTTGTCTGTGCTCCCTGATCTTCAGGTCACCACCATGAAGTTCTTAGCAGTCCTG
GTACTCTTGGGAGTTTCCATCTTCTGGTCTCTGCCCAGAATCCGACAACAGCTGCTCCA
GCTGACACGTATCCAGCTACTGGTCCTGCTGATGATGAAGCCCCTGATGCTGAAACCACT
GCTGCTGCAACCACTGCGACCACTGCTGCTCCTACCACTGCAACCACCGCTGCTTCTACC
ACTGCTCGTAAAGACATTCCAGTTTTACCCAAATGGGTTGGGGATCTCCCGAATGGTAGA
GTGTGTCCCTGAGATGGAATCAGCTTGAGTCTTCTGCAATTGGTCACAACTATTCATGCT
TCCTGTGATTTTCATCCAACCTACTTACCTTGCTACGATATCCCCTTTATCTCTAATCAGT
TTATTTTCTTTCAAATAAAAAATAACTATGAGCAACATAAAAAAAAAAAAAA

197/392

FIGURE 188

MKFLAVLVLLGVSI FLVSAQNPTTAAPADTYPATGPADDEAPDAETTAAATTATTAAPT
ATTAASTTARKDIPVLPKWVGDL PNGRVCP

198/392

FIGURE 189

GAGCGAACATGGCAGCGCGTTGGCGGTTTTGGTGTGTCTCTGTGACCATGGTGGTGGCGC
TGCTCATCGTTTTGCGACGTTCCCTCAGCCTCTGCCCCAAAGAAAGAAGGAGATGGTGTAT
CTGAAAAGGTTAGTCAGCTGATGGAATGGACTAACAAAAGACCTGTAATAAGAATGAATG
GAGACAAGTTCGGTCGCCCTTGTGAAAGCCCCACCGAGAAATTACTCCGTTATCGTCATGT
TCACTGCTCTCCAAGTGCATAGACAGTGTGTGCTTTGCAAGCAAGCTGATGAAGAATTCC
AGATCCTGGCAAACTCCTGGCGATACTCCAGTGCATTACCAACAGGATATTTTTTGCCA
TGGTGGATTTTGATGAAGGCTCTGATGTATTTAGATGCTAAACATGAATTCAGCTCCAA
CTTTCATCAACTTTCTGCAAAAGGGAAACCCAAACGGGGTGATACATATGAGTTACAGG
TGCGGGGTTTTTTCAGCTGAGCAGATTGCCCGGTGGATCGCCGACAGAACTGATGTCAATA
TTAGAGTGATTAGACCCCAATTATGCTGGTCCCTTATGTTGGGATTGCTTTTGGCTG
TTATTGGTGGACTTGTGTATCTTCGAAGAAGTAATATGGAATTTCTCTTTAATAAACTG
GATGGGCTTTTGCAGCTTTGTGTTTTGTGCTTGCTATGACATCTGGTCAAATGTGGAACC
ATATAAGAGGACCACCATATGCCATAAGAATCCCCACACGGGACATGTGAATTATATCC
ATGGAAGCAGTCAAGCCAGTTTGTAGCTGAAACACACATTGTTCTTCTGTTTAAATGGTG
GAGTTACCTTAGGAATGGTGCTTTTATGTGAAGCTGCTACCTCTGACATGGATATTGGAA
AGCGAAAGATAATGTGTGTGGCTGGTATTGGACTTGTGTATTATTCTTCAGTTGGATGC
TCTCTATTTTTAGATCTAAATATCATGGCTACCCATACAGCTTTCTGATGAGTTAAAAAG
GTCCCAGAGATATATAGACACTGGAGTACTGGAAATTGAAAAACGAAAAATCGTGTGTGT
TGAAAAGAAGAATGCAACTTGTATATTTTGTATTACCTCTTTTTTTCAAGTGATTTAAAT
AGTTAATCATTTAACCAGAAAGATGTGTAGTGCCTTAACAAGCAATCCTCTGTCAAAAT
CTGAGGTATTTGAAAATAATTATCCTCTTAACCTTCTCTTCCAGTGAACCTTATGGAAC
ATTTAATTTAGTACAATTAAGTATATTATAAAAATTGTAAACTACTACTTTGTTTTAGT
TAGAACAAAGCTCAAACTACTTTAGTTAACTTGGTCATCTGATTTTATATTGCCTTATC
CAAAGATGGGGAAAGTAAGTCCTGACCAGGTGTTCCACATATGCCTGTTACAGATAACT
ACATTAGGAATTCATTCTTAGCTTCTTCATCTTTGTGTGGATGTGTATACTTTACGCATC
TTTCTTTTGAGTAGAGAAATTATGTGTGTCATGTGGTCTTCTGAAAATGGAACACCATT
CTTCAGAGCACACGTCTAGCCCTCAGCAAGACAGTTGTTTCTCCTCCTCCTTGCATATTT
CCTACTGCGCTCCAGCCTGAGTGATAGAGTGAGACTCTGTCTCAAAAAAAGTATCTCTA
AATACAGGATTATAATTTCTGCTTGAGTATGGTGTAACTACCTTGTATTTAGAAAGATT
TCAGATTCATTCCATCTCCTTAGTTTTCTTTAAGGTGACCCATCTGTGATAAAAATATA
GCTTAGTGCTAAAATCAGTGTAACCTTATACATGGCCTAAAATGTTTCTACAAATTAGAGT
TTGTCACTTATTCATTTGTACCTAAGAGAAAAATAGGCTCAGTTAGAAAAGGACTCCCT
GGCCAGGCGCAGTGACTTACGCCTGTAATCTCAGCACTTTGGGAGGCCAAGGCAGGCAGA
TCACGAGGTCAGGAGTTCGAGACCATCCTGGCCAACATGGTGAAACCCCGTCTCTACTAA
AAATATAAAAATTAGCTGGGTGTGGTGGCAGGAGCCTGTAATCCCAGCTACACAGGAGGC
TGAGGCACGAGAATCACTTGAACCTCAGGAGATGGAGGTTTCAGTGAGCCGAGATCACGCC
ACTGCACTCCAGCCTGGCAACAGAGCGAGACTCCATCTCAAAAAAAAAAAAAA

199/392

FIGURE 190

MAARWRFWCVSVTMVVALLVCDVPSASAQRKKEMVLSEKVSQLMWETNKRFPVIRMNGDK
FRRLVKAPPRNYSVIVMFTALQLHRQCVVCKQADEEFQILANSWRYSSAFTNRIFFAMVD
FDEGSDVFQMLNMNSAPTFINFPKAGKPKRGDTYELQVRGFSAEQIARWIADRTDVNIRV
IRPPNYAGPLMLGLLLAVIGGLVYLRRSNMEFLFNKTGWAFALCFVLAMTSGQMWNHIR
GPPYAHKNPHTGHVNYIHGSSQAQFVAETHIVLLFNGGVTLGMVLLCEAATSDMDIGKRK
IMCVAGIGLVVLFFSWMLSIFRSKYHGYPSFLMS

Signal peptide:
amino acids 1-29

Transmembrane domains:
amino acids 183-205, 217-237, 217-287, 301-321

200/392

FIGURE 191

GAGAGAAGTCAGCCTGGCAGAGAGACTCTGAAATGAGGGATTAGAGGTGTTCAAGGAGCA
AGAGCTTCAGCCTGAAGACAAGGGAGCAGTCCCTGAAGACGCTTCTACTGAGAGGTCTGC
CATGGCCTCTCTTGGCCTCCAACTTGTGGGCTACATCCTAGGCCTTCTGGGGCTTTTGGG
CACACTGGTTGCCATGCTGCTCCCCAGCTGGAAAACAAGTTCTTATGTCGGTGCCAGCAT
TGTGACAGCAGTTGGCTTCTCCAAGGGCCTCTGGATGGAATGTGCCACACACAGCACAGG
CATCACCCAGTGTGACATCTATAGCACCTTCTGGGCCTGCCCGCTGACATCCAGGCTGC
CCAGGCCATGATGGTGACATCCAGTGCAATCTCCTCCCTGGCCTGCATTATCTCTGTGGT
GGGCATGAGATGCACAGTCTTCTGCCAGGAATCCCGAGCCAAAGACAGAGTGGCGGTAGC
AGGTGGAGTCTTTTTCATCCTTGGAGGCCTCCTGGGATTCAATCCTGTTGCCTGGAATCT
TCATGGGATCCTACGGGACTTCTACTCACCCTGGTGCCTGACAGCATGAAATTTGAGAT
TGGAGAGGCTCTTTACTTGGGCATTATTTCTTCCCTGTTCTCCCTGATAGCTGGAATCAT
CCTCTGCTTTTCTGCTCATCCCAGAGAAATCGCTCCAACCTACTACGATGCCTACCAAGC
CCAACCTCTTGCCACAAGGAGCTCTCCAAGGCCTGGTCAACCTCCCAAAGTCAAGAGTGA
GTTCAATTCCTACAGCCTGACAGGGTATGTGTGAAAGAACCAGGGGCCAGAGCTGGGGGGT
GGCTGGGTCTGTGAAAAACAGTGGACAGCACCCCGAGGGCCACAGGTGAGGGACACTACC
ACTGGATCGTGTGAGAAGGTGCTGCTGAGGATAGACTGACTTTGGCCATTGGATTGAGCA
AAGGCAGAAATGGGGGCTAGTGTAACAGCATGCAGGTTGAATTGCCAAGGATGCTCGCCA
TGCCAGCCTTTCTGTTTTCTCACCTTGCTGCTCCCCTGCCCTAAGTCCCCAACCCTCAA
CTTGAAACCCCATTCCTTAAGCCAGGACTCAGAGGATCCCTTTGCCCTCTGGTTTACCT
GGGACTCCATCCCCAAACCCACTAATCACATCCCCTGACTGACCCTCTGTGATCAAAGA
CCCTCTCTCTGGCTGAGGTTGGCTCTTAGCTCATTGCTGGGGATGGGAAGGAGAAGCAGT
GGCTTTTGTGGGCATTGCTCTAACCTACTTCTCAAGCTTCCCTCCAAAGAACTGATTGG
CCCTGGAACCTCCATCCCCTCTTGTATGACTCCACAGTGTCCAGACTAATTTGTGCAT
GAACTGAAATAAAACCATCCTACGGTATCCAGGGAACAGAAAGCAGGATGCAGGATGGGA
GGACAGGAAGGCAGCCTGGGACATTTAAAAAATA

201/392

FIGURE 192

MASLGLQLVGYILGLLGLLGTLVAMLLPSWKTSSYVGASIVTAVGFSKGLWMECATHSTG
ITQCDIYSTLLGLPADIQAAQAMMTSSAISSLACIISVVGMRCTVFCQESRAKDRVAVA
GGVFFILGGLLGFIPIVAVNLHGILRDFYSPLVPDSMKFEIGEALYLGIISSLFSLIAGII
LCFSCSSQRNRSNYYDAYQAQPLATRSSPRPGQPPKVKSEFNYSYSLTGYV

Important features of the protein:

Signal peptide:

amino acids 1-24

Transmembrane domains:

amino acids 82-102, 117-140, 163-182

N-glycosylation site:

amino acids 190-193

PMP-22 / EMP / MP20 family proteins:

amino acids 46-59

202/392

FIGURE 193

CTCCACTGCAACCACCCAGAGCCATGGCTCCCCGAGGCTGCATCGTAGCTGTCTTTGCCA
TTTTCTGCATCTCCAGGCTCCTCTGCTCACACGGAGCCCCAGTGGCCCCCATGACTCCTT
ACCTGATGCTGTGCCAGCCACACAAGAGATGTGGGGACAAGTTCTACGACCCCCCTGCAGC
ACTGTTGCTATGATGATGCCGTCGTGCCCTTGGCCAGGACCCAGACGTGTGGAACTGCA
CCTTCAGAGTCTGCTTTGAGCAGTGCTGCCCCCTGGACCTTCATGGTGAAGCTGATAAACC
AGAACTGCGACTCAGCCCGGACCTCGGATGACAGGCTTTGTGCGCAGTGTCAGCTAATGGA
ACATCAGGGGAACGATGACTCCTGGATTCTCCTTCCTGGGTGGGCCTGGAGAAAGAGGCT
GGTGTTACCTGAGATCTGGGATGCTGAGTGGCTGTTTGGGGGCCAGAGAAAACACACTC
AACTGCCCACTTCATTCTGTGACCTGTCTGAGGCCCACCCTGCAGCTGCCCTGAGGAGGC
CCACAGGTCCCCCTTCTAGAATTCTGGACAGCATGAGATGCGTGTGCTGATGGGGGCCAG
GGACTCTGAACCCTCCTGATGACCCCTATGGCCAACATCAACCCGGCACCACCCCAAGGC
TGGCTGGGGAACCCCTCACCCCTTCTGTGAGATTTTCCATCATCTCAAGTTCTCTTCTATC
CAGGAGCAAAGCACAGGATCATAATAAATTTATGTACTTTATAAATGAAA

203/392

FIGURE 194

MAPRGCIVAVFAIFCISRLLC SHGAPVAPMTPYLM LCQPHKRCGDKFYDPLQHCCYDDAV
VPLARTQTCGNCTFRVCFEQCCPWTFMVKLINQNCD SARTSDDRLCRSVS

Signal peptide:
amino acids 1-24

204/392

FIGURE 195

CATTTCCAACAAGAGCACTGGCCAAGTCAGCTTCTTCTGAGAGAGTCTCTAGAAGACATG
ATGCTACACTCAGCTTTGGGTCTCTGCCTCTTACTCGTCACAGTTTCTTCCAACCTTGCC
ATTGCAATAAAAAAGGAAAAGAGGCCTCCTCAGACACTCTCAAGAGGATGGGGAGATGAC
ATCACTTGGGTACAACTTATGAAGAAGGTCTCTTTTATGCTCAAAAAAGTAAGAAGCCA
TTAATGGTTATTTCATCACCTGGAGGATTGTCAATACTCTCAAGCACTAAAGAAAGTATTT
GCCCCAAATGAAGAAATACAAGAAATGGCTCAGAATAAGTTCATCATGCTAAACCTTATG
CATGAAACCACTGATAAGAATTTATCACCTGATGGGCAATATGTGCCTAGAATCATGTTT
GTAGACCCTTCTTTAACAGTTAGAGCTGACATAGCTGGAAGATACTCTAACAGATTGTAC
ACATATGAGCCTCGGGATTTACCCCTATTGATAGAAAACATGAAGAAAGCATTAAAGACTT
ATTCAGTCAGAGCTATAAGAGATGATGGAAAAAGCCTTCACTTCAAAGAAGTCAAATTT
CATGAAGAAAACCTCTGGCACATTGACAAATACTAAATGTGCAAGTATATAGATTTTGTA
ATATTACTATTTAGTTTTTTTAAATGTGTTTGCAATAGTCTTATTAAAATAAATGTTTTTT
AAATCTGA

205/392

FIGURE 196

MMLHSALGLCLLLVTVSSNLAIKKEKRPPQTL SRGWGDDITWVQTYEEGLFYAQSKK
PLMVIHHLED CQYSQALKKVFAQNEEIQEMAQNK FIMLNLMHETTDKNLSPDGQYVPRIM
FVDPSLTVRADIAGRYSNRLYTYEPRDLPLLIENMKKALRLIQSEL

Important features:

Signal peptide:

amino acids 1-23

N-myristoylation site:

amino acids 51-57

206/392

FIGURE 197

GGGGGCGGGTGCCTGGAGCACGGCGCTGGGGCCGCCCCGAGCGCTCACTCGCTCGCACTC
AGTCGCGGGAGGCTTCCCCGCGCCGGCCGCGTCCCCGCCGCTCCCCGGCACCAGAAGTTC
CTCTGCGCGTCCGACGGCGACATGGGCGTCCCCACGGCCCTGGAGGCCGGCAGCTGGCGC
TGGGGATCCCTGCTCTTCGCTCTCTTCCTGGCTGCGTCCCTAGGTCCGGTGGCAGCCTTC
AAGGTCGCCACGCCGTATTCCCTGTATGTCTGTCCCGAGGGGCAGAACGTCAACCTCACC
TGCAGGCTCTTGGGCCCCTGTGGACAAAGGGCACGATGTGACCTTCTACAAGACGTGGTAC
CGCAGCTCGAGGGGCGAGGTGCAGACCTGCTCAGAGCGCCGGCCCATCCGCAACCTCACG
TTCCAGGACCTTCACCTGCACCATGGAGGCCACCAGGCTGCCAACACCAGCCACGACCTG
GCTCAGCGCCACGGGCTGGAGTCGGCCTCCGACCACCATGGCAACTTCTCCATCACCATG
CGCAACCTGACCCCTGCTGGATAGCGGCCTCTACTGCTGCCTGGTGGTGGAGATCAGGCAC
CACCATCCGAGCACAGGGTCCATGGTGCATGGAGCTGCAGGTGCAGACAGGCAAAGAT
GCACCATCCAACGTGTGTGGTGTACCCATCCTCTCCAGGATAGTGAAAACATCACGGCT
GCAGCCCTGGCTACGGGTGCCTGCATCGTAGGAATCCTCTGCCTCCCCCTCATCCTGCTC
CTGGTCTACAAGCAAAGGCAGGCAGCCTCCAACCGCCGTGCCAGGAGCTGGTGC GGATG
GACAGCAACATTCAAGGGATTGAAAACCCCGGCTTTGAAGCCTCACCACCTGCCAGGGG
ATACCCGAGGCCAAAGTCAGGCACCCCTGTCTATGTGGCCAGCGGCAGCCTTCTGAG
TCTGGGCGGCATCTGCTTTCGGAGCCCAGCACCCCTGTCTCCTCCAGGCCCGGAGAC
GTCTTCTTCCCATCCCTGGACCTGTCCCTGACTCTCCAAACTTTGAGGTCTCTAGCCC
AGCTGGGGGACAGTGGGCTGTTGTGGCTGGGTCTGGGGCAGGTGCATTTGAGCCAGGGCT
GGCTCTGTGAGTGGCCTCCTTGGCCTCGGCCCTGGTTCCTCCCTCCTGCTCTGGGCTCA
GATACTGTGACATCCAGAACCCAGCCCTCAACCCCTCTGGATGCTACATGGGGATGC
TGGACGGCTCAGCCCCGTGTTCCAAGGATTTTGGGGTGCTGAGATTCTCCCCTAGAGACCT
GAAATTCACCAGCTACAGATGCCAAATGACTTACATCTTAAGAAGTCTCAGAACGTCCAG
CCCTTCAGCAGCTCTCGTTCTGAGACATGAGCCTTGGGATGTGGCAGCATCAGTGGGACA
AGATGGACACTGGGGCCACCCTCCCAGGCACCAGACACAGGGCACGGTGGAGAGACTTCTC
CCCCGTGGCCGCCTTGGCTCCCCCGTTTTGCCCGAGGCTGCTCTTCTGTGAGACTTCTC
TTTGTACCACAGTGGCTCTGGGGCCAGGCCTGCCTGCCCACTGGCCATCGCCACCTTCCC
CAGCTGCCTCCTACCAGCAGTTTCTCTGAAGATCTGTCAACAGGTTAAGTCAATCTGGGG
CTTCCACTGCCTGCATTCCAGTCCCCAGAGCTTGGTGGTCCCGAAACGGGAAGTACATAT
TGGGGCATGGTGGCCTCCGTGAGCAATGGTGTCTTGGGCAATCTGAGGCCAGGACAGAT
GTTGCCCCACCCACTGGAGATGGTGTCTGAGGGAGGTGGGTGGGGCCTTCTGGGAAGGTGA
GTGGAGAGGGGCACCTGCCCCCGCCCTCCCCTACTCCCCTGCTCAGCGCGGG
CCATTGCAAGGGTGCCACACAATGTCTTGTCCACCCTGGGACACTTCTGAGTATGAAGCG
GGATGCTATTAAAACTACATGGGGAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAGA

207/392

FIGURE 198

MGVPTALEAGSWRWGSLLFALFLAASLGPVAAFQVATPYSLYVCPEGQNVTLTCRLLGPV
DKGHDVTFYKTYWRSSRGEVQTCSERRPIRNLTFQDLHLHHGGHQAANTSHDLAQRHGLE
SASDHHGNFSITMRNLTLDSGLYCCLVVEIRHHHSEHRVHGAMELQVQTGKDAPSNCVV
YPSSSQDSENITAAALATGACIVGILCLPLILLLVYKQRQAASNRRRAQELVRMDSNIQGI
ENPGFEASPPAQGIPEAKVRHPLSYVAQRQPSESGRHLLSEPSTPLSPPGPGDVFFPSLD
PVPDSPNFEVI

Signal peptide:
amino acids 1-28

Transmembrane domain:
amino acids 190-216

208/392

FIGURE 199

CTAGCCTGCGCCAAGGGGTAGTGAGACCGCGCGGCAACAGCTTGCGGCTGCGGGGAGCTC
CCGTGGGCGCTCCGCTGGCTGTGCAGGCGGCCATGGATTCTTGCGGAAAATGCTGATCT
CAGTCGCAATGCTGGGCGCAGGGGCTGGCGTGGGCTACGCGCTCCTCGTTATCGTGACCC
CGGGAGAGCGGCGGAAGCAGGAAATGCTAAAGGAGATGCCACTGCAGGACCCAAGGAGCA
GGGAGGAGGCGGCCAGGACCCAGCAGCTATTGCTGGCCACTCTGCAGGAGGCAGCGACCA
CGCAGGAGAACGTGGCCTGGAGGAAGAACTGGATGGTTGGCGGCGAAGGCGGCGCCAGCG
GGAGGTCACCGTGAAGACCGGACTTGCCTCCGTGGGCGCCGGACCTTGGCTTGGGCGCAGG
AATCCGAGGCAGCCTTTCTCCTTCGTGGGCCAGCGGAGAGTCCGACCGAGATACCATG
CCAGGACTCTCCGGGGTCCTGTGAGCTGCCGTCCGGTGAGCACGTTTCCCCCAAACCCTG
GACTGACTGCTTTAAGGTCCGCAAGGCGGGCCAGGGCCGAGACGCGAGTCGGATGTGGTG
AACTGAAAGAACCAATAAAATCATGTTCTCCAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

209/392

FIGURE 200

MDSLRLKMLISVAMLGAGAGVGYALLVIVTPGERRKQEMLKEMPLQDPRSREEAARTQQLL
LATLQEAATTQENVAWRKNNMVGEGGASGRSP

Signal peptide:
amino acids 1-18

210/392

FIGURE 201

GACAGCTGTGTCTCGATGGAGTAGACTCTCAGAACAGCGCAGTTTGGCCCTCCGCTCACGC
AGAGCCTCTCCGTGGCTTCCGCACCTTGAGCATTAGGCCAGTTCTCCTCTTCTCTAAT
CCATCCGTACCTCTCCTGTTCATCCGTTTCCATGCCGTGAGGTCCATTACAGAACACAT
CCATGGCTCTCATGCTCAGTTTGGTTCTGAGTCTCCTCAAGCTGGGATCAGGGCAGTGGC
AGGTGTTTGGGCCAGACAAGCCTGTCCAGGCCCTTGGTGGGGGAGGACGCAGCATTCTCCT
GTTTCTGTCTCCTAAGACCAATGCAGAGGCCATGGAAGTGCAGTTCTTTCAGGGGCCAGT
TCTCTAGCGTGGTCCACCTCTACAGGGACGGGAAGGACCAGCCATTTATGCAGATGCCAC
AGTATCAAGGCAGGACAAAACCTGGTGAAGGATTCTATTGCGGAGGGGCGCATCTCTCTGA
GGCTGGAAAACATTACTGTGTTGGATGCTGGCCTCTATGGGTGCAGGATTAGTTCCAGT
CTTACTACCAGAAGGCCATCTGGGAGCTACAGGTGTGAGCACTGGGCTCAGTTCTCTCA
TTTCCATCACGGGATATGTTGATAGAGACATCCAGCTACTCTGTGAGTCTCGGGCTGGT
TCCCCCGGGCCACAGCGAAGTGGAAAGGTCCACAAGGACAGGATTTGTCCACAGATCCA
GGACAAACAGAGACATGCATGGCCTGTTTGTGATGTGGAGATCTCTCTGACCGTCCAAGAGA
ACGCCGGGAGCATATCCTGTTCCATGCGGCATGCTCATCTGAGCCGAGAGGTGGAATCCA
GGGTACAGATAGGAGATACCTTTTTTCGAGCCTATATCGTGGCACCTGGCTACCAAAGTAC
TGGAATACTCTGCTGTGGCCTATTTTTTGGCATTGTTGGACTGAAGATTTTCTTCTCCA
AATTCAGTGGAATAATCCAGGCGGAACCTGGACTGGAGAAGAAAGCACGGACAGGCAGAAT
TGAGAGACGCCCCGAAACACGCAGTGGAGGTGACTCTGGATCCAGAGACGGCTCACCCGA
AGCTCTGCGTTTCTGATCTGAAAACCTGTAACCCATAGAAAAGCTCCCCAGGAGGTGCCTC
ACTCTGAGAAGAGATTTACAAGGAAGAGTGTGGTGGCTTCTCAGAGTTTCCAAGCAGGGA
AACATTACTGGGAGGTGGACGGAGGACACAATAAAAGGTGGCGCTGGGAGTGTGCCGGG
ATGATGTGGACAGGAGGAAGGAGTACGTGACTTTGTCTCCCGATCATGGGTACTGGGTCC
TCAGACTGAATGGAGAACATTTGTATTTTACATTAAATCCCCGTTTATCAGCGTCTTCC
CCAGGACCCACCTACAAAAATAGGGGTCTTCTGGACTATGAGTGTGGGACCATCTCCT
TCTTCAACATAAATGACCAGTCCCTTATTTATACCTGACATGTCCGTTTGAAGGCTTAT
TGAGGCCCTACATTGAGTATCCGTCTTATAATGAGCAAAATGGAACCTCCCATAGTCATCT
GCCCAGTCACCCAGGAATCAGAGAAAGAGGCCTCTTGGCAAAGGGCCTCTGCAATCCCAG
AGACAAGCAACAGTGAGTCTCTCTCACAGGCAACCACGCCCTTCTCCCCAGGGGTGAAA
TGTAAGGATGAATCACATCCCACATTCTTCTTTAGGGATATTAAGGTCTCTCTCCAGATC
CAAAGTCCCGCAGCAGCCGGCCAAGGTGGCTTCCAGATGAAGGGGGACTGGCCTGTCCAC
ATGGGAGTCAGGTGTCTATGGCTGCCCTGAGCTGGGAGGGAAGAAGGCTGACATTACATTT
AGTTTGCTCTCACTCCATCTGGCTAAGTGATCTTGAAATACCACCTCTCAGGTGAAGAAC
CGTCAGGAATTCCCATCTCACAGGCTGTGGTGTAGATTAAGTAGACAAGGAATGTGAATA
ATGCTTAGATCTTATTGATGACAGAGTGTATCCTAATGGTTTGTTCATTATATTACACTT
TCAGTAAAAAAA

211/392

FIGURE 202

MALMLSLVLSLLKLGSGQWQVFGPDKPVQALVGEDAAFSCFLSPKTNAEAMEVRFFRGQF
SSVVHLYRDGKDQPFMQMPQYQGRTKLVKDSIAEGRISLRLENITVLDAGLYGCRISQS
YYQKAIWELQVSALGSVPLISITGYVDRDIQLLCQSSGWFFRPTAKWKGPQGQDLSTDSR
TNRDMHGLFDVEISLTVQENAGSISCSMRHAHLSREVESRVQIGDTFFEPISWHLATKVL
GILCCGLFFGIVGLKIFFSKFQWKIQAELDWRRKHGQAEIRDARKHAVEVTLDPETAHPK
LCVSDLKTVTHRKAPEVPHSEKRFTKSVVASQSFQAGKHYWEVDGGHNKRWRVGVCRD
DVDRRKEYVTLSPDHGYWVLRLNGEHLYFTLNPRFISVFPRTPTKIGVFLDYECGTISF
FNINDQSLIYTLTCRFEGLLRPYIEYPSYNEQNGTPIVICPVTQESEKEASWQRASAIPE
TSNSESSSQATTPLPRGEM

Signal peptide:
amino acids 1-17

Transmembrane domain:
amino acids 239-255

212/392

FIGURE 203

TGCGGCGCAGTGCTAGACCTGGGAGGATGGGCGGCCTGCTGCTGGCTGCTTTTCTGGCTTT
GGTCTCGGTGCCCAGGGCCCAGGCCGTGTGGTTGGGAAGACTGGACCCTGAGCAGCTTCT
TGGGCCCTGGTACGTGCTTGCGGTGGCCTCCCGGAAAAGGGCTTTGCCATGGAGAAGGA
CATGAAGAACGTCGTGGGGGTGGTGGTGACCCTCACTCCAGAAAACAACCTGCGGACGCT
GTCCTCTCAGCACGGGCTGGGAGGGTGTGACCAGAGTGTCATGGACCTGATAAAGCGAAA
CTCCGGATGGGTGTTTGAGAATCCCTCAATAGGCGTGCTGGAGCTCTGGGTGCTGGCCAC
CAACTTCAGAGACTATGCCATCATCTTCACTCAGCTGGAGTTCGGGGACGAGCCCTTCAA
CACCGTGGAGCTGTACAGTCTGACGGAGACAGCCAGCCAGGAGGCCATGGGGCTCTTCAC
CAAGTGGAGCAGGAGCCTGGGCTTCTGTGTCACAGTAGCAGGCCCAGCTGCAGAAGGACCT
CACCTGTGCTCACAAGATCCTTCTGTGAGTGCTGCGTCCCCAGTAGGGATGGCGCCCACA
GGGTCTGTGACCTCGGCCAGTGTCACCCACCTCGCTCAGCGGCTCCCGGGGCCAGCA
CCAGCTCAGAATAAAGCGATTCCACAGCA

213/392

FIGURE 204

MGGLLLAAFLALVSVPRQAVWLGRDLPEQLLGPWYVLAVASREKGFAMEKDMKNVVGVV
VTLTPENNLRTLSSQHGLGGCDQSVMDLIKRNSGWVFENPSIGVLELWVLATNFRDYAI I
FTQLEFGDEPFNTVELYSITETASQEAMGLFTKWSRSLGFLSQ

Signal peptide:
amino acids 1-20

214/392

FIGURE 205

GACGCCCAGTGACCTGCCGAGGTCCGGCAGCACAGAGCTCTGGAGATGAAGACCCTGTTCC
TGGGTGTCACGCTCGGCCTGGCCGCTGCCCTGTCCTTCACCCTGGAGGAGGAGGATATCA
CAGGGACCTGGTACGTGAAGGCCATGGTGGTCGATAAGGACTTTCGGAGGACAGGAGGC
CCAGGAAGGTGTCCCCAGTGAAGGTGACAGCCCTGGGCGGTGGGAAGTTGGAAGCCACGT
TCACCTTCATGAGGGAGGATCGGTGCATCCAGAAGAAAATCCTGATGCGGAAGACGGAGG
AGCCTGGCAAATACAGCGCCTATGGGGGCAGGAAGCTCATGTACCTGCAGGAGCTGCCCCA
GGAGGGACCACTACATCTTTTACTGCAAAGACCAGCACCATGGGGGCCTGCTCCACATGG
GAAAGCTTGTGGGTAGGAATTCTGATACCAACCGGGAGGCCCTGGAAGAATTTAAGAAAT
TGGTGCAGCGCAAGGGACTCTCGGAGGAGGACATTTTCACGCCCCTGCAGACGGGAAGCT
GCGTTCCCGAACACTAGGCAGCCCCCGGGTCTGCACCTCCAGAGCCCACCCTACCACCAG
ACACAGAGCCCCGGACCACCTGGACCTACCCTCCAGCCATGACCCTTCCCTGCTCCCACCC
ACCTGACTCCAAATAAAGTCCTTTTCCCCCAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAA

215/392

FIGURE 206

MKTLFLGVTLGLAAALSFTLEEDITGTWYVKAMVVDKDFPEDRRPRKVSPVKVTALGGG
KLEATFTFMREDRCIQKKILMRKTEEPGKYSAYGGRKLMYLQELPRRDHYIFYCKDQHHG
GLLHMGKLVGRNSDTNREALEEFKKLVQRKGLSEEDIFTPLQTGSCVPEH

Important features:

Signal peptide:

amino acids 1-17

216/392

FIGURE 207

GTTCCGCAGATGCAGAGGTTGAGGTGGCTGCGGGACTGGAAGTCATCGGGCAGAGGTCTC
ACAGCAGCCAAGGAACCTGGGGCCCGCTCCTCCCCCTCCAGGCCATGAGGATTCTGCAG
TTAATCCTGCTTGCTCTGGCAACAGGGCTTGTAGGGGGAGAGACCAGGATCATCAAGGGG
TTCGAGTGCAAGCCTCACTCCCAGCCCTGGCAGGCAGCCCTGTTTCGAGAAGACGCGGCTA
CTCTGTGGGGCGACGCTCATCGCCCCAGATGGCTCCTGACAGCAGCCCACTGCCTCAAG
CCCCGCTACATAGTTCACCTGGGGCAGCACAACTCCAGAAGGAGGAGGGCTGTGAGCAG
ACCCGGACAGCCACTGAGTCCTTCCCCACCCCGGCTTCAACAACAGCCTCCCCAACAAA
GACCACCGCAATGACATCATGCTGGTGAAGATGGCATCGCCAGTCTCCATCACCTGGGCT
GTGCGACCCCTCACCCCTCTCCTCACGCTGTGTCACTGCTGGCACCAGCTGCCTCATTTCC
GGCTGGGGCAGCACGTCCAGCCCCAGTTACGCCTGCCTCACACCTTGCGATGCGCCAAC
ATCACCATCATTGAGCACCAGAAGTGTGAGAACGCCTACCCCGGCAACATCACAGACACC
ATGGTGTGTGCCAGCGTGCAGGAAGGGGGCAAGGACTCCTGCCAGGGTGACTCCGGGGGC
CCTCTGGTCTGTAAACCAGTCTCTTCAAGGCATTATCTCCTGGGGCCAGGATCCGTGTGCG
ATCACCCGAAAGCCTGGTGTCTACACGAAAGTCTGCAAATATGTGGACTGGATCCAGGAG
ACGATGAAGAACAATTAGACTGGACCCACCCACCACAGCCCATCACCTCCATTTCCACT
TGGTGTGTTGGTTCTGTTCACTCTGTTAATAAGAAACCCTAAGCCAAGACCTCTACGAA
CATTCTTTGGGCCTCCTGGACTACAGGAGATGCTGTCACTTAATAATCAACCTGGGGTTC
GAAATCAGTGAGACCTGGATTCAAATTCTGCCTTGAAATATTGTGACTCTGGGAATGACA
ACACCTGGTTTGTCTCTGTTGTATCCCCAGCCCCAAAGACAGCTCCTGGCCATATATCA
AGGTTTCAATAAATATTTGCTAAATGAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAA

217/392

FIGURE 208

MRILQLILLALATGLVGGETRIIKGFECKPHSQPWQAALFEKTRLLCGATLIAPRWLLTA
AHCLKPRYIVHLGQHNLQKEEGCEQTRTATESFPHPGFNNSLPNKDHRNDIMLVKMASPV
SITWAVRPLTLSSRCVTAGTSCLISGWGSTSSPQLRLPHTLRCANITIIHQKCENAYPG
NITDTMVCASVQEGGKSDSCQGDSSGGLVCNQSLQGIISWGQDPCAITRKPGVYTKVCKYV
DWIQETMKNN

Important features:**Signal peptide:**

amino acids 1-18

Serine proteases, trypsin family, histidine active site:

amino acids 58-63

N-glycosylation sites:

amino acids 99-102, 165-168, 181-184, 210-213

Glycosaminoglycan attachment site:

amino acids 145-148

Kringle domain proteins:

amino acids 197-209, 47-64

Serine proteases, trypsin family, histidine protein:

amino acids 199-209, 47-63, 220-243

Apple domain proteins:

amino acids 222-249, 189-222

218/392

FIGURE 209

GCGGCCACACGCAGCTAGCCGGAGCCCGGACCAGGCGCCTGTGCCTCCTCCTCGTCCCTC
GCCGCGTCCGCGAAGCCTGGAGCCGGCGGGAGCCCCGCGCTCGCCATGTCGGGCGAGCTC
AGCAACAGGTTCCAAGGAGGGAAGGCGTTCGGCTTGCTCAAAGCCCGGCAGGAGAGGAGG
CTGGCCGAGATCAACCGGGAGTTTCTGTGTGACCAGAAGTACAGTGATGAAGAGAACCTT
CCAGAAAAGCTCACAGCCTTCAAAGAGAAGTACATGGAGTTTGACCTGAACAATGAAGGC
GAGATTGACCTGATGTCTTTAAAGAGGATGATGGAGAAGCTTGGTGTCCCCAAGACCCAC
CTGGAGATGAAGAAGATGATCTCAGAGGTGACAGGAGGGGTGACGTGACACTATATCCTAC
CGAGACTTTGTGAACATGATGCTGGGGAAACGGTCGGCTGTCCTCAAGTTAGTCATGATG
TTTGAAGGAAAAGCCAACGAGAGCAGCCCCAAGCCAGTTGGCCCCCTCCAGAGAGAGAC
ATTGCTAGCCTGCCCTTGAGGACCCCGCCTGGACTCCCAGCCTTCCCACCCCATACCTCC
CTCCCGATCTTGCTGCCCTTCTTGACACACTGTGATCTCTCTCTCTCTCATTGTGTTGGT
CATTGAGGGTTTGTGTTGTGTTTTTCATCAATGTCTTTGTAAAGCACAAATTATCTGCCTTA
AAGGGGCTCTGGGTGCGGGAATCCTGAGCCTTGGGTCCCCTCCCTCTCTTCTTCCCTCCT
TCCCCGCTCCCTGTGCAGAAGGGCTGATATCAAACCAAAAAGTAGAGGGGGCAGGGCCAG
GGCAGGGAGGCTTCCAGCCTGTGTTCCCCTCACTTGGAGGAACCAGCACTCTCCATCCTT
TCAGAAAGTCTCCAAGCCAAGTTCAGGCTCACTGACCTGGCTCTGACGAGGACCCCAGGC
CACTCTGAGAAGACCTTGGAGTAGGGACAAGGCTGCAGGGCCTCTTTCGGGTTTCTTGG
ACAGTGCCATGGTTCCAGTGCTCTGGTGTCACCCAGGACACAGCCACTCGGGGCCCCGCT
GCCCCAGCTGATCCCCACTCATTCCACACCTCTTCTCATCCTCAGTGATGTGAAGGTGGG
AAGGAAAGGAGCTTGGCATTGGGAGCCCTTCAAGAAGGTACCAGAAGGAACCCTCCAGTC
CTGCTCTCTGGCCACACCTGTGCAGGCAGCTGAGAGGCAGCGTGCAGCCCTACTGTCCCT
TACTGGGGCAGCAGAGGGCTTCGGAGGCAGAAGTGAGGCCTGGGGTTTGGGGGGAAAGGT
CAGCTCAGTGCTGTTCCACCTTTTAGGGAGGATACTGAGGGGACCAGGATGGGAGAATGA
GGAGTAAAATGCTCACGGCAAAGTCAGCAGCACTGGTAAGCCAAGACTGAGAAATACAAG
GTTGCTTGTCTGACCCCAATCTGCTTGAAAAAAAAAAAAAAAAAAAAA

219/392

FIGURE 210

MSGELSNRFQGGKAFGLLKARQERRLAELNREFLCDQKYSDEENLPEKLTAFKEKYMED
LNNEGEIDLMSLKRMMEKLGVPKTHLEMKKMISEVTGGVSDTISYRDFVNMMLGKRSAVL
KLVMFEGKANESSPKPVGPPPERDIASLP

220/392

FIGURE 211

CTGGGATCAGCCACTGCAGCTCCCTGAGCACTCTCTACAGAGACGCGGACCCAGACATG
AGGAGGCTCCTCCTGGTCACCAGCCTGGTGGTTGTGCTGCTGTGGGAGGCAGGTGCAGTC
CCAGCACCCAAGGTCCCTATCAAGATGCAAGTCAAACACTGGCCCTCAGAGCAGGACCCA
GAGAAGGCCTGGGGCGCCCGTGTGGTGGAGCCTCCGGAGAAGGACGACCAGCTGGTGGTG
CTGTTCCCTGTCCAGAAGCCGAAACTCTTGACCACCGAGGAGAAGCCACGAGGTCAGGGC
AGGGGCCCCATCCTTCCAGGCACCAAGGCCTGGATGGAGACCGAGGACACCCTGGGCCGT
GTCCTGAGTCCCGAGCCCGACCATGACAGCCTGTACCACCCTCCGCCTGAGGAGGACCAG
GGCGAGGAGAGGCCCCGGTTGTGGGTGATGCCAAATCACCAGGTGCTCCTGGGACCGGAG
GAAGACCAAGACCACATCTACCACCCCACTAGGGCTCCAGGGGCCATCACTGCCCCCGC
CCTGTCCCAAGGCCAGGCTGTTGGGACTGGGACCCCTCCCTACCCTGCCCCAGCTAGACA
AATAAACCCAGCAGGCAAAAAAAAAAAAAAAAAAAAA

221/392

FIGURE 212

MRRLLLVTSLVVVLLWEAGAVPAPKVPIKMQVKHWPSEQDPEKAWGARVVEPPEKDDQLV
VLFPVQKPKLLTTEEKPRGQGGRGPILPGTKAWMETEDTLGRVLSPEPDHDSLYHPPPEED
QGEERPRLWVMPNHQVLLGPPEEDQDHIYHPQ

222/392

FIGURE 213

CAGGCAGAAGCGAACAAAGACCCAGCAAGAGAAGGCAGAGGCTAAGACCCATCCCGTATC
TGCTCTCCTGAAATAATTCTGGAGTCATGCCTGAAATGCCAGAGGACATGGAGCAGGAGG
AAGTTAACATCCCTAATAGGAGGGTTCTGGTTACTGGTGCCACTGGGCTTCTTGGCAGAG
CTGTACACAAAGAAATTTAGCAGAATAATTGGCATGCAGTTGGCTGTGGTTTCAGAAGAG
CAAGACCAAAATTTGAACAGGTTAATCTGTTGGATTCTAATGCAGTTCATCACATCATT
ATGATTTTCAGCCCCATGTTATAGTACATTGTGTCAGCAGAGAGAAGACCAGATGTTGTAG
AAAATCAGCCAGATGCTGCCTCTCAACTTAATGTGGATGCTTCTGGGAATTTAGCAAAGG
AAGCAGCTGCTGTTGGAGCATTCTCATCTACATTAGCTCAGATTATGTATTTGATGGAA
CAAATCCACCTTACAGAGAGGAAGACATACCAGCTCCCCTAAATTTGTATGGCAAAACAA
AATTAGATGGAGAAAAGGCTGTCTGGAGAACATCTAGGAGCTGCTGTTTGGAGGATTCT
TATTCTGTATGGGGAAGTTGAAAAGCTCGAAGAAAGTGCTGTGACTGTTATGTTTGATA
AAGTGCAGTTCAGCAACAAGTCAGCAAAACATGGATCACTGGCAGCAGAGGTTCCCCACAC
ATGTCAAAGATGTGGCCACTGTGTGCCGAGCTAGCAGAGAAGAGAATGCTGGATCCAT
CAATTAAGGGAACCTTTCACTGGTCTGGCAATGAACAGATGACTAAGTATGAAATGGCAT
GTGCAATTGCAGATGCCTTCAACCTCCCCAGCAGTCACTTAAGACCTATTACTGACAGCC
CTGTCTTAGGAGCACACGTCGAGAAATGCTCAGCTTGACTGCTCCAAATGGAGACCT
TGGGCATTGGCCAACGAACACCATTTTGAATTTGGAATCAAAGAATCACTTTGGCCTTTCC
TCATTGACAAGAGATGGAGACAAACGGTCTTTTCACTTAGTTTATTTGTGTTGGGTTCTTTT
TTTTTTTAAATGAAAAGTATAGTATGTGGCACTTTTTTAAAGAACAAAGGAAATAGTTTTG
TATGAGTACTTTAATTGTGACTCTTAGGATCTTTCAGGTAAATGATGCTCTTGCACTAGT
GAAATTGTCTAAAGAACTAAAGGGCAGTCATGCCCTGTTTGCAGTAATTTTCTTTT
TCATTTTGTGTTGCTGCTAACTTGGAGTTTGAGTATAGTAAATTATGATCCTTAAAT
ATTTGAGAGTCAGGATGAAGCAGATCTGCTGTAGACTTTTCAGATGAAATTGTTTATTCT
CGTAACCTCCATATTTTCAGGATTTTGAAGCTGTTGACCTTTTCATGTTGATTATTTTA
AATTGTGTGAAATAGTATAAAATCATTTGGTGTTCATTATTTGCTTTGCCTGAGCTCAGA
TCAAAATGTTTGAAGAAAGGAACCTTTATTTTGAAGTTACGTACAGTTTTTATGCTTGA
GATATTTCAACATGTTATGTATATTGGAACCTTCTACAGCTTGATGCCTCCTGCTTTTATA
GCAGTTTATGGGGAGCACCTTGAAGAGCGTGTGTACATGTATTTTTTTCTAGGCAACA
TTGAATGCAAACGTGTATTTTTTTAATATAAATATATAACTGTCCTTTTCATCCCATGTT
GCCGCTAAGTGATATTTTATATGTGTGGTTATACTCATAAATGGGCCTTGTAAGTCTT
TTCACCATTCATGAATAATAAATATGTACTGCTGGCATGTAATGCTTAGTTTTCTTG
TATTTACTTCTTTTTTTTAAATGTAAGGACCAACTTCTAACTAATTGTTCTTTGTTGC
TTTAATTTTAAAAATTACATTCTTCTGATGTAACATGTGATACATACAAAAGAATATAG
TTTAATATGTATTGAAATAAAACACAATAAAAT

223/392

FIGURE 214

MPEMPEDMEQEEVNIPNRRVLVTGATGLLGRAVHKEFQQNNWHAVGCGFRRARPKFEQVN
LLDSNAVHHI IHDFQPHVIVHCAAERRPDVVENQPDAAASQLNVDAAGNLAKEAAAVGAFL
IYISSDYVFDGTNPPYREEDI PAPLNLYGKTKLDGEKAVLENNLGA AVLRIPILYGEVEK
LEESAVTVMFDKVQFSNKSANMDHWQQRFP THVKDVATVCRQLAEKRMLDPSIKGTFHWS
GNEQMTKYEMACAIADAFNLPSSHLRPITDSPVLGAQRPRNAQLDCSKLET LGIGQRTPF
RIGIKESLWPFLIDKRWRQTVFH

Signal peptide:
amino acids 1-30

Transmembrane domain:
amino acids 105-127

N-glycosylation site:
amino acids 197-201

N-myristoylation site:
amino acids 303-309

Short-chain dehydrogenases/reductases family proteins:
amino acids 18-30

224/392

FIGURE 215

GTGAATGTGAGGGTTTGATGACTTTCAGATGTCTAGGAACCAGAGTGGGTGCAGGGGCCC
CAGGCAGGGCTGATTCTTGGGCGGAGGAGAGTAGGGTAAAGGGTTCTGCATGAGCTCCTT
AAAGGACAAAGGTAACAGAGCCAGCGAGAGAGCTCGAGGGGAGACTTTGACTTCAAGCCA
CAGAATTGGTGGAAGTGTGCGCGCCGCCGCCGCTCGCTCCTGCAGCGCTGTTCGACCTA
GCCGCTAGCATCTTCCCGAGCACCGGGATCCCGGGGTAGGAGGCGACGCGGGCGAGCACC
AGCGCCAGCCGGCTGCGGCTGCCCACACGGCTCACCATGGGCTCCGGGCGCCGGGCGCTG
TCCGCGGTGCCGGCCGTGCTGCTGGTCCACGCTGCCGGGGCTGCCCGTCTGGGCACAG
AACGACACGGAGCCCATCGTGCTGGAGGGCAAGTGTCTGGTGGTGTGCGACTCGAACCCG
GCCACGGACTCCAAGGGCTCCTCTTCTCCCGCTGGGGATATCGGTCCGGGCGGCCAAC
TCCAAGCTCGCCTTCTCGGCGGTGCGGAGCACCAACCACGAGCCATCCGAGATGAGCAAC
AAGACGCGCATCTTTACTTCGATCAGATCCTGGTGAATGTGGGTAAATTTTTCACATTG
GAGTCTGTCTTTGTAGCACCAAGAAAAGGAATTTACAGTTTCAGTTTTCACGTGATTAAA
GTCTACCAGAGCCAAACATCCAGGTTAACTTGATGTTAAATGGAAAACAGTAATATCT
GCCTTTGCGGGGGACAAAGATGTTACTCGTGAAGCTGCCACGAATGGTGTCTGCTCTAC
CTAGATAAAGAGGATAAGGTTTACCTAAAACCTGGAGAAAGGTAATTTGGTTGGAGGCTGG
CAGTATTCACGTTTTCTGGCTTTCTGGTGTTCCTCCCTATAGGATTCAATTTCTCCATGA
TGTTTCATCCAGGTGAGGGATGACCCACTCCTGAGTTATTGGAAGATCATTTTTTTCATCAT
TGGATTGATGTCTTTTATTGGTTTCTCATGGGTGGATATGGATTCTAAGGATTCTAGCCT
GTCTGAACCAATACAAAATTTACAGATTATTTGTGTGTGTCTGTTTCAGTATATTTGGA
TTGGGACTCTAAGCAGATAATACTATGCTTAAATGTAACAGTCAAAGCTGTCTGCAAG
ACTTATTCTGAATTTCAATTCCTGGGATTACTGAATTAGTTACAGATGTGGAATTTTATT
TGTTTAGTTTTTAAAAGACTGGCAACCAGGTCTAAGGATTAGAAAACCTAAAGTTCTGAC
TTCAATCAACGGTTAGTGTGATACTGCCAAAGAACTGTATACTGTGTTAATATATTGATT
ATATTTGTTTTTATTCCTTTGGAATTAGTTTGTGTTGGTTCTTGTAATAAACTGGATTTT
TTTTTTCAGTAACTGGTATTATGTTTTCTCTTAAATAAGGTAATGAATGGCTTGCCAC
AAATTTACCTTGACTACGATATCATCGACATGACTTCTCTCAAAAAAAAAAGAATGCTTCA
TAGTTGTATTTTAATTGTATATGTGAAAGAGTCATATTTTCCAAGTTATATTTTCTAAGA
AGAAGAATAGATCATAAATCTGACAAGGAAAAAGTTGCTTACCCAAAATCTAAGTGCTCA
ATCCCTGAGCCTCAGCAAAACAGCTCCCCCTCCGAGGGAAATCTTATACTTTATTGCTCAA
CTTTAATTAAAATGATTGATAATAACCACTTTATTAAAAACCTAAGGTTTTTTTTTTTTC
CGTAGACATGACCACTTTATTAACGGTGGTGGGATGCTGTTGTTTCTAATTATACCTAT
TTTTCAAGGCTTCTGTTGTATTTGAAGTATCATCTGGTTTTGCCTTAACTCTTTAAATTG
TATATATTTATCTGTTTAGCTAATATTAAATTCAAATATCCCATATCTAAATTTAGTGCA
ATATCTTGTCCTTTTGTATAGGTCATATGAATTCATAAAATTATTTATGTCTGTTATAGAA
TAAAGATTAATATATGTTAAAAAAA

225/392

FIGURE 216

MSGRRALSAVPAVLLVLTLPGLPVWAQNDTEPIVLEGKCLVVCDSPATDSKGSSSSPL
GISVRAANSKVAFAVRSTNHEPSEMSNKTRIIYFDQILVNVGNFFTTLESV FVAPRKGIY
SFSFHVIK VYQSQTIQVNLMLNGKPVISAFAGDKDVTREAA TNGVLLYLDKEDKVYLKLE
KGNLVGGWQYSTFSGFLVFPL

Signal peptide:
amino acids 1-27

226/392

FIGURE 217

CGGCAACCAGCCGCCGCCACCACCGCTGCCACTGCCGCCCTGCCGGGGCCATGTTTCGCTC
TGGGCTTGCCCTTCTTGGTGCTCTTGGTGGCCTCGGTCGAGAGCCATCTGGGGGTTCGG
GGCCCAAGAACGTCTCGCAGAAAGACGCCGAGTTTGAGCGCACCTACGTGGACGAGGTCA
ACAGCGAGCTGGTCAACATCTACACCTTCAACCATACTGTGACCCGCAACAGGACAGAGG
GCGTGCGTGCTGTCTGTGAACGTCCCTGAACAAGCAGAAGGGGGCGCCGTTGCTGTTTGTGG
TCCGCCAGAAGGAGGCTGTGGTGTCCTTCCAGGTGCCCCAATCCTGCGAGGGATGTTTC
AGCGCAAGTACCTCTACCAAAAGTGGAACGAACCCTGTGTGTCAGCCCCCACCAGAAGATG
AGTCGGAGATTCACTTCTTCTACGTGGATGTGTCCACCCTGTCAACAGTCAACACCACAT
ACCAGCTCCGGGTGAGCCGCATGGACGATTTTGTGCTCAGGACTGGGGAGCAGTTCAGCT
TCAATACCACAGCAGCACAGCCCCAGTACTTCAAGTATGAGTTCCTGAAGGCGTGACT
CGGTAATTGTCAAGGTGACCTCCAACAAGGCCTTCCCCTGCTCAGTCATCTCCATTCAGG
ATGTGCTGTGCTCTATGACCTGGACAACAACGTAGCCTTCATCGGCATGTACCAGA
CGATGACCAAGAAGGCGGCCATCACCGTACAGCGCAAAGACTTCCCCAGCAACAGCTTTT
ATGTGGTGGTGGTGGTGAAGACCGAAGACCAAGCCTGCCGGGGCTCCCTGCCTTTCTACC
CCTTCGCAGAAGATGAACCGGTGATCAAGGGCACCGCCAGAAAACCCCTGTCACTGCTGG
TGTCTCAAGCAGTCACGTCTGAGGCATACGTCACTGAGGGATGCTCTTTTGCCTGGGTATAT
TTCTCTCCTTTTACCTGCTGACCGTCTCCTGGCCTGCTGGGAGAACTGGAGGCAGAAGA
AGAAGACCCTGCTGGTGGCCATTGACCGAGCCTGCCAGAAAGCGGTCACCCTCGAGTCC
TGGCTGATTCTTTTCTGGCAGTTCCTTATGAGGGTTACAACATATGGCTCCTTTGAGA
ATGTTTCTGGATCTACCGATGGTCTGGTTGACAGCGCTGGCACTGGGGACCTCTCTTACG
GTTACCAGGGCCGCTCCTTTGAACCTGTAGGTACTCGGCCCGAGTGGACTCCATGAGCT
CTGTGGAGGAGGATGACTACGACACATTGACCGACATCGATTCCGACAAGAATGTCATTC
GCACCAAGCAATACCTCTATGTGGCTGACCTGGCACGGAAGGACAAGCGTGTTCTGCGGA
AAAAGTACCAGATCTACTTCTGGAACATTGCCACCATTGCTGTCTTCTATGCCCTTCCTG
TGGTGCAGCTGGTGATCACCTACCAGACGGTGGTGAATGTCACAGGGAATCAGGACATCT
GCTACTACAACCTCCTCTGCGCCCACTGGGCAATCTCAGCGCCTTCAACAACATCC
TCAGCAACCTGGGGTACATCCTGCTGGGGCTGCTTTTCTGCTCATCATCCTGCAACGGG
AGATCAACCACAACCGGGCCCTGCTGCGCAATGACCTCTGTGCCCTGGAATGTGGGATCC
CCAAACACTTTGGGCTTTTCTACGCCATGGGCACAGCCCTGATGATGGAGGGGCTGCTCA
GTGCTTGCTATCATGTGTGCCCCAACTATAACCAATTTCCAGTTTGACACATCGTTTCATGT
ACATGATCGCCGACTCTGCATGCTGAAGCTCTACCAGAAGCGGCACCCGACATCAACG
CCAGCGCCTACAGTGCCCTACGCCTGCCTGGCCATTGTCTCTCTCTGTGCTGGGCG
TGGTCTTTGGCAAAGGGAACACGGCGTTCTGGATCGTCTTCTCCATCATTACATCATCG
CCACCTTGCTCCTCAGCACGCAGCTCTATTACATGGGCGGGTGGAACTGGACTCGGGGA
TCTTCCGCCGCATCCTCCACGTGCTCTACACAGACTGCATCCGGCAGTGCAGCGGGCCGC
TCTACGTGGACCGCATGGTGCTGCTGGTCACTGGGCAACGTCATCAACTGGTTCGCTGGCTG
CCTATGGGCTTATCATGCGCCCCAATGATTTGCTTCTTCTTCTTGTGGCCATTGGCATCT
GCAACCTGCTCCTTTACTTCGCCTTCTACATCATCATGAAGCTCCGGAGTGGGGAGAGGA
TCAAGCTCATCCCCCTGCTCTGCATCGTTTGACCTCCGTGGTCTGGGGCTTCGCGCTCT
TCTTCTTCTTCCAGGGACTCAGCACCTGGCAGAAAACCCCTGCAGAGTGGAGGGAGACA
ACCGGGACTGCATCCTCCTCGACTTCTTTGACGACCACGACATCTGGCACTTCTCTCCT
CCATCGCCATGTTTCGGGTCTTCTGCTGTTGCTGACACTGGATGACGACCTGGATACTG
TGCAGCGGGACAAGATCTATGTCTTCTAGCAGGAGCTGGGCCCTTCGCTTCACTCAAGG
GGCCCTGAGCTCCTTTGTGTCATAGACCGGTCACTCTGTGCTGCTGTGGGGATGAGTCCC
AGCACCGCTGCCAGCACTGGATGGCAGCAGGACAGCCAGGTCTAGCTTAGGCTTGGCCT
GGGACAGCCATGGGGTGGCATGGAACCTTGCAGCTGCCCTCTGCCGAGGAGCAGGCCTGC
TCCCCTGGAACCCCCAGATGTTGGCCAAATTGCTGCTTTCTTCTCAGTGTTGGGGCCTTC

227/392

CATGGGCCCCTGTCCTTTGGCTCTCCATTTGTCCCTTTGCAAGAGGAAGGATGGAAGGGA
CACCCCTCCCCATTTATGCCTTGCATTTTGCCCGTCCTCCTCCCCACAATGCCCCAGCCT
GGGACCTAAGGCCTCTTTTCTCCTCCATACTCCCACTCCAGGGCCTAGTCTGGGGCCTGA
ATCTCTGTCCTGTATCAGGGCCCCAGTTCTCTTTGGGCTGTCCCTGGCTGCCATCACTGC
CCATTCCAGTCAGCCAGGATGGATGGGGGTATGAGATTTTGGGGGTGGCCAGCTGGTGC
CAGACTTTTGGTGCTAAGGCCTGCAAGGGGCCTGGGGCAGTGCGTATTCTCTTCCCTCTG
ACCTGTGCTCAGGGCTGGCTCTTTAGCAATGCGCTCAGCCCAATTTGAGAACCGCCTTCT
GATTCAAGAGGCTGAATTCAGAGGTCACCTCTTCATCCCATCAGCTCCCAGACTGATGCC
AGCACCAGGACTGGAGGGAGAAGCGCCTCACCCCTTCCCTTCCTTCTTTCCAGGCCCTTA
GTCTTGCCAAACCCCAGCTGGTGGCCTTTAGTGCCATTGACACTGCCCCAAGAATGTCCA
GGGGCAAAGGAGGGATGATACAGAGTTCAGCCCGTTCTGCCTCCACAGCTGTGGGCACCC
CAGTGCCTACCTTAGAAAGGGGCTTCAGGAAGGGATGTGCTGTTTCCCTCTACGTGCCCA
GTCCTAGCCTCGCTCTAGGACCCAGGGCTGGCTTCTAAGTTTCCGTCCAGTCTTCAGGCA
AGTTCTGTGTTAGTCATGCACACACATACCTATGAAACCTTGGAGTTTACAAAGAATTGC
CCCAGCTCTGGGCACCCCTGGCCACCCCTGGTCCTTGGATCCCCCTTCGTCCCACCTGGTCCA
CCCCAGATGCTGAGGATGGGGGAGCTCAGGCGGGGCCTCTGCTTTGGGGATGGGAATGTG
TTTTTCTCCAACTTGTTTTTATAGCTCTGCTTGAAGGGCTGGGAGATGAGGTGGGTCT
GGATCTTTTCTCAGAGCGTCTCCATGCTATGGTTGCATTTCCGTTTTCTATGAATGAATT
TGCATTCAATAAACCAACCAGACTCAAAAAAAAAAAAAA

228/392

FIGURE 218

MFALGLPFLVLLVASVESHLGVLGPKNVSQKDAEFERTYVDEVNSELVNIYTFNHTVTRN
RTEGVRVSVNVLNKQKGAPLLFVVVRQKEAVVSFQVPLILRGMFORKYLYQKVERTLCQPP
TKNESEIQFFYVDVSTLSPVNTTYQLRVSRMDDFVLRTGEQFSFNTTAAQPQYFKYEFPE
GVDSVIVKVTSNKAFFPCSVISIQDVLCPVYDLNNAFIGMYQTMTKKAAITVQRKDFPS
NSFYVVVVVKTEDQACGGSLPFYFPAEDEPVDQGHRQKTL SVLVSQAVTSEAYVSGMLFC
LGIFLSFYLLTVLLACWENWRQKKKTLLVAIDRACPESGHPRVLADSFPGSSPYEGYNYG
SFENVSGSTDGLVDSAGTGDLSYGYQGRSFEPVGTRPRVDSMSSVEEDDYDTLTDIDSDK
NVIRTKQYLYVADLARKDKRVLRRKKYQIYFWNIATIAVFYALPVVQLVITYQTVVNVNVTGN
QDICYYNFLCAHPLGNLSAFNNILSNLGYILLGLLFLLIILQREINHNRRALLRNDLCALC
CGIPKHFGLFYAMGTALMMEGLLSACYHVCNPTNFQFDTSEFMYMIAGLCMLKLYQKRHP
DINASAYSAYACLAIVIFFSVLGVVFGKGNATFWIVFSIIHI IATLLLSTQLYYMGRWKL
DSGIFRRILHVLYTDCIRQCSGPLYVDRMVLLVMGNVINWSLAAYGLIMRPNDFASYLLA
IGICNLLLYFAFYIIMKLRSGERIKLIPLLCIVCTSVVWGFALFFFFQGLSTWQKTPAES
REHNRDCILLDFDDHDIWHFLSSIAMFGSFLVLLTLDDDLDTVQRDKIYVF

Important features of the protein:**Signal peptide:**

amino acids 1-18

Transmembrane domains:

amino acids 292-317, 451-470, 501-520, 607-627, 751-770

Leucine zipper pattern:

amino acids 497-518

N-glycosylation sites:amino acids 27-30, 54-57, 60-63, 123-126, 141-144, 165-168,
364-367, 476-479, 496-499, 572-575, 603-606, 699-702

229/392

FIGURE 219

[illegible]

230/392

FIGURE 220

MRSTILLFCLLGSTRSLPQLKPALGLPPTKLAPDQGTLPNQQQSNQVFPSLSLIPLTQML
TLGPDHLHLLNPAAGMTPGTQTHPLTLGGLNVQQQLHPHVLPIFVTQLGAQGTLISSEELP
QIFTSLIIHSLFPGGILPTSQAGANPDVQDGSLPAGGAGVNPATQGTPAGRLPTPSGTDD
DFAVTTPAGIQRSTHAIEEATTESANGIQ

Signal peptide:
amino acids 1-16

231/392

FIGURE 221

GACTTTGCTTGAATGGTTTACATATTTCTGCTCGCTGTCTACATATCACAAATATAGTCTT
ACGTTTTTGTTAAAACTTTGGGGTGTGAGGAGTTGAGCTTGCTCAGCAAGCCAGCATGGCT
AGGATGAGCTTTGTTATAGCAGCTTGCCAATTGGTGCTGGGCCTACTAATGACTTTCATTA
ACCGAGTCTTCCATACAGAATAGTGAGTGCCACAACCTTTGCGTATGTGAAATTCGTCCC
TGGTTTACCCACAGTCAACTTACAGAGAAGCCACCACTGTTGATTGCAATGACCTCCGC
TTAACAAGGATTCCCAGTAACCTCTCTAGTGACACACAAGTGCTTCTCTTACAGAGCAAT
AACATCGCGAAGACTGTGGATGAGCTGCAGCAGCTTTTCAACTTGACTGAACTAGATTTT
TCCCAAAACAACCTTTACTAACATTAAGGAGGTGCGGCTGGCAAACCTAACCCAGCTCAC
ACGCTGCATTTGGAGGAAAATCAGATTACCGAGATGACTGATTACTGTCTACAAGACCT
AGCAACCTTCAAGAACTCTACATCAACCACAACCAAATTAGCACTATTTCTGCTCATGCT
TTTGACAGGCTTAAAAAATCTATTAAGGCTCCACCTGAACTCCAACAAATTGAAAGTTATT
GATAGTCGCTGGTTTGATTCTACACCCAACCTGGAAATTCTCATGATCGGAGAAAACCT
GTGATTGGAATTCTGGATATGAACTTCAAACCCCTCGCAAATTTGAGAAGCTTAGTTTTG
GCAGGAATGTATCTCACTGATATTCCTGGAAATGCTTTGGTGGGTCTGGATAGCCTTGAG
AGCCTGTCTTTTTATGATAACAACTGGTTAAAAGTCCCTCAACTTGCCCTGCAAAAAGTT
CCAAATTTGAAATTCTTAGACCTCAACAAAAACCCATTCAAAAAATCCAAGAAGGGGAC
TTCAAAAATATGCTTCGGTTAAAAGAACTGGGAATCAACAATATGGGCGAGCTCGTTTCT
GTCGACCGCTATGCCCTGGATAACTTGCCTGAACTCACAAAGCTGGAAGCCACCAATAAC
CCTAACTCTCTTACATCCACCGCTTGGCTTTCGGAAGTGTCCCTGCTCTGGAAAGCTTG
ATGCTGAACAACAATGCCCTGAATGCCATTTACCAAAGACAGTCGAATCCCTCCCCAAT
CTGCGTGAGATCAGTATCCATAGCAATCCCTCAGGTGTGACTGTGTGATCCACTGGATT
AACTCCAACAAAACCAACATCCGCTTCATGGAGCCCTGTCCATGTCTGTGCCATGCGG
CCCGAATATAAAGGGCACCAGGTGAAGGAAGTTTAAATCCAGGATTCGAGTGAACAGTGC
CTCCCAATGATATCTCACGACAGCTTCCCAAATCGTTTTAAACGTGGATATCGGCACGACG
GTTTTCTTAGACTGTGAGCCATGGCTGAGCCAGAACCTGAAATTTACTGGGTCACTCCC
ATTGGAATAAGATAACTGTGGAAACCTTTTCAGATAAATACAAGCTAAGTAGCGAAGGT
ACCTTGGAATATCTAACATACAAATTGAAGACTCAGGAAGATACACATGTGTTGCCAG
AATGTCCAAGGGGCAGACACTCGGGTGGCAACAATTAAGGTTAACGGGACCTTCTGGAT
GGTACCCAGGTGCTAAAAATATACGTCAAGCAGACAGAATCCCATTCATCTTAGTGTCC
TGGAAAGTTAATTCCAATGTGATGACGTCAAACCTTAAATGGTGTCTGCCACCATGAAG
ATTGATAACCCCTCACATAACATATACTGCCAGGGTCCCAGTCGATGTCCATGAATACAAC
CTAACGCATCTGCAGCCTTCCACAGATTATGAAGTGTGTCTCACAGTGTCCAATATTCAT
CAGCAGACTCAAAAGTCATGCGTAAATGTCAACAACCAAAAATGCCGCTTCGCAGTGGAC
ATCTCTGATCAAGAAACCAGTACAGCCCTTGCTGCAGTAATGGGGTCTATGTTTGCCGTC
ATTAGCCTTGCGTCCATTGCTGTGTACTTTGCCAAAAGATTTAAGAGAAAAAACTACCAC
CACTCATTAACCTCTGGGAAGGTGACAGCGAGAAAGACAAAGATGGTTCTGCAGACACC
AAGCCAACCCAGGTGACACATCCAGAAGCTATTACATGTGGTAACTCAGAGGATATTTT
GCTTCTGGTAGTAAGGAGCACAAAGACGTTTTTGTCTTTATTCTGCAAAAGTGAACAAGTT
GAAGACTTTTGTAATTTTGTACTTTGCTAGTTTGTGGCAGAGTGGAGAGGACGGGTGGATA
TTTTCAATTTTGTAGTATAGCGTATCGCAAGGGTTTGACACGGCTGCCAGCGACTCTAG
GCTTCCAGTCTGTGTTTGGTTTTATTCTTATCATTATTATGATTGTTATTATATTATTA
TTTTATTTTGTGTTGTGTGCTTAACTCAATAATGCTGTTCTAACTACAGTGCTCAATAAA
ATGATTAATGACAGGAAAAAAGAAAAAAGAAAAAAGAAAAAAGAAAAAAGAAAAAAG
AAAAAAGAAAAAAGAAAAAAGAAAAAAGAAAAAAGAAAAAAGAAAAAAGAAAAAAG

232/392

FIGURE 222

MARMSFVIAACQLVLGLLMTSLTESSIONSECPQLCVCEIRPWFTPQSTYREATTVDCND
LRLTRIPSNLSSDTQVLLQLSQNNIAKTVDLQQLFNLTELDQSQNNFTNIKEVGLANLTQ
LTTLHLEENQITEMTDYCLQDLNLQELYINHNQISTISAHAFAGLKNLLRLHLNSNKLK
VIDSRWFDSTPNLEILMIGENPVIGILDMNFKPLANLRSLVLAGMYLTDIPGNALVGLDS
LESLSFYDNKLVKVPQLALQKVPNLKFLDLNKNPIHKIQEGDFKNMLRLKELGINNMGEL
VSVDRYALDNLPELTKLEATNNPKLSYIHRLAFRSVPALSLMLNNAALNAIYQKTVESL
PNLREISIHNSNPLRCDLCVIHWINSNKTNIREFMEPLSMFCAMPPEYKGHVKEVLIQDSSE
QCLPMISHDSFPNRLNVDIGTTVFLDCRAMAEPEPEIYWVTPIGNKITVETLSDKYKLSS
EGTLEISNIQIEDSGRYTCVAQNVQGADTRVATIKVNGTLLDGTQVLKIYVKQTESHSL
VSWKVNNSVMTSNLKWSSATMKIDNPHITYTARVPVDVHEYNLTHLQPSTDYEVCLTVSN
IHQQTQKSCVNVTTKNAFAVDISDQETSTALAAVMGSMFAVISLASIAVYFAKRFRKRN
YHSLKLYMQKTSSIPLNELYPPLINLWEGDSEKDKDGSADTKPTQVDTSRSYMW

Important features:**Signal peptide:**

Amino acids 1-25

Transmembrane domain:

Amino acids 508-530

N-glycosylation sites:Amino acids 69-73;96-100;106-110;117-121;385-389;517-521;
582-586;611-615**Tyrosine kinase phosphorylation site:**

Amino acids 573-582

N-myristoylation sites:

Amino acids 16-22;224-230;464-470;637-643;698-704

FIGURE 223

[illegible]

234/392

FIGURE 224

MASYLYGVLFVGLCAPIYCVSPANAPSAYPRPSSTKSTPASQVYSLNTDFAFRLYRRLV
LETPSQNIFFSPVSVSTSLAMLSLGAHSVTKTQILQGLGFNLTHTPESAIHQGFQHLVHS
LTVPSKDLTLKMGSALFVKKELQLQANFLGNVKRLYEAEVFSTDFSNPSIAQARINSHVK
KKTQGKVVDIIQGLDLLTAMVLVNHIFKAKWEKPFHLEYTRKNFPFLVGEQVTVQVPM
HQKEQFAFGVDTLNCFLVQMDYKGDVAFFVLPSKGKMRQLEQALSARTLIKWSHSLQK
RWIEVFIPRFSISASYNLETILPKMGIQNAFDKNADFSGIKRDSLQVSKATHKAVLDVS
EEGTEATAATTTKFIVRSKDGPSYFTVSFNRTFLMMITNKATDGILFLGKVENPTKS

Signal peptide:
amino acids 1-20

235/392

FIGURE 225

GGGAGAGAGGATAAATAGCAGCGTGGCTTCCCTGGCTCCTCTCTGCATCCTTCCCGACCT
TCCCAGCAATATGCATCTTGACAGTCTGGTCGGCTCCTGCTCCCTCCTTCTGCTACTGGG
GGCCCTGTCTGGATGGGCGGCCAGCGATGACCCCATTGAGAAGGTCATTGAAGGGATCAA
CCGAGGGCTGAGCAATGCAGAGAGAGAGGTGGGCAAGGCCCTGGATGGCATCAACAGTGG
AATCACGCATGCCGGAAGGGAAGTGGAGAAGGTTTTCAACGGACTTAGCAACATGGGGAG
CCACACCGGCAAGGAGTTGGACAAAGGCGTCCAGGGGCTCAACCACGGCATGGACAAGGT
TGCCCATGAGATCAACCATGGTATTGGACAAGCAGGAAAGGAAGCAGAGAAGCTTGGCCA
TGGGGTCAACAACGCTGCTGGACAGGCCGGAAGGAAGCAGACAAAGCGGTCCAAGGGTT
CCACACTGGGGTCCACCAGGCTGGGAAGGAAGCAGAGAACTTGGCCAAGGGGTCAACCA
TGCTGCTGACCAGGCTGGAAAGGAAGTGGAGAAGCTTGGCCAAGGTGCCACCATGCTGC
TGGCCAGGCCGGAAGGAGCTGCAGAATGCTCATAATGGGGTCAACCAAGCCAGCAAGGA
GGCCAACCAGCTGCTGAATGGCAACCATCAAAGCGGATCTTCCAGCCATCAAGGAGGGGC
CACAACCACGCCGTTAGCCTCTGGGGCCTCAGTCAACACGCCTTTCATCAACCTTCCCGC
CCTGTGGAGGAGCGTCGCCAACATCATGCCCTAAACTGGCATCCGGCCTTGCTGGGAGAA
TAATGTCGCCGTTGTCACATCAGCTGACATGACCTGGAGGGGTGGGGGTGGGGACAGG
TTTCTGAAATCCCTGAAGGGGGTTGTAAGGGATTTGTGAATAAACTTGATACACCA

236/392

FIGURE 226

MHLARLVGSCSLLLLLLGALSGWAASDDPIEKVIEGINRGLSNAEREVKGALDGINSGITH
AGREVEKVFNGLSNMGSHGTGKELDKGVQGLNHGMDKVAHEINHGIGQAGKEAEKLGHG
VN
NAAGQAGKEADKAVQGFHTGVHQAGKEAEKLGQGVNHAADQAGKEVEKLGQGAHHAAGQA
GKELQNAHNGVNVQASKEANQLLNGNHQSGSSSHQGGATTTPLASGASVNTPFINLPALWR
SVANIMP

Important features of the protein:

Signal peptide:

amino acids 1-25

Homologous region to circumsporozoite (CS) repeats:

amino acids 35-225

237/392

FIGURE 227

GAAGTAGAGGTGTTGTGCTGAGCGGCGCTCGGCGAACTGTGTGGACCGTCTGCTGGGACT
CCGGCCCTGCGTCCGCTCAGCCCCGTGGCCCCGCGCACCTACTGCCATGGAGACGCGGCC
TCGTCTCGGGGCCACCTGTTTGCTGGGCTTCAGTTTCCTGCTCCTCGTCATCTCTTCTGA
TGGACATAATGGGCTTGGAAGGGTTTTGGAGATCATATTCATTGGAGGACACTGGAAGA
TGGGAAGAAAGAAGCAGCTGCCAGTGGACTGCCCCGTGATGGTGATTATTCATAAATCCTG
GTGTGGAGCTTGCAAAGCTCTAAAGCCCCAAATTTGCAGAATCTACGGAAATTTCAGAACT
CTCCCATAAATTTTGTTATGGTAAATCTTGAGGATGAAGAGGAACCCAAAGATGAAGATTT
CAGCCCTGACGGGGGTATATTCACGAATCCTTTTTCTGGATCCCAGTGGCAAGGTGCA
TCCTGAAATCATCAATGAGAAATGGAAACCCAGCTACAAGTATTTTTATGTCAGTGCCGA
GCAAGTTGTTCAAGGGATGAAGGAAGCTCAGGAAAGGCTGACGGGTGATGCCTTCAGAAA
GAAACATCTTGAAGATGAATTGTAACATGAATGTGCCCCCTTCTTTCATCAGAGTTAGTGT
TCTGGAAGGAAAGCAGCAGGGAAGGGAATATTGAGGAATCATCTAGAACAATTAAGCCGA
CCAGGAAACCTCATTCCTACCTACACTGGAAGGAGCGCTCTCACTGTGGAAGAGTTCTGC
TAACAGAAGCTGGTCTGCATGTTTGTGGATCCAGCGGAGAGTGGCAGACTTTCTTCTCCT
TTTCCCTCTCACCTAAATGTCAACTTGTCATTGAATGTAAAGAATGAAACCTTCTGACAC
AAAA

238/392

FIGURE 228

METRPRLGATCLLGFSFLLLVISSDGHNGLGKGFGDHIHWRTLEDGKKEAAASGLPLMVI
IHKSWCGACKALKPKFAESTEISELSHNFVMVNLEDEEPEKDEDFSPDGGYIPRILFLDP
SGKVHPEIINENGNP SYKYFYVSAEQVVQGMKEAQERLTGDAFRKKHLEDEL

Signal peptide:

Amino acids 1-23

Thioredoxin family proteins Homology Block:

Amino acids 58-75

239/392

FIGURE 229

CCCACGCGTCCGCCCACGCGTCCGGGTGCCACTCGCGCGCCGGCCGCGCTCCGGGCTTCT
CTTTTCCCTCCGACGCGCCACGGCTGCCCAGACATTCCGGCTGCCGGGTCTGGAGAGCTC
CCCGAACCCCTCCGCGGAGAGGAGCGAGGCGGCGCCAGGGTGGCCCCCGGGGCGCGCTTG
GTCTCGGAGAAGCGGGGACGAGGCCGGAGGATGAGCGACTGAGGGCGACGCGGGCACTGA
CGCGAGTTGGGGCCGCGACTACCGGCAGCTGACAGCGCGATGAGCGACTCCCCAGAGACG
CCCTAGCCCCGGTGTGCGCGCCAGGCGGAGCGCGCAGGTGGGGCTGGGCTGTTAGTGGTCC
GCCCCACGCGGGTCGCCGGCCGGCCAGGATGGGCGCTGGCAACCCGGGCCCCGCGCCCCG
CGCTGCTACCCCTGCGCCCCGCTGCGAGCCCCGGCGTCCGGCCCCGCGCCCTGCGCTCATGGA
CGGCGGCTCCCGGCTGGCGGGCGGCGGCCCCCCGGGCTGTGAATGCGACTCGCCCCCTCGGC
CGCGCTCCCCGCCCCGCCCCCGCCGGGCTGGTAGGGGATGCCAGCTCCACTGCGAT
GGCAGTTGGCGCGCTCTCCAGTTCCCTCCTGCTCACCTGCTGCCTGATGGTGGCTCTGTG
CAGTCCGAGCATCCCGCTGGAGAAGCTGGCCCAGGCACCGAGCAGCGGGCCAGAGAGAA
GCGTGAGCACGCCACTCGGGACGGCCCCGGGGCGGGTGAACGAGCTCGGGCGCCCCGGCGAG
GGACGAGGGCGGCAGCGGCCGGGACTGGAAGAGCAAGAGCGGCCGTGGGCTCGCCGGCCG
TGAGCCGTGGAGCAAGCTGAAGCAGGCCCTGGGTCTCCAGGGCGGGGGCGCCAAGGCCGG
GGATCTGCAGGTCCGGCCCCGCGGGACACCCCGCAGGCGGAAGCCCTGGCCGCAGCCGC
CCAGGACGCGATTGGCCCCGGAATCGCGCCACGCCCCGAGCCACCCGAGGAGTACGTGTA
CCCGGACTACCGTGGAAGGGCTGCGTGACGAGAGCGGCTTCGTGTACGCGATCGGGGA
GAAGTTCGCGCCGGGGCCCCCTCGGCCCTGCCGTGCCTGTGCACCGAGGAGGGCCGCTGTG
CGCGCAGCCCCGAGTGCCCGAGGCTGCACCCGCGCTGCATCCACGTGACACGAGCCAGTG
CTGCCCCGAGTGCAAGGAGAGGAAGAATACTGCGAGTTCCGGGGCAAGACCTATCAGAC
TTTGAGGAGTTTCGTGGTGTCTCCATGCGAGAGGTGTCGCTGTGAAGCCAACGGTGAGGT
GCTATGCACAGTGTGACGCTGTCCCCAGACGGAGTGTGTGGACCCTGTGTACGAGCCTGA
TCAGTGCTGTCCCATCTGCAAAAATGGTCCAACTGCTTTGCAGAAACCGCGGTGATCCC
TGCTGGCAGAGAAGTGAAGACTGACGAGTGCACCATATGCCACTGTACTTATGAGGAAGG
CACATGGAGAATCGAGCGGCAGGCCATGTGCACGAGACATGAATGCAGGCAAAATGTAGAC
GCTTCCCAGAACACAACTCTGACTTTTTCTAGAACATTTTACTGATGTGAACATTCTAG
ATGACTCTGGGAACTATCAGTCAAAGAAGACTTTTGATGAGGAATAATGGAAAATTGTTG
GTACTTTTCTTTTCTTGATAACAGTTACTACAACAGAAGGAAATGGATATATTTCAAAA
CATCAACAAGAACTTTGGGCATAAAATCCTTCTCTAAATAAATGTGCTATTTTCACAGTA
AGTACACAAAAGTACACTATTATATATCAAATGTATTTCTATAATCCCTCCATTAGAGAG
CTTATATAAGTGTTTTCTATAGATGCAGATTAAAAATGCTGTGTTGTCAACCGTCAAAAA
AAAAAAAAAAAAAAAAAAAAA

240/392

FIGURE 230

MPSSTAMAVGALSSSLVTCCLMVALCSPSIPLEKLAQAPEQPGQEKREHATRDGPGRVN
ELGRPARDEGGSGRDWKSXSGRGLAGREPWSKLKQAWVSQGGGAKAGDLQVRPRGDTPOA
EALAAAAQDAIGPELAPTPEPPPEEYVYPDYRGKGCVDSESGFVYAIGKFAFGPSACPCLC
TEEGPLCAQPECPRLHPRCIHVDTSQCCPQCKERKNYCEFRGKTYQTLEEFVVSPCERCR
CEANGEVLCTVSACPQTECVDPVYEPDQCCPICKNGPNCFAETAVIPAGREVKTDCTIC
HCTYEETWRIERQAMCTRHECRQM

Important features of the protein:

Signal peptide:

amino acids 1-27

Transmembrane domain:

amino acids 11-30

Glycosaminoglycan attachment site:

amino acids 80-83

N-myristoylation sites:

amino acids 10-15, 102-107, 103-108

Cell attachment sequence:

amino acids 114-117

EGF-like domain cysteine pattern signature:

amino acids 176-187

241/392

FIGURE 231

GGCCGGACGCCTCCGCGTTACGGGATGAATTAACGGCGGGTTCCGCACGGAGGTTGTGAC
CCCTACGGAGCCCCAGCTTGCCACGCACCCCACTCGGCGTCGCGCGGCGTGCCCTGCTT
GTCACAGGTGGGAGGCTGGAACTATCAGGCTGAAAAACAGAGTGGGTACTCTCTTCTGGG
AAGCTGGCAACAAATGGATGATGTGATATATGCATTCCAGGGGAAGGGAAATTGTGGTG
TTCTGAACCCATGGTCAATTAACGAGGCAGTTTCTAGCTACTGCACGTACTTCATAAAGC
AGGACTCTAAAAGCTTTTGAATCATGGTGTGATGGAAAGGGATTTACTTTTATACTGACTC
TGTTTTGGGGAAGCTTTTTTGAAGCATTTCATGCTGAGTCCCTTTTTTACCTTTGATGT
TTGTAAACCCATCTTGGTATCGCTGGATCAACAACCGCCTTGTGGCAACATGGCTCACCC
TACCTGTGGCATTATTGGAGACCATGTTTGGTGTAAAAGTGATTATAACTGGGGATGCAT
TTGTTTCTGGAGAAAGAAGTGTCATTATCATGAACCATCGGACAAGAATGGACTGGATGT
TCCTGTGGAATTGCCTGATGCGATATAGCTACCTCAGATTGGAGAAAATTTGCCTCAAAG
CGAGTCTCAAAGGTGTTTCTGGATTGGTTGGGCCATGCAGGCTGCTGCCTATATCTTCA
TTCATAGGAAATGGAAGGATGACAAGAGCCATTTCGAAGACATGATTGATTACTTTTGTG
ATATTACGAACCACTTCAACTCCTCATATTTCCAGAAGGGACTGATCTCACAGAAAACA
GCAAGTCTCGAAGTAATGCATTTGCTGAAAAAAATGGACTTCAGAAATATGAATATGTTT
TACATCCAAGAACTACAGGCTTTACTTTTTGTGGTAGACCGTCTAAGAGAAGGTAAGAACC
TTGATGCTGTCCATGATATCACTGTGGCGTATCCTCACAACATTTCCTCAATCAGAGAAGC
ACCTCCTCCAAGGAGACTTTCCAGGGAAATCCACTTTCACGTCCACCGGTATCCAATAG
ACACCCTCCCCACATCCAAGGAGGACCTTCAACTCTGGTGCCACAAACGGTGGGAAGAGA
AAGAAGAGAGGCTGCGTTCTTCTATCAAGGGGAGAAGAATTTTATTTTACCGGACAGA
GTGTCATTCCACCTTGCAAGTCTGAACTCAGGGTCTTGTGGTCAAATTGCTCTCTATAC
TGTATTGGACCTGTTTCAAGCTGCAATGTGCTTACTCATATATTTGTACAGTCTTGTTA
AGTGGTATTTTATAATCACCATTGTAATCTTTGTGCTGCAAGAGAGAATATTTGGTGGAC
TGGAGATCATAGAACTTGATGTTACCGACTTTTACACAAACAGCCACATTTAAATTCAA
AGAAAAATGAGTAAGATTATAAGGTTTGCCATGTGAAAACCTAGAGCATATTTTGGAAAT
GTTCTAAACCTTTCTAAGCTCAGATGCATTTTTCATGACTATGTCGAATATTTCTTACT
GCCATCATTATTTGTTAAAGATATTTTGCACCTTAATTTTGTGGGAAAAATATTGCTACAA
TTTTTTTTTAATCTCTGAATGTAATTTTCGATACTGTGTACATAGCAGGGAGTGATCGGGT
GAAATAACTTGGGCCAGAATATTATTAAACAATCATCAGGCTTTTAAA

242/392

FIGURE 232

MHSRGREIVVLLNPWSINEAVSSYCTYFIKQDSKSF GIMVSWKGIYFILTLFWGSFFGSI
FMLSPFLPLMFVNPSWYRWINNRLVATWLTLPVALLETMFGVKVIITGDAFVPGERSVII
MNHRTMRMDWMFLWNCLMRYSYLRLEKICLKASLKGVPFGFGWAMQAAAYIFIHRKWKDDKS
HFEDMIDYFCDIHEPLQLLIFPEGTDLTENSKSRSNFAEKNGLQKYEYVLHPRTTGFTF
VVDRLREGKNLDAVHDITVAYPHNIPQSEKHLLOGDFPREIH FHVHRYPIDTLPTS KEDL
QLWCHKRWEEKERLRSFYQGEKNFYFTGQSVIPPCKSELRLVVKLLSILYWTLFSPAM
CLLIYLYSLVKWYFIITIVIFVLQERIFGGLEIIELACYRLLHKQPHLNSKKNE

Important features of the protein:

Signal peptide:

amino acids 1-22

Transmembrane domains:

amino acids 44-63, 90-108, 354-377

243/392

FIGURE 233

CGGCTCGAGTGCAGCTGTGGGGAGATTTCACTGCATTGCCTCCCCCTGGGTGCTCTTCATC
TTGGATTTGAAAAGTTGAGAGCAGCATGTTTTGCCCCACTGAAACTCATCCTGCTGCCAGTG
TTACTGGATTATTCCTTGGGCCTGAATGACTTGAATGTTTTCCCCGCCTGAGCTAACAGTC
CATGTGGGTGATTTCAGCTCTGATGGGATGTGTTTTCCAGAGCACAGAAGACAAATGTATA
TTCAAGATAGACTGGACTCTGTCACCAGGAGAGCACGCCAAGGACGAATATGTGCTATAC
TATTACTCCAATCTCAGTGTGCCTATTGGGCGCTTCCAGAACC GCGTACACTTGATGGGG
GACATCTTATGCAATGATGGCTCTCTCCTGCTCCAAGATGTGCAAGAGGCTGACCAGGGA
ACCTATATCTGTGAAATCCGCCTCAAAGGGGAGAGCCAGGTGTTCAAGAAGGCGGTGGTA
CTGCATGTGCTTCCAGAGGAGCCCAAAGAGCTCATGGTCCATGTGGGTGGATTGATTCAG
ATGGGATGTGTTTTCCAGAGCACAGAAGTGAAACACGTGACCAAGGTAGAATGGATATTT
TCAGGACGGCGCGCAAAGGAGGAGATTGTATTTCGTTACTACCACAACTCAGGATGTCT
GTGGAGTACTCCCAGAGCTGGGGCCACTTCCAGAATCGTGTGAACCTGGTGGGGGACATT
TTCCGCAATGACGGTTCCATCATGCTTCAAGGAGTGAGGGAGTCAGATGGAGGAACTAC
ACCTGCAGTATCCACCTAGGGAACCTGGTGTTCAGAAAACCATTTGTGCTGCATGTGAGC
CCGGAAGAGCCTCGAACACTGGTGACCCCGGCAGCCCTGAGGCCTCTGGTCTTGGGTGGT
AATCAGTTGGTGATCATTGTGGGAATTGTCTGTGCCACAATCCTGCTGCTCCCTGTTCTG
ATATTGATCGTGAAGAAGACCTGTGGAAATAAGAGTTCAGTGAATTCTACAGTCTTGGTG
AAGAACACGAAGAAGACTAATCCAGAGATAAAAGAAAAACCTGCCATTTTGAAAGATGT
GAAGGGGAGAAACACATTTACTCCCCAATAATTGTACGGGAGGTGATCGAGGAAGAAGAA
CCAAGTGAAAAATCAGAGGCCACCTACATGACCATGCACCCAGTTTGGCCTTCTCTGAGG
TCAGATCGGAACAACTCACTTGAAAAAAAGTCAGGTGGGGGAATGCCAAAAACACAGCAA
GCCTTTTGAGAAGAATGGAGAGTCCCTTCATCTCAGCAGCGGTGGAGACTCTCTCCTGTG
TGTGTCCTGGGCCACTCTACCAGTGATTTCAGACTCCCGCTCTCCAGCTGTCTCCTGT
CTCATTTGTTTGGTCAATACACTGAAGATGGAGAATTTGGAGCCTGGCAGAGAGACTGGAC
AGCTCTGGAGGAACAGGCCTGCTGAGGGGAGGGGAGCATGGACTTGGCCTCTGGAGTGGG
ACACTGGCCCTGGGAACCAGGCTGAGCTGAGTGGCCTCAAACCCCCCGTTGGATCAGACC
CTCCTGTGGGCAGGGTTCTTAGTGGATGAGTTACTGGGAAGAATCAGAGATAAAACCAA
CCCAAATCAA

244/392

FIGURE 234

MFCPLKLILLPVLLDYSGLNDLNVSPPELTVHVGDSALMGCVFQSTEDKCIFKIDWTLS
PGEHAKDEYVLYYYSNLSVPIGRFQNRVHLMGDILCNDGSLLLQDVQEADQGTYICEIRL
KGESQVFKKAVVLHVLPEEPKELMVHVGGLIQMGCVFQSTEVKHVTKVEWIFSGRRAKEE
IVFRYYHKL RMSVEYSQSWGHFQNRVNLVGDI FRNDGSIMLQGVRES DGGNYTCSIHLGN
LVFKKTIVLHVSPEEPRTLVT PAALRPLVLGGNQLV IIVGIVCATILLLPVLILIVKKTC
GNKSSVNSTVLVKNTKKTNPEIKEKPCHFERCEGEKHIYSPIIVREVIEEEEPSEKSEAT
YMTMHPVWPSLRSDRNN SLEKKSGGMPKTQQAF

245/392

FIGURE 235

TAAACAGCTACAATATTCAGGGCCAGTCACTTGCCATTTCTCATAACAGCGTCAGAGA
GAAAGAACTGACTGAAACGTTTGAGATGAAGAAAGTTCTCCTCCTGATCACAGCCATCTT
GGCAGTGGCTGTTGGTTTCCCACTCTCTCAAGACCAGGAACGAGAAAAAGAAGTATCAG
TGACAGCGATGAATTAGCTTCAGGGTTTTTTGTGTTCCCTTACCCATATCCATTTGCCC
ACTTCCACCAATTCCATTTCCAAGATTTCCATGGTTTAGACGTAATTTTCTATTCCAAT
ACCTGAATCTGCCCCTACAACCTCCCTTCCTAGCGAAAAGTAAACAAGAAGGATAAGTCA
CGATAAACCTGGTCACCTGAAATTGAAATTGAGCCACTTCCTTGAAGAATCAAAATTCCT
GTTAATAAAAGAAAAACAAATGTAATTGAAATAGCACACAGCATTCTCTAGTCAATATCT
TTAGTGATCTTCTTTAATAAACATGAAAGCAAAGATTTTGGTTTCTTAATTTCCACA

246/392

FIGURE 236

MKKVLLLLITAILAVAVGFPVSQDQEREKRSISDSDELASGFFVFPYPYPFRPLPPIFFPR
FPWFRRNFPIPIPIESAPTTPLPSEK

Important features of the protein:

Signal peptide:

amino acids 1-17

Homologous region to B3-hordein:

amino acids 47-85

247/392

FIGURE 237

TCGCCATGGCCTCTGCCGGAATGCAGATCCTGGGAGTCGTCCTGACACTGCTGGGCTGGG
TGAATGGCCTGGTCTCCTGTGCCCTGCCCATGTGGAAGGTGACCGCTTTCATCGGCAACA
GCATCGTGGTGGCCCAGGTGGTGTGGGAGGGCCTGTGGATGTCCTGCGTGGTGCAGAGCA
CCGGCCAGATGCAGTGCAAGGTGTACGACTCACTGCTGGCGCTGCCACAGGACCTGCAGG
CTGCACGTGCCCTCTGTGTCATCGCCCTCCTTGTGGCCCTGTTGGCTTGCTGGTCTACC
TTGCTGGGGCCAAGTGTACCACCTGTGTGGAGGAGAAGGATTCCAAGCCCCGCTGGTGC
TCACCTCTGGGATTGTCTTTGTCTCATCTCAGGGGTCTGACGCTAATCCCCGTGTGCTGGA
CGGCGCATGCCATCATCCGGGACTTCTATAACCCCTGGTGGCTGAGGCCCAAAAGCGGG
AGCTGGGGGCCCTCCCTCTACTTGGGCTGGGCGGCCTCAGGCCTTTTGTGTGCTGGGTGGGG
GGTTGCTGTGCTGCACTTGCCCCCTCGGGGGGTCCCAGGGCCCCAGCCATTACATGGCCC
GCTACTCAACATCTGCCCCCTGCCATCTCTCGGGGGCCCTCTGAGTACCCTACCAAGAATT
ACGTCTGACGTGGAGGGGAATGGGGGCTCCGCTGGCGCTAGAGCCATCCAGAAGTGGCAG
TGCCCAACAGCTTTGGGATGGGTTCGTACCTTTTGTCTGCTCCTGCTATTTTTCTTT
TGACTGAGGATATTTAAATTCATTTGAAAACAGGCAAGGTGTTGACTCAGACTCTCA
CTTAGGCTCTGCTGTTTCTCACCCCTTGGATGATGGAGCCAAAGAGGGGATGCTTTGAGAT
TCTGGATCTTGACATGCCATCTTAGAAGCCAGTCAAGCTATGGAACATAATGCCGAGGCT
GCTTGCTGTGCTGGCTTTGCAACAAGACAGACTGTCCCCAAGAGTTCTGCTGCTGCTGG
GGGCTGGGCTTCCCTAGATGTCACTGGACAGCTGCCCCCATCCTACTCAGGTCTCTGGA
GCTCCTCTCTTCACCCCTGGAAAAACAAATCATCTGTTAACAAGGACTGCCACCTCCG
GAACTTCTGACCTCTGTTTCTCCGTCCTGATAAGACGTCCACCCCCAGGGCCAGGTCC
CAGCTATGTAGACCCCCGCCCCACCTCCAACACTGCACCCTTCTGCCCTGCCCCCTCG
TCTCACCCCTTTACACTCACATTTTTATCAAATAAAGCATGTTTTGTTAGTGCA

248/392

FIGURE 238

MASAGMQILGVVLTLLGWVNGLVSCALPMWKVTAFIGNSIVVAQVVWEGLWMSCVVQSTG
QMCKVYDSLALLPQDLQAARALCVIALLVALLFGLLVYLAGAKCTTCVEEKDSKARLVLT
SGIVFVISGVLTLIPVCWTAHAIIIRDFYNPLVAEAQKRELGASLYLGWAASGLLLLGGGL
LCCTCPSGGSQGPHSHYMARYSTSAPAIISRGPSYPTKNYV

Transmembrane domains:

amino acids 8-30 (type II), 82-102, 121-140, 166-186

249/392

FIGURE 239

AGTGACAATCTCAGAGCAGCTTCTACACCACAGCCATTTCCAGCATGAAAGATCACTGGGG
GTCTCCTTCTGCTCTGTACAGTGGTCTATTTCTGTAGCAGCTCAGAAGCTGCTAGTCTGT
CTCCAAAAAAGTGGACTGCAGCATTTACAAGAAGTATCCAGTGGTGGCCATCCCCTGCC
CCATCACATACCTACCAGTTTGTGGTTCTGACTACATCACCTATGGGAATGAATGTCACT
TGTGTACCGAGAGCTTGAAAAGTAATGGAAGAGTTCAGTTTCTTCACGATGGAAGTTGCT
AAATTCTCCATGGACATAGAGAGAAAAGGAATGATATTCTCATCATCATCTTCATCATCCC
AGGCTCTGACTGAGTTTCTTTCAGTTTTACTGATGTTCTGGGTGGGGGACAGAGCCAGAT
TCAGAGTAATCTTGACTGAATGGAGAAAGTTTCTGTGCTACCCCTACAAACCCATGCCTC
ACTGACAGACCAGCATTTTTTTTTTAAACACGTCAATAAAAAAATAATCTCCCAGA

250/392

FIGURE 240

MKITGGLLLLCTVVYFCSSSEAASLSPKKVDCSIYKKYPVVAIPCPITYLPVCGSDYITY
GNECHLCTESLKSNQGRVQFLHDGSC

Signal peptide:
amino acids 1-19

251/392

FIGURE 241

CCCGCGCCCGGTTCTCCCTCGCAGCACCTCGAAGTGCGCCCCCTCGCCCTCCTGCTCGCGC
CCCGCCGCCATGGCTGCCTCCCCCGCGCGGCCTGCTGTCTGGCCCTGACCGGGCTGGCG
CTGCTCCTGCTCCTGTGCTGGGGCCCAGGTGGCATAAGTGGAAATAAACTCAAGCTGATG
CTTCAAAAACGAGAAGCACCTGTTCCAATAAGACTAAAGTGCCGTTGATGAGAATAAA
GCCAAGAATTCCCTTGGCAGCCTGAAGCGCCAGAAGCGGCAGCTGTGGGACCGGACTCGG
CCCGAGGTGCAGCAGTGGTACCAGCAGTTTCTCTACATGGGCTTTGATGAAGCGAAATTT
GAAGATGACATCACCTATTGGCTTAACAGAGATCGAAATGGACATGAATACTATGGCGAT
TACTACCAACGTCACCTATGATGAAGACTCTGCAATTGGTCCCCGGAGCCCCTACGGCTTT
AGGCATGGAGCCAGCGTCAACTACGATGACTACTTAACCATGACTTGCCACACGCTGTACA
AGAAGCAAATAGCGATTCTCTTCATGTATCTCCTAATGCCTTACACTACTTGGTTTCTGA
TTTGCTCTATTTTCAGCAGATCTTTTCTACCTACTTTGTGTGATCAAAAAGAAGAGTTAA
ACAACACATGTAAATGCCTTTTGATATTTTCATGGGAATGCCTCTCATTTAAAAATAGAA
ATAAAGCATTTTGTAAAAAGA

252/392

FIGURE 242

MAASPARPAVLALTGLALLLLLCWGPGGISGNKLLMLQKREAPVPTKTKVAVDENKAKE
FLGSLKRQKRQLWDRTRPEVQQWYQQFLYMGFDEAKFEDDITYWLNDRDRNGHEYYGDDYYQ
RHYDEDSAIGPRSPYGFRHGASVNYDDY

Signal peptide:
amino acids 1-30

253/392

FIGURE 243

CTCCATTAAACCACCACCAGCTCCCCAAGCCACCCCTTCAGCCATGAAGTTCCTGCTCCT
GGTCTTGGCAGCCCTCGGATTCTTGACCCAGGTGATCCCAGCCAGTGCAGGTGGGTCAAA
ATGTGTGAGTAACACCCAGGATACTGCAGGACATGTTGCCACTGGGGGGAGACAGCATT
GTTTCATGTGCAACGCTTCCAGAAAATGCTGCATCAGCTACTCCTTCCTGCCGAAGCCTGA
CCTACCACAGCTCATCGGTAACCACTGGCAATCAAGGAGAAGAAACACACAAAGGAAAGA
CAAGAAGCAACAAACGACCGTAACATCATAAATAACCACTGCTATCGCCTCCACCAACTCA
GAGAAATATCATTTCCACAGTTCCAATTCCTCCTACATTGCTGAGTACTAGCCAAGGCTC
CTCTTTATGGGGCAGATATCTATAGCCAACCCCAAACTTCTGTCTTCTATCATTCTGTC
ATTCATCTAGTAACATAATTTGGAGTTTGTATCTATCTTACGAGAACAATCATCATGCAGA
TTCGTCCACAGGGGATCTGTCTAGTTTGGGTCCTCCAAATGAAAAATGTCAAGACAGAATT
GGACATGCAAAAGATTGACTGGGAGAACACACCTCTGATGGACAAAGGTGAGAGAGCA
GCCACAGGCAGGGAGAGCCTTCAGACTGCAACGCTGGCCTGATACGTGTCAAAGGAGAGCA
GGGATAGAGGAGGATTGAATAGAAGGAGACTAAGACTGCAGCTCTAAGAAAGTCTCAGCC
AAACAGATGGGGAGGCCCAAAGCAAGGCTTGCCCCCTCAGAGGAGCTCACGCAGGGCAGGA
ATAGCCAGGTTCTCATATCCCAGGGGTTTCAGACTTGGCTGAGAACAGCCCCCTGGAGAACA
TGGGGTGAAGTCTACCATAGGTCTGGAAGTATGAGGCTGTCCACCAACTATCCCCCTTGAA
GCAAGTTCTCTTGAAAGGAAATCTAAACAGTGCACCCCCATGGCTGCCACGGAGTATAAG
GAGGGAGAGAAAGGAGCTGAAAGTCTAGGTTTGGCCAGCTAGGTAGACTGACTTGTGAGG
TATTTATTTATTCATTTGAGTAACAAAGCAGACAGAATACATAGCCACCATTGGTAGTAC
ACCCCAAAGCAAGGATGGCATGATGCTGGTGAAGTCAACGTCCTACTCATGGTGTCAA
ATTGGCATAATCCTCTTGGAAGCTGTGTGGAAATAAGCACAGAGAAGCAGAACTCTAAT
TGCTTAATCCACTAAACATTACTTCTGGGAATTGGCTCATCATAAATTATCCAAGAGAAA
GCACAAAGTTATGGGCACAAAGGTTTTCATATAATATTATTTAAATGCTGAGAAAATG
AAAAATCTAAATGGTGAAATATATACTAATGCCATCTATAAATACAAACAAATAGAATG
TTTATAGAATAATGGAACATAATAACATTATTCAAAATTGCATTTATGCTATAGTTGTCA
AAATTGTCTCCTTATATGATACAAAACCTCATGAAAATTATGACTTTTTTGTGTTGGTGGG
AAGCAGAATTATGCATAAATTTCTCTTACAGTTCGATGCCCATTAGTTTATATAACAT
TTATTTGACACGTACTGACTTCTATCTGAGAAGAACAAACCAAAACACTCAGGCCTAAAT
AATTAACAAACGGTCCCTAAACACTAGCAAACCAGATAAGAAAAGATGTTAATGCCCATTC
CTAACTTATGTCTTAGACCAAAATTAATTCTAGATGGTTTAAATGACAGTGTAAGAGT
AAAGTATTAAGAGATTGTGTGGTCAAATATTCAATTTAAGAGCAAGGAAATTCTTATAAA
TATAACAATAGAGGCAGAACTCATGTAAGAATAAATTGATTAGGTGGTATTAAATATTAA
GTTCTTATGTATGTCAAAGATATCATTTTGAAATTCATCCATCTTATTGGGTATTGCAG
GAGTTCATTCTTTTGTGTTATAAATACTCTCCGTCATATGAATAGTATTCAATTTGTAT
ACTGGTTTGTGATGGACATTGGGTTGTCCAGTTTATGGCTATTACAAATAAGCTT
CTATGAACATTTATGTACA

254/392

FIGURE 244

MKFLLLVLAALGFLTQVIPASAGGSKCVSNTPGYCRTCCHWGETALFMCNASRKCCISYS
FLPKPDLPLIGNHWQSRRRNTQRKDKKQQTTVTS

Important features of the protein:

Signal peptide:

amino acids 1-16

Transmembrane domain:

amino acids 1-22

N-glycosylation site:

amino acids 50-53

cAMP- and cGMP-dependent protein kinase phosphorylation site:

amino acids 79-82

N-myristoylation site:

amino acids 23-28

255/392

FIGURE 245

GGAGAGAGGCGCGCGGGTGAAAGGCGCATTGATGCAGCCTGCGGCGGCCTCGGAGCGCGG
CGGAGCCAGACGCTGACCACGTTCTCTCCTCGGTCTCCTCCGCCTCCAGCTCCGCGCTG
CCCGGCAGCCGGGAGCCATGCGACCCCAGGGCCCCGCGCCTCCCCGCAGCGGCTCCGCG
GCCTCCTGCTGCTCCTGCTGCTGCAGCTGCCCCGCGCGTTCGAGCGCCTCTGAGATCCCCA
AGGGGAAGCAAAAGGCGCAGCTCCGGCAGAGGGAGGTGGTGGACCTGTATAATGGAATGT
GCTTACAAGGGCCAGCAGGAGTGCTGGTTCGAGACGGGAGCCCTGGGGCCAATGTTATTC
CGGGTACACCTGGGATCCCAGGTCCGGATGGATTCAAAGGAGAAAAGGGGGAATGTCTGA
GGGAAAGCTTTGAGGAGTCCTGGACACCCAACTACAAGCAGTGTTTATGGAGTTCATTGA
ATTATGGCATAGATCTTGGGAAAATTGCGGAGTGACATTTACAAAGATGCGTTCAAATA
GTGCTCTAAGAGTTTTGTTTCAAGTGGCTCACTTCGGCTAAAATGCAGAAATGCATGCTGTC
AGCGTTGGTATTTTACATTCAATGGAGCTGAATGTTTCAAGACCTCTTCCCATTTGAAGCTA
TAATTTATTTGGACCAAGGAAGCCCTGAAATGAATTAACAATTAATATTCATCGCACTT
CTTCTGTGGAAGGACTTTGTGAAGGAATTGGTGGCTGGATTAGTGGATGTTGCTATCTGGG
TTGGCACTTGTTTCAAGATTACCCAAAAGGAGATGCTTCTACTGGATGGAATTCAGTTTCTC
GCATCATTATTGAAGAACTACCAAAATAAATGCTTTAATTTTTCATTTGCTACCTCTTTTT
TTATTATGCCTTGGAATGGTTCACTTAAATGACATTTTAAATAAGTTTATGTATACATCT
GAATGAAAAGCAAAGCTAAATATGTTTACAGACCAAAGTGTGATTTTCACTGTTTTTAA
ATCTAGCATTATTATTTTGGCTTCAATCAAAGTGGTTTCAATATTTTTTTTAGTTGGTT
AGAATACTTTCTTCATAGTCACATTCTCTCAACCTATAATTTGGAATATTGTTGTGGTCT
TTTGTTTTTTCTCTTAGTATAGCATTTTAAAAAAATATAAAAGCTACCAATCTTTGTAC
AATTTGTAAATGTAAAGAATTTTTTTTTATATCTGTAAATAAAAATTATTTCCAACA

256/392

FIGURE 246

MRPQGPAASPQRLRGLLLLLLLQLPAPSSASEIPKGKQKAQLRQREVVDLYNGMCLQGPA
GVPGRDGSPGANVIPGTPGIPGRDGFKEKGECLRESFEESWTPNYKQCSWSSLNYGIDL
GKIAECTFTKMRSNSALRVLFSGSLRLKCRNACCQRWYFTFNGAECSGPLPIEAIYLDQ
GSPEMNSTINIHRTSSVEGLCEGIGAGLVDVAIWVGTCSDYPKGDASTGWNSVSRIEEE
LPK

Signal peptide:
amino acids 1-30

Transmembrane domain:
amino acids 195-217

257/392

FIGURE 247

GGCCGTTGGTTGGTGC GCGGCTGAAGGGTGTGGCGCGAGCAGCGTCGTTGGTTGGCCGGC
GGCGGGCCGGGACGGGCATGGCCCTGCTGCTGTGCCTGGTGTGCCTGACGGCGGCGCTGG
CCCACGGCTGTCTGCACTGCCACAGCAACTTCTCCAAGAAGTTCTCCTTCTACCGCCACC
ATGTGAACTTCAAGTCCTGGTGGGTGGGCGACATCCCCGTGTCAGGGGCGCTGCTCACCG
ACTGGAGCGACGACACGATGAAGGAGCTGCACCTGGCCATCCCCGCCAAGATCACCCGGG
AGAAGCTGGACCAAGTGGCGACAGCAGTGTACCAGATGATGGATCAGCTGTACCAGGGGA
AGATGTACTTCCCCGGGTATTTCCCCAACGAGCTGCGAAACATCTTCCGGGAGCAGGTGC
ACCTCATCCAGAACGCCATCATCGAAAGGCACCTGGCACCAGGCAGCTGGGGAGGAGGGC
AGCTCTCCAGGGAGGGACCCAGCCTAGCACCTGAAGGATCAATGCCATCACCCCGCGGGG
ACCTCCCCTAAGTAGCCCCCAGAGGCGCTGGGAGTGTGCCACCGCCCTCCCCTGAAGTT
TGCTCCATCTCACGCTGGGGGTCAACCTGGGGACCCCTTCCCTCCGGGCCATGGACACAC
ATACATGAAAACCAGGCCGCATCGACTGTCAGCACCGCTGTGGCATCTTCCAGTACGAGA
CCATCTCCTGCAACAACCTGCACAGACTCGCACGTGCCTGCTTTGGCTATAACTGCGAGT
AGGGCTCAGGCATCACACCCACCCGCTGCCAGGGCCCTACTGTCCCTGGGGTCCCAGGCTC
TCCTTGAGGGGGCTCCCCGCCTTCCACCTGGCTGTCATCGGGTAGGGCGGGGCCGTGGG
TTCAGGGGCGCACCACTTCCAAGCCTGTGTCCACAGGTCCTCGGCGCAGTGGGAAGTCAG
CTGTCCAGGGCCTCCTGAACTACATAAATAACTGGCACAAGTAAGTCCCTCCTCAAACC
AACACAGGCAGTGTGTGTATGTGAGCACCTCGTGGGTGAGTATGTGTGGGGCACAGGCTG
GCTCCCTCAGCTCCACGTCTTAGAGGGGCTCCCGAGGAGGTGGAACCTCAACCCAGCTC
TGCGCAGGAGGCGGCTGCAGTCTTTTCTCCCTCAAAGGTCTCCGACCCCTCAGCTGGAGG
CGGGCATCTTTCTAAAGGGTCCCCATAGGGTCTGGTTCCACCCCATCCAGGTCTGTGG
TCAGAGCCTGGGAGGGTTCCCTACGATGGTTAGGGGTGCCCCATGGAGGGGCTGACTGCC
CCACATTGCCTTTTACAGACAGGACACGAGCATGAGGTAAGGCCGCCCTGACCTGGACTTCA
GGGGGAGGGGGTAAAGGGAGAGAGGAGGGGGGCTAGGGGGTCTCTAGATCAGTGGGGGC
ACTGCAGGTGGGGCTCTCCCTATACCTGGGACACCTGCTGGATGTACCTCTGCAACCAC
ACCCATGTGGTGGTTTCATGAACAGACCACGCTCCTCTGCCTTCTCCTGGCCTGGGACAC
ACAGAGCCACCCCGGCCTTGTGAGTGACCCAGAGAAGGGAGGCCTCGGGAGAAGGGGTGC
TCGTAAGCCAACACCAGCGTGCCGCGGCCTGCACACCCCTTCGGACATCCAGGCACGAGG
GTGTCGTGGATGTGGCCACACATAGGACCACACGTCCCAGCTGGGAGGAGAGGCCTGGGG
CCCCCAGGGAGGGAGGCAGGGGTGGGGGACATGGAGAGCTGAGGCAGCCTCGTCTCCCC
GCAGCCTGGTATCGCCAGCCTTAAGGTGTCTGGAGCCCCACACTTGGCCAACCTGACCT
TGGAAGATGCTGCTGAGTGTCTCAAGCAGCACTGACAGCAGCTGGGCCTGCCCCAGGGCA
ACGTGGGGGCGGAGACTCAGCTGGACAGCCCCTGCCTGTCACTCTGGAGCTGGGCTGCTG
CTGCCTCAGGACCCCTCTCCGACCCCGACAGAGCTGAGCTGGCCAGGGCCAGGAGGGC
GGGAGGGAGGGAATGGGGTGGGCTGTGCGCAGCATCAGCGCCTGGGCAGGTCCGCAGAG
CTGCGGGATGTGATTAAAGTCCCTGATGTTTCTC

258/392

FIGURE 248

MALLLCLVCLTAALAHGCLHCHSNFSKKFSFYRHHVNFKSWVVDIPVSGALLTDWSDDT
MKELHLAIPAKITREKLDQVATAVYQMMDQLYQGKMYFPGYFPNELRNIFREQVHLIQNA
IIERHLAPGSWGGGQLSREGPSLAPEGSMPSPRGDLP

Signal peptide:
amino acids 1-15

259/392

FIGURE 249

CGACGATGCTACGCGCGCCCGGCTGCCTCCTCCGGACCTCCGTAGCGCCTGCCGCGGCCC
TGGCTGCGGCGCTGCTCTCGTCGCTTGCGCGCTGCTCTCTTCTAGAGCCGAGGGACCCGG
TGGCCTCGTCGCTCAGCCCCCTATTTCCGGCACCAAGACTCGCTACGAGGATGTCAACCCCG
TGCTATTGTGCGGGCCCCGAGGCTCCGTGGCGGGACCCCTGAGCTGCTGGAGGGGACCTGCA
CCCCGGTGACGCTGGTCGCCCTCATTGCCACGGCACCCGCTACCCACGGTCAAACAGA
TCCGCAAGCTGAGGCAGCTGCACGGGTGCTGCAGGCCCGCGGGTCCAGGGATGGCGGGG
CTAGTAGTACCGGCAGCCGCGACCTGGGTGCAGCGCTGGCCGACTGGCCTTTGTGGTACG
CGGACTGGATGGACGGGCAGCTAGTAGAGAAGGGACGGCAGGATATGCGACAGCTGGCGC
TGCGTCTGGCCTCGCTCTTCCCGGCCCTTTTCAGCCGTGAGAACTACGGCCGCTGCGGC
TCATCACCAGTTCGAAGCACCCTGCATGGATAGCAGCGCCGCTTCTGTCAGGGGCTGT
GGCAGCACTACCACCTGGCTTGCCGCCGCGGACGTCGCAGATATGGAGTTTGGACCTC
CAACAGTTAATGATAAACTAATGAGATTTTTTGATCACTGTGAGAAGTTTTTAATGAAG
TAGAAAAAATGCTACAGCTCTTTATCAGTGGAAGCCTTCAAACTGGACCAGAAATGC
AGAACATTTTAAAAAAGTTGCAGCTACTTTGCAAGTGCCAGTAAATGATTTAAATGCAG
ATTTAATTCAAGTAGCCTTTTTCACCTGTTCAATTTGACCTGGCAATTAAAGGTGTTAAAT
CTCCTTGGTGTGATGTTTTTGACATAGATGATGCAAAGGTATTAGAATATTTAAATGATC
TGAAACAATATTGGAAAAGAGGATATGGGTATACTATTAACAGTCGATCCAGCTGCACCT
TGTTTCAGGATATCTTTAGCACTTGACAAAGCAGTTGAACAGAAACAAAGGTCTCAGC
CAATTTCTTCTCCAGTCATCTCCAGTTTGGTCATGCAGAGACTCTTCTTCCACTGCTTT
CTCTCATGGGCTACTTCAAAGACAAGGAACCCCTAACAGCGTACAATTACAAAAACAAA
TGCATCGGAAGTTCCGAAGTGGTCTCATTTGTACCTTATGCCTCGAACCTGATATTTGTGC
TTTACCCTGTGAAAAATGCTAAGACTCCTAAAGAACAATTCCGAGTGCAGATGTTATTAA
ATGAAAAGGTGTTACCTTTGGCTTACTCACAAGAACTGTTTCATTTTATGAAGATCTGA
AGAACCACTACAAGGACATCCTTCAGAGTTGTCAAACAGTGAAGAATGTGAATTAGCAA
GGGCTAACAGTACATCTGATGAACTATGAGTAACTGAAGAACATTTTTAATTCCTTTAGGA
ATCTGCAATGAGTGATTACATGCTTGTAATAGGTAGGCAATTCCTTGATTACAGGAAGCT
TTTATATTACTTGAGTATTTCTGTCTTTTCACAGAAAAACATTGGGTTTCTCTCTGGGTT
TGGACATGAAATGTAAGAAAAGATTTTCACTGGAGCAGCTCTCTTAAGGAGAAACAAAT
CTATTTAGAGAAACAGCTGGCCCTGCAAATGTTTACAGAAATGAAATTCCTTCTACTTAT
ATAAGAAATCTCAGCTGAGATAGAATTGTGATTTTATAATAACACTTGAAAAGTGCTGG
AGTAACAAAATATCTCAGTTGGACCATCCTTAACCTTGATTGAACTGTCTAGGAACCTTAC
AGATTGTTCTGCAGTTCTCTCTTTTCTCCTCAGGTAGGACAGCTCTAGCATTTTCTTAA
TCAGGAATATTGTGGTAAGCTGGGAGTATCACTCTGGAAGAAAGTAACATCTCCAGATGA
GAATTTGAAACAAGAAACAGAGTGTTGTAAAAGGACACCTTCACTGAAGCAAGTCGGA
GTACAATGAAAATAAATATTTTTGGTATTTATTTATGAAATATTTGAACATTTTTTCAAT
AATTCCTTTTTACTTCTAGGAAGTCTCAAAAAGACCATCTTAAATTATTATATGTTTGGAC
AATTAGCAACAAGTCAGATAGTTAGAATCGAAGTTTTTCAAATCCATTGCTTAGCTAACT
TTTTCATTTCTGTCACTTGGCTTCGATTTTTATATTTTCTATTATATGAAATGTATCTTT
TGGTTGTTTGATTTTCTTTCTTTCTTTGTAAATAGTTCTGAGTTCTGTCAAATGCCGTG
AAAGTATTTGCTATAATAAAGAAATTCTTGTGACTTTAAAAA

260/392

FIGURE 250

MLRAPGCLLRTSVAPAAALAAALLSSLARCSLLEPRDPVASSLSFYFGTKTRYEDVNPVL
LSGPEAPWRDPELLEGTCTPVQLVALIRHGTRYPTVKQIRKLRQLHGLLQARGSRDGGAS
STGSRDLGAALADWPLWYADWMDGQLVEKGRQDMRQLALRLASLFPALFSRENYGRLRLI
TSSKHRCMDSSAAFLOGLWQHYPGLPPPDVADMEFGPPTVNDKLMRFFDHCEKFLTEVE
KNATALYHVEAFKTGPEMQNILKKVAATLQVPVNDLNADLIQVAFFTCSFDLAIKGVKSP
WCDVFDIDDAKVLEYLNDLKQYWKRGYGYTTINSRSSCTLFQDIFQHLDKAVEQKQRSQPI
SSPVILQFGHAETLLPLLSLMGYFKDKEPLTAYNYKKQMRKFRSGLIVPYASNLI FVLY
HCENAKTPKEQFRVQMLLNEKVLPLAYSQETVSFYEDLKNHYKDILQSCQTSEECELARA
NSTSDEL

Important features:**Signal sequence**

amino acids 1-30

N-glycosylation sites:

amino acids 242-246, 481-485

N-myristoylation sites.

amino acids 107-113, 113-119, 117-123, 118-124, 128-134

Endoplasmic reticulum targeting sequence:

amino acids 484-489

FIGURE 251

GGAGAGCCGCGGCTGGGACCGGAGTGGGGAGCGCGGCTGGAGGTGCCACCGCGCGGCT
TGGCGGAGAGATCAGAAGCCTCTTCCCCAAGCCGAGCCTCAGCGGGGACCCGGGCT
CAGGGACGCGGCGGCGGCGGCGGCGACTGCAGTGGCTGGACGATGGCAGCGTCCGCCGGA
GCCGGGGCGGTGATTGCAGCCCCAGACAGCCGGCGCTGGCTGTGGTTCGGTGCTGGCGGCG
GCGCTTGGGCTCTTGACAGCTGGAGTATCAGCCTTGGAAAGTATATACGCCAAAAGAAATC
TTCGTGGCAAATGGTACACAAGGGAAGCTGACCTGCAAGTTCAAGTCTACTAGTACGACT
GGCGGGTTGACCTCAGTCTCCTGGAGCTTCCAGCCAGAGGGGGCCGACACTACTGTGTCTG
TTTTTCCACTACTCCCAAGGGCAAGTGTACCTTGGGAATTATCCACCATTAAAGACAGA
ATCAGCTGGGCTGGAGACCTTGACAAGAAAGATGCATCAATCAACATAGAAAATATGCAG
TTTATACACAATGGCACCTATATCTGTGATGTCAAAAACCCCTCTGACATCGTTGTCCAG
CCTGGACACATTAGGCTCTATGTCTGTAGAAAAAGAGAATTTGCCTGTGTTTTCCAGTTTGG
GTAGTGGTGGGCATAGTTACTGCTGTGGTCTAGGTCTCACTCTGCTCATCAGCATGATT
CTGGCTGTCTCTATAGAAGGAAAACTCTAAACGGGATTACACTGGCTGCAGTACATCA
GAGAGTTTGTCAACAGTTAAGCAGGCTCCTCGGAAGTCCCCCTCCGACACTGAGGGTCTT
GTAAAGAGTCTGCCTTCTGGATCTCACCAGGGCCAGTCATATATGCACAGTTAGACCA
TCCGGCGGACATCACAGTGACAAGATTAAACAAGTCAGAGTCTGTGGTGTATGCGGATATC
CGAAAGAATTAAGAGAATACCTAGAACATATCCTCAGCAAGAAACAAAACCAAACCTGGAC
TCTCGTGCAAAAATGTAGCCCATTAACCATGTAGCCTTGGAGACCCAGGCAAGGACAA
GTACACGTGTACTCACAGAGGGAGAGAAAAGATGTGTACAAAGGATATGTATAAATATTCT
ATTTAGTCATCCTGATATGAGGAGCCAGTGTTCATGATGAAAAGATGGTATGATTCTAC
ATATGTACCCATTGTCTTGCTGTTTTTGTACTTTCTTTTCAGGTCATTTACAATTGGGAG
ATTTCAGAAACATTCTTTTACCATCATTTAGAAATGGTTTGCCTTAATGGAGACAATAG
CAGATCCTGTAGTATTTCCAGTAGACATGGCCTTTTAATCTAAGGGCTTAAGACTGATTA
GTCTTAGCATTTACTGTAGTTGGAGGATGGAGATGCTATGATGGAAGCATACCCAGGGTG
GCCTTTAGCACAGTATCAGTACCATTATTTGTCTGCCGCTTTTAAAAAATACCCATTGG
CTATGCCACTTGAAAAACAATTTGAGAAGTTTTTTTGAAGTTTTTCTCACTAAAATATGG
GCAATTGTTAGCCTTACATGTTGTGTAGACTTACTTTAAGTTTGCACCCTTGAAATGTGT
CATATCAATTTCTGGATTCAATATAGCAAGATTAGCAAAAGGATAAATGCCGAAGGTCAC
TCATTCTGGACACAGTTGGATCAATACTGATTAAAGTAGAAAATCCAAGCTTTGCTTGAGA
ACTTTTGTAAACGTGGAGAGTAAAAAGTATCGGTTTTTA

262/392

FIGURE 252

MAASAGAGAVIAAPDSRRWLWSVLAAALGLLTAGVSALEVYTPKEIFVANGTQGKLTCKF
KSTSTTGGLTSVSWSFQPEGADTTVSFFHYSQGQVYLGNYPPFKDRISWAGDLDKKDASI
NIENMQFIHNGTYICDVKNPPDIVVQPGHIRLYVVEKENLPVFPVWVVVGIVTAVVLGLT
LLISMILAVLYRRKNSKRDTGTCSTSESLSPVKQAPRKSPSDTEGLVKSLPSGSHQGPVI
YAQLDHSGGHHSDKINKSESVVYADIRKN

Signal peptide:
amino acids 1-37

Transmembrane domain:
amino acids 161-183

263/392

FIGURE 253

GTGACACTATAGAAGAGCTATGACGTCGCATGCACGCGTACGTAAGCTCGGAATTCGGCT
CGAGGCTGGTGGGAAGAAGCCGAGATGGCGGCAGCCAGCGCTGGGGCAACCCGGCTGCTC
CTGCTCTTGCTGATGGCGGTAGCAGCGCCAGTCGAGCCCGGGGCAGCGGCTGCCGGGCC
GGGACTGGTGC GCGAGGGGCTGGGGCGGAAGGTGAGAGGGCGAGGCCTGTGGCACGGTG
GGGCTGCTGCTGGAGCACTCATTTGAGATCGATGACAGTGCCAACTTCCGGAAGCGGGGC
TCACTGCTCTGGAACCAGCAGGATGGTACCTTGTCCCTGTCACAGCGGCAGCTCAGCGAG
GAGGAGCGGGCCGACTCCGGGATGTGGCAGCCCTGAATGGCCTGTACCGGGTCCGGATC
CCAAGGCGACCCGGGGCCCTGGATGGCCTGGAAGCTGGTGGCTATGTCTCCTCCTTTGTC
CCTGCGTGCTCCCTGGTGGAGTCGCACCTGTGCGACCAGCTGACCCTGCACGTGGATGTG
GCCGGCAACGTGGTGGGCGTGTCGGTGGTGACGCACCCCGGGGGCTGCCGGGGCCATGAG
GTGGAGGACGTGGACCTGGAGCTGTTCAACACCTCGGTGCAGCTGCAGCCGCCACCACA
GCCCCAGGCCCTGAGACGGCGGCCTTCATTGAGCGCCTGGAGATGGAACAGGCCCCAGAAG
GCCAAGAACCCCCAGGAGCAGAAGTCCTTCTTCGCCAAATACTGGATGTACATCATTCCC
GTCGTCCTGTTCTCATGATGTCAGGAGCGCCAGACACCGGGGGCCAGGGTGGGGGTGGG
GGTGGGGGTGGTGGTGGGGGTAGTGGCCTTTGCTGTGTGCCACCCTCCCTGTAAGTCTAT
TTAAAAACATCGACGATACATTGAAATGTGTGAACGTTTTGAAAAGCTACAGCTTCCAGC
AGCCAAAAGCAACTGTTGTTTTGGCAAGACGGTCCTGATGTACAAGCTTGATTGAAATTC
ACTGCTCACTTGATACGTTATTCAGAAACCCAAGGAATGGCTGTCCCCATCCTCATGTGG
CTGTGTGGAGCTCAGCTGTGTTGTGTGGCAGTTTATTAACTGTCCCCCAGATCGACACG
CAAAAAAAAA

264/392

FIGURE 254

MAAASAGATRLLLLLLLMAVAAPSRARGSGCRAGTGARGAGAEGREGEACGTVGLLLEHSF
EIDDSANFRKRGSLLWNQQDGTLSLSQRQLSEEERGLRDVAALNGLYRVRI PRRPGALD
GLEAGGYVSSFVPACSLVESHLSQDLTLHVDVAGNVVGVSVVTHPGGCRGHEVEDVDLEL
FNTSVQLQPPTTAPGPETA AFIERLEMEQAQKAKNPQEQKSFFAKYWMIIPVVLFLMMS
GAPDTGGQGGGGGGGGGGGSGLCVPPSL

Signal peptide:
amino acids 1-24

Transmembrane domain:
amino acids 226-243

265/392

FIGURE 255

GCGACGCGCGGCGGGGCGGCGAGAGGAAACGCGGCGCCGGGCGGGCCCGGCCCTGGAGA
TGGTCCCCGCGCGCCGCGGGCTGGTGTTGTCTCGTGCTCTGGCTCCCCGCGTGCGTCGCGG
CCCACGGCTTCCGTATCCATGATTATTTGTACTTTCAAGTGCTGAGTCCTGGGGACATTC
GATACATCTTCACAGCCACACCTGCCAAGGACTTTGGTGGTATCTTTCACACAAGGTATG
AGCAGATTCACCTTGTCCCCGCTGAACCTCCAGAGGCCTGCGGGGAACCTCAGCAACGGTT
TCTTCATCCAGGACCAGATTGCTCTGGTGGAGAGGGGGGGCTGCTCCTTCCTCTCCAAGA
CTCGGGTGGTCCAGGAGCACGGCGGGCGGGCGGTGATCATCTCTGACAACGCAGTTGACA
ATGACAGCTTCTACGTGGAGATGATCCAGGACAGTACCCAGCGCACAGCTGACATCCCCG
CCCTCTTCCTGCTCGGCCGAGACGGCTACATGATCCGCCGCTCTCTGGAACAGCATGGGC
TGCCATGGGCCATCATTCCATCCCAGTCAATGTCACCAGCATCCCCACCTTTGAGCTGC
TGCAACCGCCCTGGACCTTCTGGTAGAAGAGTTTGTCCCACATTCAGCCATAAGTGA
CTGAGCTGGGAAGGGGAAACCCAGGAATTTTGCTACTTGGAATTTGGAGATAGCATCTGG
GGACAAGTGGAGCCAGGTAGAGGAAAAGGTTTGGGCGTTGCTAGGCTGAAAGGGAAGCC
ACACCACTGGCCTTCCCTTCCCCAGGGCCCCCAAGGTGTCTCATGCTACAAGAAGAGGC
AAGAGACAGGCCCCAGGGCTTCTGGCTAGAACCCGAAACAAAAGGAGCTGAAGGCAGGTG
GCCTGAGAGCCATCTGTGACCTGTCACACTCACCTGGCTCCAGCCTCCCCTACCCAGGGT
CTCTGCACAGTGACCTTCACAGCAGTTGTTGGAGTGGTTTAAAGAGCTGGTGTTTGGGGA
CTCAATAAACCTCACTGACTTTTTAGCAATAAAGCTTCTCATCAGGGTTGCAAAAAAA
AAAAAAAAAAAAAAAAAAAA

266/392

FIGURE 256

MVPGAAGWCCLVLWLPACVAAHGFRIDYLYFQVLSPGDIRYIFTATPAKDFGGIFHTRY
EQIHLVPAEPPEACGELSNGFFIQDQIALVERGGCSFLSKTRVVQEHGGRAVIISDNAV
NDSFYVEMIQDSTQRTADIPALFLLGRDGYMIRRSLEQHGLPWAIISIPVNVTSIPTFEL
LQPPWTFW

Signal peptide:
amino acids 1-20

267/392

FIGURE 257

CTCGCTTCTTCCTTCTGGATGGGGGCCCAGGGGGCCCAGGAGAGTATAAAGGCGATGTGG
AGGGTGCCCGGCACAACCAGACGCCCAGTCACAGGCGAGAGCCCTGGGATGCACCGGCCA
GAGGCCATGCTGCTGCTGCTCACGCTTGCCCTCCTGGGGGGCCCCACCTGGGCAGGGAAG
ATGTATGGCCCTGGAGGAGGCAAGTATTTTCAGCACCCTGAAGACTACGACCATGAAATC
ACAGGGCTGCGGGTGTCTGTAGGTCTTCTCCTGGTGAAAAGTGTCCAGGTGAAACTTGGA
GACTCCTGGGACGTGAAACTGGGAGCCTTAGGTGGGAATACCCAGGAAGTCACCCCTGCAG
CCAGGCGAATACATCACAAAAGTCTTTGTGCGCCTTCCAAGCTTTCCTCCGGGGTATGGTC
ATGTACACCAGCAAGGACCGCTATTTCTATTTTGGGAAGCTTGATGGCCAGATCTCCTCT
GCCTACCCCAGCCAAGAGGGGCAGGTGCTGGTGGGCATCTATGGCCAGTATCAACTCCTT
GGCATCAAGAGCATTGGCTTTGAATGGAATTATCCACTAGAGGAGCCGACCACTGAGCCA
CCAGTTAATCTCACATACTCAGCAAACCTACCCGTGGGTGCTAGGGTGGGGTATGGGGC
CATCCGAGCTGAGGCCATCTGTGTGGTGGTGGCTGATGGTACTGGAGTAACTGAGTCGGG
ACGCTGAATCTGAATCCACCAATAAATAAAGCTTCTGCAGAAAA

268/392

FIGURE 258

MHRPEAMLLLLTLALLGGPTWAGKMYGPGGGKYFSTTEDYDHEITGLRVSVGLLLVKSQ
VKLGDSWDVKLGALGGNTQEVTLQPGYITKVFAFQAFLRGMVMTSKDRYFYFGKLDG
QISSAYPSQEGQVLVGIYGQYQLLGIKSIGFEWNYPLEEPTTEPPVNLTYSANSPVGR

Signal peptide:
amino acids 1-22

269/392

FIGURE 259

CAGACATGGCTCAGTCACTGGCTCTGAGCCTCCTTATCCTGGTTCTGGCCTTTGGCATCC
CCAGGACCCAAGGCAGTGATGGAGGGGCTCAGGACTGTTGCCTCAAGTACAGCCAAAGGA
AGATTCCCGCCAAGGTTGTCCGCAGCTACCGGAAGCAGGAACCAAGCTTAGGCTGCTCCA
TCCCAGCTATCCTGTTCTTGCCCCGCAAGCGCTCTCAGGCAGAGCTATGTGCAGACCCAA
AGGAGCTCTGGGTGCAGCAGCTGATGCAGCATCTGGACAAGACACCATCCCCACAGAAAC
CAGCCCAGGGCTGCAGGAAGGACAGGGGGGCCTCCAAGACTGGCAAGAAAGGAAAGGGCT
CCAAAGGCTGCAAGAGGACTGAGCGGTCACAGACCCCTAAAGGGCCATAGCCCAGTGAGC
AGCCTGGAGCCCTGGAGACCCACACAGCCTCACCAGCGCTTGAAGCCTGAACCCAAGATG
CAAGAAGGAGGCTATGCTCAGGGGCCCTGGAGCAGCCACCCCATGCTGGCCTTGCCACAC
TCTTTCTCCTGCTTTAACCACCCCATCTGCATTCCCAGCTCTACCCTGCATGGCTGAGCT
GCCACAGCAGGCCAGGTCCAGAGAGACCGAGGAGGGAGAGTCTCCAGGGAGCATGAGA
GGAGGCAGCAGGACTGTCCCCTTGAAGGAGAATCATCAGGACCCTGGACCTGATACGGCT
CCCCAGTACACCCACCTCTTCCTTGTAATATGATTATACCTAACTGAATAAAAAGCT
GTTCTGTCTTCCCNCCCA

270/392

FIGURE 260

MAQSLALSLLILVLAFGIPRTQGS DGG AQDCCLKYSQRKIPAKVVRSYRKQEPSLGCSIP
AILFLPRKRSQAELCADPKELWVQQLMQHLDKTPSPQKPAQGCRKDRGASKTGKKKGKGSK
GCKRTERSQTPKGP

Important features of the protein:

Signal peptide:

amino acids 1-17

cAMP- and cGMP-dependent protein kinase phosphorylation site:

amino acids 67-71

N-myristoylation sites:

amino acids 17-23, 23-29, 27-33, 108-114, 118-124, 121-127

Amidation site:

amino acids 112-116

Small cytokines:

amino acids 51-91

271/392

FIGURE 261

GGGACTACAAGCCGCGCCGCGCTGCCGCTGGCCCCCTCAGCAACCCCTCGACATGGCGCTGA
GGCGGCCACCGCGACTCCGGCTCTGCGCTCGGCTGCCTGACTTCTTCCTGCTGCTGCTTT
TCAGGGGCTGCCTGATAGGGGCTGTAAATCTCAAATCCAGCAATCGAACCCCACTGGTAC
AGGAATTTGAAAGTGTGGAACGTCTTGCATCATTACGGATTTCGAGACAAGTGACCCCA
GGATCGAGTGGAAGAAAATTCAAGATGAACAAACCACATATGTGTTTTTTGACAACAAAA
TTCAGGGGAGACTTGGCGGGTCGTGCAGAAATACTGGGGAAGACATCCCTGAAGATCTGGA
ATGTGACACGGAGAGACTCAGCCCTTTATCGCTGTGAGGTCGTTGCTCGAAATGACCGCA
AGGAAATTGATGAGATTGTGATCGAGTTAACTGTGCAAGTGAAGCCAGTGACCCCTGTCT
GTAGAGTGCCGAAGGCTGTACCAGTAGGCAAGATGGCAACACTGCACTGCCAGGAGAGTG
AGGGCCACCCCGGCCTCACTACAGCTGGTATCGCAATGATGTACCCTGCCACGGATT
CCAGAGCCAATCCCAGATTTTCGAATTTCTTCTTTCCACTTAAACTCTGAAACAGGCACCT
TGGTGTTCACTGCTGTTTCAAGGACGACTCTGGGCAGTACTACTGCATTGCTTCCAATG
ACGCAGGCTCAGCCAGGTGTGAGGAGCAGGAGATGGAAGTCTATGACCTGAACATTGGCG
GAATTATTGGGGGGGTTCTGGTTGTCTTGGCTGTACTGGCCCTGATCACGTTGGGCATCT
GCTGTGCATACAGACGTGGCTACTTCATCAACAATAAACAGGATGGAGAAAGTTACAAGA
ACCCAGGGAACCCAGATGGAGTTAACTACATCCGCACTGACGAGGAGGGCGACTTCAGAC
ACAAGTCATCGTTTGTGATCTGAGACCCGCGGTGTGGCTGAGAGCGCACAGAGCGCACGT
GCACATACCTCTGCTAGAACTCCTGTCAAGGCAGCGAGAGCTGATGCACTCGGACAGAG
CTAGACACTCATTGAGAAGCTTTTCGTTTTGGCCAAAGTTGACCACTACTCTTCTTACTC
TAACAAGCCACATGAATAGAAGAAATTTTCTCAAGATGGACCCGGTAAATATAACCACAA
GGAAGCGAAACTGGGTGCGTTCACTGAGTTGGGTTCTTAATCTGTTTCTGGCCTGATTCC
CGCATGAGTATTAGGGTGATCTTAAAGAGTTTGCTCACGTAAACGCCCGTGTGGGCCCT
GTGAAGCCAGCATGTTTACCCTGGTTCGTTGAGCAGCCACGACAGCACCATTGTGAGATGG
CGAGGTGGCTGGACAGCACAGCAGCGCATCCCGCGGGAACCCAGAAAGGCTTCTTAC
ACAGCAGCCTTACTTCATCGGCCCCACAGACACCACCGCAGTTTCTTCTTAAAGGCTCTGC
TGATCGGTGTTGCAGTGTCCATTGTGGAGAAGCTTTTTGGATCAGCATTTTGTAAAAACA
ACCAAAATCAGGAAGGTAAATTGGTTGCTGGAAGAGGGATCTTGCTGAGGAACCTGCT
TGTCCAACAGGGTGTGAGGATTTAAGGAAAACCTTCGTTTAGGCTAAGTCTGAAATGGT
ACTGAAATATGCTTTTCTATGGGTCTGTTTATTTTATAAAATTTTACATCTAAATTTTT
GCTAAGGATGTATTTGATTATTGAAAAGAAAATTTCTATTTAACTGTAAATATATTGT
CATACAATGTTAAATAACCTATTTTTTTTAAAAAAGTTCAACTTAAGGTAGAAGTTCCAAG
CTACTAGTGTTAAATTGGAAAATATCAATAATTAAGAGTATTTTACCAAGGAATCCTCT
CATGGAAGTTTACTGTGATGTTTCTTTCTCACACAAGTTTTAGCCTTTTTTACAAGGGA
ACTCATACTGTCTACACATCAGACCATAGTTGCTTAGGAAACCTTTAAAAATTCAGTTA
AGCAATGTTGAAATCAGTTTGCATCTCTTCAAAAGAAACCTCTCAGGTTAGCTTTGAACT
GCCTCTTCTGAGATGACTAGGACAGTCTGTACCCAGAGGCCACCCAGAAGCCCTCAGAT
GTACATACAGATGCCAGTCAGCTCCTGGGGTTGCGCCAGGCGCCCCCGCTCTAGCTCA
CTGTTGCCTCGCTGTCTGCCAGGAGGCCCTGCCATCCTTGGGCCCTGGCAGTGGCTGTGT
CCCAGTGAGCTTTACTCAGTGGCCCTTGCTTCATCCAGCACAGCTCTCAGGTGGGCACT
GCAGGGACACTGGTGTCTTCCATGTAGCGTCCCAGCTTTGGGCTCCTGTAACAGACCTCT
TTTTGGTTATGGATGGCTCACAAAATAGGGCCCCCAATGCTATTTTTTTTTTTAAGTTT
GTTTAATTATTTGTTAAGATTGTCTAAGGCCAAAGGCAATTGCGAAATCAAGTCTGTCAA
GTACAATAACATTTTAAAGAAAATGGATCCCCTGTTTCTTTGCCACAGAGAAAGC
ACCCAGACGCCACAGGCTCTGTGCGATTTCAAAACAAACCATGATGGAGTGGCGGCCAGT
CCAGCCTTTTAAAGAACGTCAGGTGGAGCAGGTCAGGTGAAAGGCCTGGCGGGGAGGAAAG
TGAAACGCCTGAATCAAAAGCAGTTTTCTAATTTTGACTTTAAATTTTTTCATCCGCCGGA

272/392

GACACTGCTCCCATTTGTGGGGGGACATTAGCAACATCACTCAGAAGCCTGTGTTCTTCA
AGAGCAGGTGTTCTCAGCCTCACATGCCCTGCCGTGCTGGACTCAGGACTGAAGTGCTGT
AAAGCAAGGAGCTGCTGAGAAGGAGCACTCCACTGTGTGCCTGGAGAATGGCTCTCACTA
CTCACCTTGTCTTTCAGCTTCCAGTGTCTTGGGTTTTTTATACTTTGACAGCTTTTTTTTT
AATTGCATACATGAGACTGTGTTGACTTTTTTTAGTTATGTGAAACACTTTGCCGCAGGC
CGCCTGGCAGAGGCAGGAAATGCTCCAGCAGTGGCTCAGTGCTCCCTGGTGTCTGCTGCA
TGGCATCCTGGATGCTTAGCATGCAAGTTCCCTCCATCATTGCCACCTTGGTAGAGAGGG
ATGGCTCCCCACCCTCAGCGTTGGGGATTACGCTCCAGCCTCCTTCTTGGTTGTCATAG
TGATAGGGTAGCCTTATTGCCCCCTCTTCTTATACCCTAAAACCTTCTACACTAGTGCCA
TGGGAACCAGGTCTGAAAAAGTAGAGAGAAGTGAAAGTAGAGTCTGGGAAGTAGCTGCCT
ATAACTGAGACTAGACGGAAAAGGAATACTCGTGTATTTTAAGATATGAATGTGACTCAA
GACTCGAGGCCGATACGAGGCTGTGATTCTGCCTTTGGATGGATGTTGCTGTACACAGAT
GCTACAGACTTGTAATAACACACCGTAATTTGGCATTTGTTTAACCTCATTTATAAAAGC
TTCAAAAAAACCCA

273/392

FIGURE 262

MALRRPPRLRLCARLPDFFLLLLFRGCLIGAVNLKSSNRTPVVQEFESVELSCIITDSQT
SDPRIEWKKIQDEQTTYVFFDNKIQGDLAGRAEILGKTSLSKIWNVTRRDSALYRCEVVAR
NDRKEIDEIVIELTVQVKPVPVPCRVPKAVPVGKMATLHCQESEGHPRPHYSWYRNDVPL
PTDSRANPRFRNSSFHLNSETGTLVFTAVHKDDSGQYYCIASNDAGSARCEEQEMEVDL
NIGGIIGGVLVVLAVLALITLGICCAYYRRGYFINNKQDGESYKNPGKPDGVNYIRTDEEG
DFRHKSSFVI

Important features of the protein:

Signal peptide:

amino acids 1-30

Transmembrane domain:

amino acids 243-263

N-glycosylation sites:

amino acids 104-107, 192-195

cAMP- and cGMP-dependent protein kinase phosphorylation site:

amino acids 107-110

Casein kinase II phosphorylation site:

amino acids 106-109, 296-299

Tyrosine kinase phosphorylation site:

amino acids 69-77

N-myristoylation sites:

amino acids 26-31, 215-220, 226-231, 243-248, 244-249, 262-267

274/392

FIGURE 263

CCAGGACCAGGGCGCACCGGCTCAGCCTCTCACTTGT CAGAGGCCGGGGAAGAGAAGCAA
AGCGCAACGGTGTGGTCCAAGCCGGGGCTTCTGCTTCGCCTCTAGGACATACACGGGACC
CCCTAACTTTCAGTCCCCCAAACGCGCACCCCTCGAAGTCTTGAAGTCCAGCCCCGCACATC
CACGCGCGGCACAGGCGCGGCAGGCGGCAGGTCCCGGCCGAAGGCGATGCGCGCAGGGGG
TCGGGCAGCTGGGCTCGGGCGGCGGGAGTAGGGCCCCGGCAGGGAGGCAGGGAGGCTGCAT
ATTCAGAGTCGCGGGCTGCGCCCTGGGCAGAGGCCGCCCTCGCTCCACGCAACACCTGCT
GCTGCCACCGCGCCGCGATGAGCCGCGTGGTCTCGCTGCTGCTGGGCGCCGCGCTGCTCT
GCGGCCACGGAGCCTTCTGCCGCCGCGTGGTCAGCGGCCAAAAGGTGTGTTTGTGCTGACT
TCAAGCATCCCTGCTACAAAATGGCCTACTTCCATGAAGTGTCCAGCCGAGTGAGCTTTC
AGGAGGCACGCCTGGCTTGTGAGAGTGAGGGAGGAGTCTCCTCAGCCTTGAGAATGAAG
CAGAACAGAAGTTAATAGAGAGCATGTTGCAAAACCTGACAAAACCCGGGACAGGGATTT
CTGATGGTGATTTCTGGATAGGGCTTTGGAGGAATGGAGATGGGCAAAACATCTGGTGCCT
GCCAGATCTCTACCACTGGTCTGATGGAAGCAATTCCAGTACCGAAACTGGTACACAG
ATGAACCTTCTGCGGAAGTGAAAAGTGTGTTGTGATGTATCACCAACCAACTGCCAATC
CTGGCCTTGGGGGTCCCTACCTTTACCAGTGGAATGATGACAGGTGTAACATGAAGCACA
ATTATATTTGCAAGTATGAACCAGAGATTAATCCAACAGCCCCTGTAGAAAAGCCTTATC
TTACAAATCAACCAGGAGACACCCATCAGAATGTGGTTGTTACTGAAGCAGGTATAATTC
CCAATCTAATTTATGTTGTTATACCAACAATACCCCTGCTCTTACTGATACTGGTTGCTT
TTGGAACCTGTTGTTTCCAGATGCTGCATAAAAGTAAAGGAAGAACAAAACTAGTCCAA
ACCAGTCTACACTGTGGATTTCAAAGAGTACCAGAAAAGAAAGTGGCATGGAAGTATAAAT
AACTCATTTGACTTGGTTCCAGAATTTTGTAAATCTGGATCTGTATAAGGAATGGCATCAG
AACAATAGCTTGAATGGCTTGAAATCACAAAGGATCTGCAAGATGAAGTGAAGCTCCC
CCTTGAGGCAAATATTAAAGTAATTTTATATGTCTATTATTTTCAATTAAGAATATGCT
GTGCTAATAATGGAGTGAGACATGCTTATTTTGTCTAAAGGATGCACCCAACTTCAAAT
TCAAGCAAATGAAATGGACAATGCAGATAAAGTTGTTATCAACACGTGCGGAGTATGTGT
GTTAGAAGCAATTCCTTTTATTTCTTTACCTTTTCATAAGTTGTTATCTAGTCAATGTAA
TGTATATTGTATTGAAATTTACAGTGTGCAAAAGTATTTTACCTTTGCATAAGTGTGTTGA
TAAAAATGAAGTGTCTAATATTTATTTTATGGCATCTCATTTTTCAATACATGCTCTT
TTGATTAAAGAACTTATTACTGTTGTCAACTGAATTCACACACACACAAATATAGTACC
ATAGAAAAAGTTTGTCTTCGAAATAATTCATCTTTCAGCTTCTCTGCTTTTGGTCAAT
GTCTAGGAAATCTCTTCAGAAATAAGAAGCTATTTCAATTAAGTGTGATATAAACCTCCTC
AAACATTTTACTTAGAGGCAAGGATTGTCTAATTTCAATTGTGCAAGACATGTGCCTTAT
AATTATTTTACTTAAATTTAAACAGATTTTGTAAATAATGTAAGTGTGTTAATAGGTGC
ATAAACACTAATGCAGTCAATTTGAACAAAAGAGTGACATACACAATATAAATCATATG
TCTTCACACGTTGCCTATATAATGAGAAGCAGCTCTCTGAGGGTTCTGAAATCAATGTGG
TCCCTCTCTTGCCCACTAAACAAAGATGGTTGTTGCGGGGTTGGGATTGACACTGGAGGC
AGATAGTTGCAAAGTTAGTCTAAGGTTTCCCTAGCTGTATTTAGCCTCTGACTATATTAG
TATACAAAGAGGTGATGTGGTTGAGACCAGGTGAATAGTCACTATCAGTGTGGAGACAAG
CACAGCACACAGACATTTTAGGAAGGAAAGGAAGTACGAAATCGTGTGAAAATGGGTGG
AACCCATCAGTGATCGCATATTCATTGATGAGGGTTTGTCTGAGATAGAAAATGGTGGCT
CCTTTCTGTCTTATCTCCTAGTTTCTTCAATGCTTACGCTTGTCTCTCAAGAGAAAG
TTGTAAGTCTCTGGTCTTCATATGTCCCTGTGCTCCTTTTAACCAATAAAGAGTCTTG
TTTCTGGGGGAA

275/392

FIGURE 264

MSRVVSLLLGAALLCGHGAFRRVVSQGKVCFAFKHPCYKMAFHELSRVSFQEARLA
CESEGGVLLSLENAEQKLIESMLQNLTKPGTGISDGDFWIGLWRNGDQTSGACPDLYQ
WSDGSNSQYRNWYTDEPSCGSEKCVVMYHQPTANPGLGGPYLYQWNDDRCNMKHNYICKY
EPEINPTAPVEKPYLTNQPGDTHQNVVTEAGIIPNLIYVVIPTIPLLLLILVAFGTCCF
QMLHKSKGRTKTSPNQSTLWISKSTRKESGMEV

Important features of the protein:

Signal peptide:

amino acids 1-21

Transmembrane domain:

amino acids 214-235

N-glycosylation sites:

amino acids 86-89 and 255-258

cAMP- and cGMP-dependent protein kinase phosphorylation site:

amino acids 266-269

N-myristoylation sites:

amino acids 27-32, 66-71, 91-96, 93-98, 102-107, 109-114, 140-145 and 212-217

276/392

FIGURE 265

GGAGAATGGAGAGAGCAGTGAGAGTGGAGTCCGGGGTCCTGGTCGGGGTGGTCTGTCTGC
TCCTGGCATGCCCTGCCACAGCCACTGGGCCCGAAGTTGCTCAGCCTGAAGTAGACACCA
CCCTGGGTCGTGTGCGAGGCCGGCAGGTGGGCGTGAAGGGCACAGACCGCCTTGTGAATG
TCTTTCTGGGCATTCCATTTGCCAGCCGCCACTGGGCCCTGACCGGTTCTCAGCCCCAC
ACCCAGCACAGCCCTGGGAGGGTGTGCGGGATGCCAGCACTGCGCCCCAATGTGCCTAC
AAGACGTGGAGAGCATGAACAGCAGCAGATTTGTCTCAACGGAAAACAGCAGATCTTCT
CCGTTTCAGAGGACTGCCTGGTCCTCAACGTCTATAGCCCAGCTGAGGTCCCCGAGGGT
CCGGTAGGCCGGTCATGGTATGGGTCCATGGAGGCGCTCTGATAACTGGCGCTGCCACCT
CCTACGATGGATCAGCTCTGGCTGCCCTATGGGGATGTGGTCGTGGTTACAGTCCAGTACC
GCCTTGGGGTCCTTGGCTTCTTCAGCACTGGAGATGAGCATGCACCTGGCAACCAGGGCT
TCCTAGATGTGGTAGCTGCTTTGCGCTGGGTGCAAGAAAACATCGCCCCCTTCGGGGGTG
ACCTCAACTGTGTCACTGTCTTTGGTGGATCTGCCGGTGGGAGCATCATCTCTGGCCTGG
TCCTGTCCCCAGTGGCTGCAGGGCTGTTCCACAGAGCCATCACACAGAGTGGGGTCATCA
CCACCCCAGGGATCATCGACTCTCACCCCTTGGCCCCTAGCTCAGAAAATCGCAAACACCT
TGGCCTGCAGCTCCAGCTCCCCGGCTGAGATGGTGCAGTGCCTTCAGCAGAAAAGAAGGAG
AAGAGCTGGTCCTTAGCAAGAAGCTGAAAAATACTATCTATCCTCTCACCGTTGATGGCA
CTGTCTTCCCCAAAAGCCCCAAGGAATCCTGAAGGAGAAGCCCTTCCACTCTGTGCCCT
TCCTCATGGGTGTCAACAACCATGAGTTTCAGCTGGCTCATCCCCAGGGGCTGGGGTCTCC
TGGATACAATGGAGCAGATGAGCCGGGAGGACATGCTGGCCATCTCAACACCCGTCTTGA
CCAGTCTGGATGTGCCCCCTGAGATGATGCCACCCGTATAGATGAATACCTAGGAAGCA
ACTCGGACGCACAAGCCAAATGCCAGGCGTTCAGGAATTCATGGGTGACGTATTTCATCA
ATGTTCCACCCGTCACTTTTTTCAAGATACCTTCGAGATTCTGGAAGCCCTGTCTTTTTCT
ATGAGTTCCAGCATCGACCCAGTTCTTTTGCAGAGATCAAACCTGCCTGGGTGAAGGCTG
ATCATGGGGCCGAGGGTGCTTTTTGTGTTTCGGAGGTCCCTTCCTCATGGACGAGAGCTCCC
GCCTGGCCTTTTCCAGAGGCCACAGAGGAGGAGAAGCAGCTAAGCCTCACCATGATGGCCC
AGTGGACCCACTTTTGGCCGGACAGGGGACCCCAATAGCAAGGCTCTGCCTCCTTGGCCCC
AATTCAACCAGGCGGAACAATATCTGGAGATCAACCCAGTGCCACGGGCCGGACAGAAGT
TCAGGGAGGCCTGGATGCAGTTCTGGTCAGAGACGCTCCCCAGCAAGATACAACAGTGGC
ACCAGAAGCAGAAGAACAGGAAGGCCCAGGAGGACCTCTGAGGGCCAGGCCTGAACCTTCT
TGGCTGGGGCAAACCACTCTTCAAGTGGTGGCAGAGTCCCAGCACGGCAGCCCGCCTCTC
CCCCTGCTGAGACTTTAATCTCCACCAGCCCTTAAAGTGTGCGCCGCTCTGTGACTGGAG
TTATGCTCTTTTTGAAATGTCACAAGGCCGCTCCACCTCTGGGGCATTGTACAAGTTCT
TCCCTCTCCCTGAAGTGCCTTTTCTGCTTCTTTCGTGGTAGGTTCTAGCACATTCCTCTA
GCTTCCCTGGAGGACTCACTCCCCAGGAAGCCTTCCCTGCCTTCTCTGGGCTGTGCGGCCC
CGAGTCTGCGTCCATTAGAGCACAGTCCACCCGAGGCTAGCACCGTGTCTGTGTCTGTCT
CCCCCTCAGAGGAGCTCTCTCAAAATGGGGATTAGCCTAACCCCACTCTGTCACCCCACAC
CAGGATCGGGTGGGACCTGGAGCTAGGGGGTGTGTTGCTGAGTGAGTGAGTGAAACACAGA
ATATGGGAATGGCAGCTGCTGAACCTTGAACCCAGAGCCTTCAGGTGCCAAAGCCATACTC
AGGCCCCCACCAGCATTGTCCACCCTGGCCAGAAGGGTGCATGCCAATGGCAGAGACCTG
GGATGGGAGAAGTCTTGGGGCGCCAGGGGATCCAGCCTAGAGCAGACCTTAGCCCCCTGAC
TAAGGCCTCAGACTAGGGCGGGAGGGGTCTCCTCCTCTCTGCTGCCAGTCTTGGCCCCCT
GCACAAGACAACAGAATCCATCAGGGCCATGAGTGTCAACCAGACCTGACCCTACCAAT
TCCAGCCCCCTGACCCTCAGGACGCTGGATGCCAGCTCCCAGCCCCAGTGCCGGGTCTCTC
CTCCCTTCTGGCTTGGGGAGACCAGTTTCTGGGGAGCTTCCAAGAGCACCCACCAAGAC
ACAGCAGGACAGGCCAGGGGAGGGCATCTGGACCAGGGCATCCGTGCGGCTATTGTCA
GAGAAAAGAAGAGACCCACCCACTCGGGCTGCAAAAGGTGAAAAGCACCAAGAGGTTTTT

277/392

AGATGGAAGTGAGAGGTGACAGTGTGCTGGCAGCCCTCACAGCCCTCGCTTGCTCTCCCT
GCCGCTCTGCCTGGGCTCCCACTTTGGCAGCACTTGAGGAGCCCTTCAACCCGCCGCTG
CACTGTAGGAGCCCCCTTCTGGGCTGGCCAAGGCCGGAGCCAGCTCCCTCAGCTTGCGGG
GAGGTGCGGAGGGAGAGGGGCGGGCAGGAACCGGGGCTGCGCGCAGCGCTTGCGGGCCAG
AGTGAGTTCGGGTGGGCGTGGGCTCGGCGGGGCCCCACTCAGAGCAGCTGGCCGGCCCC
AGGCAGTGAGGGCCTTAGCACCTGGGCCAGCAGCTGCTGTGCTCGATTTCTCGCTGGGCC
TTAGCTGCCTCCCCGCGGGCAGGGCTCGGGACCTGCAGCCCTCCATGCCTGACCCCTCC
CCCACCCCCCGTGGGCTCCTGTGCGGCCGGAGCCTCCCCAAGGAGCGCCGCCCTGCTC
CACAGCGCCCACTCCCATCGACCACCCAAGGGCTGAGGAGTGCGGGTGACAGCGCGGGA
CTGGCAGGCAGCTCCACCTGCTGCCCCAGTGCTGGATCCACTGGGTGAAGCCAGCTGGGC
TCCTGAGTCTGGTGGGACTTGGAGAACCTTTATGTCTAGCTAAGGGATTGTAAATACAC
CGATGGGCACTCTGTATCTAGCTCAAGGTTTGTAAACACACCAATCAGCACCCCTGTGTCT
AGCTCAGTGTTTGTGAATGCACCAATCCACACTCTGTATCTGGCTACTCTGGTGGGGACT
TGGAGAACCTTTGTGTCCACACTCTGTATCTAGCTAATCTAGTGGGGATGTGGAGAACCT
TTGTGTCTAGCTCAGGGATCGTAAACGCACCAATCAGCACCCCTGTCAAAACAGACCACTT
GACTCTCTGTAAATGGACCAATCAGCAGGATGTGGGTGGGGCGAGACAAGAGAATAAAA
GCAGGCTGCCTGAGCCAGCAGTGACAACCCCCCTCGGGTCCCCTCCCACGCCGTGGAAGC
TTTGTTCTTTCGCTCTTTGCAATAAATCTTGCTACTGCCCAAAA

278/392

FIGURE 266

MERAVRVESGVLVGVVCLLLACPATATGPEVAQPEVDTTLGRVVRGRQVGVKGTDLNVNF
LGIPFAQPPLGPDRFSAPHPAQPWEGVRDASTAPPMCLQDVESMNSSRFVLNGKQQIFSV
SEDCLVLNVYSPAIEVPAGSGRPVMVWVHGGALITGAATSYDGSALAAYGDVVVVTVQYRL
GVLGFFSTGDEHAPGNQGFLLDVVAALRWVQENIAPFGGDLNLCVTVFGGSAGGSIISGLVL
SPVAAGLFHRAITQSGVITTPGIIDSHPWPLAQKIANLTLACSSSSPAEMVQCLQQKEGEE
LVLSKKLKNTIYPLTVDGTVFPKSPKELLKEKPFHSVPFLMGVNNHEFSWLI PRGWGLLD
TMEQMSREDMLAISTPVLTSLDVPEMMPTVIDEYLGSNSDAQAKCQAFQEFMGDVFINV
PTVSFSRYLRDSGSPVFFYEFQHRPSSF AKIKPAWVKADHGAEGAFVFGGPFLMDESSL
AFPEATEEEKQLSLTMMAQWTHFARTGDPNSKALPPWPQFNQAEQYLEINPVPRAGQKFR
EAWMQFWSETLPSKIQQWHQKQKNRKAQEDL

Important features of the protein:**Signal peptide:**

amino acids 1-27

Transmembrane domain:

amino acids 226-245

N-glycosylation site:

amino acids 105-109

N-myristoylation sites:amino acids 10-16, 49-55, 62-68, 86-92, 150-156, 155-161,
162-168, 217-223, 227-233, 228-234, 232-238, 262-268, 357-363,
461-467**Prokaryotic membrane lipoprotein lipid attachment site:**

amino acids 12-23

Carboxylesterases type-B serine active site:

amino acids 216-232

279/392

FIGURE 267

TGTCGCCTGGCCCTCGCCATGCAGACCCCGCGAGCGTCCCCTCCCCGCCCCGGCCCTCCTG
CTTCTGCTGCTGCTACTGGGGGGCGCCACGGCCTCTTTCCTGAGGAGCCGCCGCCGCTT
AGCGTGGCCCCCAGGGACTACCTGAACCACTATCCCGTGTTTGTGGGCAGCGGGCCCGGA
CGCCTGACCCCCGAGAAGGTGCTGACGACCTCAACATCCAGCGAGTCTTGCGGGTCAAC
AGGACGCTGTTTATTGGGGACAGGGACAACCTTACC CGTAGAGCTGGAGCCCCCACG
TCCACGGAGCTGCGGTACCAGAGGAAGCTGACCTGGAGATCTAACCCAGCGACATAAAC
GTGTGTGCGATGAAGGGCAAACAGGAGGGCGAGTGTGAAACTTCGTAAAGGTGCTGCTC
CTTCGGGACGAGTCCACGCTCTTTGTGTGCGGTTCCAACGCCTTCAACCCGGTGTGCGCC
AACTACAGCATAGACACCCTGCAGCCCGTCGGAGACAACATCAGCGGTATGGCCCGCTGC
CCGTACGACCCCAAGCACGCCAATGTTGCCCTCTTCTCTGACGGGATGCTCTTACAGCT
ACTGTTACCGACTTCTAGCCATTGATGCTGTCTATACC CGCAGCCTCGGGGACAGGCC
ACCCTGCGCACCGTGAAACATGACTCCAAGTGGTTCAAAGAGCCTTACTTTGTCCATGCG
GTGGAGTGGGGCAGCCATGTCTACTTCTTCTTCCGGGAGATTGCGATGGAGTTAACTAC
CTGGAGAAGGTGGTGGTGTCCCGCGTGGCCCGAGTGTGCAAGAACGACGTGGGAGGCTCC
CCCCGCGTGCTGGAGAAGCAGTGGACGTCCTTCTGAAGGCGCGGCTCAACTGCTCTGTA
CCCGGAGACTCCCATTTCTACTTCAACGTGCTGCAGGCTGTCACGGGCGTGGTCAGCCTC
GGGGGCCCGCCCGTGGTCTGGCCGTTTTTTCCACGCCCAGCAACAGCATCCCTGGCTCG
GCTGTCTGCGCCTTTGACCTGACACAGGTGGCAGCTGTGTTTGAAGGCCGCTTCCGAGAG
CAGAAGTCCCCGAGTCCATCTGGACGCCGGTGCCGGAGGATCAGGTGCCTCGACCCCGG
CCCGGGTGCTGCGCAGCCCCCGGGATGCAGTACAATGCCTCCAGCGCCTTGCCGGATGAC
ATCCTCAACTTTGTCAAGACCCACCCTCTGATGGACGAGGCGGTGCCCTCGCTGGGCCAT
GCGCCCTGGATCCTGCGGACCCTGATGAGGCACCAGCTGACTCGAGTGGCTGTGGACGTG
GGAGCCGGCCCCCTGGGGCAACCAGACCCTTGTCTTCTGGGTTCTGAGGCGGGGACGGTC
CTCAAGTTCCTCGTCCGGCCCCAATGCCAGCACCTCAGGGACGTCTGGGCTCAGTGTCTTC
CTGGAGGAGTTTGAGACCTACCGGCCGGACAGGTGTGGACGGCCCCGGCGGTGGCGAGACA
GGGCAGCGGCTGTGAGCTTGGAGCTGGACGAGCTTCGGGGGGCCTGCTGGCTGCCTTC
CCCCGCTGCGTGGTCCGAGTGCTGTGGCTCGCTGCCAGCAGTACTCGGGGTATGAAG
AACTGTATCGGCAGTCAGGACCCCTACTGCGGGTGGGCCCCCGACGGCTCCTGCATCTTC
CTCAGCCCCGGGCACCAGAGCCGCTTTTGAGCAGGACGTGTCCGGGGCCAGCACCTCAGGC
TTAGGGGACTGCACAGGACTCCTGCGGGCCAGCCTCTCCGAGGACCGCGCGGGGCTGGTG
TCGGTGAACCTGCTGTTAACGTGTCGGTGGCGGCCTTCGTGGTGGGAGCCGTGGTGTCC
GGCTTCAGCGTGGGCTGGTTCGTGGGCCCTCCGTGAGCGGCGGGAGCTGGCCCGGCGCAAG
GACAAGGAGGCCATCCTGGCGCACGGGGCGGGCGAGGCGGTGCTGAGCGTCAGCCGCTG
GGCGAGCGCAGGGCGCAGGGTCCCGGGGGCGGGGCGGAGGCGGTGGCGGTGGCGCCGGG
GTTCCCCCGGAGGCCCTGCTGGCGCCCCCTGATGCAGAACGGCTGGGCCAAGGCCACGCTG
CTGCAGGGCGGGCCCCACGACCTGGACTCGGGGCTGCTGCCACGCCCGAGCAGACGCCG
CTGCCGCGAGAAGCGCCTGCCCACTCCGCACCCGCACCCCCACGCCCTGGGCCCCCGCGCC
TGGGACCACGGCCACCCCTGCTCCCGCCTCCGCTTCATCCTCCCTCCTGCTGCTGGCG
CCCGCCCGGGCCCCCGAGCAGCCCCCGCGCCTGGGGAGCCGACCCCCGAGGCCGCGCTC
TATGCTGCCCCGGCCCCGGCGCGCCTCCACGGCGACTTCCCGCTCACCCCCACGCCAGC
CCGGACCGCCGGCGGGTGGTGTCCGCGCCACGGGCCCCCTTGACCCAGCCTCAGCCGCC
GATGGCCTCCCGCGGCCCTGGAGCCCCCGCCCGACGGGCAGCCTGAGGAGGCCACTGGGC
CCCCACGCCCTCCGGCCGCCACCCTGCGCCGCACCCACACGTTCAACAGCGGCGAGGCC
CGGCCTGGGGACCGCCACCGCGGCTGCCACGCCCCGGCCGGGCACAGACTTGGCCACCTC
CTCCCCATATGGGGGGGCGGACAGGACTGCGCCCCCGTGCCCTAGGCCGGGGGCCCCCG
ATGCCTTGGCAGTGCCAGCCACGGGAACAGGAGCGAGAGACGGTGCCAGAACGCCGGGG
CCCCGGGCAACTCCGAGTGGGTGCTCAAGTCCCCCCCCGCGACCCACCCGCGAGTGGGGG

280/392

GCCCCCTCCGCCACAAGGAAGCACAAACCAGCTCGCCCTCCCCCTACCCGGGGCCGCAGGA
CGCTGAGACGGTTTGGGGGTGGGTGGGCGGGAGGACTTTGCTATGGATTTGAGGTTGACC
TTATGCGCGTAGGTTTTGGTTTTTTTTTGCAGTTTTGGTTTCTTTTGCGGTTTTCTAACC
AATTGCACAACCTCCGTTCTCGGGGTGGCGGCAGGCAGGGGAGGCTTGGACGCCGGTGGGG
AATGGGGGGCCACAGCTGCAGACCTAAGCCCTCCCCACCCCTGGAAAGGTCCCTCCCCA
ACCCAGGCCCCCTGGCGTGTGTGGGTGTGCGTGCGTGTGCGTGCCGTGTTTCGTGTGCAAGG
GGCCGGGGAGGTGGGCGTGTGTGTGCGTGCCAGCGAAGGCTGCTGTGGGCGTGTGTGTCA
AGTGGGCCACGCGTGCAGGGTGTGTGTCCACGAGCGACGATCGTGGTGGCCCCAGCGGCC
TGGGCGTTGGCTGAGCCGACGCTGGGGCTTCCAGAAGGCCCGGGGGTCTCCGAGGTGCCG
GTTAGGAGTTTGAACCCCCCCCCACTCTGCAGAGGGAAGCGGGGACAATGCCGGGGTTTCA
GGCAGGAGACACGAGGAGGGCCTGCCCCGAAGTCACATCGGCAGCAGCTGTCTAAAGGGC
TTGGGGGCCTGGGGGGCGGCGAAGGTGGGTGGGGCCCCCTCTGTAAATACGGCCCCAGGGT
GGTGAGAGAGTCCCATGCCACCCGTCCCCTTGTGACCTCCCCCTATGACCTCCAGCTGA
CCATGCATGCCACGTGGCTGGCTGGGTCTCTGCCCTCTTTGGAGTTTGCCTCCCCCAGC
CCCCCTCCCCATCAATAAACTCTGTTTACAACCAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

281/392

FIGURE 268

MQTPRASPPRPALLLLLLLLLLGGAHGLFPPEPPPLSVAPRDYLNHYPVFVSGSGPGRLTPAE
GADDLNIQVLRVNRTLFIGDRDNLRYVELEPPPTSTELRYQRKLTWRSNPSDINVCRMKG
KQEGECRNFVKVLLLRDESTLFVCGSNAFNPVCANYSIDTLQPVGDNISGMARCPYDPKH
ANVALFSDGMLFTATVTDFLAIDAVIYRSLGDRPTLRTVKHDSKWFKEPYFVHAVEWGSH
VYFFFREIAMEFNYLEKVVVSRVARVCKNDVGGSPRVLEKQWTSFLKARLNCSVPGDSHF
YFNVLQAVTGVVSLGGRPVLAVFSTPSNSIPGSAVCAFDLTQVAAVFEGRFREQKSPES
IWTPVPEDQVPRPRPGCCAAPGMQYNASSALPDDILNFVKTHPLMDEAVPSLGHAPWILR
TLMRHQLTRVAVDVGAGPWGNQTVVFLGSEAGTVLKFLVRPNASTSGTSGLSVFLEEFET
YRPDRCGRPGGGETGQRLLSLELDAASGGLLAAPRCVVRVPVARCQQYSGCMKNCIGSQ
DPYCGWAPDGCIFLSPGTRAAFEQDVSGASTSGLGDCGTGLLRASLSEDRAGLVSVNLLV
TSSVAAFVVGAVVSGFSVGVFVGLRERRELARRKDKEAILAHGAGEAVLSVSRLEGERRAQ
GPGGRGGGGGGGAGVPPEALLAPLMQNGWAKATLLQGGPHDLDSGLLPTPEQTPLPQKRL
PTPHPHPHALGPRAWDHGHPLLPASASSSLLLLAPARAPEQPPAPGEPTPDGRLYAARPG
RASHGDFPLTPHASPDRRRVVSAPTGPLDPASAADGLPRPWSPPTGSLRRPLGPHAPPA
ATLRRTHTFNSGEARPGDRHRGCHARPGTDLAHLPLPYGGADRTAPPVP

Important features of the protein:

Signal peptide:

amino acids 1-25

Transmembrane domains:

amino acids 318-339, 598-617

N-glycosylation sites.

amino acids 74-78, 155-159, 167-171, 291-295, 386-390,
441-445, 462-466

Glycosaminoglycan attachment sites:

amino acids 51-55, 573-577

cAMP- and cGMP-dependent protein kinase phosphorylation site:

amino acids 102-106

N-myristoylation sites:

amino acids 21-27, 50-56, 189-195, 333-339, 382-388, 448-454,
490-496, 491-497, 508-514, 509-515, 531-537, 558-564, 569-575,
574-580, 580-586, 610-616, 643-649, 663-669, 666-672, 667-673,
668-674, 669-675, 670-676, 868-874, 879-885

282/392

FIGURE 269

ATCTGAGTGAGCTAACTGACACAATGAACTGTCAGGCATGTTTCTGCTCCTCTCTCTGG
CTCTTTTCTGCTTTTAAACAGGTGTCTTCAGTCAGGGAGGACAGGTTGACTGTGGTGAGT
TCCAGGACCCCAAGGTCTACTGCACTCGGGAATCTAACCCACACTGTGGCTCTGATGGCC
AGACATATGGCAATAAATGTGCCTTCTGTAAGGCCATAGTGAAAAGTGGTGGAAAGATTA
GCCTAAAGCATCCTGGAAAATGCTGAGTAAAGCCAATGTTTCTTGGTGACTTGCCAGCT
TTTGCAGCCTTCTTTTCTCACTTCTGCTTATACTTTTGCTGGTGGATTCCTTTAATTCAT
AAAGACATACCTACTCTGCCTGGGTCTTGAGGAGTTCAATGTATGTCTATTTCTCTTGAT
TCACTTGTCAATAAAGTACATTCTGCAAAAGCAAAA

283/392

FIGURE 270

MKLSGMFLLLSLALFCFLTGVFSQGGQVDCGEFQDPKVYCTRESNPHCGSDGQTYGNKCA
FCKAIVKSGGKISLKHPGKC

Important features of the protein:

Signal peptide:

amino acids 1-23

N-myristoylation sites:

amino acids 26-32, 52-58, 56-62, 69-75

Kazal serine protease inhibitors family signature:

amino acids 40-63

284/392

FIGURE 271

AACTTCTACATGGGCCTCCTGCTGCTGGTGCTCTTCCTCAGCCTCCTGCCGGTGGCCTAC
ACCATCATGTCCCTCCCACCCTCCTTTGACTGCGGGCCGTTTCAGGTGCAGAGTCTCAGTT
GCCCCGGAGCACCTCCCCTCCCGAGGCAGTCTGCTCAGAGGGCCTCGGCCCAGAATTCCA
GTTCTGGTTTCATGCCAGCCTGTAAAAGGCCATGGAACTTTGGGTGAATCACCGATGCCA
TTTAAGAGGGTTTTCTGCCAGGATGGAAATGTTAGGTCGTTCTGTGTCTGCGCTGTTTCAT
TTCAGTAGCCACCAGCCACCTGTGGCCGTTGAGTGCTTGAAATGAGGAACTGAGAAAATT
AATTTCTCATGTATTTTTCTCATTATTTATTTAATTTTTAACTGATAGTTGTACATATTT
GGGGGTACATGTGATATTTGGATACATGTATACAATATATAATGATCAAATCAGGGTAAC
TGGGATATCCATCACATCAAACATTTATTTTTTATTCTTTTATAGACAGAGTCTCACTCTG
TCACCCAGGCTGGAGTGCAGTGGTGCCATCTCAGCTTACTGCAACCTCTGCCTGCCAGGT
TCAAGCGATTCTCATGCCTCCACCTCCCAAGTAGCTGGGACTACAGGCATGCACCACAAT
GCCCAACTAATTTTTGTATTTTTAGTAGAGACGGGGTTTTGCCATGTTGCCAGGCTGGC
CTTGAACTCCTGGCCTCAAACAATCCACTTGCCTCGGCCTCCCAAAGTGTTATGATTACA
GGCGTGAGCCACCGTGCCTGGCCTAAACATTTATCTTTTCTTTGTGTTGGGAACTTTGAA
ATTATACAATGAATTATTGTAACTGTCATCTCCCTGCTGTGCTATGGAACACTGGGACT
TCTTCCCTCTATCTAACTGTATATTTGTACCAGTTAACCAACCGTACTTCATCCCCACTC
CTCTCTATCCTTCCCAACCTCTGATCACCTCATTTACTCTCTACCTCCATGAGATCCAC
TTTTTTTAGCTCCCATGTGAGTAAGAAAATGCAATATTTGTCTTTCTGTGCCTGGCTTA
TTTCACTTAACATAATGACTTCCTGTTCCATCCATGTTGCTGCAAATGACAGGATTTTCGT
TCTTAATTTCAATTAAAATAACCACACATGGCAAAAA

285/392

FIGURE 272

MGLLLLVLFLSLLPVAYTIMSLPPSFDCGPFRCRVSVAREHLPSRGSLLRGPRPRIPVLV
SCQPVKGHGTLGESPMFPKRVFCQDGNVRSFCVCAVHFSSHQPPVAVECLK

Important features of the protein:

Signal peptide:

amino acids 1-18

N-myristoylation site:

amino acids 86-92

Zinc carboxypeptidases, zinc-binding region 2 signature:

amino acids 68-79

286/392

FIGURE 273

TTCTGAAGTAACGGAAGCTACCTTGTATAAAGACCTCAACACTGCTGACCATGATCAGCG
CAGCCTGGAGCATCTTCCTCATCGGGACTAAAATTGGGCTGTTCCCTTCAAGTAGCACCTC
TATCAGTTATGGCTAAATCCTGTCCATCTGTGTGTCGCTGCGATGCGGGTTTCATTTACT
GTAATGATCGCTTTCTGACATCCATTCCAACAGGAATACCAGAGGATGCTACAACCTCTCT
ACCTTCAGAACAACCAAATAAATAATGCTGGGATTCCCTTCAGATTTGAAAACTTGCTGA
AAGTAGAAAGAATATACCTATAACCACAACAGTTTAGATGAATTTCCCTACCAACCTCCCAA
AGTATGTAAAAGAGTTACATTTGCAAGAAAATAACATAAGGACTATCACTTATGATTAC
TTTCAAAAATTCCTTATCTGGAAGAATTACATTTAGATGACAACTCTGTCTCTGCAGTTA
GCATAGAAGAGGGAGCATTCGAGACAGCAACTATCTCCGACTGCTTTTCCTGTCCCGTA
ATCACCTTAGCACAAATTCCTGCGGTTTGCCAGGACTATAGAAGAACTACGTTGGATG
ATAATCGCATATCCACTATTTTCATCACCATCTCTTCAAGGTCTCACTAGTCTAAAACGCC
TGGTTCTAGATGGAAACCTGTTGAACAATCATGGTTTAGGTGACAAAGTTTCTTCAACC
TAGTTAATTTGACAGAGCTGTCCCTGGTGCGGAATTCCCTGACTGCTGCACCAGTAAACC
TTCCAGGCACAAACCTGAGGAAGCTTTATCTTCAAGATAACCACATCAATCGGGTGCCCC
CAAATGCTTTTTCTTATCTAAGGCAGCTCTATCGACTGGATATGTCCAATAATAACCTAA
GTAATTTACCTCAGGGTATCTTTGATGATTTGGACAATATAACACAACCTGATTCTTCGCA
ACAATCCCTGGTATTGCGGGTGCAAGATGAAATGGGTACGTGACTGGTTACAATCACTAC
CTGTGAAGGTCAACGTGCGTGGGCTCATGTGCCAAGCCCCAGAAAAGGTTTCGTGGGATGG
CTATTAAGGATCTCAATGCAGAACTGTTTGATTGTAAGGACAGTGGGATTGTAAGCAGCA
TTCAGATAACCACTGCAATACCCAACACAGTGTATCCTGCCCAAGGACAGTGGCCAGCTC
CAGTGACCAAACAGCCAGATATTAAGAACCCCAAGCTCACTAAGGATCAACAAACCACAG
GGAGTCCCTCAAGAAAAACAATTACAATTACTGTGAAGTCTGTACCTCTGATACCATTCT
ATATCTCTTGGAACCTTGCTCTACCTATGACTGCTTTGAGACTCAGCTGGCTTAAACTGG
GCCATAGCCCGGCATTTGGATCTATAACAGAAACAATTGTAACAGGGGAACGCAGTGAGT
ACTTGGTCACAGCCCTGGAGCCTGATTCACCCCTATAAAGTATGCATGGTTCCCATGGAAA
CCAGCAACCTCTACCTATTTGATGAAACTCCTGTTTGTATTGAGACTGAACTGCACCCC
TTCGAATGTACAACCCTACAACCACCCCTCAATCGAGAGCAAGAGAAAGAACCTTACAAA
ACCCCAATTTACCTTTGGCTGCCATCATTTGGTGGGGCTGTGGCCCTGGTTACCATTGCCC
TTCTTGCTTTAGTGTGTGGTATGTTTCATAGGAATGGATCGCTCTTCTCAAGGAACCTGTG
CATATAGCAAAGGGAGGAGAAGAAAGGATGACTATGCAGAAGCTGGCACTAAGAAGGACA
ACTCTATCCTGGAAATCAGGGAAACTTCTTTTCAGATGTTACCAATAAGCAATGAACCCA
TCTCGAAGGAGGAGTTTGTAAATACACACCATATTTCCCTCCTAATGGAATGAATCTGTACA
AAAACAATCACAGTGAAAGCAGTAGTAACCGAAGCTACAGAGACAGTGGTATTCCAGACT
CAGATCACTCACACTCATGATGCTGAAGGACTCACAGCAGACTTGTGTTTTGGGTTTTTTT
AAACCTAAGGGAGGTGATGGT

287/392

FIGURE 274

MISAAWSIFLIGTKIGLFLQVAPLSVMAKSCPSVCRC DAGFIYCND RFLTSIPTGIPEDA
TTLYLQNNQINNAGIPSDLKNLLKVERIYLYHNSLDEFPTNL PKYVKELHLQENNI RTIT
YDSLSKIPYLEELHDDNSVSAVSIIEGAFRDSNYLRLLFLSRNHLSTIPWGLPRTIEEL
RLDDNRISTISSPSLQGLTSLKRLVLDGNLLNNHGLGDKVFFNLVNLTELSLVRNSLTAA
PVNLPGTNLRKLYLQDNHINRVPPNAFSYLRLQLYRLDMSNNLSNLPQGIFDDL DITQL
ILRNNPWYCGCKMKWVRDWLQSLPVKVNVRGLMCQAPEKVRGMAIKDLNAELFDCKDSGI
VSTIQITTAIPNTVYPAQGQWPAPVTKQPDIKNPKLTKDQQTGSPSRKTITITVKS VTS
DTIHISWKLALPMTALRLSWLKLGHSPAFGSITETIVTGERSEYLVTALEPDSPYKVC MV
PMETS NLYLFDETPVC IETETAPLRMYNPTTTLNREQEKEPYKNPNLPLAAIIGGA VALV
TIAL LALVCWYVHRNGSLFSRNCAYS KGRRRKDDYAEAGTKKDNSILEIRETSFQMLPIS
NEPISKEEFVIHTIFPPNGMNL YKNNHSESSSNRSYRDSGIPDS DSHSHS

Important features of the protein:**Signal peptide:**

amino acids 1-28

Transmembrane domain:

amino acids 531-552

N-glycosylation sites:

amino acids 226-229, 282-285, 296-299, 555-558, 626-629, 633-636

Tyrosine kinase phosphorylation site:

amino acids 515-522

N-myristoylation sites:

amino acids 12-17, 172-177, 208-213, 359-364, 534-539, 556-561, 640-645

Amidation site:

amino acids 567-570

Leucine zipper pattern:

amino acids 159-180

Phospholipase A2 aspartic acid active site:

amino acids 34-44

288/392

FIGURE 275

AGGGCCCCGCGGGTGGAGAGAGCGACGCCCCGAGGGGATGGCGGCAGCGTCCCCGAGCGCCT
CTGGCTGGGCGCTACTGCTGCTGGTGGCACTTTGGCAGCAGCGCGCGGCCGCTCCGGCG
TCTTCCAGCTGCAGCTGCAGGAGTTTCATCAACGAGCGCGGCGTACTGGCCAGTGGGCGGC
CTTGCGAGCCCCGGCTGCCGACTTTCTTCCGCGTCTGCCTTAAGCACTTCCAGGCGGTG
TCTCGCCCCGACCTGCACCTTCGGGACCGTCTCCACGCCGGTATTGGGCACCAACTCCT
TCGCTGTCCGGGACGACAGTAGCGGCGGGGGCGCAACCTCTCCAAGTGCCTTCAATT
TCACCTGGCCGGGTACCTTCTCGCTCATCATCGAAGCTTGGCACGCGCCAGGAGACGACC
TGCGGCCAGAGGCCTTGCCACCAGATGCACTCATCAGCAAGATCGCCATCCAGGGCTCCC
TAGCTGTGGGTGAGAACTGGTTATTGGATGAGCAAACCAGCACCTCACAAGGCTGCGCT
ACTCTTACCGGGTCATCTGCAGTGACAATACTATGGAGACAAGTCTCCCGCTGTGCA
AGAAGCCCAATGACCACTTCGGCCACTATGTGTGCCAGCCAGATGGCAACTGTCTGCTCC
TGCCCCGTTGGACTGGGGAATATTGCCAACCCCTATCTGTCTTTTCGGGCTGTGATGAAC
AGAATGGCTACTGCAGCAAGCCAGCAGAGTGCCTCTGCCGCCAGGCTGGCAGGGCCGGC
TGTGTAACGAATGCATCCCCACAATGGCTGTGCGCCACGGCACCTGCAGCACTCCCTGGC
AATGTACTTGTGATGAGGGCTGGGGAGGCCTGTTTTGTGACCAAGATCTCAACTACTGCA
CCCACCACTCCCCATGCAAGAATGGGGCAACGTGCTCCAACAGTGGGCAGCGAAGCTACA
CCTGCACCTGTGCGCCAGGCTACACTGGTGTGGACTGTGAGCTGGAGCTCAGCGAGTGTG
ACAGCAACCCCTGTGCAATGGAGGCAGCTGTAAGGACCAGGAGGATGGCTACCACTGCC
TGTGTCTCCGGGCTACTATGGCCTGCAGTGTGAACACAGCACCTTGAGCTGCGCCGACT
CCCCCTGCTTCAATGGGGGCTCCTGCCGGGAGCGCAACCAGGGGGCCAAGTATGCTTGTG
AATGTCCCCCAACTTCACCGGCTCCAAGTGCAGAGAAGAAAGTGGACAGGTGCACCAGCA
ACCCCTGTGCCAACGGGGACAGTGCCTGAACCGAGGTCCAAGCCGCATGTGCCGCTGCC
GTCCTGGATTACAGGGCACCTACTGTGAAGTCCACGTCAGCGACTGTGCCCCGTAACCTT
GCGCCACGGTGGCACTTGCCATGACCTGGAGAATGGGCTCATGTGCACCTGCCCTGCCG
GCTTCTCTGGCCGACGCTGTGAGGTGCGGACATCCATCGATGCCTGTGCCTCGAGTCCCT
GCTTCAACAGGGCCACCTGCTACACCGACCTCTCCACAGACACCTTTGTGTGCAACTGCC
CTTATGGCTTTGTGGGCAGCCGCTGCGAGTTCCCCGTGGGCTTGCCGCCAGCTTCCCCCT
GGGTGGCCGTCTCGCTGGGTGTGGGGCTGGCAGTGTGCTGGTACTGCTGGGCATGGTGG
CAGTGGCTGTGCGGCAGCTGCGGCTTCGACGGCCGGACGACGGCAGCAGGGAAGCCATGA
ACAACTTGTGCGACTTCCAGAAGGACAACCTGATTCCTGCGCCAGCTTAAAAACACAA
ACCAGAAGAAGGAGCTGGAAGTGGACTGTGGCCTGGACAAGTCCAAGTGTGGCAAACAGC
AAAACCACACATTGGACTATAATCTGGCCCCAGGGCCCCCTGGGGCGGGGACCATGCCAG
GAAAGTTTCCCCACAGTGACAAGAGCTTAGGAGAGAAGGCGCCACTGCGGTTACACAGTG
AAAAGCCAGAGTGTGCGATATCAGCGATATGCTCCCCAGGGACTCCATGTACCAGTCTG
TGTGTTTGATATCAGAGGAGAGGAATGAATGTGTGTCATTGCCACGGAGGTATAAGGCAGGA
GCCTACCTGGACATCCCTGCTCAGCCCCGCGGCTGGACCTTCCTTCTGCATTGTTTACA

289/392

FIGURE 276

MAAASRSASGWALLLLVALWQORAAGSGVFQLQLQEFINERGVLASGRPCEPGCRTLFFRV
CLKHFQAVVSPGPCTFGTVSTPVLGTNSFAVRDSSGGGRNPLQLPFNFTWPGTFSLIIE
AWHAPGDDLRPEALPPDALISKIAIQGSLAVGQNWLLDEQTSTLTRLRYSYRVICSDNY
GDNC SRLCKKRNDHFHGYVCQPDGNLSCLPGWTGEYCQQPICLSGCHEQNGYCSKPAECL
CRPGWQGRLCNECIPHNGCRHGTCTSPWQCTCDEGWGGLFCDQDLNYCTHHSPCKNGATC
SNSGQRSYTCTCRPGYTGVDCELELSECDSPNCRNGGSKDQEDGYHCLCPPGYGLHCE
HSTLSCADSPCFNGGSCRERNQGANACECPNFTGSNCEKKVDRCTSNPCANGGQCLNR
GPSRMCRCRPGFTGTCELVSDCARNPCAAGGTCHDLENGLMCTCPAGFSGRRCEVRTS
IDACASSPCFN RATCYTDLSTDTFVCNCPYGFVGSRCFVPVGLPPSFPWVAVSLGVGLAV
LLVLLGMVAVAVRQLRLRRPDDGSREAMNNLSDFQKDNLI PAAQLKNTNQKKELEVDCGL
DKSNC GKQQNHTLDYNLAPGPLGRGTM PGKFPHSDKSLGEKAPLRLHSEKPECRISAICS
PRDSMYQSVCLISEERNECVIATEV

Important features of the protein:**Signal peptide:**

amino acids 1-26

Transmembrane domain:

amino acids 530-552

N-glycosylation sites:amino acids 108-112, 183-187, 205-209, 393-397, 570-574,
610-614**Glycosaminoglycan attachment site:**

amino acids 96-100

Tyrosine kinase phosphorylation site:

amino acids 340-347

N-myristoylation sites:amino acids 42-48, 204-210, 258-264, 277-283, 297-303,
383-389, 415-421, 461-467, 522-528, 535-541, 563-569,
599-605, 625-631**Amidation site:**

amino acids 471-475

Aspartic acid and asparagine hydroxylation site:

amino acids 339-351

EGF-like domain cysteine pattern signature:amino acids 173-185, 206-218, 239-251, 270-282, 310-322,
348-360, 388-400, 426-438, 464-476, 506-518**Calcium-binding EGF-like:**

amino acids 224-245, 255-276, 295-316, 333-354, 373-394,

290/392

FIGURE 277

GGCAGTGCAGCCGCCTCACAGGTGCGCGGACGGGCCAGGCGGGCGGCCTCCTGAACCGAA
CCGAATCGGCTCCTCGGGCCGTGCTCCTCCCGCCCCCTCCTCGCCCGCCGCGGAGTTTTC
TTTCGGTTTCTTCCAAGATTCTTGGCCTTCCCTCGACGGAGCCGGGCCCAGTGCAGGGGGC
GCAGGGCGCGGGAGCTCCACCTCCTCGGCTTTCCCTGCGTCCAGAGGCTGGCATGGCGCG
GGCCGAGTACTGAGCGCACGGTCGGGGCACAGCAGGGCCGGGGGTGCAGCTGGCTCGCG
CCTCCTCTCCGGCCGCGTCTCCTCCGGTCCCTGGCGAAAGCCATTGAGACACCAGCTGG
ACGTCACGCGCCGGAGCATGTCTGGGAGTCAGAGCGAGGTGGCTCCATCCCCGCAGAGTC
CGCGGAGCCCCGAGATGGGACGGGACTTGCGGGCCGGGTCCCGCGTGCTCCTGCTCCTGC
TTCTGCTCCTGCTGGTGTACCTGACTCAGCCAGGCAATGGCAACGAGGGCAGCGTCACTG
GAAGTTGTTATTGTGGTAAAAGAATTTCTTCCGACTCCCCGCCATCGGTTCACTTCATGA
ATCGTCTCCGGAAACACCTGAGAGCTTACCATCGGTGTCTATACTACACGAGGTTCCAGC
TCCTTTCTGAGCGTGTGTGGGGGCAACAAGGACCCATGGGTTCAAGGAATTGATGAGCT
GTCTTGATCTCAAAGAATGTGGACATGCTTACTCGGGGATTGTGGCCCACCAGAAGCATT
TACTTCCTACCAGCCCCCAATTTCTCAGGCCTCAGAGGGGGCATCTTCAGATATCCACA
CCCCTGCCCAGATGCTCCTGTCCACCTTGCACTCAGCGCCCCACCCTCCCAGTAG
GATCACTGTCCTCGGACAAAGAGCTCACTCGTCCCAATGAAACCACCATTCACTGCGG
GCCACAGTCTGGCAGCTGGGCCTGAGGCTGGGGAGAACCAGAAGCAGCCGGAAAAAATG
CTGGTCCCACAGCCAGGACATCAGCCACAGTGCCAGTCCTGTGCCTCCTGGCCATCATCT
TCATCCTCACCGCAGCCCTTTCTATGTGCTGTGCAAGAGGAGGAGGGGGCAGTCACCGC
AGTCCTCTCCAGATCTGCCGGTTCATTATATACCTGTGGCACCTGACTCTAATACCTGAG
CCAAGAATGGAAGCTTGTGAGGGTAACTGTGGCTTATTCTTACAAAAGTGAATAAAG
GAGACTGACCCCTGACAACATGGTAGGCACTGTAAAAA

291/392

FIGURE 278

MGRDLRPGSRVLLLLLLLLLVYLTQPGNGNEGSVTGSCYCGKRISSDSPPSVQFMNRLRK
HLRAYHRCLYYTRFQLLSWSVCGGNKDPWVQELMSCLDLKECGHAYSGIVAHQKHLPTS
PPISQASEGASSDIHTPAQMLLSTLQSTQRPTLPVGSLSDDKELTRPNETTIHTAGHSLA
AGPEAGENQKQPEKNAGPTARTSATVPVLCLLAIFILTAALSYVLCKRRRGQSPQSSPD
LPVHYIPVAPDSNT

Important features of the protein:

Signal peptide:

1-26

Transmembrane domain:

204-223

N-glycosylation site:

168-172

cAMP- and cGMP-dependent protein kinase phosphorylation site:

42-46

N-myristoylation site:

29-35, 32-38, 36-42, 156-162

Amidation site:

40-44

292/392

FIGURE 279

CGCGAGGCGCGGGGAGCCTGGGACCAGGAGCGAGAGCCGCTACCTGCAGCCGCCGCCCA
CGGCACGGCAGCCACCATGGCGCTCCTGCTGTGCTTCGTGCTCCTGTGCGGAGTAGTGGA
TTTCGCCCAGAAAGTTTGAGTATCACTACTCCTGAAGAGATGATTGAAAAAGCCAAAGGGGA
AACTGCCTATCTGCCATGCAAATTTACGCTTAGTCCCGAAGACCAGGGACCGCTGGACAT
CGAGTGGCTGATATCACCAGCTGATAATCAGAAGGTGGATCAAGTGATTATTTTATATTC
TGGAGACAAAATTTATGATGACTACTATCCAGATCTGAAAGGCCGAGTACATTTTACGAG
TAATGATCTCAAATCTGGTGATGCATCAATAAATGTAACGAATTTACAACGTGCAGATAT
TGGCACATATCAGTGCAAAGTGAAAAAGCTCCTGGTGTTGCAAATAAGAAGATTTCATCT
GGTAGTTCTTGTTAAGCCTTCAGGTGCGAGATGTTACGTTGATGGATCTGAAGAAATTGG
AAGTGACTTTAAGATAAAATGTGAACCAAAAGAAGGTTCACTTCCATTACAGTATGAGTG
GCAAAAATTTGCTGACTCACAGAAAATGCCCACTTCATGGTTAGCAGAAATGACTTCATC
TGTTATATCTGTAAAAAATGCCTCTTCTGAGTACTCTGGGACATACAGCTGTACAGTCAG
AAACAGAGTGGGCTCTGATCAGTGCCTGTTGCGTCTAAACGTTGTCCCTCCTTCAAATAA
AGCTGGACTAATTGCAGGAGCCATTATAGGAACCTTTGCTTGCTCTAGCGCTCATTGGTCT
TATCATCTTTTGCTGTCGTAAAAAGCGCAGAGAAGAAAAATATGAAAAGGAAGTTCATCA
CGATATCAGGGAAGATGTGCCACCTCCAAAGAGCCGTACGTCCACTGCCAGAAGCTACAT
CGGCAGTAATCATTCATCCCTGGGGTCCATGTCTCCTTCCAACATGGAAGGATATTCCAA
GACTCAGTATAACCAAGTACCAAGTGAAGACTTTGAACGCACCTCCTCAGAGTCCGACTCT
CCCACCTGCTAAGTTCAAGTACCCTTACAAGACTGATGGAATTACAGTTGTATAAATATG
GACTACTGAAGAATCTGAAGTATTGTATTATTTGACTTTATTTTAGGCCTCTAGTAAAGA
CTTAAATGTTTTTTAAAAAAGCACAGGCACAGAGATTAGAGCAGCTGTAAGAACACAT
CTACTTTATGCAATGGCATTAGACATGTAAGTCAGATGTCATGTCAAAATTAGTACGAGC
CAAATTCCTTTGTTAAAAAACCTATGTATAGTGACACTGATAGTTAAAGATGTTTTATT
ATATTTTCAATAACTACCCTAACAAATTTTAACTTTTCATATGCATATCTGATATGT
GGTCTTTTAGGAAAAAGTATGGTTAATAGTTGATTTTTTCAAAGGAAATTTTAAATTCCTTA
CGTTCTGTTTAAATGTTTTTGCTATTTAGTTAAATACATTGAAGGGAATACCCGTTCTTT
TCCCTTTTATGCACACAACAGAAACACGCGTTGTCATGCCTCAAACCTATTTTTTATTG
CAACTACATGATTTTACACAATTCTCTTAAACAACGACATAAAATAGATTTCTTGTATA
TAAATAACTTACATACGCTCCATAAAGTAAATTTCTCAAAGGTGCTAGAACAAATCGTCCA
CTTCTACAGTGTTCTCGTATCCAACAGAGTTGATGCACAATATATAAATACTCAAGTCCA
ATATTAAAACTTAGGCACCTTGACTAACTTTAATAAAATTTCTCAAACCTATATCAATATC
TAAAGTGATATATTTTTTAAAGAAAGATTATTTCTCAATAACTTCTATAAAAAATAAGTTTG
ATGGTTTGGCCCATCTAACTTCACTACTATTAGTAAGAACTTTTAACTTTTAAATGTGTAG
TAAGGTTTATTCTACCTTTTTCTCAACATGACACCAACACAATCAAAAACGAAGTTAGTG
AGGTGCTAACATGTGAGGATTAATCCAGTGATTCCGGTCACAATGCATTCCAGGAGGAGG
TACCCATGTCACTGGAATTGGGCGATATGGTTTTATTTTTCTTCCCTGATTTGGATAACC
AAATGGAACAGGAGGAGGATAGTGATTCTGATGGCCATTCCCTCGATACATTCCCTGGCTT
TTTTCTGGGCAAAGGGTGCCACATTGGAAGAGGTGGAAATATAAGTTCTGAAATCTGTAG
GGAAGAGAACACATTAAGTTAATTCAAAGGAAAAAATCATCATCTATGTTCCAGATTTCT
CATTAAAGACAAAGTTACCCACAACACTGAGATCACATCTAAGTGACACTCCTATTGTCA
GGTCTAAATACATTAAAAACCTCATGTGTAATAGGCGTATAATGTATAACAGGTGACCAA
TGTTTTCTGAATGCATAAAGAAATGAATAAACTCAAACACAGTACTTCCTAAACAACCTTC
AACCAAAAAAGACCAAAACATGGAACGAATGGAAGCTTGTAAGGACATGCTTGTTTTAGT
CCAGTGGTTTCCACAGCTGGCTAAGCCAGGAGTCACTTGAGGCTTTTAAATACAAAACA
TTGGAGCTGGAGGCCATTATCCTTAGCAAACTAATGCAGAAACAGAAAATCAACTACCGC
ATGTTCTCACTTATAAGTGGGAGGTAATGATAAGAACTTATGAACACAAAGAAGGAAACA
ATAGACATTGGAGTCTATTTGAGAGGGGAGGGTGGGAGAAGGAAAAGGAGCAGAAAAGAT
AACTATTGAGTACTGCCTTCACACCTGGGTGATGAAATAATATGTACAACAAATCCCTGT
GACACATGTTTACCTATGGAACAAACCTTCATGTGTATCCCTAAACCTAAAAATAAAGTT

293/392

FIGURE 280

MALLLCFVLLCGVVDFARSLSITTPPEEMIEKAKGETAYLPCKFTLSPEDQGPLDIEWLIS
PADNQKVDQVIILYSGDKIYDDYYPDLKGRVHFTSNDLKSGDASINVTNLQLSDIGTYQC
KVKKAPGVANKKIHLVVLVKPSGARCIVDGSEEIGSDFKIKCEPKEGSLPLQYEWQKLS
SQKMPTSWLAEMTSSVISVKNASSEYSGTYSCTVRNRVGSQCLLRNLNVPPSNKAGLIA
GAIIGTLLALALIGLIIFCCRKKRREEKYEKEVHHDIREDVPPPKSRTSTARSYIGSNHS
SLGSMSPSNMEGYSKTQYNQVPSEDFERTPQSPTLPPAKFKYPYKTDGITVV

Signal sequence.

amino acids 1-19

Transmembrane domain:

amino acids 236-257

N-glycosylation sites:

amino acids 106-110, 201-205, 298-302

Tyrosine kinase phosphorylation sites:

amino acids 31-39, 78-85, 262-270

N-myristoylation sites:amino acids 116-122, 208-214, 219-225, 237-243, 241-247,
245-251, 296-302**Myelin P0 protein:**

amino acids 96-125

294/392

FIGURE 281

TGCATCAGTGCCCAGGCAAGCCCAGGAGTTGACATTTCTCTGCCCAGCCATGGGCCTCAC
CCTGCTCTTGCTGCTGCTCCTGGGACTAGAAGGTCAGGGCATAGTTGGCAGCCTCCCTGA
GGTGCTGCAGGCACCCGTGGGAAGCTCCATTCTGGTGCAAGTCCACTACAGGCTCCAGGA
TGTCAAAGCTCAGAAGGTGTGGTGCCGGTTCTTGCCGGAGGGGTGCCAGCCCCCTGGTGTC
CTCAGCTGTGGATCGCAGAGCTCCAGCGGGCAGGCGTACGTTTCTCACAGACCTGGGTGG
GGGCCTGCTGCAGGTGGAAATGGTTACCCTGCAGGAAGAGGATGCTGGCGAGTATGGCTG
CATGGTGGATGGGGCCAGGGGGCCCCAGATTTTGCACAGAGTCTCTCTGAACATACTGCC
CCCAGAGGAAGAAGAAGAGACCCATAAGATTGGCAGTCTGGCTGAGAACGCATTCTCAGA
CCCTGCAGGCAGTGCCAACCCTTTGGAACCCAGCCAGGATGAGAAGAGCATCCCCCTTGAT
CTGGGGTGCTGTGCTCCTGGTAGGTCTGCTGGTGGCAGCGGTGGTGCTGTTTGCTGTGAT
GGCCAAGAGGAAACAAGAATCCCTCCTCAGTGGTCCACCACGTCAGTGACTCTGGACCGG
CTGCTGAATTGCCTTTGGATGTACCACACATTAGGCTTGACTCACCACCTTCATTTGACA
ATACCACCTACACCAGCCTACCTCTTGATTCCCCATCAGGAAAACCTTCACTCCCAGCTC
CATCCTCATTGCCCCCTCTACCTCCTAAGGTCCTGGTCTGCTCCAAGCCTGTGACATATG
CCACAGTAATCTTCCCGGGAGGGAACAAGGGTGGAGGGACCTCGTGTGGGCCAGCCCAGA
ATCCACCTAACAATCAGACTCCATCCAGCTAAGCTGCTCATCACACTTTAAACTCATGAG
GACCATCCCTAGGGGTCTGTGCATCCATCCAGCCAGCTCATGCCCTAGGATCCTTAGGA
TATCTGAGCAACCAGGGACTTTAAGATCTAATCCAATGTCCTAACTTTACTAGGGAAAGT
GACGCTCAGACATGACTGAGATGTCTTGGGGAAGACCTCCCTGCACCCAACTCCCCCACT
GGTTCTTCTACCATTACACACTGGGCTAAATAAACCTAATAATGATGTGCAAAAAAAAAA
AA

295/392

FIGURE 282

MGLTLLLLLLLLLGLGQGIVGSLPEVLQAPVGSSILVQCHYRLQDVKAQKVWCRFLPEGCQ
PLVSSAVDRRAPAGRRTFLTDLGGGLLQVEMVTLQEEDAGEYGCMVDGARGPQILHRVSL
NILPPEEEEEETHKIGSLAENAFSDPAGSANPLEPSQDEKSIPLIWGAVLLVGLLVAAVVL
FAVMAKRKQESLLSGPPRQ

Important features of the protein:

Signal peptide:

amino acids 1-15

Transmembrane domain:

amino acids 161-181

N-myristoylation sites:

amino acids 17-23, 172-178

Amidation site:

amino acids 73-79

296/392

FIGURE 283

GTAGCATAGTGTGCAGTTCACCTGGACCAAAGCTTTGGCTGCACCTCTTCTGGAAAGCTG
GCCATGGGGCTCTTCATGATCATTGCAATTCTGCTGTTCCAGAAACCCACAGTAACCGAA
CAACTTAAGAAGTGCTGGAATAACTATGTACAAGGACATTGCAGGAAAATCTGCAGAGTA
AATGAAGTGCCTGAGGCACTATGTGAAAATGGGAGATACTGTTGCCTCAATATCAAGGAA
CTGGAAGCATGTAAAAAATTACAAAGCCACCTCGTCCAAAGCCAGCAACACTTGCACTG
ACTCTTCAAGACTATGTTACAATAATAGAAAATTTCCAAGCCTGAAGACACAGTCTACA
TAAATCAAATACAATTTTCGTTTTCACTTGCTTCTCAACCTAGTCTAATAAACTAAGGTGA
TGAGATATACATCTTCTTCCTTCTGGTTTCTTGATCCTTAAAATGACCTTCGAGCATATT
CTAATAAAGTGCATTGCCAGTTAAAAA

297/392

FIGURE 284

MGLFMIIAILLFQKPTVTEQLKKCWNNYVQGHCRKICRVNEVPEALCENGRYCCLNIKEL
EACKKITKPPRPKPATLALTLQDYVTIIEFSLKTQST

Important features of the protein:

Signal peptide:

None

Transmembrane domain:

None

cAMP- and cGMP-dependent protein kinase phosphorylation site:

64-68

298/392

FIGURE 285

GATGGCGCAGCCACAGCTTCTGTGAGATTTCGATTTCTCCCCAGTTCCCCTGTGGGTCTGA
GGGGACCAGAAGGGTGAGCTACGTTGGCTTTCTGGAAGGGGAGGCTATATGCGTCAATTC
CCCAAAACAAGTTTTGACATTTCCCCTGAAATGTCATTCTCTATCTATTCACTGCAAGTG
CCTGCTGTTCCAGGCCTTACCTGCTGGGCACCTAACGGCGGAGCCAGGATGGGGACAGAAT
AAAGGAGCCACGACCTGTGCCACCAACTCGCACTCAGACTCTGAACTCAGACCTGAAATC
TTCTCTTCACGGGAGGCTTGGCAGTTTTTCTTACTCCTGTGGTCTCCAGATTTCAAGGCCT
AAGATGAAAGCCTCTAGTCTTGCCTTCAGCCTTCTCTCTGCTGCGTTTTATCTCCTATGG
ACTCCTTCCACTGGACTGAAGACACTCAATTTGGGAAGCTGTGTGATCGCCACAAACCTT
CAGGAAATACGAAATGGATTTTTCTGAGATACGGGGCAGTGTGCAAGCCAAAGATGGAAAC
ATTGACATCAGAATCTTAAGGAGGACTGAGTCTTTGCAAGACACAAAGCCTGCGAATCGA
TGCTGCCTCCTGCGCCATTTGCTAAGACTCTATCTGGACAGGGTATTTAAAAACTACCAG
ACCCCTGACCATTATACTCTCCGGAAGATCAGCAGCCTCGCCAATTCCTTTCTTACCATC
AAGAAGGACCTCCGGCTCTCTCATGCCCACATGACATGCCATTGTGGGGAGGAAGCAATG
AAGAAATACAGCCAGATTCTGAGTCACTTTGAAAAGCTGGAACCTCAGGCAGCAGTTGTG
AAGGCTTTGGGGGAACTAGACATTCTTCTGCAATGGATGGAGGAGACAGAATAGGAGGAA
AGTGATGCTGCTGCTAAGAATATTGAGGTCAGAGCTCCAGTCTTCAATACCTGCAGAG
GAGGCATGACCCCAAACCACCATCTCTTTACTGTACTAGTCTTGTGCTGGTCAAGTGTA
TCTTATTTATGCATTACTTGCTTCCTTGCAATGATTGTCTTTATGCATCCCCAATCTTAAT
TGAGACCATACTTGTATAAGATTTTTTGTAATATCTTTCTGCTATTGGATATATTTATTAG
TTAATATATTTATTTATTTTTTGCTATTTAATGTATTTATTTTTTTACTTGGACATGAAA
CTTTAAAAAAATTACAGATTATATTTATAACCTGACTAGAGCAGGTGATGTATTTTTAT
ACAGTAAAAAATAACCTTGTAATTTCTAGAAGAGTGGCTAGGGGGGTATTTCATTTG
TATTCAACTAAGGACATATTTACTCATGCTGATGCTCTGTGAGATATTTGAAATGAACC
AATGACTACTTAGGATGGGTGTGGAATAAGTTTTGATGTGGAATTGCACATCTACCTTA
CAATTACTGACCATCCCAGTAGACTCCCAGTCCCATAATTGTGTATCTTCCAGCCAGG
AATCCTACACGGCCAGCATGTATTTCTACAAATAAAGTTTTCTTGCATACCAAAAAAA
AAAAA

299/392

FIGURE 286

MKASSLAFSLLSAAFYLLWTPSTGLKTLNLGSCVIATNLQEIRNGFSEIRGSVQAKDGN
DIRILRRTESLQDTKPANRCCLLRHLLRLYLDRVFKNYQTPDHYTLRKISSLANSTLIK
KDLRLSHAHMTCHCGEAMKKYSQILSHFEKLEPQAAVVKALGELDILLQWMEETE

300/392

FIGURE 287

AATGCCCCATGCGCACCCACAGCTCGCGCTCCTGCAAGTGTTCTTTCTGGTGTTCCCCG
ATGGCGTCCGGCCTCAGCCCTCTTCTCCCCATCAGGGGCAGTGCCACGTCTTTGGAGC
TGCAGCGAGGGACGGATGGCGGAACCTCCAGTCCCCTTCAGAGGCGACTGCAACTCGCC
CGGCCGTGCCTGGACTCCCTACAGTGGTCCCTACTCTCGTGACTCCCTCGGCCCCCTGGGA
ATAGGACTGTGGACCTCTTCCCAGTCTTACCGATCTGTGTCTGTGACTTGACTCCTGGAG
CCTGCGATATAAATTGCTGCTGCGACAGGGACTGCTATCTTCTCCATCCGAGGACAGTTT
TCTCCTTCTGCCTTCCAGGCAGCGTAAGGTCTTCAAGCTGGGTTTGTGTAGACAACTCTG
TTATCTTCAGGAGTAATTCCCCGTTTCTTCAAGAGTTTTCATGGATTCTAATGGAATCA
GGCAGTTTTGTGTCCATGTGAACAACCTCAAACCTTAACTATTTCCAGAAGCTTCAAAAGG
TCAATGCAACCAACTTCCAGGCCCTGGCTGCAGAGTTTGGAGGCGAATCATTCACTTCAA
CATTCCAAACTCAATCACCACCATCTTTTTTACAGGGCTGGGGACCCCATTTCTACTTACT
TCCCCAAGTGGTCTGTAATAAGCTTGCTGAGACAACCTGCAGGAGTTGGAGCTGGGGGAC
TCTGTGCTGAAAGCAATCCTGCAGGTTTCTAGAGAGTAAAAGTACAACCTGCACTCGTT
TTTTCAAGAACCTGGCTAGTAGCTGTACCTTGGATTGAGCCCTCAATGCTGCCTCTTACT
ATAACTTCACAGTCTTAAAGGTTCCAAGAAGCATGACTGATCCACAGAATATGGAGTTCC
AGGTTCTGTAAATACTTACCTCACAGGCTAATGCTCCTCTGTTGGCTGGAAACACTTGTC
AGAATGTAGTTTCTCAGGTCACCTATGAGATAGAGACCAATGGGACTTTTGGAATCCAGA
AAGTTTCTGTGAGTTTGGGACAAACCAACCTGACTGTTGAGCCAGGCGCTTCCTTACAGC
AACACTTCATCCTTCGCTTCAGGGCTTTTCAACAGAGCACAGCTGCTTCTCTCACCAGTC
CTAGAAGTGGGAATCCTGGCTATATAGTTGGGAAGCCACTCTTGGCTCTGACTGATGATA
TAAGTTACTCAATGACCCTCTTACAGAGCCAGGGTAATGGAAGTTGCTCTGTTAAAAGAC
ATGAAGTGCAGTTTGGAGTGAATGCAATATCTGGATGCAAGCTCAGGTTGAAGAAGGCAG
ACTGCAGCCACTTGCAAGCAGGAGATTTATCAGACTCTTCATGGAAGGCCAGACCAGAGT
ATGTTGCCATCTTTGGTAATGCTGACCCAGCCAGAAAGGAGGGTGGACCAGGATCCTCA
ACAGGCACTGCAGCATTTAGCTATAAACTGTACTTCTGCTGTCTCATAACAGTTTCCC
TGGAGATCCAGGTATTGTGGGCATATGTAGGTCTCCTGTCCAACCCGCAAGCTCATGTAT
CAGGAGTTCGATTCTTATACCAGTGCCAGTCTATACAGGATTCTCAGCAAGTTACAGAAG
TATCTTTGACAACTCTTGTGAACCTTGTGGACATTACCCAGAAGCCACAGCCTCCAAGGG
GCCAACCCAAAATGGACTGGAAATGGCCATTGCACTTCTTCCCTTCAAAGTGGCATTCA
GCAGAGGAGTATTCTCTCAAAAATGCTCAGTCTCTCCATCCTTATCCTGTGCCTCTTAC
TACTTGGAGTTCTCAACCTAGAGACTATGTGAAGAAAAGAAAATAATCAGATTTCAGTTT
TCCCTATGAGAACTCTGAGGCAGCCACTTATCTTGGCTAAATAGAACCTCACCTGCTCA
TGACCAGAGAGCATTTAGGATAATAGATGACCTAACTGAAGGAATCCTTGTATATGAAAG
GAGTTATTTAGAAAAGCAATAAAAATATTTTATTCATCNTAAAAAAAAA

301/392

FIGURE 288

M RTPQLALLQVFFLVFPDGV RPQPSSSPSGAVPTSLELQRGTDGGTLQSPSEATATRP AV
PGLPTVVPTLVTPSAPGNRTVDLFPVLPICVCDLTPGACDINCCCDRDCYLLHPRTVFSF
CLPGSVRSSSWVCVDNSVIFRSNSPFPSRVFMD SNGIRQFCVHVNN SNLNYFQKLQKVNA
TNFQALAAEFGGESFTSTFQTQSPPSFYRAGDPILTYFPKWSVISLLRQPAGVGAGGLCA
ESNPAGFLESKSTTCTRFFKNLASSCTLDSALNAASYNFTVLKVPRSM TDPQNMEFQVP
VILTSQANAPLLAGNTCQNVVSQVTYE IETNGTFGIQKVS VSLGQTNLTVEPGASLQQHF
ILRFRAFQQSTAASLTSPRSGNPGYIVGKPLLALTDDISYSMTLLQSQNGSCSVKRHEV
QFGVNAISGCKLRLKKADC SHLQQEIYQTLHGRPRPEYVAIFGNADPAQKGGWTRILNRH
CSISAINCTSCCLIPVSLEIQVLWAYVGLLSNPQAHVSGVRFLYQCQSIQDSQQVTEVSL
TTLVNFVDITQKPQPPRGQPKMDWKWPFDFFPFKVAFSRGVFSQKCSVSPILILCLLLLG
VLNLETM

Important features of the protein:**Signal peptide:**

amino acids 1-22

Transmembrane domains:

amino acids 484-505, 581-600

N-glycosylation sites:amino acids 78-82, 165-169, 179-185, 279-285, 331-337,
347-351, 410-414, 487-491**N-myristoylation sites:**

amino acids 30-36, 41-47, 124-130, 232-238, 236-242, 409-415

Prokaryotic membrane lipoprotein lipid attachment site:

amino acids 420-431

302/392

FIGURE 289

CGCGGAGCCCTGCGCTGGGAGGTGCACGGTGTGCACGCTGGACTGGACCCCCATGCAACC
CCGCGCCCTGCGCCTTAACCAGGACTGCTCCGCGCGCCCCCTGAGCCTCGGGCTCCGGCCC
GGACCTGCAGCCTCCCAGGTGGCTGGGAAGAACTCTCCAACAATAAATACATTTGATAAG
AAAGATGGCTTTAAAAGTGCTACTAGAACAAGAGAAAAACGTTTTTCACTCTTTTAGTATT
ACTAGGCTATTTGTGCTAAAGTGACTTGTGAATCAGGAGACTGTAGACAGCAAGAATT
CAGGGATCGGTCTGGAACCTGTGTTCCCTGCAACCAGTGTGGGCCAGGCATGGAGTTGTC
TAAGGAATGTGGCTTCGGCTATGGGGAGGATGCACAGTGTGTGACGTGCCGGCTGCACAG
GTTCAAGGAGGACTGGGGCTTCAGAAATGCAAGCCCTGTCTGGAAGTGCAGTGGTGAA
CCGCTTTCAGAAGGCAAATTGTTACGCCACCAAGTGTGCCATCTGCGGGGACTGCTTGCC
AGGATTTTATAGGAAGACGAACTTGTTCGGCTTTCAAGACATGGAGTGTGTGCCTTGTGG
AGACCCCTCCTCCTTACGAACCGCACTGTGCCAGCAAGGTCAACCTCGTGAAGATCGC
GTCCACGGCCTCCAGCCCACGGGACACGGCGCTGGCTGCCGTTATCTGCAGCGCTCTGGC
CACCGTCTGTGGCCCTGCTCATCTCTGTGTGTCATCTATTGTAAGAGACAGTTTATGGA
GAAGAAACCCAGCTGGTCTCTGCGGTCTCAGGACATTCAGTACAACGGCTCTGAGCTGTC
GTGTTTTGACAGACCTCAGCTCCACGAATATGCCACAGAGCCTGCTGCCAGTGCCGCCG
TGAAGTCAAGTGCAGACCTGCGGGCCGGTGCCTTCTCCATCCATGTGCTGTGAGGAGGC
CTGCAGCCCCAACCCGGCGACTCTTGGTTGTGGGGTGCATTCTGCAGCCAGTCTTCAGGC
AAGAAACGCAGGCCAGCCGGGGAGATGGTGCCGACTTTCTTCGGATCCCTCACGCAGTC
CATCTGTGGCGAGTTTTCAGATGCCTGGCCTCTGATGCAGAATCCCATGGGTGGTGACAA
CATCTCTTTTTGTGACTCTTATCCTGAACCTCACTGGAGAAGACATTCATCTCTCAATCC
AGAACTTGAAAGCTCAACGTCTTTGGATTCAAATAGCAGTCAAGATTTGGTTGGTGGGGC
TGTTCCAGTCCAGTCTCATTCTGAAAACCTTTACAGCAGCTACTGATTTATCTAGATATAA
CAACACACTGGTAGAATCAGCATCAACTCAGGATGCACTAACTATGAGAAGCCAGCTAGA
TCAGGAGAGTGGCGCTGTCATCCACCCAGCCACTCAGACGTCCCTCCAGGAAGCTTAAAG
AACCTGCTTCTTTCTGTCAGTAGAAGCGTGTGCTGGAACCCAAAGAGTACTCCTTTGTAG
GCTTATGGACTGAGCAGTCTGGACCTTGCATGGCTTCTGGGGCAAAAATAAATCTGAACC
AACTGACGGCATTTGAAGCCTTTCAGCCAGTTGCTTCTGAGCCAGACCAGCTGTAAGCT
GAAACCTCAATGAATAACAAGAAAAGACTCCAGGCCGACTCATGATACTCTGCATCTTTC
CTACATGAGAAGCTTCTCTGCCACAAAAGTGACTTCAAAGACTGATGGGTTGAGCTGGCA
GCCTATGAGATTGTGGACATATAACAAGAAACAGAAATGCCCTCATGCTTATTTTCATGG
TGATTGTGGTTTTACAAGACTGAAGACCCAGAGTATACTTTTTCTTTCAGAAATAATTT
CATACCGCCTATGAAATATCAGATAAATTACCTTAGCTTTTATGTAGAATGGGTTCAAAA
GTGAGTGTTCCTATTTGAGAAGGACACTTTTTCATCATCTAAACTGATTCGCATAGGTGG
TTAGAATGGCCCTCATATTGCCTGCCTAAATCTTGGGTTTATTAGATGAAGTTTACTGAA
TCAGAGGAATCAGACAGAGGAGGATAGCTCTTCCAGAATCCACACTTCTGACCTCAGCC
TCGGTCTCATGAACACCCGCTGATCTCAGGAGAACACCTGGGCTAGGGAATGTGGTCGAG
AAAGGGCAGCCCATTGCCCAGAATTAACACATATTGTAGAGACTTGTATGCAAAGGTTGG
CATATTTATATGAAAATTAGTTGCTATAGAAACATTTGTTGCATCTGTCCCTCTGCCTGA
GCTTAGAAGGTTATAGAAAAGGGTATTTATAAACATAAATGACCTTTTACTTGCAATTGT
ATCTTATACTAAAGGCTTTAGAAATTACAACATATCAGGTTCCCCTACTACTGAAGTAGC
CTTCCGTGAGAACACACCACATGTTAGGACTAGAAGAAAATGCACAATTTGTAGGGGTTT
GGATGAAGCAGCTGTAAGTCCCTAGTGTAGTTTGACCAGGACATTGTGCTGCTCCTTCC
AATTGTGTAAGATTAGTTAGCACATCATCTCCTACTTTAGCCATCCGGTGTGGATTTAA
GAGGACGGTGCTTCTTCTATTAAAGTGCTCCATCCCTACCATCTACACATTAGCATTG
TCTCTAGAGCTAAGACAGAAATTAACCCCGTTCAGTCACAAAGCAGGGAATGGTTCAATT
ACTCTTAATCTTTATGCCCTGGAGAAGACCTACTTGAACAGGGCATATTTTTTAGACTTC
TGAACATCAGTATGTTTCGAGGGTACTATGATATTTTGGTTTGAATTGCCCTGCCCAAGT
CACTGTCTTTTAACTTTTAACTGAATATTAATAATGTATCTGTCTTTCCT

303/392

FIGURE 290

MALKVLLLEQEKTFFTLLVLLGYLSCKVTCESGDCRQQEFRDRSGNCVPCNQCGPGMELSK
ECGFGYGEDAQCVTCLHRFKEDWGFQKCKPCLDCAVVNRFAQKANCSTSDAICGDCLPG
FYRKTCLVGFQDMECVPCGDP PPPPYEPHCASKVNLVKIASTASSPRDTALAAVICSALAT
VLLALLILCVIYCKRQFMKKPSWSLRSQDIQYNGSELSCFDRPQLHEYAHRACCQCRD
SVQTCGPVRLLLPSMCCEEACSPNPATLGCGVHSAASLQARNAGPAGEMVPTFFGSLTQSI
CGEFSDAWPLMQNPMGGDNISFCDSYPELTGEDIHSLNPELESSTSLDSNSSQDLVGGAV
PVQSHSENFATAATDLRYNNTLVESASTQDALTMRSQLDQESGAVIHPATQTSLQEA

Important features of the protein:**Signal peptide:**

Amino acids 1-25

Transmembrane domain:

Amino acids 169-192

N-glycosylation sites:

Amino acids 105-109; 214-218; 319-323; 350-354; 368-372; 379-383

cAMP- and cGMP-dependent protein kinase phosphorylation sites:

Amino acids 200-204; 238-242

Tyrosine kinase phosphorylation site:

Amino acids 207-214

N-myristoylation sites:

Amino acids 55-61; 215-221; 270-276

Prokaryotic membrane lipoprotein lipid attachment site:

Amino acids 259-270

TNFR/NGFR family cysteine-rich region proteins:

Amino acids 89-96

304/392

FIGURE 291

CCTGGAGCCGGAAGCGCGGCTGCAGCAGGGCGAGGCTCCAGGTGGGGTCCGGTTCCGCATC
CAGCCTAGCGTGTCACGATGCGGCTGGGCTCCGGGACTTTTCGCTACCTGTTGCGTAGCG
ATCGAGGTGCTAGGGATCGCGGTCTTCCTTCGGGGATTCTTCCCGGCTCCCGTTCCGTTCC
TCTGCCAGAGCGGAACACGGAGCGGAGCCCCAGCGCCCGAACCCTCGGCTGGAGCCAGT
TCTAACTGGACCACGCTGCCACCACCTCTCTTCAGTAAAGTTGTTATTGTTCTGATAGAT
GCCTTGAGAGATGATTTTGTGTTTGGGTCAAAGGGTGTGAAATTTATGCCCTACACAAC
TACCTTGTGGAAAAAGGAGCATCTCACAGTTTTGTGGCTGAAGCAAAGCCACCTACAGTT
ACTATGCCTCGAATCAAGGCATTGATGACGGGGAGCCTTCCTGGCTTTGTGACGTCATC
AGGAACCTCAATTCTCTGCACTGCTGGAAGACAGTGTGATAAGACAAGCAAAGCAGCT
GGAAAAAGAATAGTCTTTTATGGAGATGAAACCTGGGTAAATTTATCCCAAAGCATT
GTGGAATATGATGGAACAACCTCATTTTTTCGTGTGAGATTACACAGAGGTGGATAATAAT
GTCACGAGGCATTTGGATAAAGTATTAAAAAGAGGAGATTGGGACATATTAATCCTCCAC
TACCTGGGGCTGGACCACATTGGCCACATTTTCAGGGCCCAACAGCCCCCTGATTGGGCAG
AAGCTGAGCGAGATGGACAGCGTGCTGATGAAGATCCACACCTCACTGCAGTCGAAGGAG
AGAGAGACGCCCTTACCCAATTTGCTGGTTCTTTGTGGTGACCATGGCATGTCTGAAACA
GGAAGTCACGGGGCCTCCTCCACCGAGGAGGTGAATACACCTCTGATTTTAATCAGTTCT
GCGTTTGAAAGGAAACCCGGTGATATCCGACATCCAAAGCACGTCCAATAGACGGATGTG
GCTGCGACACTGGCGATAGCACTTGGCTTACCGATTCCAAAAGACAGTGTAGGGAGCCTC
CTATTCCCAGTTGTGGAAGGAAGACCAATGAGAGAGCAGTTGAGATTTTACATTTGAAT
ACAGTGCAGCTTAGTAACTGTTGCAAGAGAATGTGCCGTGATATGAAAAAGATCCTGGG
TTTGAGCAGTTTAAATGTGAGAAAGATTGCATGGGAACTGGATCAGACTGTACTTGGAG
GAAAAGCATTGAGAGTCTATTCAACCTGGGCTCCAAGGTTCTCAGGCAGTACCTGGAT
GCTCTGAAGACGCTGAGCTTGTCCCTGAGTGCACAAGTGGCCCAGTTCTCACCCTGCTCC
TGCTCAGCGTCCCACAGGCACTGCACAGAAAGGCTGAGCTGGAAGTCCCCTGTCTATCTC
CTGGGTTTTCTCTGCTCTTTTATTTGGTGATCCTGGTTCTTTTCGGCCGTTTACGTCATTG
TGTGCACCTCAGCTGAAAGTTGCTGCTACTTCTGTGGCCTCTCGTGGCTGGCGGCAGGCT
GCCTTTCGTTTACCAGACTCTGGTTGAACACCTGGTGTGTGCCAAGTGTGGCAGTGCCC
TGGACAGGGGGCCTCAGGGAAGGACGTGGAGCAGCCTTATCCCAGGCCTCTGGGTGTCCC
GACACAGGTGTTACATCTGTGCTGTCAGGTGAGATGCCTCAGTTCTTGAAAGCTAGGT
TCCTGCGACTGTTACCAAGGTGATTGTAAAGAGCTGGCGGTACAGAGGAACAAGCCCC
CAGCTGAGGGGGTGTGTGAATCGGACAGCCTCCAGCAGAGGTGTGGGAGCTGCAGCTGA
GGGAAGAAGAGACAATCGGCCTGGACACTCAGGAGGGTCAAAGGAGACTTGGTCGCACC
ACTCATCCTGCCACCCCCAGAATGCATCCTGCCTCATCAGGTCCAGATTTCTTTCCAAGG
CGGACGTTTTCTGTTGGAATTCTTAGTCTTGGCCTCGGACACCTTCATTTCGTTAGCTGG
GGAGTGGTGGTGAGGCAGTGAAGAAGAGGCGGATGGTCACACTCAGATCCACAGAGCCCA
GGATCAAGGGACCCACTGCAGTGGCAGCAGGACTGTTGGGCCCCCACCCTGCAC
AGCCCTCATCCCCTCTGGCTTGAGCCGTGAGAGGCCCTGTGCTGAGTGTCTGACCGAGA
CACTCACAGCTTTGTCATCAGGGCACAGGCTTCCTCGGAGCCAGGATGATCTGTGCCACG
CTTGACCTCGGGCCCATCTGGGCTCATGCTCTCTCTCTGCTATTGAATTAGTACCTAG
CTGCACACAGTATGTAGTTACCAAAGAATAAACGGCAATAATTGAGAAAAAAA

305/392

FIGURE 292

MRLGSGTFATCCVAIEVLGIAVFLRGFFPAPVRSSARAEGHGAEPPEPSAGASSNWTTL
PPPLFSKVIVLIDALRDDFVFGSKGVKFMPTTYLVEKGASHSFVAEAKPPTVTMPRIK
ALMTGSLPGFVDVIRNLNSPALLEDVIRQAKAAGKRIVFYGDETWVKLFPKHFEYDGT
TSFFVSDYTEVDNNVTRHLDKVLKRGDWDILILHYLGLDGHIGHISGPNSPLIGKLSEMD
SVLMKIHTSLQSKERETPLPNLLVLCGDHGMSETGSHGASSTEEVNTPLILISSAFERKP
GDIRHPKHVQ

Important features of the protein:**Signal peptide:**

amino acids 1-34

Transmembrane domain:

amino acids 58-76

N-glycosylation sites:

amino acids 56-60, 194-198

N-myristoylation sites:amino acids 6-12, 52-58, 100-106, 125-131, 233-239, 270-276,
275-281, 278-284**Amidation site:**

amino acids 154-158

Cell attachment sequence:

amino acids 205-208

306/392

FIGURE 293

AGCCAGGCAGCACATCACAGCGGGAGGAGCTGTCCCAGGTGGCCCAGCTCAGCAATGGCA
ATGGGGGTCCCCAGAGTCATTCTGCTCTGCCTCTTTGGGGCTGCGCTCTGCCTGACAGGG
TCCCAAGCCCTGCAGTGCTACAGCTTTGAGCACACCTACTTTGGCCCCCTTGACCTCAGG
GCCATGAAGCTGCCCAGCATCTCCTGTCCTCATGAGTGCTTTGAGGCTATCCTGTCTCTG
GACACCGGGTATCGCGCGCCGGTGACCCCTGGTGCGGAAGGGCTGCTGGACCGGGCCTCCT
GCGGGCCAGACGCAATCGAACCCGGACGCGCTGCCGCCAGACTACTCGGTGGTGCGCGGC
TGCACAACTGACAAATGCAACGCCACCTCATGACTCATGACGCCCTCCCCAACCTGAGC
CAAGCACCCGACCCGCCGACGCTCAGCGGCGCCGAGTGCTACGCCTGTATCGGGGTCCAC
CAGGATGACTGCGCTATCGGCAGGTCCCGACGAGTCCAGTGTCACCAGGACCAGACCGCC
TGCTTCCAGGGCAGTGGCAGAATGACAGTTGGCAATTTCTCAGTCCCTGTGTACATCAGA
ACCTGCCACCGGCCCTCCTGCACCACCGAGGGCACCACCAGCCCCCTGGACAGCCATCGAC
CTCCAGGGCTCCTGCTGTGAGGGGTACCTCTGCAACAGGAAATCCATGACCCAGCCCTTC
ACCAGTGCTTCAGCCACCACCCCTCCCCGAGCACTACAGGTCCCTGGCCCTGCTCCTCCCA
GTCCTCCTGCTGGTGCGGCTCTCAGCATAGACCGCCCCTCCAGGATGCTGGGGACAGGGC
TCACACACCTCATTCTTGCTGCTTCAGCCCCTATCACATAGCTCACTGGAAAATGATGTT
AAAGTAAGAATTGCAAAA

307/392

FIGURE 294

MAMGVPRVILLCLFGAALCLTGSQALQCYSFEHTYFGPFDLRAMKLPSISCPHECFEAIL
SLDTGYRAPVTILVRKGCWTGPPAGQTQSNPDALPPDYSVVRGCTTDKCNAHLMTHDALPN
LSQAPDPPTLSGAECYACIGVHQDDCAIGRSRRVQCHQDQTACFQSGRMTVGNFSPVY
IRTCHRPSCCTTEGTTSPWTAIDLQSCCEGYLCNRKSMTQPFTSASATTPPRALQVLALL
LPVLLLVGLSA

Important features of the protein:**Signal peptide:**

amino acids 1-19

Transmembrane domain:

amino acids 233-251

N-glycosylation sites:

amino acids 120-124, 174-178

N-myristoylation sites:

amino acids 15-21, 84-90

308/392

FIGURE 295

AATCGGCTGATTCTGCATCTGGAACTGCCTTCATCTTGAAAGAAAAGCTCCAGGTCCCT
TCTCCAGCCACCCAGCCCCAAGATGGTGATGCTGCTGCTGCTGCTTTCCGCACTGGCTGG
CCTCTTCGGTGCGGCAGAGGGACAAGCATTTTCATCTTGGGAAGTCCCCAATCCTCCGGT
GCAGGAGAATTTTGACGTGAATAAGTATCTCGGAAGATGGTACGAAATTGAGAAGATCCC
AACAAACCTTTGAGAATGGACGCTGCATCCAGGCCAACTACTCACTAATGGAAAACGGAAA
GATCAAAGTGTTAAACCAGGAGTTGAGAGCTGATGGAAGTGAATCAAATCGAAGGTGA
AGCCACCCAGTTAACCTCACAGAGCCTGCCAAGCTGGAAGTTAAGTTTTCTGGTTTTAT
GCCATCGGCACCGTACTGGATCCTGGCCACCGACTATGAGAAGTATGCCCTCGTGTATTC
CTGTACCTGCATCATCCAACTTTTTCACGTGGATTTTGCTTGGATCTTGGCAAGAAACCC
TAATCTCCCTCCAGAAACAGTGGACTCTCTAAAAAATATCCTGACTTCTAATAACATTGA
TGTCAGAAAATGACGGTCACAGACCAGGTGAACTGCCCCAAGCTCTCGTAACCAGGTTTC
TACAGGGAGGCTGCACCCACTCCATGTTACTTCTGCTTCGCTTTCCCCTACCCACCCCC
CCCCATAAGACAAACCAATCAACCACGACAAAGGAAGTTGACCTGAACATGTAACCAT
GCCCTACCCTGTTACCTTGCTAGCTGCAAAATAAACTTGTTGCTGACCTGCTGTGCTCGC
AAAAAA

309/392

FIGURE 296

MVMLLLLLLSALAGLFGAAEGQAFHLGKCPNPPVQENFDVNKYLGRWYEIEKIPITTFENG
RCIQANYSLMENGKIKVLNQELRADGTVNQIEGEATPVNLTEPAKLEVKFSWFMPSPAY
WILATDYENYALVYSCTCIIQLFHVDFAWILARNPNLPPETVDSLKNILTSNNIDVKKM
TVTDQVNCPKLS

Signal sequence:

1-16

N-glycosylation site:

65-68

98-101

cAMP- and cGMP-dependent protein kinase phosphorylation site:

175-178

N-myristoylation site:

13-18

16-21

Lipocalin proteins:

36-47

120-130

Lipocalin / cytosolic fatty-acid binding proteins:

41-185

310/392

FIGURE 297

GGGTGATTGAACTAAACCTTCGCCGCACCGAGTTTGACGTACGGCCGTCACCCGCACCGC
TGCCTGCTTGCGGTTGGAGAAATCAAGGCCCTACCGGGCCTCCGTAGTCACCTCTCTATA
GTGGGCGTGGCCGAGGCCGGGGTGACCCTGCCGGAGCCTCCGCTGCCAGCGACATGTTCA
AGGTAATTCAGAGGTCCGTGGGGCCAGCCAGCCTGAGCTTGCTCACCTTCAAAGTCTATG
CAGCACCAAAAAAGGACTCACCTCCCAAAAATTCCGTGAAGGTTGATGAGCTTTCCTCT
ACTCAGTTCCTGAGGGTCAATCGAAGTATGTGGAGGAGGCAAGGAGCCAGCTTGAAGAAA
GCATCTCACAGCTCCGACACTATTGCGAGCCATACACAACCTGGTGTGAGGAAACGTACT
CCCAAACCTAAGCCCAAGATGCAAAGTTTGGTTCAATGGGGGTTAGACAGCTATGACTATC
TCCAAAATGCACCTCCTGGATTTTTTCCGAGACTTGGTGTATTGGTTTTGCTGGCCTTA
TTGGACTCCTTTTGGCTAGAGGTTCAAAAATAAAGAAGCTAGTGTATCCGCCTGGTTTTCA
TGGGATTAGCTGCCTCCCTCTATTATCCACAACAAGCCATCGTGTTTGCCAGGTCAGTG
GGGAGAGATTATATGACTGGGGTTTACGAGGATATATAGTCATAGAAGATTTGTGGAAGG
AGAACTTTCAAAGCCAGGAAATGTGAAGAATTCACCTGGAACCTAAGTAGAAAACCTCCAT
GCTCTGCCATCTTAATCAGTTATAAGGTAAACATTGGAACTCCATAGAATAAATCAGTAT
TTCTACAGAAAAATGGCATAGAAGTCAGTATTGAATGTATTAAATTGGCTTTCCTTCTTCA
GGAAAACTAGACCAGACCTCTGTTATCTTCTGTGAAATCATCCTACAAGCAAACCTAACC
TGGAATCCCTTCACCTAGAGATAATGTACAAGCCTTAGAACTCCTCATTCTCATGTTGCT
ATTTATGTACCTAATTAAAACCAAGTTTAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAA

311/392

FIGURE 298

MFKVIQRSVGPASLSLLTFKVYAAPKKDSPPKNSVKVDELSTLYSVPEGQSKYVEEARSQLE
EESISQLRHYCEPYTTWCQETYSQTKPKMQSLVQWGLDSYDYLQNAPPGFFPRLGVIGFA
GLIGLLLARGSKIKKLVYPPGFMGLAASLYYPQQAIVFAQVSGERLYDWGLRGYIVIEDL
WKENFQKPGNVKNSPGTK

Important features:

Signal peptide:

Amino acids 1-23

Transmembrane domain:

Amino acids 111-130

cAMP- and cGMP-dependent protein kinase phosphorylation site:

Amino acids 26-30

Tyrosine kinase phosphorylation site:

Amino acids 36-44

N-myristoylation sites:

Amino acids 124-130;144-150;189-195

312/392

FIGURE 299

CCGCTGAGATGTACGAACTTCCGGTTCTCCGGGCAGCTGCCACTGCTGTAGCTTCTGCCA
CCTGCCACGACCGGGCCTCTCCCTGGCGTTTGGTCACCTCTGCTTCATTCTCCACCGCGC
CTATGGTCCCTCTTGAGCCAGCGTGGCGGGCCTGGCGGCTCCCGGGTGGTGAGAGAGCG
GTCCGGGAACGATGAAGGCCTCGCAGTGCTGCTGCTGTCTCAGCCACCTCTTGGCTTCCG
TCCTCCTCCTGCTGTTGCTGCCTGAACTAAGCGGGCCCCTGGCAGTCCTGCTGCAGGCAG
CCGAGGCCGCGCCAGGTCTTGGGCCTCCTGACCCTAGACCACGGACATTACCGCCGCTGC
CACCGGGCCCTACCCCTGCCCAGCAGCCGGGCGGTGGTCTGGCTGAAGCTGCGGGGCCG
GGGGCTCCGAGGGAGGCAATGGCAGCAACCCTGTGGCCGGGCTTGAGACGGACGATCACG
GAGGGAAGGCCGGGAAGGCTCGGTGGGTGGCGGCCCTTGCTGTGAGCCCCAACCTGGCG
ACAAGCCCATGACCCAGCGGGCCCTGACCGTGTTGATGGTGGTGAGCGGCGCGGTGCTGG
TGTACTTCGTGGTCAGGACGGTCAGGATGAGAAGAAGAAACCGAAAGACTAGGAGATATG
GAGTTTTGGACACTAACATAGAAAATATGGAATTGACACCTTTAGAACAGGATGATGAGG
ATGATGACAACACGTTGTTTGATGCCAATCATCCTCGAAGATAAGAATGTGCCTTTTGAT
GAAAGAAC'TTTATCTTTCTACAATGAAGAGTGGAATTTCTATGTTTAAGGAATAAGAAGC
CACTATATCAATGTTGGGGGGGTATTTAAGTTACATATATTTTAACAACCTTTAATTTGC
TGTTGCAATAAATACCGTATCCTTTTATTATATCTTTATATGTATAGAAGTACTCTATTA
ATGGGCTCAGAGATGTTGGGGATAAAGTATACTGTAATAATTTATCTGTTTGAAAATTAC
TATAAAACGGTGTTTTCTGGTGGTTTTTGTTCCTGCTTACCATATGATTGTAAATTGT
TTTATGTATTAATCAGTTAATGCTAATTATTTTGTCTGATGTCATATGTTAAAGAGCTAT
AAATTCCAACAACCAACTGGTGTGTAATAATAATTTAAATTTCCTTTACTGAAAGGTAT
TTCCCATTTTTGTGGGGAAGAAGCCAAATTTATTACTTTGTGTTGGGGTTTTTAAAT
ATTAAGAAATGTCTAAGTTATTGTTTGCAAAACAATAAATATGATTTTAAATTCTCTAA
AAAAAAA

313/392

FIGURE 300

MKASQCCCCLSHLLASVLLLLLLPELSGPLAVLLQAEEAAPGLGPPDPRPRTLPPPLPPGP
TPAQQPGRGLAEAAAGPRGSEGGNGSNPVAGLETDDHGGKAGEGSVGGLAVSPNPGDKPM
TQRALTVLMVVSGAVLVYFVVRTVRMRRNRKTRRYGVLDTNIE NMELTPLEQDDEDDDN
TLFDANHPRR

Signal peptide:
amino acids 1-28

Transmembrane domain:
amino acids 124-140

N-glycosylation site:
amino acids 83-87

N-myristoylation sites:
amino acids 69-75, 78-84, 81-87, 97-103, 103-109, 106-112,
157-160

314/392

FIGURE 301

CTCGGCTGGATTTAAGGTTGCCGCTAGCCGCTGGGAATTTAAGGGACCCACACTACCTT
CCCGAAGTTGAAGGCAAGCGGTGATTGTTTGTAGACGGCGCTTTGTCATGGGACCTGTGC
GGTTGGGAATATTGCTTTTCCTTTTTTGGCCGTGCACGAGGCTTGGGCTGGGATGTTGA
AGGAGGAGGACGATGACACAGAACGCTTGCCAGCAAATGCGAAGTGTGTAAGCTGCTGA
GCACAGAGCTACAGGCGGAAGTGAAGTGCACCGGTGATCTCGAGAGGTGCTGGAGCTGG
GGCAGGTGCTGGATACAGGCAAGAGGAAGAGACACGTGCCCTTACAGCGTTTCAGAGACAA
GGCTGGAAGAGGCCCTTAGAGAATTTATGTGAGCGGATCCTGGACTATAGTGTTACGCTG
AGCGCAAGGGCTCACTGAGATATGCCAAGGGTCAGAGTCAGACCATGGCAACACTGAAAG
GCCTAGTGAGCAAGGGGGTGAAGGTGGATCTGGGGATCCCTCTGGAGCTTTGGGATGAGC
CCAGCGTGGAGGTCACATACCTCAAGAAGCAGTGTGAGACCATGTTGGAGGAGTTTGAAG
ACATTGTGGGAGACTGGTACTTCCACCATCAGGAGCAGCCCCTACAAAATTTTCTCTGTG
AAGGTCATGTGCTCCCAGCTGCTGAAACTGCATGTCTACAGGAACTTGGACTGGAAAGG
AGATCACAGATGGGGAAGAGAAAACAGAAGGGGAGGAAGAGCAGGAGGAGGAGGAGGAAG
AGGAGGAAGAGGAAGGGGGAGACAAGATGACCAAGACAGGAAGCCACCCCAAACCTTGACC
GAGAAGATCTTTTGACCCTTGCTTTGAGCCCCCAGGAGGGGAAGGGATCATGGAGAGCCC
TCTAAAGCCTGCACCTCTCCCTGCTCCACAGCTTTCAGGGTGTGTTTATGAGTGAATCCAC
CCAAGCTTGTAGCTGTTCTCTCCCATCTAACCTCAGGCAAGATCCTGGTGAAACAGCATG
ACATGGCTTCTGGGGTGGAGGGTGGGGGTGGAGGTCTGCTCCTAGAGATGAACTCTATC
CAGCCCCCTTAATTGGCAGGTGTATGTGCTGACAGTACTGAAAGCTTTCCTCTTTAACTGA
TCCCCCCCCACCCAAAAGTCAGCAGTGGCACTGGAGCTGTGGGCTTTGGGGAAGTCACT
TAGCTCCTTAAGGTCTGTTTTTAGACCCTTCCAAGGAAGAGGCCAGAACGGACATTCTCT
GCGATCTATATACATTGCCTGTATCCAGGAGGCTACACACCAGCAAACCGTGAAGGAGAA
TGGGACACTGGGTGATGGCCTGGAGTTGCTGATAATTTAGGTGGGATAGATACTTGGTCT
ACTTAAGCTCAATGTAACCCAGAGCCCACCATATAGTTTTATAGGTGCTCAACTTTCTAT
ATCGCTATTAACTTTTTTCTTTTTTTCTA

315/392

FIGURE 302

MGPVRLGILLFLFLAVHEAWAGMLKEEDDDTERLPSKCEVCKLLSTELQAELSRTGRSRE
VLELGQVLDTGKRKRHPYPSVSETRLEEALENLCERILDYSVHAERKGSRLRYAKGQSQTM
ATLKGLVQKGVKVDLGIPLELWDEPSVEVTYLKKQCETMLEEFEDIVGDWYFHHQEQLQ
NFLCEGHVLPAAETACLQETWTGKEITDGEEKTEGEEEQEEEEEEEEEGGDKMTKTGSH
PKLDREDL

Important features of the protein:

Signal peptide:

amino acids 1-21

cAMP- and cGMP-dependent protein kinase phosphorylation site:

amino acids 106-110

N-myristoylation site:

amino acids 115-121

Amidation site:

amino acids 70-74

316/392

FIGURE 303

CTCCTGCACTAGGCTCTCAGCCAGGGATGATGCGCTGCTGCCGCCGCCGCTGCTGCTGCC
GGCAACCACCCCATGCCCTGAGGCCGTTGCTGTTGCTGCCCCCTCGTCCTTTTACCTCCCC
TGGCAGCAGCTGCAGCGGGCCCAAACCGATGTGACACCATATACCAGGGCTTCGCCGAGT
GTCTCATCCGCTTGGGGGACAGCATGGGCCGCGGAGGCGAGCTGGAGACCATCTGCAGGT
CTTGGAATGACTTCCATGCCTGTGCCTCTCAGGTCCTGTCAGGCTGTCCGGAGGAGGCAG
CTGCAGTGTGGGAATCACTACAGCAAGAAGCTCGCCAGGCCCCCCCGTCCGAATAACTTGC
ACACTCTGTGCGGTGCCCCGGTGCA TGTTGGGAGCGCGGCACAGGCTCCGAAACCAACC
AGGAGACGCTGCGGGCTACAGCGCCTGCACTCCCCATGGCCCCCTGCGCCCCCACTGCTGG
CGGCTGCTCTGGCTCTGGCCTACCTCCTGAGGCCTCTGGCCTAGCTTGTTGGGTTGGGTA
GCAGCGCCCGTACCTCCAGCCCTGCTCTGGCGGTGGTTGTCCAGGCTCTGCAGAGCGCAG
CAGGGCTTTTCATTAAAGGTATTTATATTTGTA

317/392

FIGURE 304

MMRCCRRRCCCRQPPHALRPLLLLPLVLLPPLAAAAAGPNRCDTIYQGFAECLIRLGDSM
GRGGELETICRSWNDFHACASQVLSGCPEEAAAVWESLQQEARQAPRPNNLHTLCGAPVH
VRERGTGSETNQETLRATAPALPMAPAPPLLAAALALAYLLRPLA

Signal peptide:
Amino acids 1-35

Transmembrane domain:
Amino acids 141-157

N-myristoylation site:
Amino acids 127-133

Prokaryotic membrane lipoprotein lipid attachment site:
Amino acids 77-88

318/392

FIGURE 305

AAGTACTTGTGTCCGGGTGGTGGACTGGATTAGCTGCGGAGCCCTGGAAGCTGCCTGTCC
TTCTCCCTGTGCTTAACCAGAGGTGCCCATGGGTTGGACAATGAGGCTGGTCACAGCAGC
ACTGTTACTGGGTCTCATGATGGTGGTCACTGGAGACGAGGATGAGAACAGCCCGTGTGC
CCATGAGGCCCTCTTGGACGAGGACACCCTCTTTTGCCAGGGCCTTGAAGTTTTCTACCC
AGAGTTGGGGAACATTGGCTGCAAGGTTGTTCCCTGATTGTAACAACTACAGACAGAAGAT
CACCTCCTGGATGGAGCCGATAGTCAAGTTCCCGGGGGCCGTGGACGGCGCAACCTATAT
CCTGGTGATGGTGGATCCAGATGCCCCCTAGCAGAGCAGAACCAGACAGAGATTCTGGAG
ACATTGGCTGGTAACAGATATCAAGGGCGCCGACCTGAAGAAAGGAAGATTCAAGGCCA
GGAGTTATCAGCCTACCAGGCTCCCTCCCCACCGGCACACAGTGGCTTCCATCGCTACCA
GTTCTTTGTCTATCTTCAGGAAGGAAAAGTCATCTCTCTCCTTCCCAAGGAAAACAAAAC
TCGAGGCTCTTGGAAAATGGACAGATTTCTGAACCGCTTCCACCTGGGCGAACCTGAAGC
AAGCACCAGTTTCATGACCCAGAACTACCAGGACTACCAACCCTCCAGGCTCCAGAGG
AAGGGCCAGCGAGCCCAAGCACAAAACCAGGCAGAGATAGCTGCCTGCTAGATAGCCGGC
TTTGCCATCCGGGCATGTGGCCACACTGCTCACCACCGACGATGTGGGTATGGAACCCCC
TCTGGATACAGAACCCTTCTTTTCCAAATTAATAAAAAAAAAAATCATCAA

319/392

FIGURE 306

MGWTMRLVTAALLLGLMMVVTGDEDENSPCAHEALLDEDTLFCQGLEVFYPELGNIGCKV
VPDCNNYRQKITSWMEPIVKFPGAVDGATYILVMVDPDAPSRAEPRQRFWRHWLVTDIKG
ADLKKGKIQQGELSAYQAPSPPAHSGFHRYQFFVYLQEGKVISLLPKENKTRGSWKMDRF
LNRFHLGEPEASTQFMTQNYQDSPTLQAPRGRASEPKHKTRQR

Important features of the protein:

Signal peptide:

amino acids 1-22

N-glycosylation site:

amino acids 169-173

Tyrosine kinase phosphorylation site:

amino acids 59-68

N-myristoylation sites:

amino acids 54-60, 83-89, 130-136

Phosphatidylethanolamine signature:

amino acids 113-157

320/392

FIGURE 307

AAGGAGCAGCCCGCAAGCACCAAGTGAGAGGCATGAAGTTACAGTGTGTTTCCCTTTGGC
TCCTGGGTACAATACTGATATTGTGCTCAGTAGACAACCACGGTCTCAGGAGATGTCTGA
TTTCCACAGACATGCACCATATAGAAGAGAGTTTCCAAGAAATCAAAGAGCCATCCAAG
CTAAGGACACCTTCCCAAATGTCACCTATCCTGTCCACATTGGAGACTCTGCAGATCATT
AGCCCTTAGATGTGTGCTGCGTGACCAAGAACCTCCTGGCGTTCTACGTGGACAGGGTGT
TCAAGGATCATCAGGAGCCAAACCCCAAATCTTGAGAAAAATCAGCAGCATTGCCAACT
CTTTCCTCTACATGCAGAAAACCTGCGGCAATGTGAGAACAGAGGCAGTGTCACTGCA
GGCAGGAAGCCACCAATGCCACCAGAGTCATCCATGACAACCTATGATCAGCTGGAGGTCC
ACGCTGCTGCCATTAAATCCCTGGGAGAGCTCGACGTCTTTCTAGCCTGGATTAATAAGA
ATCATGAAGTAATGTTCTCAGCTTGATGACAAGGAACCTGTATAGTGATCCAGGGATGAA
CACCCCCTGTGCGGTTTACTGTGGGAGACAGCCACCTTGAAGGGGAAGGAGATGGGGAA
GGCCCCCTTGACAGCTGAAAGTCCCACTGGCTGGCCTCAGGCTGTCTTATTCCGCTTGAAAA
TAGGCAAAAAGTCTACTGTGGTATTTGTAATAAACTCTATCTGCTGAAAGGGCCTGCAGG
CCATCCTGGGAGTAAAGGGCTGCCTTCCCATCTAATTTATTGTAAAGTCATATAGTCCAT
GTCTGTGATGTGAGCCAAGTGATATCCTGTAGTACACATTGTACTGAGTGGTTTTTCTGA
ATAAATTCCATATTTTACCTATGA

321/392

FIGURE 308

MKLQCVSLWLLGTILILCSVDNHGLRRCLISTDMHHIEESFQEIKRAIQAKDTFPNVTIL
STLETLQIIKPLDVCCVTKNLLAFYVDRVFKDHQEPNPKILRKISSIANSFLYMQKTLRQ
CQEQRQCHCRQEATNATRVIHNDYDQLEVHAAAIKSLGELDVFLAWINKNHEVMFSA

Signal sequence:

amino acids 1-18

N-glycosylation sites:

amino acids 56-60, 135-139

cAMP- and cGMP-dependent protein kinase phosphorylation site:

amino acids 102-106

N-myristoylation site:

amino acids 24-30

Actinin-type actin-binding domain signature 1:

amino acids 159-169

322/392

FIGURE 309

GTCGACCCACGCGTCCGAAGCTGCTGGAGCCACGATTTCAGTCCCCTGGACTGTAGATAAA
GACCCTTTCTTGCCAGGTGCTGAGACAACCACACTATGAGAGGCACTCCAGGAGACGCTG
ATGGTGGAGGAAGGGCCGTCTATCAATCAATCACTGTTGCTGTTATCACATGCAAGTATC
CAGAGGCTCTTGAGCAAGGCAGAGGGGATCCCATTTATTTGGGAATCCAGAATCCAGAAA
TGTGTTTGTATTGTGAGAAGGTTGGAGAACAGCCACATTGCAGCTAAAAGAGCAGAAGA
TCATGGATCTGTATGGCCAACCCGAGCCCGTGAAACCCCTTCTTTTCTACCGTGCCAAGA
CTGGTAGGACCTCCACCCTTGAGTCTGTGGCCTTCCCGGACTGGTTCATTGCCTCCTCCA
AGAGAGACCAGCCCATCATTCTGACTTCAGAACTTGGGAAGTCATACAACACTGCCTTTG
AATTAAATATAAATGACTGA¹ACTCAGCCTAGAGGTGGCAGCTTGGTCTTTGTCTTAAAGT
TTCTGGTTCCCAATGTGTTTTCGTCTACATTTTCTTAGTGTCAATTTTCACGCTGGTGCTG
AGACAGGAGCAAGGCTGCTGTTATCATCTCATTTTATAATGAAGAAGAAGCAATTACTTC
ATAGCAACTGAAGAACAGGATGTGGCCTCAGAAGCAGGAGAGCTGGGTGGTATAAGGCTG
TCCTCTCAAGCTGGTGCTGTGTAGGCCACAAGGCATCTGCATGAGTGACTTTAAGACTCA
AAGACCAAACACTGAGCTTTCTTCTAGGGGTGGGTATGAAGATGCTTCAGAGCTCATGCG
CGTTACCCACGATGGCATGACTAGCACAGAGCTGATCTCTGTTTCTGTTTTGCTTTATTC
CCTCTTGGGATGATATCATCCAGTCTTTATATGTTGCCAATATACCTCATTGTGTGTAAT
AGAACCTTCTTAGCATTAAAGACCTTGTAACAAAAATAATTCCTGGGGTGGGTATGAAGA
TGCTTCAGAGCTCATGCGCGTTACCCACGATGGCATGACTAGCACAGAGCTGATCTCTGT
TTCTGTTTTGCTTTATTCCTCTTGGGATGATATCATCCAGTCTTTATATGTTGCCAATA
TACCTCATTGTGTGTAATAGAACCTTCTTAGCATTAAAGACCTTGTAACAAAAATAATTC
TTGTGTAAAGTTAAATCATTTTTGTCTAATTGTAATGTGTAATCTTAAAGTTAAATAAA
CTTTGTGTATTTATATAATAATAAGCTAAAACCTGATATAAAATAAAGAAAGAGTAAACTG

323/392

FIGURE 310

MRGTPGDADGGGRAVYQSITVAVITCKYPEALEQGRGDPIYLGIONPEMCLYCEKVGEQP
TLQLKEQKIMDLYGQPEPVKPFIFYRAKTGRTSTLESVAFPDWFIASSKRDPPIILTSEL
GKSYNTAFELNIND

Signal sequence:
amino acids 1-17

N-myristoylation site:
amino acids 10-16

Cell attachment sequence:
amino acids 36-39

324/392

FIGURE 311

GCGAGGCTGCACCAGCGCCTGGCACCATGAGGACGCCTGGGCCTCTGCCCCGTGCTGCTGC
TGCTCCTGGCGGGAGCCCCGCGCGCGGCCACTCCCCGACCTGCTACTCCCGCATGC
GGGCCCTGAGCCAGGAGATCACCCGCGACTTCAACCTCCTGCAGGTCTCGGAGCCCTCGG
AGCCATGTGTGAGATACCTGCCCAGGCTGTACCTGGACATACACAATTACTGTGTGCTGG
ACAAGCTGCGGGACTTTGTGGCCTCGCCCCGTGTTGGAAAGTGGCCCAGGTAGATTTCCT
TGAAGGACAAAGCACGGAAGCTGTACACCATCATGAACTCGTTCTGCAGGAGAGATTGG
TATTCCTGTTGGATGACTGCAATGCCTTGGAATACCCAATCCCAGTGACTACGGTCCTGC
CAGATCGTCAGCGCTTAAGGGAAGTGAAGACCAGAGAAAGAACCCAAGAGAACTAAAGTTAT
GTCAGCTACCCAGACTTAATGGGCCAGAGCCATGACCCTCACAGGTCTTGTGTTAGTTGT
ATCTGAAACTGTTATGTATCTCTACCTTCTGGAAAACAGGGCTGGTATTCCTACCCAG
GAACCTCCTTTGAGCATAGAGTTAGCAACCATGCTTCTCATTCCCTTGACTCATGTCTTG
CCAGGATGGTTAGATACACAGCATGTTGATTTGGTCACTAAAAAGAAGAAAGGACTAAC
AAGCTTCACTTTTATGAACAACTATTTTGAGAACATGCACAATAGTATGTTTTTTATTACT
GGTTTAATGGAGTAATGGTACTTTTTATTCTTTCTTGATAGAAACCTGCTTACATTTAACC
AAGCTTCTATTATGCCTTTTTTCTAACACAGACTTTCTTCACTGTCTTTCATTTAAAAAGA
AATTAATGCTCTTAAGATATATATTTTACGTAGTGCTGACAGGACCCACTCTTTCATTGA
AAGGTGATGAAAATCAAATAAAGAATCTCTTCACATGGA

325/392

FIGURE 312

MRTPGPLPVLLLLLAGAPAAPPTPPTCYSRMRALSQEITRDFNLLQVSEPSEPCVRYLPR
LYLDIHNYCVLDKLRDFVASPPCWKVAQVDSLKDKARKLYTIMNSFCRRDLVFLDDCNA
LEYPIPVTTVLPDRQR

Important features of the protein:

Signal peptide:

amino acids 1-19

Tyrosine kinase phosphorylation site:

amino acids 60-69

N-myristoylation site:

amino acids 16-22

326/392

FIGURE 313

GAGCGACGCTGTCTCTAGTCGCTGATCCCAAATGCACCGGCTCATCTTTGTCTACACTCT
AATCTGCGCAAACCTTTTGCAGCTGTCTGGGACACTTCTGCAACCCCGCAGAGCGCATCCAT
CAAAGCTTTGCGCAACGCCAACCTCAGGCGAGATGACTTGTACCGAAGAGATGAGACCAT
CCAGGTGAAAGGAAACGGCTACGTGCAGAGTCCTAGATTCCCGAACAGCTACCCCAGGAA
CCTGCTCCTGACATGGCGGCTTCACTCTCAGGAGAATACACGGATACAGCTAGTGTGTTGA
CAATCAGTTTGGATTAGAGGAAGCAGAAAATGATATCTGTAGGTATGATTTTGTGGAAGT
TGAAGATATATCCGAAACCAGTACCATTATTAGAGGACGATGGTGTGGACACAAGGAAGT
TCCTCCAAGGATAAAATCAAGAACGAACCAAATTAATACATTCAAGTCCGATGACTA
CTTTGTGGCTAAACCTGGATTCAAGATTTATTATTCTTTGCTGGAAGATTTCCAACCCGC
AGCAGCTTCAGAGACCAACTGGGAATCTGTCACAAGCTCTATTTCAAGGGTATCCTATAA
CTCTCCATCAGTAACGGATCCCACTCTGATTGCGGATGCTCTGGACAAAAAATTCAGAG
ATTTGATACAGTGAAGATCTGCTCAAGTACTTCAATCCAGAGTCATGGCAAGAAGATCT
TGAGAATATGTATCTGGACACCCCTCGGTATCGAGGCAGGTCATACCATGACCGGAAGTC
AAAAGTTGACCTGGATAGGCTCAATGATGATGCCAAGCGTTACAGTTGCACTCCCAGGAA
TTACTCGGTCAATATAAGAGAAGAGCTGAAGTTGGCCAATGTGGTCTTCTTTCCACGTTG
CCTCCTCGTGCAGCGCTGTGGAGGAAATTGTGGCTGTGGAAGTGTCAACTGGAGGTCCTG
CACATGCAATTCAGGGAAAACCGTGAAAAAGTATCATGAGGTATTACAGTTTGAGCCTGG
CCACATCAAGAGGAGGGGTAGAGCTAAGACCATGGCTCTAGTTGACATCCAGTTGGATCA
CCATGAACGATGCGATTGTATCTGCAGCTCAAGACCACCTCGATTAAGAGAATGTGCACAT
CCTTACATTAAGCCTGAGAGAA

327/392

FIGURE 314

MHRLIFVYTLCANFCSCRDTSATPQSASIKALRNANLRRDDLRYRDETIQVKNGYVQS
PRFPNSYPRNLLLTWRLHSQENTRIQLVFDNQFGLLEEAENDICRYDFVEVEDISETSTII
RGRWCGHKEVPPRIKSRTNQIKITFKSDDYFVAKPGFKIYYSLLEDFQPAAASETNWESV
TSSISGVSYNPSVTDPTLIADALDKKIAEFDTVEDLLKYFNPESWQEDLENMYLDTPRY
RGRSYHDRKSKVDLDRLNDDAKRYSCTPRNYSVNIREELKLANVVFFPRCLLVQRCGGNC
GCGTVNWRSTCNSGKTVKKYHEVLQFEPGHIKRRGRAKTMALVDIQLDHHERCDCICSS
RPPR

Signal peptide:
amino acids 1-18

N-glycosylation site:
amino acids 270-274

cAMP- and cGMP-dependent protein kinase phosphorylation site:
amino acids 262-266

Tyrosine kinase phosphorylation site:
amino acids 256-265

N-myristoylation sites:
amino acids 94-100, 186-192, 297-303, 298-304

TonB-dependent receptor proteins signature 1:
amino acids 1-56

328/392

FIGURE 315

CGGCTCGAGGCTCCCGCCAGGAGAAAGGAACATTCTGAGGGGAGTCTACACCCTGTGGAG
CTCAAGATGGTCCTGAGTGGGGCGCTGTGCTTCCGAATGAAGGACTCGGCATTGAAGGTG
CTTTATCTGCATAATAACCAGCTTCTAGCTGGAGGGCTGCATGCAGGGAAGGTCATTAAA
GGTGAAGAGATCAGCGTGGTCCCCAATCGGTGGCTGGATGCCAGCCTGTCCCCCGTCATC
CTGGGTGTCCAGGGTGGAAAGCCAGTGCCTGTCTGTGGGGTGGGGCAGGAGCCGACTCTA
ACACTAGAGCCAGTGAACATCATGGAGCTCTATCTTGGTGCCAAGGAATCCAAGAGCTTC
ACCTTCTACCGGCGGGACATGGGGCTCACCTCCAGCTTCGAGTCGGCTGCCTACCCGGGC
TGGTTCCTGTGCACGGTGCCTGAAGCCGATCAGCCTGTCTCAGACTCACCCAGCTTCCCGAG
AATGGTGGCTGGAATGCCCCCATCACAGACTTCTACTTCCAGCAGTGTGACTAGGGCAAC
GTGCCCCCAGAACTCCCTGGGCAGAGCCAGCTCGGGTGAGGGGTGAGTGGAGGAGACCC
ATGGCGGACAATCACTCTCTCTGCTCTCAGGACCCCCACGTCTGACTTAGTGGGCACCTG
ACCATTCTTGTCTTCTGCTTCCAGTTTGGATAAAATCTGAGATTGAGGCTCAGTCCACG
GTCTTCCCCACTGGATGGTGTCTACTGCTGTGGAACCTTGTA AAAACCATGTGGGGTAAA
CTGGGAATAACATGAAAAGATTTCTGTGGGGGTGGGGTGGGGGAGTGGTGGGAATCATTC
CTGCTTAATGGTAAGTACGACAAGTGTACCCCTGAGCCCCGAGGCCAACCCATCCCCAGTT
GAGCCTTATAGGGTCACTAGCTCTCCACATGAAGTCCTGTCACTACCACTGTGCAGGAG
AGGGAGGTGGTCATAGAGTCAGGGATCTATGGCCCTTGGCCAGCCCCACCCCTTCCCT
TTAATCCTGCCACTGTCTATGCTACCTTTCTCTATCTCTTCCCTCATCATCTTGTGTGG
GCATGAGGAGGTGGTGATGTCTCAGAAGAAATGGCTCGAGCTCAGAAGATAAAAGATAAGTA
GGGTATGCTGATCCTCTTTTAAAAACCCAAGATACAATCAAAATCCCAGATGCTGGTCTC
TATTTCCCATGAAAAAGTGTCTCATGACATATTGAGAAGACCTACTTACAAAGTGGCATATA
TTGCAATTTATTTTAAATTAAGATACCTATTTATATATTTCTTTATAGAAAAAGTCTG
GAAGAGTTTACTTCAATTGTAGCAATGTCTAGGGTGGTGGCAGTATAGGTGATTTTCTTT
TAATCTGTGTTAATTTATCTGTATTTCTTAATTTTCTACAATGAAGATGAATTCCTGTA
TAAAAATAAGAAAAGAAATTAATCTTGAGGTAAGCAGAGCAGACATCATCTCTGATTGTC
CTCAGCCTCCACTTCCCCAGAGTAAATTCAAATTGAATCGAGCTCTGCTGCTCTGGTTGG
TTGTAGTAGTGATCAGGAAACAGATCTCAGCAAAGCCACTGAGGAGGAGGCTGTGCTGAG
TTTGTGTGGCTGGAATCTCTGGGTAAAGGAATTAAGAAACAAAAATCATCTGGTAATCT
TTCTTAGAAGGATCACAGCCCCCTGGGATTCCAAGGCATTGGATCCAGTCTCTAAGAAGGC
TGCTGTACTGGTTGAATTTGTGTCCCCCTCAAATTCACATCCTTCTTGGAATCTCAGTCTG
TGAGTTTATTTGGAGATAAGGTCTCTGCAGATGTAGTTAGTTAAGACAAGGTCTATGCTGG
ATGAAGGTAGACCTAAATTCATATGACTGGTTTCTTGTATGAAAAGGAGAGGACACAG
AGACAGAGGAGACGCGGGGAAGACTATGTAAAGATGAAGGCAGAGATCGGAGTTTTCAG
CCACAAGCTAAGAAACACCAAGGATTGTGGCAACCATCAGAAGCTTGGAAGAGGCAAGA
AGAATCTTCCCTAGAGGCTTTAGAGGGATAACGGCTCTGCTGAAACCTTAATCTCAGAC
TTCCAGCCTCCTGAACGAAGAAAGATAAATTTGGCTGTTTTAAGCCACCAAGGATAAT
TGGTTACAGCAGCTCTAGGAACTAATACAGCTGCTAAAATGATCCCTGTCTCTCGTGT
TTACATTCTGTGTGTGTCCCCCTCCACAATGTACCAAAGTTGTCTTGTGACCAATAGAA
TATGGCAGAAGTGATGGCATGCCACTTCCAAGATTAGGTTATAAAAGACACTGCAGCTTC
TACTTGAGCCCTCTCTCTCTGCCACCCACCGCCCCCAATCTATCTTGGCTCACTCGCTCT
GGGGGAAGCTAGCTGCCATGCTATGAGCAGGCCTATAAAGAGACTTACGTGGTA AAAAAT
GAAGTCTCCTGCCACAGCCACATTAGTGAACCTAGAAGCAGAGACTCTGTGAGATAATC
GATGTTTGTGTTGTTTAAAGTTGCTCAGTTTGGTCTAACTTGTTATGCAGCAATAGATAAA
TAATATGCAGAGAAAGAG

329/392

FIGURE 316

MVLSGALCFRMKDSALKVLYLHNNQLLAGGLHAGKVIKGEESVVPNRWLDASLSPVILG
VQGSQCLSCGVGQEPTLTLEPVNIMELYLGAKESKSFTFYRRDMGLTSSFESAAYPGWF
LCTVPEADQPVRLTQLPENGGWNAPIITDFYFQQCD

N-myristoylation sites:

amino acids 29-34, 30-35, 60-65, 63-68, 73-78, 91-96, 106-111

Interleukin-1 signature:

amino acids 111-131

Interleukin-1 proteins:

amino acids 8-29, 83-120, 95-134, 64-103

330/392

FIGURE 317

ATGGAACTTGGACTTGGAGGCCTCTCCACGCTGTCCCACTGCCCCCTGGCCTAGGCGGCAG
CCTGCCCTGTGGCCCACCCTGGCCGCTCTGGCTCTGCTGAGCAGCGTCGCAGAGGCCTCC
CTGGGCTCCGCGCCCCGAGCCCTGCCCCCGCGAAGGCCCCCGCCTGTCTTGGCGTCC
CCCGCCGGCCACCTGCCGGGGGGACGCACGGCCCGCTGGTGCAGTGGAAGAGCCCGGCGG
CCGCCCGCGCAGCCTTCTCGGCCCCGCGCCCCGCGCCCTGCACCCCCATCTGCTCTTCCC
CGCGGGGGCCGCGCGGCGCGGGCTGGGGGCCCGGGCAGCCGCGCTCGGGCAGCGGGGGCG
CGGGGCTGCCGCTGCGCTCGCAGCTGGTGC CGGTGCGCGCGCTCGGCCTGGGCCACCGC
TCCGACGAGCTGGTGC GTTTCCGCTTCTGCAGCGGCTCCTGCCGCCGCGCGCTCTCCA
CACGACCTCAGCCTGGCCAGCCTACTGGGCGCCGGGGCCCTGCGACCGCCCCGGGCTCC
CGGCCCCGTCAGCCAGCCCTGCTGCCGACCCACGCGCTACGAAGCGGTCTCCTTCATGGAC
GTCAACAGCACCTGGAGAACCGTGGACCGCCTCTCCGCCACCGCCTGCGGCTGCCTGGGC
TGA

331/392

FIGURE 318

MELGLGGLSTLSHCPWPRRQPALWPTLAALALLSSVAEASLGSAPRSPAPREGPPPPVLAS
PAGHLPGGRTARWCSEGRARRPPPPQPSRPAPPPPPAPPSALPRGGRAARAGGPGSRARAAGA
RGCRLRSQ LVPVRALGLGHRSEDLVRFRFCSGSCRRARSPHDLSLASLLGAGALRPPPGS
RPVSQPCCRPTRYEAVSFMDVNSTWRTVDRLSATAACGCLG

signal sequence:

Amino acids 1-39

N-glycosylation site:

Amino acids 202-206

N-myristoylation sites:

Amino acids 6-12;67-73;102-108;109-115;119-125

332/392

FIGURE 319

GTTGCTATGTTGCCAGGCTGGTCTTGAAGTGCCTTGACCTCCTAAAGTGTGGAACCAC
AGACGTGAGCCACTCCACCCAGCCTAAACTTCATCTTCTTTGGATGAGATGAACACTTT
TAACAAGAGAACAGGACTCTATATAAATCGCTGTGGGCTCACCACCTCTAAGGAGGAGCA
CTGACTGAAGACAGAAAAATTGATGAACTGAAGAAGACATGGTCCATTATGCCTTACAAA
CTTACACAGTGCTTTGGGAATTCCAAAGTACTCAGTGGAGAGAGGTGTTTCAGGAGCCGT
AGAGCCAGATCGTCATCATGTCTGCATTGTGGCTGCTGCTGGGCCTCCTTGCCCTGATGG
ACTTGTCTGAAAGCAGCAACTGGGGATGCTATGGAAACATCCAAAGCCTGGACACCCCTG
GAGCATCTTGTGGGATTGGAAGACGTCACGGCCTGAACTACTGTGGAGTTTCGTGCTTCTG
AAAGGCTGGCTGAAATAGACATGCCATACCTCCTGAAATATCAACCCATGATGCAAACCA
TTGGCCAAAAGTACTGCATGGATCCTGCCGTGATCGCTGGTGTCTTGTCCAGGAAGTCTC
CCGGTGACAAAATTCTGGTCAACATGGGCGATAGGACTAGCATGGTGCAGGACCCCTGGCT
CTCAAGCTCCACATCCTGGATTAGTGAGTCTCAGGTTTCCCAGACAACTGAAGTTCTGA
CTACTAGAATCAAAGAAATCCAGAGGAGGTTTTCCAACCTGGACCCCTGACCAGTACCTGA
GAGGTGGACTCTGTGCCTACAGTGGGGGTGCTGGCTATGTCCGAAGCAGCCAGGACCTGA
GCTGTGACTTCTGCAATGATGTCCTTGCACGAGCCAAGTACCTCAAGAGACATGGCTTCT
AACATCTCAGATGAAACCCAAGACCATGATCACATATGCAGCCTCAAATGTTACACAGAT
AAAACTAGCCAAGGGCACCTGTAACCTGGGAATCTGAGTTTGACCTAAAAGTCATTAAAAT
AACATGAATCCCATTTAAAAAAAAAAAAAA

333/392

FIGURE 320

MSALWLLLGLLLALMDLSESSNWGCYGNIQSLDTPGASCGIGRRHGLNYCGVRASERLAEI
DMPYLLKYQPMQTIGQKYCMDPAVIAGVLSRKSPGDKILVNMGDRTSMVQDPGSQAPTS
WISESQVSQTTEVLTTRIKEIQRRFPTWTPDQYLRGGLCAYSGGAGYVRSSQDLSCDFCN
DVLARAKYLKRHGF

Important features of the protein:

Signal peptide:

amino acids 1-19

N-myristoylation sites:

amino acids 23-29, 26-32, 35-41, 45-51, 50-56, 76-82, 156-162

Amidation site:

amino acids 40-44

334/392

FIGURE 321

GCCTTATAAAGTAGCCTCTGCATCTGCCTGCCTCGGGCAGAGGAGGGCTACCCCTGGGGCT
GAGAGTTCACCTGTCTCAGGAACCACCTGAGCCACAGATCCTGTGGGCAGCGGCCAGGG
CAGCCATGGCTTGGGCAAGTAGGCTGGGCCTGCTGCTGGCACTGCTGCTGCCCGTGGTCG
GTGCCTCCACGCCAGGCACCGTGGTCCGACTCAACAAGGCAGCATTGAGCTACGTGTCTG
AAATTGGGAAAGCCCCCTCTCCAGCGGGCCCTGCAGGTCACTGTCCCTCATTTCTGGACT
GGAGTGGAGAGGCGCTTCAGCCCACCAGGATCCGGATTCTGAATGTCCATGTGCCCCGCC
TCCACCTGAAATTCATTGCTGGTTTCGGAGTGCGCCTGCTGGCAGCAGCTAATTTTACTT
TCAAGGTCTTTTCGCGCCCCAGAGCCCCCTGGAGCTGACGCTGCCTGTGGAAGTGTGGCTG
ACACCCGCGTGACCCAGAGCTCCATCAGGACCCCTGTGGTCAGCATCTCTGCCTGCTCTT
TATTCTCGGGCCACGCCAACGAGTTTGATGGCAGTAACAGCACCTCCCACGCGCTGCTGG
TCCTGGTGAGAGCACATTAAAGCTGTCTTGAGTAACAAGCTGTGCCTGAGCATCTCCA
ACCTGGTGAGGGTGTCAATGTCCACCTGGGCACCTTAATTGGCCTCAACCCCGTGGGTCT
CTGAGTCCCAGATCCGCTATTCCATGGTCAGTGTGCCCACTGTCAACAGTGACTACATTT
CCCTGGAAGTCAATGCTGTTCTCTTCTGCTGGGCAACCCCATCATCCTGCCCACGGATG
CCACCCCTTTTGTGTTGCCAAGGCATGTGGGTACCGAGGGCTCCATGGCCACCGTGGGCC
TCTCCCAGCAGCTGTTTGAAGTCTGCGCTCCTGCTGCTGCAGAAGGCCGGTGCCCTCAACC
TGGACATCACAGGGCAGCTGAGGTGCGATGACAACCTGCTGAACACCTCTGCTCTGGGCC
GGCTCATCCCGAGGTGGCCCGCCAGTTTCCCGAGCCCATGCCTGTGGTGCTCAAGGTGC
GGCTGGGTGCCACACCTGTGGCCATGCTCCACACAAACAACGCCACCCTGCGGCTGCAGC
CCTTCGTGGAGGTCCCTGGCCACAGCCTCCAACCTCGGCTTTCCAGTCCCTCTTCTCCCTGG
ATGTGGTAGTGAAGTTGAGACTCCAGCTCTCTGTGTCCAAGGTGAAGCTTCAGGGGACCA
CGTCTGTGCTGGGGGATGTCCAGCTCACGGTGGCCTCCTCCAACGTGGGCTTCATTGATA
CAGATCAGGTGCGCACACTGATGGGCACCGTTTTTGAGAAGCCCCCTGCTGGACCATCTCA
ATGCTCTCTTGGCCATGGGAATTGCCCTCCCTGGTGTGGTCAACCTCCACTATGTTGCC
CTGAGATCTTTGTCTATGAGGGCTACGTGGTGATATCCAGTGGACTCTTCTACCAGAGCT
GAGGCAAGACCACTGGGAGGCCTGAGAGTGGGCCAGCTCGCTGCTCAGGCGAATTTCTCA
TTTCAAGCCACTGGGGAAACTGAGGCAAAACCATACTTAGTCATCACCAACAAGCTGGAC
TGCTTAGCTGGGCTGTTTTATCTTCCCTGAGTGCCTGGGTCTCCCTCCCTCACTTCTGCC
CTTTCCCTTCCCTCCTCCTTCTCCTCCCTCTTCCCTCATCTCCCCCTCCTTCCCTCTGC
CCCACCCAGGGGGAGCAGACTGCTCCTCCAGGCTGTATAGACCTGCCCTCTTGCAATTA
AACAACTTCTCTTGAGCTGC

335/392

FIGURE 322

MAWASRLGLLLALLLPVVGASTPGTVVRLNKAALSYVSEIGKAPLQRALQVTVPHFLDWS
GEALQPTRIRILNVHVPRLHLKFIAGFGVRLLAANFTFKVFRAPEPLELTPVELLADT
RVTQSSIRTPVVSISACSLFSGHANEFDGSNSTSHALLVLVQKHIKAVLSNKLCLSISNL
VQGVNVHLGTLIGLNPVGPESQIRYSMVSVPVTSDYISLEVNAVLFLLGNPIILPTDAT
PFVLPRHVGTEGSMATVGLSQQLFDSALLLLQKAGALNLDITGQLRSDDNLLNTSALGRL
IPEVARQFPEPMPVVLKVRLGATPVAMLHTNNATLRLQPFVEVLATASNSAFQSLFSLDV
VVLNRLQLSVSKVKLQGTTSVLGDVQLTVASSNVGFIDTDQVRTLMGTVFEKPLLDHLNA
LLAMGIALPGVVNLHYVAPEIFVYEGYVVISSGLFYQS

Important features of the protein:**Signal peptide:**

Amino acids 1-20

Transmembrane domain:

Amino acids 217-236

N-glycosylation sites:

Amino acids 96-100;151-155;293-297;332-336

N-myristoylation sites:

Amino acids 8-14;149-155;189-195;249-255;252-258;283-289

LBP / BPI / CETP family proteins:

Amino acids 22-50; 251-287

336/392

FIGURE 323

TTGAAAATCTACTCTATCAGCTGCTGTGGTTGCCACCATTCTCAGGACCCTCGCCATGAA
AGCCCTTATGCTGCTCACCTGTCTGTTCTGCTCTGCTGGGTCTCAGCTGACATTGCTG
TCACTCCTGCTACAAGGTCCCTGTGCTGGGCTGTGTGGACCGGCAGTCCTGCCGCCTGGA
GCCAGGACAGCAATGCCTGACAACACATGCATACCTTGGTAAGATGTGGGTTTTCTCCAA
TCTGCGCTGTGGCACACCAGAAGAGCCCTGTCAGGAGGCCTTCAACCAAACCAACCGCAA
GCTGGGTCTGACATATAACACCACCTGCTGCAACAAGGACAAGTCAACAGCGCAGGACC
CCGGCCCACTCCAGCCCTGGGCCTTGTCTTCCTTACCTCCTTGGCTGGCCTTGGCCTCTG
GCTGCTGCACTGAGACTCATTCCATTGGCTGCCCCCTCCTCCACCTGCCTTGGCCTGAGC
CTCTCTCCCTGTGTCTCTGTATCCCCTGGCTTTACAGAATCGTCTCTCCCTAGCTCCCAT
TTCTTTAATTAAACACTGTTCCGAGTGGTCTCCTCATCCATCCTTCCCACCTCACACCCT
TCACTCTCCTTTTTCTGGGTCCCTTCCCACTTCTTCCAGGACCTCCATTGGCTCCTAGA
AGGGCTCCCCACTTTGCTTCCCTATACTCTGCTGTCCCCTACTTGAGGAGGGATTGGGATC
TGGGCCTGAAATGGGGCTTCTGTGTTGTCCCCAGTGAAGGCTCCCACAAGGACCTGATGA
CCTCACTGTACAGAGCTGACTCCCCAAACCCAGGCTCCCATATGTACCCCATCCCCATA
CTCACCTCTTTCCATTTTGAGTAATAAATGTCTGAGTCTGGAAAAAAAAAAAAAAAAAAAA

337/392

FIGURE 324

MKALMLLTLSVLLCWVSADIRCHSCYKVPVLGCVDROSCRLEPGQQCLTTHAYLGKMWVF
SNLRCGTPEEPCQEAFNQTNRKLGTYNTTCCNKDNCNSAGPRPTPALGLVFLTSLAGLG
LWLLH

Important features of the protein:

Signal peptide:

amino acids 1-18

N-glycosylation sites:

amino acids 77-81, 88-92

N-myristoylation site:

amino acids 84-90

Ly-6 / u-PAR domain protein signature:

amino acids 85-98

338/392

FIGURE 325

ACGGGCCG CAGCGGCAGTGACGTAGGGTTGGCGCAGCGATCCGTTGCGGCTGCAGCTCTG
CAGTCGGGCCGTTCCCTTCGCCGCCGCCAGGGGTAGCGGTGTAGCTGCGCAGCGTCGCGCG
CGCTACCGCACCCAGGTTCCGGCCCGTAGGCGTCTGGCAGCCCGGCCATCTTCATCGAG
CGCCATGGCCG CAGCCTGCGGGCCGGGAGCGGCCGGGTACTGCTTGCTCCTCGGCTTGCA
TTTGTTTCTGCTGACCGCGGGCCCTGCCCTGGGCTGGAACGACCCTGACAGAATGTTGCT
GCGGGATGTAAAAGCTCTTACCCTCCACTATGACCGCTATACCACCTCCCGCAGGCTGGA
TCCCATCC CACAGTTGAAATGTGTTGGAGGCACAGCTGGTGTGATTCTTATACCCAAA
AGTCATACAGTGT CAGAACAAGGCTGGGATGGGTATGATGTACAGTGGGAATGTAAGAC
GGACTTAGATATTGCATACAAATTTGGAAAACTGTGGTGAGCTGTGAAGGCTATGAGTC
CTCTGAAGACCAGTATGTACTAAGAGGTTCTTGTGGCTTGGAGTATAATTAGATTATAC
AGAACTTGGCCTGCAGAACTGAAGGAGTCTGGAAAGCAGCACGGCTTTGCCTCTTCTC
TGATTATTATTATAAGTGGTCCTCGGCGGATTCTGTAAACATGAGTGGATTGATTACCAT
CGTGGTACTCCTTGGGATCGCCTTTGTAGTCTATAAGCTGTTCTGAGTGACGGGCAGTA
TTCTCCTCCACCGTACTCTGAGTATCCTCCATTTTCCACCGTTACCAGAGATTCACCAA
CTCAGCAGGACCTCCTCCCCCAGGCTTTAAGTCTGAGTTCACAGGACCACAGAATACTGG
CCATGGTGCAACTCTGGTTTTGGCAGTGCTTTTACAGGACAACAAGGATATGAAAATTC
AGGACCAGGGTTCTGGACAGGCTTGGGAAGTGGTGAATACTAGGATATTTGTTTGGCAG
CAATAGAGCGGCAACACCCTTCTCAGACTCGTGGTACTACCCGTCCTATCCTCCCTCCTA
CCCTGGCAGTGGAATAGGGCTTACTCACCCCTTCATGGAGGCTCGGGCAGCTATTCGGT
ATGTTCAAACCTCAGACACGAAAACCAGAACTGCATCAGGATATGGTGGTACCAGGAGACG
ATAAAGTAGAAAGTTGGAGTCAAACACTGGATGCAGAAATTTTGGATTTTTCATCACTTT
CTCTTTAGAAAAAAGTACTACCTGTTAACAATTGGGAAAAGGGGATATTCAAAGTTCT
GTGGTGTTATGTCCAGTGTAGCTTTTGTATTCTATTATTTGAGGCTAAAAGTTGATGTG
TGACAAAATACTTATGTGTTGTATGTCAGTGTAAACATGCAGATGTATATTGCAGTTTTTG
AAAGTGATCATTACTGTGGAATGCTAAAAATACATTAATTTCTAAACCTGTGATGCCCT
AAGAAGCATTAAGAATGAAGGTGTTGTACTAATAGAACTAAGTACAGAAAATTTTCAGTT
TTAGGTGGTTGTAGCTGATGAGTTATTACCTCATAGAGACTATAATATTCTATTTGGTAT
TATATTATTTGATGTTTGCTGTTCTTCAAACATTTAAATCAAGCTTTGGACTAATTATGC
TAATTTGTGAGTTCTGATCACTTTTGGAGCTCTGAAGCTTTGAATCATTCAGTGGTGGAGA
TGGCCTTCTGGTAACTGAATATTACCTTCTGTAGGAAAAGGTGGAAAATAAGCATCTAGA
AGGTTGTGTGAATGACTCTGTGCTGGCAAAAATGCTTGAAACCTCTATATTTCTTTCGT
TCATAAGAGGTAAAGGTCAAATTTTTCAACAAAAGTCTTTTAATAACAAAAGCATGCAGT
TCTCTGTGAAATCTCAAATATTGTTGTAATAGTCTGTTTCAATCTTAAAAAGAATCA

339/392

FIGURE 326

MAAACGPGAAGYCLLLGLHLFLLTAGPALGWNDPDRMLLRDVKALTLHYDRYTTSRRLLDP
IPQLKCVGGTAGCDSYTPKVIQCQNKGDGYDVQWECKTDLDIAYKFGKTVVSCEGYESS
EDQYVLRGSCGLEYNLDYTELGLOKLKESGKQHGFASFSDYYYKWSSADSCNMSGELITIV
VLLGIAFVVYKLFLSDGQYSPPPYSEYPPFSHRYQRFTNSAGPPPPGFKSEFTGPQNTGH
GATSGFGSAFTGQQGYENSGPGFWTGLGTGGILGYLFGSNRAATPFSDSWYYPSYPPSY
GTWNRAYSPLHGSGSYSVCSNSDTKTRTASGYGGTRRR

Signal peptide:
amino acids 1-30

Transmembrane domain:
amino acids 171-190

N-glycosylation site:
amino acids 172-176

Glycosaminoglycan attachment sites:
amino acids 244-248, 259-263, 331-335

Tyrosine kinase phosphorylation site:
amino acids 98-106

N-myristoylation sites:
amino acids 68-74, 69-75, 131-137, 241-247, 247-253, 266-272,
270-276, 278-284, 312-318

340/392

FIGURE 327

GGCACGAGGTGGAAGGGCTTTTACAAACAGATTGCTGGCCCCACCCCCCAGAATTTCTCA
TCAGGAGTGGGCAAGACCAATCATTTGCATTTCTGACAAGTTCCCAGGAGCTGCAGCTGC
TGGCCCTGGAACCACACTTTGAGAACCACTGCTTTAGACCAAACACCAAAGGAAGATGCA
GCCACCCTCCTTTACATGTCAACAACGCTCAGGGTCCATGAGTACCTCAGGCTGTCCAGCT
GAGCTCCACCTGCAGCAGCCGAGATTCCCGACTCGCTCCACCATTTGGGGGCTAGGAGTGA
AGCGTGTCAACATGGTCAGCTCATGGCCAGCCAGGAAAGCCTCTCTGCTGTGCGTCTGTG
CAGTTCTTGTTCTTCCCTGGAGGACTCTTGATCGCCTGTGATCTTGGCCAGGAGACCAG
GTGCCTGGGTCCCTTCCTGGAAGGGGACAAGTTACACACCCAGCCCCATTTCCACCA
ACTTCTACATGCCTTGGGAGAACCCTTCTACATGTTGGCTGCCCCCTTCCCTATTTCAGC
AGTGCCAGTCCTGCTTATAAACCTGAGGCCTGCTCCCATACCTTCCCTGTGCAAGTGC
CAGCCGTTATTCCAGGCAGCCCAATGTTGTTGAGGCCAGATGGATTCCTGGAAGCAGCTG
GCCCATGGATGTGAGTCATCACAGTATTCTAGAAACAGAGAAGAGGTCTTAACCTAATGC
GCATAGAGAAATTGTTCTCATTGTAAACATAACCCCTGTCCTTAGCTGATCTAGGTGGAAG
CCCAGCTTCATGTGCTAGGGGGCATGATAATGATAATAAGGAATTGTATCTAGGACTAA

341/392

FIGURE 328

MVSSWPARKASLLCVC AVLVL PWRTL GSPVILARRPGAWVPSWKGTSYTPQPHFPTNFYM
PWENLLHVGCP LPLFQQCPVLLINLRPAPHTFPVQVPAVIPGSPMLLRPDGFLEAAGPWM

Signal peptide:
amino acids 1-27

cAMP- and cGMP-dependent protein kinase phosphorylation site:
amino acids 8-12

342/392

FIGURE 329

CAAAGAGTAGTCAGTCCCTTCTTGGCTCTGCTGACACTCGAGCCCACATTCCATCACCTG
CTCCCAATCATGCAGGTCTCCACTGCTGCCCTTGCCGTCCTCCTCTGCACCATGGCTCTC
TGCAACCAGGTCTCTCTGCACCACTTGCTGCTGACACGCCGACCGCCTGCTGCTTCAGC
TACACCTCCCGACAGATTCCACAGAATTTATAGCTGACTACTTTGAGACGAGCAGCCAG
TGCTCCAAGCCCAGTGTCATCTTCCTAACCAAGAGAGGCCGGCAGGTCTGTGCTGACCCC
AGTGAGGAGTGGGTCCAGAAATACGTCAGTGACCTGGAGCTGAGTGCCGTGAGGGGTCCAG
AAGCTTCGAGGCCCGACGACCTCAGTGGGCCAGTGGGGAGGAGCAGGAGCCTGAGCCTT
GGGAACATGCGTGTGACCTCTACAGCTACCTCTTCTATGGACTGGTTATTGCCAAACAGC
CACACTGTGGGACTCTTCTTAACTTAAATTTAATTTATTTATACTATTTAGTTTTTATA
ATTTATTTTTTGATTTACAGTGTGTTTGTGATTGTTTGCTCTGAGAGTTCCCCCTGTCCC
CTCCCCCTTCCCTCACAGTGTGTCTGGTGACAACCGAGTGGCTGTCATCGGCCTGTGTAG
GCAGTCATGGCACCAAAGCCACCAGACTGACAAATGTGTATCAAATGCTTTTGTTCAGGG
CTGTGATCGGCCTGGGGAAATAATAAGATGTTCTTTTAAACGGTAAAAAA

343/392

FIGURE 330

MQVSTAALAVLLCTMALCNQVLSAPLAADTPTACCFSYTSRQIPQNFIADYFETSSQCSK
PSVIFLTKRGRQVCADPSEEWVQKYVSDLELSA

Signal sequence:

1-23

Small cytokines (intercrine/chemokine) C-C subfamily
signature:

1-35, 2-36, 10-44, 34-74, 50-90

Small cytokines (intecrine/chemokine):

24-89

344/392

FIGURE 331

GGCACGAGGTGAGACTTTAAATGAAATGTCTCACAAAGCTAGGTGATCCAGGTTTTGTGGT
CTTTGCAACCCTTGTGGTCATTGTGGCCTTGATATTAATCTTCGTGGTGGGTCCCTCGCCA
TGGACAGACAAACATTCTTGTGTACATAACAATCTGCTCTGTAATCGGCGCGTTTTTCAGT
CTCCTGTGTGAAGGGCCTGGGCATTGCTATCAAGGAGCTGTTTGCAGGGAAGCCTGTGCT
GCGGCATCCCCTGGCTTGGATTCTGCTGCTGAGCCTCATCGTCTGTGTGAGCACACAGAT
TAATTACCTAAATAGGGCCCTGGATATATTCAACACTTCCATTGTGACTCCAATATATTA
TGTATTCTTTACAACATCAGTTTTAACTTGTTTCACTATTCTTTTTAAGGAGTGGCAAGA
TATGCCTGTTGACGATGTCATTGGTACTTTGAGTGGCTTCTTTACAATCATTGTGGGGAT
ATTCTTGTTCATGCCTTTTAAAGACGTCAGCTTTAGTCTAGCAAGTCTGCCTGTGTCTTT
TCGAAAAGACGAGAAAGCAATGAATGGCAATCTCTCTAATATGTATGAAGTCTTAATAA
TAATGAAGAAAGCTTAACCTGTGGAATCGAACAACACACTGGTGAAAATGTCTCCCGAAG
AAATGGAAATCTGACAGCTTTTTTAAGAAAGGTGTAATTAAAGGTAAATCTGTGATTGTTA
TGAAGTGAATTTGAATATCATCAGAATGTGTCTGAAAAAACATTGTCCTCAAATAATGTT
CTTTAAAGGCAATCTTTTTAAAGATTTCACTAATTTGGACCAAGAAATTACTTTTCTTGT
ATTTAAACAAACAATGGTAGCTCACTAAAATGACCTCAGCACATGACGATTTCTATTAAC
ATTTTATTGTTGTAGAAGTATTTTACATTTTTCATCCCTTCTCCAAAAGCCGAATGACTA
ATGACAGTTTTAAGTCTATGAAAATGCTTTATTTTTTTCATTGGTGATGAAAGTCTGAAAT
GTGCATTTGTTCATCCCCACTCCATCAATCCCTGACCATGTAAGGCTTTTTTATTTTAAAA
AAACAGAGTTATCCAATACATTATCCTGTGATTTACCTTACCTACAAAAGTGGCTCCTG
TTTGTGTTGATGATGATTGGTTTTATTTTTGAAATATTTATTAAGGGAAAACTAAGTTACT
GAATGAAGGAACCTCTTTCTTACAAAACAAAAAAGGGCAGAAATCACCCCAAGGAACG
ATTTCTCAGGTTGAGATGATCACCGTGAATCCGGCTTCCTCTGAGCATTGATGGCCTTA
GCACCTCATCAAGCCAGCACATCCTGCCTGCTGTTGCAGCCTGGCTGGGTTTATTCTTCA
GTTACCTAATCCCATGATGCCTGGAACCTTGATTACCGTTTTTACATCAGCTCTTGTACT
TTTCAGTATATTTTTCATAATGAGTTATATTGTCAATTTAGACTTTGAACAGCTCTGGGAAA
TAGAAGACTAGGGTTGTTTCTTAAATTTAGCTCATGTTATAATAAAAAGTTGAAATG

345/392

FIGURE 332

MSHKLGDPGFVVFATLVVIVALILIFVVGPRHGQTNILVYITICSVIGAFSVSCVKGLGI
AIKELFAGKPVLRHPLAWILLLSLIVCVSTQINYLNRLDIFNTSIVTPIYYVFFTTSVL
TCSAILFKEWQDMPVDDVIGTLSGFFTIIVGIFLLHAFKDVSFSLASLPVSFRKDEKAMN
GNLSNMYEVLNNNEESLTCGIEQHTGENVSRRNGNLTAF

Signal sequence:

1-33

Transmembrane domain:

40-60, 70-90, 103-123, 139-159

N-glycosylation site:

103-106, 182-185, 208-211, 215-218

N-myristoylation site:

57-62, 140-145, 181-186, 214-219

346/392

FIGURE 333

GTGATGGCGGCTGGTGATGGGGACGTGAAGCTAGGCACCCTGGGGAGTGGCAGCGAGAGC
AGCAACGACGGCGGCAGCGAGAGTCCAGGCGACGCGGGAGCGGCAGCGGAAGGGGGAGGC
TGGGCGGCGGCGGCGTTGGCGCTTCTGACGGGGGGCGGGGAAATGCTGCTGAACGTGGCG
CTGGTGGCTCTGGTGCTGCTGGGGGCCCTACCGGCTGTGGGTGCGCTGGGGGCGGCGGGGT
CTGGGGGCGGGGCGGGGCGGGGCGAGGAGAGCCCCGCCACCTCTCTGCCTCGCATGAAG
AAGCGGGA¹CTTCAGCTTGGAGCAGCTGCGCCAGTACGACGGCTCCCGCAACCCGCGCATC
CTGCTCGCGGTCAATGGGAAAGTCTTCGACGTGACCAAAGGCAGCAAGTTCTACGGCCCCG
GCGGGTCCATATGGAATATTTGCTGGTAGGGATGCCTCCAGAGGACTGGCCACATTTTGC
CTAGATAAAGATGCACCTTAGAGATGAATATGATGATCTCTCAGATTTGAATGCAGTACAA
ATGGAGAGTGTTGCGAGAATGGGAAATGCAGTTTAAAGAAAAATATGATTATGTAGGCAGA
CTCCTAAAACCAGGAGAAGAACCATCAGAATATACAGATGAAGAAGATACCAAGGATCAC
AATAAACAGGATTGAACTTTGTAAACAACCAAAGTCAGGGGCCTTCAGAACTGCAATTCT
TACTCCCTTTCACAGACTGTCCGGAGTCTTTGGGTTTGATTACCTGCTGCGAAAAACAT
TCAACAAATTGTGTACAAGATAAATTAATCTCACTATGAAGATTTGAATAACTAGACATT
ATTTATGCTGCCAAACTCATTGTGTCAGTTGTTTGTAAATGTCTAGTGGGGCTTCATCAT
CCTGAAAAGAAGGAGACAGGGATTTTTTTAAAGAGCAAGAAAGTCACAATATTACTTCTT
TCCTTCCTTTTTTCCTTCTTTCTTTCTTTCTTTCTTTCTTTCTTTTAAATATATTG
AAGACAACCAGATATGTATTTGCTACTCAAGTGTACAGATCTCCTCAAGAAACATCAAGG
G

347/392

FIGURE 334

MAAGDGDVKLGTLGSGSESSNDGGSESPGDAGAAAEGGGWAAAALALLTGGGEMLLNVAL
VALVLLGAYRLWVRWGRRGLGAGAGAGEESPATSLPRMKKRDFSLEQLRQYDGSRNPRIL
LAVNGKVFDVTKGSKFYGPAGPYGIFAGRDASRGLATFCLDKDALRDEYDDLSDLNAVQM
ESVREWEMQFKEKYDYVGRLLKPGEEPSEYTDDEEDTKDHNKQD

Signal sequence:

None

Transmembrane domain:

45-65

Tyrosine kinase phosphorylation site:

202-210

N-myristoylation site:

11-16, 16-21, 37-42, 38-43, 79-84, 81-86, 83-88, 144-149

Amidation site:

75-78

348/392

FIGURE 335

[illegible]

349/392

FIGURE 336

TGRGYCGDHESFSGAMEEPGATPQPYLGLLLEELRRVVAALPEGMRPDSNLYGFPWELVI
CAAVVGFFAVLFFLWRSFRSRSRLYVGREKKLALMLSGLIIEKSKLLEKFSLVQKEYEG
YEVESLKDASFEKEATEAQSLEATCEKLNRSNSELEDEILCLEKELKEEKSKHSEQDEL
MADISKRIQSLEDESKSLKSQVAEAKMTFQIFQMNEERLKIAIKDALNENSQLOESQKQL
LQEAENVWKEQVSELNKQKVTTFEDSKVHAEQVLNDKESHIKTLTERLLKMKDWAAMLGEDI
TDDDNLELEMNSESENGAYLDNPPKGALKKLIHAAKLNASLKTLEGERNQIYIQLSEVDK
TKEELTEHIKNLQTQQASLQSENFHFNENQKLQQKLKVMTELYQENEMKLHRKLTVEEN
YRLEKEEKLSKVDEKISHATEELETYRKRAKDLLEELERTIHSYQGQIISHEKKAHDNWL
AARNAERNLNDLRKENAHNRQKLTETELKFELLEKDPYALDVPNTAFGRGSRGPGNPLDH
QITNERGESSCDRLTDPHRAPSDTGSLSPWDQDRRMMFPPPGQSYPDALPPQRQDRFC
SNSGRLSGPAELRSFNMPSLDKMDGSMPSSEMRNDTKDDLGNLNVDPSSLPAENEATG
PGFVPPPLAPIRGPLFPVDARGPFLRRGPPFPFPPPGAMFGASRDYFPPRDFPGPPPAPF
AMRNVYPPRGFPFPPYLPPRPGFFPPPPHSEGRSEFFPSGLIPPSNEPATEHPEPQQET

Signal sequence:

None

Transmembrane domain:

54-74

N-glycosylation site:

150-153, 338-341, 636-639

cAMP- and cGMP-dependent protein kinase phosphorylation site:

413-416

Tyrosine kinase phosphorylation site:

414-421

N-myristoylation site:

466-417, 625-630, 697-702

Leucine zipper pattern:

142-163

350/392

FIGURE 337

GGACTGCGGTCTCGGGCAGCAATGGCCGAGAAGCGCGACACACGGGACTCCGAAGCCCAG
CGGCTCCCCGACTCCTTCAAGGACAGCCCCAGTAAGGGCCTTGGACCTTGC GGATGGATT
TTGGTGGCGTTCTCATTCTTATTCACCGTTATAACTTTCCCAATCTCAATATGGATGTGC
ATAAAGATTATAAAAGAGTATGAAAGAGCCATCATCTTTAGATTGGGTCGCATTTTACAA
GGAGGAGCCAAAGGACCTGGT TTTGTTTTTATTCTGCCATGCACTGACAGCTTCATCAAA
GTGGACATGAGAACTATTTCA TTTGATATTCTCTCCTCAGGAGATCCTGACAAAGGATTCA
GTGACAATTAGCGTGGATGGTGTGGTCTATTACCGCGTTCAGAATGCAACCCCTGGCTGTG
GCAAATATCACCAACGCTGACTCAGCAACCCGTCTTTTGGCACAACTACTCTGAGGAAT
GTTCTGGGCACCAAGAATCTTTCTCAGATCCTCTCTGACAGAGAAGAAATTGCACACAAC
ATGCAGTCTACTCTGGATGATGCCACTGATGCCTGGGGAATAAAGGTGGAGCGTGTGGAA
ATTAAGGATGTGAACTACCTGTGCAGCTCCAGAGAGCTATGGCTGCAGAACGAGGCG
TCCCGCGAGGCCCCGCGCCAAGGTTATTGCAGCCGAAGGAGAAATGAATGCATCCAGGGCT
CTGAAAGAAGCCTCCATGGTCACTGAATCTCCTGCAGCCCTTCAGCTCCGATACCTG
CAGACACTGACCACCATTGCTGCTGAGAAAACTCAACAATTGTCTTCCCTCTGCCATA
GATATGCTGCAAGGAATCATAGGGGCAAAACACAGCCATCTAGGCTAGTGTAGAGATGAG
CGCTAGCCTTCCAAGCATGAAGTCGGGGACCAAATTAGCCTTTAACTCATAAAGAGAGGG
TAGGGCTTTTCTTTTCCATATGTCAATTGTGGTGTTC CAGAATGTATAGCAGTTATAA
AAATAGGTGAAAGAATTGTTAGCTTGTAATACTGAGAGATTGGTGATTTATATAAGGTA
ATCTGTTAGTCTTAAATAGTTAAAAGTTTGATTTT TAGATTATTATGTAGTAGGTTAG
ATCCCTCTTGTTTGA CTTCCTGACTCATTCTGAACCCCTAAGCACCAGGCCACAG
GCAAGAACCTGGGCTGTA ACTGCCACCTGACACCGCTGACTGGCTAAATGCTTTGCAGAA
AGTGATGACCTTACACCACAACCAGCTTCTCCAGGT CATATGTGCCTTACCTCCAGAAGT
CTTTT TTTT TTTT TTTT TTTTCTGAGATGGAGTTTCACTCTTGTTGCCAGGCTGGAGTGCAA
TAGCATGATCTCGGCTCACTGCAACCTCCGCTCCTGGGTTCAAGAGATTCTCCTGCCTC
AGCCTCCCCAGTAGCTGGGATTACAGGCTCATGCCACCATGCCAGCTAATTTTTGTATT
ATTATTATTGTTTTTTAGTAGAGACGGGTTTCACCATGTTGGCCAGGCTAGTCACGAAC
TCCTAACCTCAGGTGATCCACCCACCTCTGCCTCCAAAGTGCTGGATTACAGGCTGAGCT
ACCACCTGGT TTTGGAGAGTCTTAATTAATTGAAATTTCCCTAATGTTTCA TTTTCT
AAATCCAGCCGTGTTTCAGAATAATCCTTACTTGAGAGTAGCCATTTTCTTGTTACTTG
TCAGAACTAGAGGAAATAGCCAAGACTAATGAAAAACATTACTCTAACCTTAAAAGACT
TTTAAATTCACTACTAGAGTGGTCATTTTAAAATACATCCATGTTTTAACTTATTTTGA
GCCTTCTTTTATGAGTAAATGATTCCTCCTTGTTCTGTCTTTCAAACCAGCTAAATATT
TGTACAAAAAGTGACTTTTTCTCACTGTGCCTATTTTCATATATCAGGTTTTAAATAG
TTTTAATTTTTTAATAAAATTTTTCTCTACGTTCTATATGCAATTGTTATATATCTATTT
GAATAGCTGAAGGACTAAAATACTTTTTTAAGAGATAA CTTCAGGAAACCATTATATTTT
ACTATCTGCATGCTGT TAACTGTGGTACACTGTGAAATATGTTGATTACAAACCCATTCA
TTACATAGTATAAGGAATTCACAGTATATGACTATATAGTGTCTAATGACTGGGCAGAT
ACTGTCAACTTACAATATCTATATAGAGAGGCTTTAAACTTACCTTACTCATTTCTCTATG
ATGTATGACTTGATGCTGAAAGAGGAAGCTGGTCAGCTCCTCATGGACAACAAATTCCTTA
GTCTATAATATTAGGAGACATCTCTAGTTTTTGCAAATGTCTGTGAATCTGAGCAACCTGG
ACTTCTGCTTACTGGCCAGAAAGCTGGCGGGTGACATTTGTAACATTTCTCTTTGAGAC
TCTGAGTTCACCTAGAGAAGTCTAAGCATAACAGCTTTCTTTCCAGCACGAGCCTTTAT
AGCTCTCTTTAGCTCAACCACTCTGTCCATCCAGCCAATGGATGTCCTTCCCTGTACCCA
ATTCAAGCTTATTTTAGGGAAGCCTTGAAACTACCATGTATCTGGCTCTAGCTGAGTTAT
TGAGGATTGAGCCAGTGCAACGTTAAACTCAGTGCACCTTACATTTGATTTAAATGATGGT
TTTATCTGTTGTGTGAAGTGGTTCACCCCTGAGGACCAGGAGCCTCCATATCTGACTGA
AAACCTTTTCTGAGACTTAGAGTAACAGTACTTTTGGTTCTTGAGTTCTCCTGTCTCCA
GATACCTAAATGACCTTGACTTTTTCTGCCTTGTGAATTCGTAGTCCAATCAGCTGAAATT
AAATCACTTGGGAGGGACGCATAGAAGGAGCTCTAGGAACACAGTGCCAGTGCAGAAATT
TCTCCAGGTGGCCTCCCTTTCCAACAATGTACATAATAAAGTGTATGCACCTTCACT

351/392

FIGURE 338

MAEKRDTRDSEAQRLPDSFKDSPSKGLGPCGWILVAFSFLFTVITFPISIWMCIKIKEY
ERAIIFRLGRILQGGAKGPGLFFILPCTDSFIKVD MRTISFDIPPQEILTKDSVTISVDG
VYYRVQNATLAVANITNADSATRLLAQTTLRNVLGTKNLSQILSDREEIAHNMQSTLDD
ATDAWGIKVERVEIKDVKLPVQLQRAMAAEAEASREARAKVIAAEGEMNASRALKEASMV
ITESPAALQLRYLQTLTTIAAEKNSTIVFPLPIDMLQGIIGAKHSHLG

Signal sequence:

1-45

Transmembrane domain:

None

N-glycosylation site:

128-131, 135-138, 159-162, 229-232, 264-267

cAMP- and cGMP-dependent protein kinase phosphorylation
site:

4-7

N-myristoylation site:

26-31, 278-283, 281-286

SPFH domain/Band 7 family:

39-230

352/392

FIGURE 339

TCTAGAGCCCTCTCCCAACATGGCGGCCTCAGCAAAAAGAAGAATAAGAAGGGGAAGAC
TATCTCCCTAACAGACTTTCTGGCTGAGGATGGGGGTACTGGTGGAGGAAGCACCTATGT
TTCCAAACCAGTCAGCTGGGCTGATGAAACGGATGACCTGGAAGGAGATGTTTCGACCAC
TTGGCACAGTAACGATGACGATGTGTATAGGGCGCCTCCAATTGACCGTTCCATCCTTCC
CACTGCTCCACGGGCTGCTCGGGAACCCAATATCGACCGGAGCCGTCTTCCCAATCGCC
ACCCTACACTGCTTTTCTAGGAAACCTACCCTATGATGTTACAGAAGAGTCAATTAAGGA
ATTCTTTCGAGGATTAAATATCAGTGCAGTGCCTTTACCACGTGAACCCAGCAATCCAGA
GAGGTTGAAAGGTTTTGGTTATGCTGAATTTGAGGACCTGGATTCCCTGCTCAGTGCCCT
GAGTCTCAATGAAGAGTCTCTAGGTAACAGGAGAATTGAGTGGACGTTGCTGATCAAGC
ACAGGATAAAGACAGGGATGATCGTTCTTTTGGCCGTGATAGAAATCGGGATTCTGACAA
AACAGATACAGACTGGAGGGCTCGTCTCTGCTACAGACAGCTTTGATGACTACCCACTAG
AAGAGGTGATGATAGCTTTGGAGACAAGTATCGAGATCGTTATGATTCAGACCGGTATCG
GGATGGGTATCGGGATGGGTATCGGGATGGCCACGCCGGGATATGGATCGATATGGTGG
CCGGGATCGCTATGATGACCGAGGCAGCAGAGACTATGATAGAGGCTATGATTCCCGGAT
AGGCAGTGGCAGAAGAGCATTTGGCAGTGGGTATCGCAGGGATGATGACTACAGAGGAGG
CGGGGACCGCTATGAAGACCGATATGACAGACGGGATGATCGGTCGTGGAGCTCCAGAGA
TGATTACTCTCGGGATGATTATAGGCGTGATGATAGAGGTCCCCCAGAACCCAACT
GAATCTAAAGCCTCGGAGTACTCCTGAAGAAGATGATTCTCTGCTAGTACCTCCAGTC
CACTCGAGCTGCTTCTATCTTTGGAGGGGCAAAGCCTGTTGACACAGCTGCTAGAGAAAG
AGAAGTAGAAGAACGGCTACAGAAGGAACAAGAGAAGTTGCAGCGTCAGTGAATGAGCC
AAAAGTAGAACGACGGCCTCGGGAGAGACACCCAAGCTGGCGAAGTGAAGAACTCAGGA
ACGGGAACGGTGCAGGACAGGAAGTGAGTCATCACAACCTGGGACCTCCACCACATCTAG
CAGAAATGCACGAAGGAGAGAGAGTGAAGTCTCTAGAAAATGAAACACTCAATAAGGA
GGAAGATTGCCACTCTCCAACCTCTAAACCTCCCAAACCTGATCAGCCCCCTAAAGGTAAT
GCCAGCCCCCTCCACCAAAGGAGAATGCTTGGGTGAAGCGAAGTTCTAACCTCCTGCTCG
ATCTCAGAGCTCAGACACAGAGCAGCAGTCCCCCTACAAGTGGTGGGGGAAAAGTAGCTCC
AGCTCAACCATCTGAGGAAGGACCAGGAAGGAAAGATGAAAATAAAGTAGATGGGATGAA
TGCCCCAAAAGGCCAACTGGGAACTCTAGCCGTGGTCCAGGAGACGGAGGGAAACAGAGA
CCACTGGAAGGAGTCAGATAGGAAAGATGGCAAAAAGGATCAAGACTCCAGATCTGCACC
TGAGCCAAAGAAACCTGAGGAAAATCCAGCTTCTAAGTTCAGTCTGCAAGCAAGTATGC
TGCTCTCTCTGTTGATGGTGAAGATGAAAATGAGGGAGAAGATTATGCCGAATAGACCTC
TACATCCTGTGCTTTTCTCCTAGTTTCTCTCCACCCTGGAAACATTCGAGAGCAAATCAA
ACCTCTATCCAGACAAGACAAAATAAACTCAACATCTCCTGAAGACCTTTCTTACCTTT
TTTTAAAAACAAAAXTGAAATTATTTTGCATGCTGCTGCAGCCTTTAAAGTATTGAAGT
AACTGGAGAATTGCCAATACAGCCAGAGAGAAAAGGACTACAGCTTTTTTAGAGGAAAAGT
TGTGGTGCGTTATGTCAACATGCAGTTGCCAGTGTGATTAGTGCCTAGGGGTCTCATTTA
GCAGAAATGGTAATGACAGTGATATAATGCCTGGAACCTGGTTGGGCAGTAGGGGAGGGA
GGTAGAAGGAAAAGTGTGAGATTTCTACCTTTTAGTTTTTATCCTATTGTGGCATATATG
AATTCTCAAACATTATCTGAATAAATTTTCCACTCTTGGAAGGTAGATTTAGCCTCAAG
TTGTTCTAGTCTCCAGGAGGCTGCCAGCCCCCTCCTCTTATTTAATTCTGAGTTTGGGGG
CCAGCCTAGAGGGAATTCCTTTTTTTTTTTTTTTTAAACCCCCAGGGGGTAGTTGGGAGT
GAGACTATAGGCCATAAAGAATGGGACTGCATTGGACCAAAATAAATGGGAAAATCGTGG
TTTGAAAAGAAGCTTTTGGGAAGTGATGAGTCATTTTGCACCAGGTAATAGGGGAAAATT
GTGTGACCTCCAGCAAACACATGAATGGTTATTTCTTGGAGCCGGAAGCACTTGGGGGTC
GTGGTAATTCCCAGTGTTTTCTGTGTCTAGTTTACCCTTTCTAAACACTGTCTTTTTT
GAAAGTTTGAATATATCCACATTTCTATTGAAACCTTGAACTAAAAATTTAGACTCTTA
TCGTCACTCTTAAGTTCTTCATGCTACTCTTAACCTCCCAAAAAGCAGTATCTAAGTCACA
TACATGATGCTTGGGCATTTTCTGAGCCATGGAGAACTCTGAAAGGAAGAATCGCTGCT
TTTCTCAAGCAAATCGGTTTCTTGATGTCTTTTGGTTCTCCTTGCTGCTGCTGATGCTT

353/392

GGACCCCTTTTATTGATCAGAGTGCTCTAGAATAATGGATGGTCTTGGATGATGGATAAA
TAGGGACAGGGACAGTTAAATTGGGAGCCTTTCTTACAACCTTGATGGGATTTTTCCCCC
CAAGTTTCCTTCTCCACTGAAATGCCACACTAATGCTTGTTGGATTCATGAGGTGGCCAG
ACCAATGTGTTGTTTTGTTGTTGTTTTTTTTTTAAGCTTCCCTTGAGAGAATAAATGGTA
ATGGAGAGAATCATTTAACAAGGTCTGGTTTTCTCTTGCAACACAGTAGCTAACTTGCC
TGCTTTTATATGCATTTTTGTAGGGATCAGCTTGGTAGACAGTATTAGCGGAGAAACACC
TTGATCTTGGTTTGCAAGCCCTTCTCCCATCAGTCCTAGATTAGGCCCTGTTTCAGCCATG
CAGGGGTGTTGGTTTATGCGTGCTGCAGCAGTGGGCATAATGAATATAATTTACCCAGTG
GACAAAGGTGTGTACCAAGTGAATTTAAATAATTGGTGTGGATTGGCCAGTAGCTAAGAA
GTGGGCTTTTAAAGAGTATTGAAGATTGAAAGGGTTTTCTTTCTTTTTTAAAAAAGAAA
AACAACTATTGATTGTAGATAATGAAAAGCTAGGGTTTGCCCTCTTCATGTCCTACTCTC
CTTCCAAATAGTTATATCCAAACTGTTTTTCCCTCTCCCCTACCTTGTCCTCCCTATTA
AAATAGAAACAGGGATTGATTAATGTCCCGCTCCTGAATACATGTAAAAATTTGTACAAAA
ATATCTTCTATGAAAATGATTTGTAATCTGTAGACTTATTACCTGGGAGATGTCTTGATG
TAAATCCCATCCTTTGGGTTGTGGGTTTTTTGTTTTCTCCAAATAAATCTGATCTTTAA
AGTTAAAAAATAAATAAATAAATAAATAAATAAATAAATAAATAAATAAATAAATAAATAA
AGTTAAAAAATAAATAAATAAATAAATAAATAAATAAATAAATAAATAAATAAATAAATAA

354/392

FIGURE 340

MAASAKKKKNGKGTISLTDFLAEDGGTGGGSTYVSKPVSWADETDLEGDVSTTWHSNDD
DVYRAPPIDRSILPTAPRAAREPNIDRSRLPKSPPYTAFLGNLPYDVTEESIKEFFRGLN
ISAVRLPREPSNPERLKGFYAEFEDLDSLLSALSLSNEESLGNRRIRVDVADQAQDKORD
DRSFGRDRNRDSDKTDTDWRARPATDSFDDYPPRRGDDSFQDKYRDRYDSDRYRDGYRDG
YRDGPRRDMDRYGGDRYDDRGSRDYDRGYDSRIGSGRRAFSGSYRRDDDYRGGGDRYED
RYDRRDRSWSSRDDYSRDDYRRDDRGPPQRPKLNKPRSTPEEDDSSASTSQSTRAASI
FGGAKPVDTAAREEREVEERLQKEQEKLRQWNEPKLERPRERHPSWRSEETQERERSRT
GSESSQTGTSTTSSRNARRRESEKSLNETLNKEEDCHSPTSKPPKPDQPLKVMPPPPK
ENAWVKRSSNPAPARSQSSDTEQQSPTSGGGKVAPAQPSEEGPGRKDENVKVDGMNAPKQQT
GNSSRGPGDGGNRDHWKESDRKDGKKDQDSRSAPEPKPEENPASKFSSASKYAALSVDG
EDENEGEDYAE

Signal Sequence:

None

Transmembrane domain:

None

N-glycosylation site:

120-123, 448-451, 542-545

Glycosaminoglycan attachment site:

507-510

cAMP- and cGMP-dependent protein kinase phosphorylation site:

439-442, 486-489

Tyrosine kinase phosphorylation site:

225-233, 264-270

N-myristoylation site:

25-30, 26-31, 28-33, 118-123, 421-426, 428-433, 538-543

Amidation site:

276-279, 522-525, 563-566

Cell attachment sequence:

215-217

Eukaryotic putative RNA-binding region RNP-1 signature:

137-144

RNA recognition motif:

98-168

355/392

FIGURE 341

CGGTGGACACCACCTCAGCCCCTGAGCAGGAGTCACAGCACGAAGACCAAGCGCAAAGC
GACCCCTGCCCTCCATCCTGACTGCTCCTCCTAAGAGAGATGGCACC GGCCAGAGCAGGA
TTCTGCCCCCTTCTGCTGCTTCTGCTGCTGGGGCTGTGGGTGGCAGAGATCCCAGTCAGT
GCCAAGCCCAAGGGCATGACCTCATCACAGTGGTTTAAAATTCAGCACATGCAGCCCAGC
CCTCAAGCATGCAACTCAGCCATGAAAAACATTAACAAGCACACAAAACGGTGCAAAGAC
CTCAACACCTTCCTGCACGAGCCTTTCTCCAGTGTGGCCGCCACCTGCCAGACCCCCAAA
ATAGCCTGCAAGAATGGCGATAAAAACTGCCACCAGAGCCACGGGCCCCGTGTCCCTGACC
ATGTGTAAGCTCACCTCAGGGAAGTATCCGAAGTGCAGGTACAAAGAGAAGCGACAGAAC
AAGTCTTACGTAGTGGCCTGTAAGCCTCCCCAGAAAAAGGACTCTCAGCAATTCCACCTG
GTTCCCTGTACACTTGGACAGAGTCCTTTTAGGTTTCCAGACTGGCTTGCTCTTTGGCTGAC
CTTCAATTCCCTCTCCAGGACTCCGCACCACTCCCCTACACCCAGAGCATTCTCTTCCCC
TCATCTCTTGGGGCTGTTCTGGTTCAGCCTCTGCTGGGAGGCTGAAGCTGACACTCTGG
TGAGCTGAGCTCTAGAGGGATGGCTTTTTCATCTTTTGTGCTGTTTTCCAGATGCTTA
TCCCCAAGAAACAGCAAGCTCAGGTCTGTGGGTTCCCTGGTCTATGCCATTGCACATGTC
TCCCCTGCCCCCTGGCATTAGGGCAGCATGACAAGGAGAGGAAATAAATGGAAAGGGGGC
AA
AAA

356/392

FIGURE 342

MAPARAGFCPLLLLLLLGLWVAEIPVSAKPKGMTSSQWFKIQHMQPSPQACNSAMKNINK
HTKRCKDLNITFLHEPFSSVAATCQTPKIAKNGDKNCHQSHGFPVSLTMCKLTSGKYPNCR
YKEKRQNKSYVVACKPPQKKDSQQFHLVPVHLDRVL

Important features of the protein

Signal peptide:

1-22

Transmembrane domain:

none

N-glycosylation site:

127-131

**cAMP- and cGMP-dependent protein kinase phosphorylation
site:**

139-143

N-myristoylation site:

18-24, 32-38

Pancreatic ribonuclease family signature:

65-72

Pancreatic ribonuclease family proteins:

49-93

357/392

FIGURE 343

GCATTTGCCACTGGTTGCAGATCAGGCGGACGAGGAGCCGGGAAGGCAGAGCCATGTGGC
TGGCCCCCTGCTCTGCTCCTTCTCAGCCTCTCAGGCTGTTTCTCCATCCAAGGCCAGAGT
CTGTGAGAGCCCCAGAGCAGGGGTCCCTGACGGTTCAATGCCACTATAAGCAAGGATGGG
AGACCTACATTAAGTGGTGGTGCCGAGGGGTGCGCTGGGATACATGCAAGATCCTCATTG
AAACCAGAGGGTCCGAGCAAGGAGAGAAGAGTGACCGTGTGTCCATCAAGGACAATCAGA
AAGACCGCACGTTCACCTGTGACCATGGAGGGGCTCAGGCGAGATGACGCAGATGTTTACT
GGTGTGGGATTGAAAGAAGAGGACCTGACCTTGGGACTCAAGTGAAAGTGATCGTTGACC
CAGAGGGAGCGGCTTCCACAACAGCAAGCTCACCTACCAACAGCAATATGGCAGTGTTCA
TCGGCTCCCACAAGAGGAACCACTACATGCTCCTGGTATTTGTGAAGGTGCCCATCTTGC
TCATCTTGGTCACTGCCATCCTCTGGTTGAAGGGTCTCAGAGGGTCCCTGAGGAGCCAG
GGGAACAGCCTATCTACATGAACTTCTCCGAACCTCTGACTAAAGACATGGCCACTTAGA
GAGATGGATCTGCAGAGCCTTCTGCCCTGGCCACGTTTCCAGAAGAGACTCGGGCTGTG
GAAGGAACATCTACGAGTCCTCGGGATGCAGTGACTGAGATAGGGGCCCTGGGCCTCCGC
CCTGGCCTTGGAGCTGGTGGGCACCTCCCTGTTCTGCACAGCTCAGGGACTTAGCCAGGT
CCTCTCCTGAGCCACCATCACCTCCTGGGGTGCCAGCACCTGTTCTCTTGGTCAGGAGCT
GTAGAGATGGAGCTCAAGCACTGGACGACTCTGTCCCCACTGCTGGAATAACTCGGGCAC
AGAGCATGGGACCAAAGTACAGAAAGAGGTTGGGGGAGACCCCCCAGCCCTAGACTTCC
ATCATTCGGGAGACCAACTCAACACCGTCTTTGCCCTGAGAACCTGATATATCCGTGTTTT
TAAATTTTTTTTTTTCTAGCAAAGTTGGGTTTTAATGACTTATGTTTCATAGGAAACCTCT
CTGATCCCACACACAAGGAGGGTGATTCTGGGATGAGTTCTTGGTTCTAGGGCATGAGGG
GCTGGATGGACCCTGTCCCCAGGGAGGACATGGCTCTGAGTCCACAGGGCTGAGGAGGCA
ATGGGAACCTCCCTGGCCCCGCGCGGTGCTTGTCTCCCCCTCCACCTCTTCTCCTCC
TAGCTCCCCAAGCTCCCTGCCTATTCCCCACCTCCGAGGGGCTGCAGCTTGGGAGCCTC
CTCAGCATGACAGCTTGGGTCTCCTCCCCAAAAGAGCCTGTCAGGCCTCAAGAACCACCT
CCAGGTGGGGAGGGCAGTAACGAAAACCATCGCAGGAAATGGCACCCCTCCCTTTTCGGTG
ATGTTGAAATCATGTTACTAATGAAAACCTGTCTAGGGAAGTGGTTCTGTCTCCTCACAG
GCTTCACCCACGGCGATGAGGCCCTTGAATGTGGTCACTTTGTGCTGTATGGTTGAGGGA
CCCTCACACCAAAGGGACCTTCCCATGTGAGATGTGCTCCCGCCCCACCTGCCACAAG
CAAACACACCACACATGTTCCGCATGTTGCCCTTTGAACACCCATGAGGACGCCTCCAAC
CTGCTCTTGGTTCTAATAGGGAGTACTGACTGTGAGCAGTGGATAAAGGAGAGGGGACCC
TCTGGTCCCTAGCATGGCACCCAGAGCCTCCCCTCTTCTTGTCTTCTCAGCCAAAGAGAAA
CTTTCTCTGACTTTGAACTGAATTTAGGTCTCTGGCCAATGATGGGCCTGAAAATTCAT
AATGGCCAGAGAGGAGAGTTCGAGCCCGGCTAAGATCCCCTGAGTCATTCTGTGAGGGAC
CAAGACCCACAGTCCACCAGCCCCAGGGCCCTACCTCCTGGAATGCTTTCTTGGATCCAG
CTTCCCGAAGATCCGACCAGACCCAGGGAGGACGGCACCGCTCCGCGGGAGGGAAAAGCCA
AAGCATGGTGCTTACCAGCTGGAATCAGGGGCGAGGGGACATGGGCGCTTGTCAACGTG
ATGTCATTCTTTTCCCACCGTTTCTTCTGTTGATATTCAATGAATCCGTCAATCTCTCT
GGGAAA

358/392

FIGURE 344

MWLPPALLLLSLSGCFSSIQGPESVRAPEQGS�TVQCHYKQGWETIYIKWWCRGVRWDTCKI
LIETRGSEQGEKSDRVSIKDNQKDRTFTVTMEGLRRDDADVWCGIERRGPDLTQVKVI
VDPEGAASTTASSPTNSNMAVFIGSHKRNHYMLLVFVKVPILLILVTAILWLKGSQRVPE
EPGEQPIYMNFSPLTKDMAT

Important features of the protein:

Signal peptide:

Amino acids 1-17

Transmembrane domain:

Amino acids 151-170

N-glycosylation site:

Amino acids 190-194

Tyrosine kinase phosphorylation site:

Amino acids 95-103

N-myristoylation sites:

Amino acids 66-72;125-131

Prokaryotic membrane lipoprotein lipid attachment site:

Amino acids 5-16

359/392

FIGURE 345

CTGAGCTCCCGGGCTCCGGCAGCGCGCTGGCGGGGCGCCGCATTGCACACTCTGGGGGCG
CCGCAGTGTTCGTGGGATGGGGCAGCGGGCTGCAGCTGGCGGCCGGAATCCGCGCGCAGC
CCGGGTGCAAGTTCTCTCCTGTTGCCCTGAGTGCCCACTCCCAGGCCCTCTGTATGAGTG
ACACTTCAGTCTGCCATGGAACCTGGCCCTGCTCTGGCCTGGCTCCTGCTCCTGAGCCTG
CTGGCGGATTGTCTGAAAGCTGCTCAGTCCCAGACTTCACAGTGAAAGACATTATCTAC
CTCCATCCTTCAACCACACCATATCCTGGTGGATTTAAATGTTTCACCTGTGAAAAGGCA
GCAGACAATTATGAGTGCAACCGATGGGCTCCAGACATCTACTGCCCTCGAGAGACCAGA
TACTGCTACACTCAGCACACAATGGAAGTCACAGGAAACAGTATCTCAGTCACCAAACGC
TGTGTCCCACTGGAAGAGTGCTTATCCACTGGCTGCAGAGACTCCGAGCATGAAGGCCAC
AAGGTCTGCACCTTCTTGTGTGAAGGAATATCTGTAACCTGCCACTGCCCCGAAATGAA
ACTGATGCCACATTTGCCACGACGTACCTATAAATCAGACAAATGGGCACCCACGCTGT
ATGTCAGTGATAGTGTCTGCTTGTGGTTGTGGTTAGGGCTCATGTTATAGTGGCTCAGT
GGCTCCATGTGTTAATAGCGATCCATGGGGATCTCGATGGTCCACAGACCTGCATGAGTC
ATTGGCCTGACAGTAATTACACATGTGAGACACAACACTCTTGGAGGTCATCACAGCCAA
GCATTGCCACTTACCATGAGGAATAAATGTTGCTTCATTGTAGCCATTTTGAGTCTAACC
GAGACTCATCAAAGCCTTCTGTCTAGTACAGCCCAAGTTCCATACCATAAACGTTGTTTT
CATTCCAAGAAGTAGTTCTGCATTTATCGAGATCTGGGGTTCTTAATTTGGAAGAATACA
TGCATGAGATGCAGTAGGTCTGAGACTGTAAGATATTAGGAGTATGTTATAGGGGCATG
TATAGATGTGGGCTTTTCAGGAGAAAAGTAACCATTGGTTTAAATATAATCATGAGTTCA
TTTGTAGCTTTAGAAATTTTAAACATTGACTCCAACTGAATGGACTATTTCTTGGAAA
TTCTGACTGAGTCCCTGGAAGAGTAGTAATTCACAATTCAGCCATTTGTTCAATTAA
TTTTCCCAACATTCTTCTCCAGTGCTGGGAATCACATTTCTCTGTTCTGTGCAGAAGA
CAAAAAGGCAATCATAAAAGTTTGTATATTTGTGGGGGTGCCTGGAGGAGGATTTTCT
CAACTTAATGGAGCCACTGTCCATAAAGTGGCTGTTATCCCTTCATATAATTGGTGAGAT
CAGCCTTCTCCTTGACTTGGCACCTAATTATGCTTCATGAGATCCTAGATTCCACCTGAG
TCAATTGTGTCCAGAGCCCCAAACCAGGATGGAGTTGTTTTCCCAGATATGGGGTTCTA
TTCAGCCATAGATAATCTAGACAGAGGATTTCAGAATGAAAGGAAAAATGTGTGGAGATT
AGTCTTAGTTTCTGAGGGCCGACTAAGTGGCTCAGCCAGCTTCTTACTCCATCTGCA
GTTTCACTAGTCCAAAGAGCTCCCACTTCCAAATCCCAGTGACTTTATGGAGAAGATTCT
GCATTAAATTGTCTTTTGAATGATGGGGAAGCAAGGCATAATATGCGATGATGAGGAGAA
AGTAGACCAGTGAGGTGATTGCAAGACTAACAAGGAGACTCAATGGGAAGTTTTTCTTTC
TTTTAGATATTGCTTTTGAAGTAGATGGTAAAATTTTGTATCCTTCTGTATTTTTTG
TACCCCAAGTTACAATTTTTCTTCTTCTTGTAAATAATTTAAACAGTATTTATTTTTGT
AAGGCATAACTAGAACTAAAATATATTCTAAAAAATTCATTATTCTGAACAAAGTGATC
AAATTAGAATACATATTTTTCAACAGTGGTAGAGCTTTTAATATATGTTTATTGAAAGTT
ATCTATAATACTTGCACCAGTGTTGAAAAAAGTTAACATGTAGGCAAGAGCAATATGTTT
GTCTCAAGGATTTTTCCATGGTTTTCTCAGTGATGGTGTCTGGAATTATTAGGTGGTG
ACCATCACTGGTCTAAGTTTGTGTGCAGGGTTTTTCAGACGTGTTTTTGTGAACTTGGTA
GAACCATGGCTAATAAAGAGGACAGTGTTGTGAGGGTCCATCTGCCCTCCATAGAAAAAT
GTCTCTGGCTCATAAAATGAGACTCCCTCAGGGACTAAATATGAACTGACAGCAGTAACT
CTGATACAGAATAATCTAAATTGCATCAAATGGCCTTAATTCAGAGTTTGTAGGCTTAT
CAGTATGTTGCTTTTAAATTGGGGTGGGAAAGTAGAGGGAGAGAAAGCAAGACATTTATTA
AGCACCTCGTATGTGCCAGGCACTATGCTAAGCACTTACATAAGTTAGGATTAATCCCT
GCAAGAATCCTATAAAGAATGTTACTAGCATTTTACACTTCCCAAATGAAGGTACCAAAGC
TCAAACGCAATGTTGTGAAGCTGTTTCTTCTCAGATTTAGGTTATGTGGGATGATGTGGGA
TTGAAGAGGAAAGAAAGGTGGGATTATCCCCCTAGGAAGACTTTTCAGGCCTGACTTCATA
GGAATTCATCCATCTTATCATGTGGAGTTTATCTCACCCTGCTGTTGCAGGATGCTATTT
GCATGTGTCCCAGGTGATGTTTTTTCTTGGGGAGTAGGGGTTTGGCTTCTCATTTCAT

360/392

CCCTCTGCTAAAAGAGGAGATAGTTGATGTTGCATCTAAGATGCTATAAGACAATGAAAGTTTGATGTTGTAC
ATACCTACAAGTACCATTTTTGTGCATGATTACACTCCACTGACATCTTCCAAGTACTAC
ATGTGATTGAATAAGAAACAAGAAAGTGACCACACCAAAGCCTCCCTGGCTGGTGTACAG
GGATCAGGTCCACAGTGGTGCAGATTCAACCACCACCCAGGGAGTGCTTGCAGACTCTGC
ATAGATGTTGCTGCATGCGTCCCATGTGCCTGTCAGAATGGCAGTGTTTAATTCTCTTGA
AAGAAAGTTATTTGCTCACTATCCCCAGCCTCAAGGAGCCAAGGAAGAGTCATTCACATG
GAAGGTCCGGGACTGGTCAGCCACTCTGACTTTTCTACCACATTAAATTCTCCATTACAT
CTCACTATTGGTAATGGCTTAAGTGTAAGAGCCATGATGTGTATATTAAGCTATGTGCC
ACATATTTATTTTTAGACTCTCCACAGCATTTCATGTCAATATGGGATTAATGCCTAAACT
TTGTAAATATTGTACAGTTTGTAATCAATGAATAAAGGTTTTGAGTGTAACAAAAAAAAA
AAAAAA

361/392

FIGURE 346

MEPGPALAWLLLLSLADCLKAAQSRDFTVKDIIYLHPSTTPYPGGFKCFTCEKAADNYE
CNRWAPDIYCPRETRYCYTQHTMEVTGNSISVTKRCVPLEECLSTGCRDSEHEGHKVCTS
CCEGNICNLPLPRNETDATFATTSPINQTNHGPRCMSVIVSCLWLWLGLML

Important features of the protein:

Signal peptide:

1-22

Transmembrane domain:

None

N-glycosylation site:

134-138, 147-151

N-myristoylation site:

45-51, 87-93, 106-112, 124-130

Ly-6 / u-PAR domain protein:

115-128

362/392

FIGURE 347

GATCAAGCGCCTTCCTTTCCCTTCCTCTCCCTACTTGGCCTTTGCCCTAAGCCAAGACCT
GGCCATCAGCCTGGCTGCAGGGGCTGCAGAGCCAGCTGCACTTTTTCAGGTATGGGGGA
GGGCCAGGCACCATGAAGCCAGTGTGGGTGCACACCCTTCTGTGGATGCTACTGCTGGTG
CCCAGGCTGGGGGCCGCCCGGAAGGGGTCCCAGAAGAGGCCTCCTTCTACTATGGAACC
TTCCCTCTTGGCTTCTCCTGGGGCGTGGGCAGTTCTGCCTACCAGACGGAGGGCGCCTGG
GACCAGGACGGGAAAGGGCCTAGCATCTGGGACGTCTTACACACAGTGGGAAGGGGAAA
GTGCTTGGGAATGAGACGGCAGATGTAGCCTGTGACGGCTACTACAAGGTCCAGGAGGAC
ATCATTTCTGCTGAGGGAAGTGCACGTCAACCACTACCGATTCTCCCTGTCTTGGCCCCGG
CTCCTGCCCACAGGCATCCGAGCCGAGCAGGTGAACAAGAAGGGAATCGAATTCTACAGT
GATCTTATCGATGCCCTTCTGAGCAGCAACATCACTCCCATCGTGACCTTGCACCACTGG
GATCTGCCACAGCTGCTCCAGGTCAAATACGGTGGGTGGCAGAATGTGAGCATGGCCAAC
TACTTCAGAGACTACGCCAACCTGTGCTTTGAGGCCTTTGGGGACCGTGTGAAGCACTGG
ATCACGTTCAGTGATCCTCGGGCAATGGCAGAAAAAGGCTATGAGACGGGGCCACCATGCG
CCGGGCCTGAAGCTCCGCGGCACCGGCCTGTACAAGGCAGCACACCACATCATTAAGGCC
CACGCCAAAACCTGGCATTCTTATAACACCACGTGGCGCAGCAAGCAGCAAGGTCTGGTG
GGAATTTCACTGAACTGTGACTGGGGGGAACCTGTGGACATTAGTAACCCCAAGGACCTA
GAGGCTGCCGAGAGATACCTACAGTTCTGTCTGGGCTGGTTTGCCAACCCCATTTATGCC
GGTGACTACCCCAAGTCATGAAGGACTACATTGGAAGAAAGAGTGCAGAGCAAGGCCTG
GAGATGTCGAGGTTACCGGTGTTCTCACTCCAGGAGAAGAGCTACATTAAAGGCACATCC
GATTTCTTGGGATTAGGTCATTTTACTACTCGGTACATCACGAAAGGAACTACCCCTCC
CGCCAGGGGCCAGCTACCAGAACGATCGTGACTTGATAGAGCTGGTTGACCCAAACTGG
CCAGATCTGGGGTCTAAATGGCTATATTCTGTGCCATGGGGATTTAGGAGGCTCCTTAAC
TTTGCTCAGACTCAATACGGTGATCCTCCCATATATGTGATGGAAAATGGAGCATCTCAA
AAATTCCACTGTACTCAATTATGTGATGAGTGGAGAATTCAATACCTTAAAGGATACATA
AATGAAATGCTAAAAGCTATAAAAGATGGTGCTAATATAAAGGGGTATACTTCCTGGTCT
CTGTTGGATAAGTTTGAATGGGAGAAAGGATACTCAGATAGATATGGATTCTACTATGTT
GAATTTAACGACAGAAATAAGCCTCGCTATCCAAAGGCTTCAGTTCAATATTACAAGAAG
ATTATCATTGCCAATGGGTTTCCCAATCCAAGAGAGGTGGAAAGTTGGTACCTCAAAGCT
TTGGAAACTTGCTCTATCAACAATCAGATGCTTGCTGCAGAGCCTTTGCTAAGTCACATG
CAAATGGTTACGGAGATCGTGGTACCCACTGTCTGCTCCCTCTGTGTCCTCATCACTGCT
GTTCTACTAATGCTCCTCCTGAGGAGGCAGAGCTGAGACAGGATTATCAATTTTGAGCT
TCATAAGAGAATCTTCAGGATCTTCCTCCCTTTTCTGCTTTGAGGGTTTCCATACATTGC
TGTTTTTCAGGTTCTACAATAATTACCTTTTTTTCTCTTTCTCTTTTGGCTTGTGCTGGG
ATTTAAGAATTAGAAAATAAAAATAAGCAGAAATTA

363/392

FIGURE 348

MKPVVWATLLWMLLLVPRLGAARKGSPPEASFYFGTFPLGFSWGVGSSAYQTEGAWDQDG
KGPSIWDVFTHSGKGKVLGNETADVACDGYKQVEDIILLRELHVNHYRFSLSWPRLPT
GIRAEQVNKKGIEFYSDLIDALLSSNITPIVTLHHWDLPQLLQVKYGGWQNVSMANYFRD
YANLCFEAFGDRVKHWITFSDPRAMAEKGYETGHHAPGLKLRGTGLYKAAHHIIKAHAKT
WHSYNTTWRSKQQGLVGISLNCDWGEPVDISNPKDLEAAERYLQFCLGWFANPIYAGDYP
QVMKDYIGRKSAEQGLEMSRLPVFSLQEKSYIKGTSDFLGLGHFTTRYITERNYPSRQGP
SYQNDRLIELVDPNWPDLGSKWLYSVPWGFRRLLNFAQTQYGDPPYVMENGASQKFHC
TQLCDEWRIQYLKGYINEMLKAIKDGANIKGYTSWSLLDKFEWEKGYSDDRYGFYYVEFND
RNKPRYPKASVQYYKKIIANGFPNPREVESWYLKALETCSINNQMLAAEPLLSHMQMVT
EIVVPTVCSLCVLITAVLLMLLLRRQS

Important features:**Signal peptide:**

amino acids 1-21

Transmembrane domain:

amino acids 541-558

N-glycosylation sites:

amino acids 80-84,171-175,245-249

Glycosaminoglycan attachment site:

amino acids 72-76

cAMP- and cGMP-dependent protein kinase phosphorylation sites:

amino acids 23-27,564-568

Tyrosine kinase phosphorylation sites:

amino acids 203-211,347-355,460-468,507-514

N-myristoylation sites:

amino acids 44-50,79-85,167-173,225-231,257-263,315-321

Amidation site:

amino acids 307-311

Glycosyl hydrolases family 1 active site:

amino acids 407-416

Glycosyl hydrolases family 1 N-terminal signature:

amino acids 41-56

Motif name Glycosyl hydrolases family:

amino acids 37- 67

364/392

FIGURE 349

CGCAAAGCCGCCCTCGGGGCGCTCATGGCGGGACGCCTCCTGGGAAAGGCTTTAGCCGCG
GTGTCTCTCTCTCTGGCCTTGGCCTCTGTGACTATCAGGTCCTCGCGCTGCCGCGGCATC
CAGGCGTTTCAAGAACTCGTTTTTCATCTTCTTGGTTTCATCTTAATACCAACGTCATGTCT
GGTTCTAATGGTTCCAAAGAAAATTCTCACAATAAGGCTCGGACGTCTCCTTACCCAGGT
TCAAAGTTGAACGAAGCCAGGTTCCCTAATGAGAAAGTGGGCTGGCTTGTTGAGTGGCAA
GACTATAAGCCTGTGGAATACACTGCAGTCTCTGTCTTGGCTGGACCCAGGTGGGCAGAT
CCTCAGATCAGTGAAAGTAATTTTTCTCCAAGTTTAACGAAAAGGATGGGCATGTTGAG
AGAAAGAGCAAGAATGGCCTGTATGAGATTGAAAATGGAAGACCGAGAAATCCTGCAGGA
CGGACTGGACTGGTGGGCGGGGGCTTTTTGGGGCGATGGGGCCCAAATCACGCTGCAGAT
CCCATTATAACCAGATGGAAAAGGGATAGCAGTGGAAATAAAATCATGCATCCTGTTTCT
GGGAAGCATATCTTACAATTTGTTGCAATAAAAAGGAAAAGACTGTGGAGAATGGGCAATC
CCAGGGGGGATGGTGGATCCAGGAGAGAAGATTAGTGCCCACTGAAAAGAGAATTTGGT
GAGGAAGCTCTCAACTCCTTACAGAAAACCAGTGCTGAGAAGAGAGAAAATAGAGGAAAAG
TTGCACAAACTCTTCAGCCAAGACCACCTAGTGATATATAAGGGATATGTTGATGATCCT
CGAAACACTGATAATGCATGGATGGAGACAGAAGCTGTGAACTACCATGACGAAACAGGT
GAGATAATGGATAATCTTATGCTAGAAGCTGGAGATGATGCTGGAAAAGTGAAATGGGTG
GACATCAATGATAAACTGAAGCTTTATGCCAGTCACTCTCAATTCATCAAACCTTGTGGCT
GAGAAACGAGATGCACACTGGAGCGAGGACTCTGAAGCTGACTGCCATGCGTTGTAGCTG
ATGGTCTCCGTGTAAGCCAAAGGCCACAGAGGAGCATATACTGAAAAGAAGGCAGTATC
ACAGAATTTTATACTATAAAAAGGGCAGGGTAGGCCACTTGGCCTATTTACTTTCAAAACA
ATTTGCATTTAGAGTGTTTCGCATCAGAATAACATGAGTAAGATGAACTGGAACACAAAA
TTTTCAGCTCTTTGGTCAAAAGGAATATAAGTAATCATATTTTGTATGTATTCGATTTAA
GCATGGCTTAAATTAAATTTAAACAACCTAATGCTCTTTGAAGAATCATAATCAGAATAAA
GATAAATCTTGTATCAGCTATA

365/392

FIGURE 350

MAGRLLGKALAAVSLSLALASVTIRSSRCRGIQAFRNSFSSSWFHLNTNVMMSGNSGSKEN
SHNKARTSPYPGSKVERSQVPNEKVGWLVEWQDYKPVEYTAVSVLAGPRWADPQISESNF
SPKFNEKDGHVERKSKNGLYEIENGRPRNPAGRTGLVGRGLLGRWGPNAADPIITRWKR
DSSGNKIMHPVSGKHILQFVAIKRKDCGEWAI PGGMVDPGEKISATLKREFGEEALNSLQ
KTSAEKREIEEKLHKLFSQDHLVIYKGYVDDPRNTDNAWMETEAVNYHDETGEIMDNMLL
EAGDDAGKVKWVDINDKLKLYASHSQFIKLVAEKRDAHWSSEDSEADCHAL

Important features of the protein:**Signal peptide:**

1-20

Transmembrane domain:

None

N-glycosylation site:

55-59

cAMP- and cGMP-dependent protein kinase phosphorylation site:

179-183

N-myristoylation site:

53-59, 56-62

mutT domain signature:

215-235

366/392

FIGURE 351

CCTCTGTCTGTGCTCCCATCCCAGGGAGTATAGGTGGAGCCTCCAGAGCCCATGGACAGG
GCATGCTGGGGCTGGGCCAGCCCCAGCGGTGTCTCTAAGGCACCCCTGGGATCCCCACTG
AGCTGGCCTACTTCAGACAGCCAGGGCCCCACCCCTCTGGCCCCCTTAGTGTCAGCTCGT
GGCCCCCTTGGCATTTCCACAAGACGCCAAGATGGAGATTCCCATGGGGACCCAGGGCTGC
TTCTCAAAGAGCCTCCTGCTCTCAGCCTCAATCCTGGTCCTCTGGATGCTCCAAGGCTCC
CAGGCAGCTCTCTACATCCAGAAGATTCCAGAGCAGCCTCAAAAGAACCAGGACCTTCTC
CTGTTCAGTCCAGGGTGTCCCAGACACCTTCCAGGACTTCAACTGGTACCTGGGGGAGGAG
ACGTACGGAGGCACGAGGCTATTTACCTACATCCCTGGGATACAACGGCCTCAGAGGGAT
GGCAGTGCCATGGGACAGCGAGACATCGTGGGCTTCCCCAATGGTTCCATGCTGCTGCGC
CGCGCCAGCCTACAGACAGTGGCACCTACCAAGTAGCCATTACCATCAACTCTGAATGG
ACTATGAAGGCCAAGACTGAGGTCCAGGTAGCTGAAAAGAATAAGGAGCTGCCCAGTACA
CACCTGCCCCACCAACGCTGGGATCCTGGCGGCCACCATCATTTGGATCTCTTGCTGCCGGG
GCCCTTCTCATCAGCTGCATTGCCTATCTCCTGGTGACAAGGAAGTGGAGGGCCAGAGC
CACAGACTGCCTGCTCCGAGGGGGCCAGGGATCTCTGTCCATCTTGTGCTCGGCTGTATCC
CCAGTGCCTTCAGTGACGCCCAGCACATGGATGGCGACCACAGAGAAGCCAGAAATTGGGC
CCTGCTCATGATGCTGGTGACAACAACATCTATGAAGTGATGCCCTCTCCAGTCCTCCTG
GTGTCCCCCATCAGTGACACAAGGTCCATAAACCCAGCCCGGCCCTGCCACACCCCCA
CACCTGCAGGCGGAGCCAGAGAACCACAGTACCAGCAGGACCTGCTAAACCCCGACCCT
GCCCCCTACTGCCAGCTGGTGCCAACTTCCTGATGGGTCTTGGGCCAGGCCAGCCAGGGA
GAAGACAAGGCCCCAGCCCTCCTCTGGGAGCCTCACACCTGAGACCAGCAGGACAAGGCC
ATTGGGGGCTGTGGGGCCGATGAGGTGGACTCAGCCAAAGACTCAGCAGCACATGGGGCA
GGTGTCTTGGCAGGGGGACAGGAGACTGTAACAGGCCCAGGTCTTGTGCAGCCCTGTAA
TGCACGCCCCGCTTCGGTCTGTTCTTCAAGCAAGCTGGCCTGGGCCATGTGCCTGTGAA
AGGCAGGCTCTGGCCCCCTTCCATGCCAAAGTCCCCCAAGATCTGGATATCTGGGGACAA
GATGGTGGCCTCAGGCCTGCCTCCCAGGCAGTTGGCTGGGCTCCCAACTGTCTGTCTCTCA
ATGCCCTACCCCAACTCCACTAGTGACCCTCAGAGTCTTCTCCCCCTTAGGACAAGGCAGA
CACCCCACCATGCGGGCCTCAGGTGGCAGAGAGGCCAGCCTCACAGGCCTGTGGCCCCA
CACACCAGTCCCAGCAAGGTGACCACGGCTGCTGGACCCCTTCCCTGTTTCAGGCAGGCCC
AGCCCCCTCTCAGAACCTGCTGCCAGCTGCTGGTCTTGGCCCCCACCCTGAATCTTACTGA
GTCCCTCTGGGCAGCAGCTCCCTTCTCCACCCACCCAGCACCCTGCCAAATGTGGCC
TCAGCTTGTCTCTCCCTTCCCCAAACTATGCATTCAATCAGCAATAAATGAGCCTTTGCT
GCA

367/392

FIGURE 352

MEIPMGQTQGCFSKSLLLSASILVLWMLQGSQAALYIQKIPEQPQKNQDLLLSVQGVPDF
QDFNWYLGEETYGGTRLFTYIPGIQRPQRDGSAMGQRDIVGFNGSMLLRRQAQPTDSGT
QVAITINSEWTMKAKTEVQVAEKNKELPSTHLPTNAGILAATIIGSLAAGALLISCIAYL
LVTRNWRGQSHRLPAPRGQGSLSILCSAVSPVPSVTPSTWMATTEKPELGPAGHDAGDNNI
YEVMPSPVLLVSPISDTRSINPARPLPTPPHLQAEPENHQYQQDLLNPDPAPYCQLVPTS

Important features of the protein:

Signal peptide:

Amino acids 1-32

Transmembrane domain:

Amino acids 159-178

N-glycosylation site:

Amino acids 104-108

N-myristoylation sites:

Amino acids 6-12; 29-35; 55-61; 91-97; 157-163; 165-171

368/392

FIGURE 353

CTTCAGAACAGGTTCTCCTTCCCCAGTCACCAAGTTGCTCGAGTTAGAATTGTCTGCAATG
GCCGCCCTGCAGAAATCTGTGAGCTCTTTCCTTATGGGGACCCTGGCCACCAGCTGCCTC
CTTCTCTTGGCCCTCTTGGTACAGGGAGGAGCAGCTGCGCCCATCAGCTCCCACTGCAGG
CTTGACAAGTCCAACTTCCAGCAGCCCTATATCACCAACCGCACCTTCATGCTGGCTAAG
GAGGCTAGCTTGGCTGATAACAACACAGACGTTTCGTCTCATTGGGGAGAACTGTTCCAC
GGAGTCAGTATGAGTGAGCGCTGCTATCTGATGAAGCAGGTGCTGAACCTTCACCTTGAA
GAAGTGCTGTTCCCTCAATCTGATAGGTTCCAGCCTTATATGCAGGAGGTGGTGCCCTTC
CTGGCCAGGCTCAGCAACAGGCTAAGCACATGTCATATTGAAGGTGATGACCTGCATATC
CAGAGGAATGTGCAAAAGCTGAAGGACACAGTGAAAAAGCTTGGAGAGAGTGAGAGATC
AAAGCAATTGGAGAACTGGATTGCTGTTTATGTCTCTGAGAAATGCCTGCATTTGACCA
GAGCAAAGCTGAAAAATGAATAACTAACCCCTTCCCTGCTAGAAATAACAATTAGATG
CCCCAAAGCGATTTTTTTTAACCAAAGGAAGATGGGAAGCCAACTCCATCATGATGGG
TGGATTCCAAATGAACCCCTGCGTTAGTTACAAAGGAAACCAATGCCACTTTTGTTTATA
AGACCAGAAGGTAGACTTTCTAAGCATAGATATTTATTGATAACATTTTCATTGTAAGTGG
TGTTCTATACACAGAAAACAATTTATTTTTTAAATAATTGTCTTTTCCATAAAAAAGAT
TACTTTCCATTCCTTTAGGGGAAAAAACCCTAAATAGCTTCATGTTCCATAATCAGTA
CTTTATATTTATAAATGTATTTATTATTATTATAAGACTGCATTTTATTTATATCATTTT
ATTAATATGGATTTATTTATAGAAACATCATTGATATTGCTACTTGAGTGTAAGGCTAA
TATTGATATTTATGACAATAATTATAGAGCTATAACATGTTTATTTGACCTCAATAACA
CTTGGATATCCC

369/392

FIGURE 354

MAALQKSVSSFLMGTLATSCLLLLALLVQGGAAAPISSHCRLDKSNFQQPYITNRTFMLA
KEASLADNNTDVRLIGEKLFGVSMSERCYLMKQVLNFTLEEVLFPPQSDRFQPYMQEVVP
FLARLSNRLSTCHIEGDDLHIQRNVQKLKDTVKKLGESGEIKAIGELDLLFMSLRNACI

Important features of the protein:

Signal peptide:

amino acids 1-33

N-glycosylation sites:

amino acids 54-58, 68-72, 97-101

N-myristoylation sites:

amino acids 14-20, 82-88

Prokaryotic membrane lipoprotein lipid attachment site:

amino acids 10-21

370/392

FIGURE 355

TGGCCTACTGGAAAAAAAAAAAAAAAAAAAAAGTCACCCGGGCCCCGCGGTGGCCACAA
CATGGCTGCGGCGCCGGGGCTGCTCTTCTGGCTGTTTCGTGCTGGGGGCGCTCTGGTGGGT
CCCGGGCCAGTCGGATCTCAGCCACGGACGGCGTTTCTCGGACCTCAAAGTGTGCGGGGA
CGAAGAGTGCAGCATGTTAATGTACCGTGGAAGCTCTTGAAGACTTCACGGGCCCTGA
TTGTCGTTTTGTGAATTTTAAAAAGGTGACGATGTATATGTCTACTACAACTGGCAGG
GGGATCCCTTGAACTTTGGGCTGGAAGTGTTGAACACAGTTTGGATATTTTCCAAAAGA
TTTGATCAAGGTACTTCATAAATACACGGAAGAAGAGCTACATATTCAGCAGATGAGAC
AGACTTTGTCTGCTTTGAAGGAGGAAGAGATGATTTTAATAGTTATAATGTAGAAGAGCT
TTTAGGATCTTTGGAACCTGGAGGACTCTGTACCTGAAGAGTCGAAGAAAGCTGAAGAAGT
TTCTCAGCACAGAGAGAAATCTCCTGAGGAGTCTCGGGGGCGTGAACCTGACCCTGTGCC
TGAGCCCGAGGCATTTCAGAGCTGATTTCAGAGGATGGAGAAGGTGCTTTCTCAGAGAGCAC
CGAGGGGCTGCAGGGACAGCCCTCAGCTCAGGAGAGCCACCCTCACACCAGCGGTCTTGC
GGCTAACGCTCAGGGAGTGCAGTCTTCGTTGGACACTTTTGAAGAAATTCTGCACGATAA
ATTGAAAGTGCCGGAAGCGAAAGCAGAACTGGCAATAGTTCTCCTGCCTCGGTGGAGCG
GGAGAAGACAGATGCTTACAAAGTCCTGAAAACAGAAATGAGTCAGAGAGGAAGTGGACA
GTGCGTTATTTCATTACAGCAAAGGATTTTCGTTGGCATCAAATCTAAGTTTGTTTTACAA
AGATTGTTTTTAGTACTAAGCTGCCTTGGCAGTTTGCATTTTTTGAGCCAAACAAAAATAT
ATTATTTTCCCTTCTAAGTAAAAAAAAAAAAAAAAAAAAA

371/392

FIGURE 356

MAAAPGLLFWLFLVLGALWWVPGQSDLSHGRRFSDLKVCGDEEC SMLMYRGKALEDFTGPD
CRFVNFKKGDDVYVYYKLAGGSLELWAGSVEHSFGYFPKDLIKVLHKYTEELHIPADET
DFVCFEGGRDDFNSYNVEELLGSLELEDSVPEESKKAEVVSQHREKSPEESRGRELDVPV
EPEAFRADSEDGEGAFSESTEGLQGQPSAQESH PHTSGPAANAQGVQSSLDTFEEILHDK
LKVPGESERTGNSSPASVEREKTDAYKVLKTEMSQRGSGQCVIHYSKGFRWHQNL SLFYK
DCF

Important features of the protein:

Signal peptide:

amino acids 1-22

N-glycosylation site:

amino acids 294-298

cAMP- and cGMP-dependent protein kinase phosphorylation site:

amino acids 30-34

Tyrosine kinase phosphorylation site:

amino acids 67-76

N-myristoylation sites:

amino acids 205-211, 225-231, 277-283

Amidation site:

amino acids 28-32

FIGURE 357

[illegible]

373/392

FIGURE 358

MEAPGPRALRTALCGGCCCLLLCAQLAVAGKGARGFGRGALIRLNIWPAVQGACKQLEVC
EHCVEGDRARNLSSCMWEQCRPEEPGHCVAQSEVVKEGCSIYNRSEACPAHHHPTYPEK
TVTTGSPPVPEAHSPGFDGASFIGGVVLVLSLQAVAFFVLHFLKAKDSTYQTL

Important features of the protein:

Signal peptide:

1-29

Transmembrane domain:

141-160

N-glycosylation site:

71-75, 103-107

Tyrosine kinase phosphorylation site:

164-171

N-myristoylation site:

15-21

374/392

FIGURE 359

TTCCAGTCAGAGTTAAGTTAAAACAGAAAAAAGGAAGATGGCAAGAATATTGTTACTTTT
CCTCCCGGGTCTTGTGGCTGTATGTGCTGTGCATGGAATATTTATGGACCGTCTAGCTTC
CAAGAAGCTCTGTGCAGATGATGAGTGTGTCTATACTATTTCTCTGGCTAGTGCTCAAGA
AGATTATAATGCCCCGGACTGTAGATTCATTAACGTTAAAAAAGGGCAGCAGATCTATGT
GTA~~CT~~CAAAGCTGGTAAAAGAAAATGGAGCTGGAGAATTTTGGGCTGGCAGTGT~~TT~~TATGG
TGATGGCCAGGACGAGATGGGAGTCGTGGGTATTTCCCAGGA~~ACT~~TGGTCAAGGAACA
GCGTGTGTACCAGGAAGCTACCAAGGAAGTTCCACACCGGATATTGACTTCTTCTGCGA
GTA~~AT~~AATAATTAGTTAAA~~ACT~~GCAAATAGAAAGAAAACACCA~~AAAA~~ATAAAGAAAAGAGCAA
AAGTGGCCAAAAAATGCATGTCTGTAATTTTGGACTGACGT

375/392

FIGURE 360

MARILLLLFLPGLVAVCAVHGIFMDRLASKKLCADDECVTYISLASAQEDYNAPDCRFINV
KKGQQIYVYSKLVKENGAGEFWAGSVYGDGQDEMGVVGYFPRNLVKEQRVYQEATKEVPT
TDIDFFCE

Important features of the protein:

Signal peptide:

1-14

Transmembrane domain:

None

N-myristoylation site:

84-90

376/392

FIGURE 361

GGCACGAGCCACCACTTACAACCACACAGCCTATCCAGAAACATGAAGATAAGAAATGCT
TGTGCTGTCTTATTGAAGTACTCCTGTTTATACTTGAAGGAGTTACAGGAGCTCGAAAA
ATTTCAACTTTCTCAGGCCCTGGCTCATGGCCGTGCAATCCCAAGTGTGATGGCAGAACT
TACAACCCCTCAGAGGAGTGTTGTGTTTCATGACACCATCCTGCCCTTTAAGCGGATTAAC
CTCTGTGGCCCTAGCTGCACCTACAGGCCCTGCTTTGAGCTCTGCTGTCTGAGTCCTAT
AGCCCCAAGAAGAAATTTATTGTCAAGCTTAAAGTTCATGGAGAGAGATCCCATTGCAGT
TCATCCCCTATCTCCAGGAAGTGTAAAAGCAACAAGATTTTTTCATGGAGAAGATATTGAA
GACAACCAACTTTCTCTTAGGAAAAAAGTGGTGACCAGCCTTGAGAGTCTGCTTTCTTC
CTGCAAGCACCAGTTCCTGAATGTTCTTACTTGAAGAATGGATACCTGAAGCATTGGGGT
GCAGTGATATATGTGTCTCATTACAATGCTCCTTTGGATATTGTTTTCTAAGCATGTGT
TGGAATGTTCCCCATAACTTTCTAAAATTATCCTATTTCAATGCAACTAAAGATAAATG
TATTCCAGCCAGAGTCCACAGAGAAGGCAAGTTATGCAAGGCAGGCATGGGGCCCTCACA
AAATTTCAAGCTGTGCGACTTATGTAGTAATTTTCTACAAACAATCCCTCCTGGATATCC
AGGAGGCTCCAGACCTGAATAAAAAACCACATGTCTGTCTAGAAAAAGGGAATGAATCAAG
ATCCACAGGACCTTTTCAAGATTTTAGAAGCAGCAAACCTATGGCTGAGAGAAAAGACTCT
CTGACCAGGCAAATTGTTCTGCAGTATTCTCCGGGCGTGTAGCTCCCCTGAGTAGTCGCC
AGGCTGGTCTTGGCTTTGTAATAATACAGCTGCCTTTGAGTCCTCCCTACCCTGTTAGTA
ACCCCTTGCTGCACTGTTGTCCTTACAACCGAAATAAACTGATTAGTTG

377/392

FIGURE 362

MKIRNACAVLIEVLLFILEGVTGARKISTFSGPGSWPCNPKCDGRTYNPSEECCVHDTIL
PFKRINLCGPSCTYRPCFELCCPESYSPKKKFIVKLKVHGERSHCSSSPISRNCKSNKIF
HGEDIEDNQLSLRKKSGDQP

Important features of the protein:

Signal peptide:

1-23

Transmembrane domain:

None

Glycosaminoglycan attachment site:

31-35

N-myristoylation site:

20-26, 34-40

378/392

FIGURE 363

ACACTGGCCAAACAAAAACGAAAGCACTCCGTGCTGGAAGTAGGAGGAGAGTCAGGACTC
CCAGGACAGAGAGTGCACAACTACCCAGCACAGCCCCCTCCGCCCCCTCTGGAGGGCTGA
AGAGGGATTCCAGCCCCCTGCCACCCACAGACACGGGGCTGACTGGGGTGTCTGCCCCCCTT
GGGGGGGGGAGCACAGGGCCTCAGGCCTGGGTGCCACCTGGCACCTAGAAGATGCTCTGT
GCCCTGGTTCTTGCTGTCTTGGCACTGGGGCCGAAGCCCAGTGGTCCTTTCTCTGGAGAG
GCTTGTGGGGCCTCAGGACGCTACCCACTGCTCTCCGGGCCTCTCCTGCCGCCTCTGGGA
CAGTGACATACTCTGCCTGCCTGGGGACATCGTGCCCTGCTCCGGGCCCCGTGCTGGCGCC
TACGCACCTGCAGACAGAGCTGGTGCTGAGGTGCCAGAAGGAGACCGACTGTGACCTCTG
TCTGCGTGTGGCTGTCCACTTGGCCGTGCATGGGCACTGGGAAGAGCCTGAAGATGAGGA
AAAGTTTGGAGGAGCAGCTGACTCAGGGGTGGAGGAGCCTAGGAATGCCTCTCTCCAGGC
CCAAGTCGTGCTCTCCTTCCAGGCCTACCCTACTGCCCGCTGCGTCCTGCTGGAGGTGCA
AGTGCCCTGCTGCCCTTGTGCAGTTTGGTCAGTCTGTGGGCTCTGTGGTATATGACTGCTT
CGAGGCTGCCCTAGGGAGTGAGGTACGAATCTGGTCCTATACTCAGCCCAGGTACGAGAA
GGAACCTCAACCACACACAGCAGCTGCCTGCCCTGGCTCAACGTGTGACGAGATGG
TGACAACGTGCATCTGGTTCTGAATGTCTCTGAGGAGCAGCACTTCGGCCTCTCCCTGTA
CTGGAATCAGGTCCAGGGCCCCCAGAACCCCGGTGGCACAAAAACCTGACTGGACCGCA
GATCATTACCTTGAACCACACAGACCTGGTTCCCTGCCTCTGTATTTCAGGTGTGGCCTCT
GGAACCTGACTCCGTTAGGACGAACATCTGCCCTTCAGGGAGGACCCCCGCGCACACCA
GAACCTCTGGCAAGCCGCCCCGACTGCGACTGCTGACCCTGCAGAGCTGGCTGCTGGACGC
ACCGTGCTCGCTGCCCGCAGAAGCGGCACTGTGCTGGCGGGCTCCGGGTGGGGACCCCTG
CCAGCCACTGGTCCCACCGCTTTCCTGGGAGAACGTCACTGTGGACAAGGTTCTCGAGTT
CCCATTGCTGAAAGGCCACCCTAACCTCTGTGTTTCAGGTGAACAGCTCGGAGAAGCTGCA
GCTGCAGGAGTGCTTGTGGGCTGACTCCCTGGGGCCTCTCAAAGACGATGTGCTACTGTT
GGAGACACGAGGCCCCCAGGACAACAGATCCCTCTGTGCCTTGGAACCCAGTGGCTGTAC
TTCACTACCCAGCAAAGCCTCCACGAGGGCAGCTCGCCTTGAGAGTACTTACTACAAGA
CCTGCAGTCAGGCCAGTGTCTGCAGCTATGGGACGATGACTTGGGAGCGCTATGGGCCTG
CCCCATGGACAAATACATCCACAAGCGCTGGGCCCTCGTGTGGCTGGCCTGCCTACTCTT
TGCCGCTGCGCTTTCCTCATCCTCCTTCTCAAAAAGGATCACGCGAAAGGGTGGCTGAG
GCTCTTGAAACAGGACGTCCGCTCGGGGGCGGCCCGCAGGGGCCGCGCGGCTCTGCTCCT
CTACTCAGCCGATGACTCGGGTTTCGAGCGCCTGGTGGGCGCCCTGGCGTCCGCCCTGTG
CCAGCTGCCGCTGCGCGTGGCCGTAGACCTGTGGAGCCGTGCTGAACCTGAGCGCGCAGGG
GCCCCGTGGCTTGGTTTCACGCGCAGCGGCGCCAGACCCTGCAGGAGGGCGGCGTGGTGGT
CTTGCTCTTCTCTCCCGGTGCGGTGGCGCTGTGCAGCGAGTGGCTACAGGATGGGGTGTG
CGGGCCCCGGGGCGCACGGCCCCGACGACGCTTCCGCGCCTCGCTCAGCTGCGTGCTGCC
CGACTTCTTGAGGGCCGGGCGCCCGCAGCTACGTGGGGGCCTGCTTCGACAGGCTGCT
CCACCCGGACGCCGTACCCGCCCTTTTCGACACCGTGCCCGTCTTCACACTGCCCTCCCA
ACTGCCAGACTTCTGGGGGCCCTGCAGCAGCCTCGCGCCCCGCGTTCCGGGGCGGCTCCA
AGAGAGAGCGGAGCAAGTGTCCCGGGCCCTTCAGCCAGCCCTGGATAGCTACTTCCATCC
CCCCGGGACTCCCGCGCCGGGACGCGGGGTGGGACCAGGGGCGGGACCTGGGGCGGGGA
CGGGACTTAAATAAAGGCAGACGCTGTTTTTCTAAAAAAA

379/392

FIGURE 364

MPVPWFLLSLALGRSPVVLSELRLVGPQDATHCSPGLSCRLWDSIDLCLPGDIVPAPGPV
LAPTHLQTELVLRCQKETDCDLCLRVAVHLAVHGHWEPEDEEKFGGAADSGVEEPRNAS
LQAQVVLVSFQAYPTARCVLLEVQVPAALVQFGQSVGSVVYDCFEAALGSEVRIWSYTQPR
YEKELNHTQQLPALPWLNVSADGDNVHLVLNVSEEQHFGLSLYWNQVQGPPKPRWHKNLT
GPQIITLNLHTDLVPCLCIQVWPLEPDSVRTNICPFREDPRAHQNLWQAARLRLTLQSWL
LDAPCSLPAAALCWRAPGGDPQCPLVPPLSWENVTVDKVLEFPLLKGHPNLCVQVNSSE
KLQLQECLWADSLGPLKDDVLLLETRGPQDNRSLEPSGCTSLPSKASTRAARLGEYL
LQDLQSGQCLQLWDDDLGALWACPMCKYIHKRWALVWLACLLFAAALSLILLKKDHAKG
WLRLKQDVRSAGAAARGRAALLLYSADDSGFERLVGALASALCQLPLRVAVDLWSRRELS
AQGPVAVFWHAQRRQTLQEGGVVLLFSPGAVALCSEWLQDGVSGPGAHGPHDAFRASLSC
VLPDFLQGRAPGSYVGACFDRLLHPDAVPALFRTVPVFTLPSQLPDLGALQQPRAPRS
RLQERAEQVSRALQPALDSYFHPPGTPAPGRGVGPGAGPGAGDGT

Signal sequence:
amino acids 1-20

Transmembrane domain:
amino acids 453-475

N-glycosylation sites:
amino acids 118-121, 186-189, 198-201, 211-214, 238-241,
248-251, 334-337, 357-360, 391-394

Glycosaminoglycan attachment site:
amino acids 583-586

cAMP- and cGMP-dependent protein kinase phosphorylation site:
amino acids 552-555

N-myristoylation sites:
amino acids 107-112, 152-157, 319-324, 438-443, 516-521,
612-617, 692-697, 696-701, 700-705

380/392

FIGURE 365

AATAGAAGTCCTCAGGACGGAGCAGAGGTGGCCGGCGGGCCCGGCTGACTGCGCCTCTGC
TTTCTTTCCATAACCTTTTCTTTTCGGAATCAGGCTGCTGCGAAGGGTCTAGTTC
CGGACACTAGGGTGCCCGAACGCGCTGATGCCCCGAGTGCTCGCAGGGCTTCCCGCTAAC
CATGCTGCCGCCGCCGCCGCCGAGCTGCCTTGGCGCTGCCCTGTGCTCCTGCTACTGCT
GGTGGTGTGACGCCGCCCGACCGGCGCAAGGCCATCCCCAGGCCCAGATTACCTGCG
GCGCGGCTGGATGCGGCTGCTAGCGGAGGGCGAGGGCTGCGCTCCCTGCCGGCCAGAAGA
GTGCGCCGCCGCCGCCGGGCTGCCTGGCGGGCAGGGTGCGCGACGCGTGCGGCTGCTGCTG
GGAATGCGCCAACCTCGAGGGCCAGCTCTGCGACCTGGACCCCAGTGCTCACTTCTACGG
GCACTGCGGCGAGCAGCTTGAGTGCCGGCTGGACACAGGCGGCGACCTGAGCCGCGGAGA
GGTGCGGGAACCTCTGTGTGCCTGTCTGTCGAGAGTCCGCTCTGCGGGTCCGACGGTCA
CACTACTCCAGATCTGCCGCTGCAGGAGGCGGCCCGCGCTCGGCCCGATGCCAACCT
CACTGTGGCACACCCGGGGCCCTGCCAATCGGGGCCCCAGATCGTGTACATCCATATGA
CACTTGGAATGTGACAGGGCAGGATGTGATCTTTGGCTGTGAAGTGTTTGCCTACCCCAT
GGCCTCCATCGAGTGAGGAAGGATGGCTTGGACATCCAGCTGCCAGGGGATGACCCCCA
CATCTCTGTGCAGTTTAGGGGTGGACCCAGAGGTTTGAGGTGACTGGCTGGCTGCAGAT
CCAGGCTGTGCGTCCAGTGATGAGGGCACTTACCGCTGCCTTGGCCGCAATGCCCTGGG
TCAAGTGGAGGCCCCTGCTAGCTTGACAGTGCTCACACCTGACCAGCTGAACTCTACAGG
CATCCCCAGCTGCGATCACTAAACCTGGTTCTTGAGGAGGAGGCTGAGAGTGAAGAGAA
TGACGATTACTACTAGGTCCAGAGCTCTGGCCCATGGGGGTGGGTGAGCGGCTATAGTGT
TCATCCCTGCTCTTGAAAAGACCTGGAAAGGGGAGCAGGGTCCCTTCATCGACTGCTTTC
ATGCTGTCAGTAGGGATGATCATGGGAGGCCTATTTGACTCCAAGGTAGCAGTGTGGTAG
GATAGAGACAAAAGCTGGAGGAGGGTAGGGAGAGAAGCTGAGACCAGGACCGGTGGGGTA
CAAAGGGGGCCATGCAGGAGATGCCCTGGCCAGTAGGACCTCCAACAGGTTGTTTCCAG
GCTGGGGTGGGGGCCTGAGCAGACACAGAGGTGCAGGCACCAGGATTCTCCACTTCTTCC
AGCCCTGCTGGGCCACAGTTCTAACTGCCCTTCTCCAGGCCCTGGTTCTTGCTATTTTC
CTGGTCCCCAACGTTTATCTAGCTTGTGTTGCCCTTCCCCAACTCATCTTCCAGAACTT
TTCCCTCTCTCCTAAGCCCCAGTTGCACCTACTAACTGCAGTCCCTTTTGCTGTCTGCCG
TCTTTTGTACAAGAGAGAGAACAGCGGAGCATGACTTAGTTTCAGTGCAGAGAGATTT

381/392

FIGURE 366

MLPPFRPAAALALPVLLLLLVLTTPPPTGARPSPGPDYLRRGWMRLLAEGEGCAPCRPEE
CAAPRGCLAGRVRDACGCCWECANLEGQLCDLDP SAHFYGHCGEQLECR LDTGGDL SRGE
VPEPLCACRSQSPLCGSDGHTYSQICRLQEAARARPDANLTV AHPGPCESGPQIVSHPYD
TWNVTGQDVIFGCEVFAYPMASIEWRKDGLDIQLPGDDPHISVQFRGGPQRFEVTGWLQI
QAVRPSDEGTYRCLGRNALGQVEAPASLTVLTPDQLNSTGIPQLRSLNLVP EEEAESEEN
DDYY

Important features of the protein:**Signal peptide:**

1-30

Transmembrane domain:

None

N-glycosylation site:

159-163, 183-187, 277-281

Tyrosine kinase phosphorylation site:

244-252

N-myristoylation site:

52-58, 66-72, 113-119, 249-255

Kazal-type serine protease inhibitor domain:

121-168

Immunoglobulin domain:

186-255

Insulin-like growth factor binding proteins:

53-90

382/392

FIGURE 367

AGACGCTACAGGATGGAGCGGGGCGCAGGAGCCAAGCTGCTGCCGCTGCTGCTGCTTCTG
CGGGCGACTGGTTTTACATGTGCACAGACAGATGGCCGGAACGGCTACACGGCGGTATC
GAAGTGACCAGCGGGGTCCCTGGGGCGACTGGGCCTGGCCTGAGATGTGTCCCGATGGA
TTCTTCGCCAGCGGGTTCTCGCTCAAGGTGGAGCCTCCCCAAGGCATTCTGGCGACGAC
ACTGCACTGAATGGGATCAGGCTGCACTGCGCGCGCGGGAACGTCCTAGGCAATACGCAC
GTGGTAGAGTCCCAGTCTGGAAGCTGGGGCGAATGGAGTGAGCCGCTGTGGTGTGCGCGC
GGCGCCTACCTAGTGGCTTTCTCGCTTCGCGTGGAGGCACCCACGACCCTCGGTGACAAC
ACAGCAGCGAACAACGTGCGCTTCGCTGTTGAGACGGCGAGGAACTGCAGGGGCCTGGG
CTGAGCTGGGGAGACTTTGGAGACTGGAGTGACCATTGCCCAAGGGCGCGTGCGGCCTG
CAGACCAAGATCCAGGGACCTAGAGGCCTCGGCGATGACACTGCGCTGAACGACGCGCGC
TTATTCTGCTGCCGCAGTGAACGGCGCCGCCGCCGCTCTCTCCCGGGCCAGGAGGC
TAGTCCCACCTCTTGCTATTAAAGCTTCTCTGAGTTG

383/392

FIGURE 368

MERGAGAKLLPLLLLLLRATGFTCAQTDGRNGYTAVIEVTSGGPWGDWAWPEMCPDGFFAS
GFSLKVEPPQGIPGDDTALNGIRLHCARGNVLGNTHVVESQSGSWGGEWSEPLWCRGGAYL
VAFSLRVEAPTTLGDNNTAANNVRFRCSDGEELQGPGLSWGDFGDWSDHCPKGACGLQTKI
QGPRGLGDDTALNDARLFCCRS

Important features of the protein:

Signal peptide:

1-24

Transmembrane domain:

None

N-myristoylation site:

41-47

89-95

156-162

Growth factor and cytokines receptors family signature 2:

103-110

384/392

FIGURE 369

GCCAACACTGGCCAAACCTCGGAGACCGTCCTGCGCTCTCTGGAGACGCGCTGTCCGCGC
CCAGGGTGGTGCCATGTGGGGCGCTCGCCGCTCGTCCGTCTCCTCATCCTGGAACGCCGC
TTCGCTCCTGCAGCTGCTGCTGGCTGCGCTGCTGGCGGCGGGGGCGAGGGCCAGCGGCGA
GTACTGCCACGGCTGGCTGGACGCGCAGGGCGTCTGGCGCATCGGCTTCCAGTGTCCTCGA
GCGCTTCGACGGCGGGCGACGCCACCATCTGCTGCGGCAGCTGCGCGTTGCGCTACTGCTG
CTCCAGCGCCGAGGCGCGCTGGACCAGGGCGGCTGCGACAATGACCGCCAGCAGGGCGC
TGGCGAGCCTGGCCGGGCGGACAAAGACGGCCCCGACGGCTCGGCAGTGCCCATCTACGT
GCCGTTCCCTCATTGTTGGCTCCGTGTTTGTGCGCTTTATCATCTTGGGGTCCCTGGTGGC
AGCCTGTTGCTGCAGATGTCTCCGGCCTAAGCAGGATCCCCAGCAGAGCCGAGCCCCAGG
GGGTAAACCGCTTGATGGAGACCATCCCCATGATCCCCAGTGCCAGCACCTCCCGGGGGTC
GTCCTCACGCCAGTCCAGCACAGCTGCCAGTTCAGCTCCAGCGCCAACTCAGGGGCCCCG
GGCGCCCCCAACAAGGTCACAGACCAACTGTTGCTTGCCGGAAGGGACCATGAACAACGT
GTATGTCAACATGCCCACGAATTTCTCTGTGCTGAACTGTCAGCAGGCCACCCAGATTGT
GCCACATCAAGGGCAGTATCTGCATCCCCCATACTGCGGGTACACGGTGCAGCACGACTC
TGTGCCCATGACAGCTGTGCCACCTTTCATGGACGGCCTGCAGCCTGGCTACAGGCAGAT
TCAGTCCCCCTTCCCTCACACCAACAGTGAACAGAAGATGTACCCAGCGGTGACTGTATA
ACCGAGAGTCACTGGTGGGTTCTTTACTGAAGGGAGACGAAGGCAGGGGTGGATTTTCG
AGGTGGAAGT

385/392

FIGURE 370

MWGARRSSVSSSWNAASLLQLLLAALLAAGARASGEYCHGWLDAQGVWRIGFQCPERFDG
GDATICCGSCALRYCCSSAEARLDQGGCDNDRQQGAGEPGRADKDGPDGSAVPIYVPFLI
VGSVFVAFIILGSLVAACCCRLRPKQDPQQSRAPGGNRLMETIPMIPSASTSRGSSSRQ
SSTAASSSSSANS GARAPPTRSQTNCCLP EGTMMNVYVNMPTNFSVLNCQQATQIVPHQG
QYLHPPYVG YTVQHDSVPMTAVPPFMDGLQPGYRQIQSPFPHTNSEQKMPAVTV

Important features of the protein:

Signal peptide:

1-33

Transmembrane domain:

54-78

N-glycosylation site:

223-226

cAMP- and cGMP-dependent protein kinase phosphorylation site:

5-8

N-myristoylation site:

3-8, 30-35, 60-65, 86-91, 132-137, 211-216, 268-273

Prokaryotic membrane lipoprotein lipid attachment site:

128-138

386/392

FIGURE 371

CACCAGACAGCACTCCAGCACTCTGTTTGGGGGGCATTCGAAACAGCAAAATCACTCATA
AAAGGCCAAAAAATTGCAAAAAAATAGTAATAACCAGCATGGCACTAAATAGACCATGA
AAAGACATGTGTGTGCAGTATGAAAAATTGAGACAGGAAGGCAGAGTGTCTAGCTTGTTCCT
CCTCAGCTGGGAATGTGCATCAGGCAACTCAAGTTTTTACCACGGCATGTGTCTGTGAA
TGTCCGCAAAACATTCTCTCTCCCCAGCCTTCATGTGTAACTGGGGATGATGTGGACC
TGGGCACTGTGGATGCTCCCTTCACTCTGCAAAATTCAGCCTGGCAGCTCTGCCAGCTAAG
CCTGAGAACATTTCCCTGTGTCTACTACTATAGGAAAAATTTAACCTGCACTTGGAGTCCA
GGAAAGGAAACCAGTTATACCCAGTACACAGTTAAGAGAACTTACGCTTTTGGAGAAAAA
CATGATAATTGTACAACCAATAGTTCTACAAGTGAAAATCGTGCTTCGTGCTCTTTTTTC
CTTCCAAGAATAACGATCCAGATAATTATACCATTTGAGGTGGAAGCTGAAAATGGAGAT
GGTGTAAATTAAATCTCATATGACATACTGGAGATTAGAGAACATAGCGAAAACCTGAACCA
CCTAAGATTTTCCGTGTGAAACCAAGTTTTGGGCATCAAACGAATGATTCAAATTGAATGG
ATAAACCTGAGTTGGCGCCTGTTTCATCTGATTAAATAACACACTTCGATTCAAGGACA
GTCAACAGTACCAGCTGGATGGAAGTCAACTTCGCTAAGAACCGTAAGGATAAAAAACCAA
ACGTACAACCTCACGGGGCTGCAGCCTTTTACAGAATATGTCATAGCTCTGCGATGTGCG
GTCAAGGAGTCAAAGTTCTGGAGTGACTGGAGCCAAGAAAAAATGGGAATGACTGAGGAA
GAAGCTCCATGTGGCCTGGAAGTGTGGAGAGTCTGAAACCAGCTGAGGCGGATGGAAGA
AGGCCAGTGCGGTTGTTATGGAAGAAGGCAAGAGGAGCCCCAGTCCTAGAGAAAACACTT
GGCTACAACATATGGTACTATCCAGAAAGCAACACTAACCTCACAGAAACAATGAACACT
ACTAACCAGCAGCTTGAAGTGCATCTGGGAGGCGAGAGCTTTTGGGTGTCTATGATTTCT
TATAATTCTCTTGGGAAGTCTCCAGTGGCCACCCTGAGGATTCCAGCTATTCAAGAAAAA
TCATTTCAAGTGCATTGAGGTGATGCAGGCCTGCGTTGCTGAGGACCAGCTAGTGGTGAAG
TGGCAAAGCTCTGCTCTAGACGTGAACACTTGGATGATTGAATGGTTTCCGGATGTGGAC
TCAGAGCCCACCACCCTTTCTGGGAATCTGTGTCTCAGGCCACGAAGTGGACGATCCAG
CAAGATAAATTTAAACCTTTCTGGTGCTATAACATCTCTGTGTATCCAATGTTGCATGAC
AAAGTTGGCGAGCCATATTCCATCCAGGCTTATGCCAAAGAAGGCGTTCCATCAGAAGGT
CCTGAGACCAAGGTGGAGAACATTGGCGTGAAGACGGTCACGATCACATGGAAAAGAGATT
CCCAAGAGTGAGAGAAAGGGTATCATCTGCAACTACACCATCTTTTACCAAGCTGAAGGT
GGAAAAGGATTCTGTAAGCACGCCCATAGCGAAGTGGAAAAAACCCTAAGCCCCAGATA
GATGCTATGGATAGACCTGTTGTAGGCATGGCTCCCCATCTCATTTGTGACTTGCAACCT
GGCATGAATCACTTAGCTTCTTTAAATCTCTCTGAAAATGGGGCCAAGAGCACCCACCTT
TTGGGGTTTTTGGGGGTAAATGAGAGTGAAGTGACAGTACCTGAGAGGAGAGTCCTGAGG
AAATGGAAGGAGTTGTTATTAATTTTGTCTGGTTAGGCCCTGAATTGACCTCCCGGGAGCT
CCCCGACCATCATTTCCAGGAATGGCGTGCCTGGCTTAAAGAGTGAGGAGGAACAGACCC
TGTCACCATGACTTCTACTGCCCCTGCCAAATCATGCTTTTGTTTTTTCAGTCCACCTTAT
CTCCTGACATCTTAAATACTGGGCAAGGCTTGGATTCTTGCTTAGGCTAAATAATTTTTT
CTTATGGTAAATAACACGTAAAAATTTTTTCCAGTTTAAACATTTGAAAGTGTACAATTT
AGTGGCATTAGAAGCATTACAAATATTGTGCAACCATCACCATTATTTCCAGAACTCTTC
TATTTCTGCCCCAAATAGAAGCCCTATACCCATTATTAGTCACTCCCATTTCTCTCCTC
CCACAGCCCCCTGGCAACTACCAAACTGCTTTGTGTCTCTATGGATTGCCTATTTTGGATA
TTTCATATACATAGAATCATAAANTAAAAA

387/392

FIGURE 372

MCIRQLKFFTTACVCECPQNILSPQPSCVNLGMMWTWALWMLPSLCKFSLAALPAKPENI
SCVYYYRKNLTCTWSPGKETSQYTVKRTYAFGEKHDNCTTNSSTSENRASCSFFLPRI
TIPDNYTIEVEAENG DGVIKSHMTYWRL ENIAKTEPPKIFRVKPV LGIKRMIQIEWIKPE
LAPVSSDLKYTLRFRTVNSTSWMEVNF AKNRKDKNQTYNLTGLQPFTEYVIALRC AVKES
KFWSDWSQEKMGMT EEEAPCGLELWRVLKPAEADGRRPVRL LWKKARGAPVLEKTLGYNI
WYYPESNTNLTETMNTTNQQLHLHGGEFWVSMISYNSLGKSPVATLRIPAIQEKSFQC
IEVMQACVAEDQLVVWKQSSALDVNTWMI EWFPDVDSEPTTLSWESVSQATNWTIQQDKL
KPFWCYNISVYPMLHDKVGEPYSIQAYAKEGVPSEGPETKVENIGVKT VTTITWKEIPKSE
RKGII CNYTIFYQAEGGKGFC KHAHSEVEKNPKPQIDAMDRPVVGMAPP SHCDLQPGMNH
LASLNLSENGAKSTHLLGFWGLNESEVTV PERRVLRKWKELL

Important features of the protein:

Signal peptide:

1-46

Transmembrane domain:

None

N-glycosylation site:

59-63, 69-73, 99-103, 103-107, 125-129, 198-202, 215-219, 219-223, 309-313, 315-319, 412-416, 427-431, 487-491, 545-549, 563-567

N-myristoylation site:

32-38, 137-143, 483-489, 550-556, 561-567

Amidation site:

274-278

Growth factor and cytokines receptors family signature 1:

62-75

Fibronectin type III domain:

54-144

154-247

388/392

FIGURE 373

CCAGGTCCAACCTGCACCTCGGTTCTATCGATTGAATTCCTCGGGGATCCTCTAGAGATCC
CTCGACCTCGACCCACGCGTCCGCCAAGCTGGCCCTGCACGGCTGCAAGGGAGGCTCCTG
TGGACAGGCCAGGCAGGTGGGCCTCAGGAGGTGCCTCCAGGCGGCCAGTGGGCCTGAGGC
CCCAGCAAGGGCTAGGGTCCATCTCCAGTCCCAGGACACAGCAGCGGCCACCATGGCCAC
GCCTGGGCTCCAGCAGCATCAGCAGCCCCCAGGACCGGGGGAGGCACAGGTGGCCCCCAC
CACCCGGAGGAGCAGCTCCTGCCCCCTGTCCGGGGGATGACTGATTCTCCTCCGCCAGGCC
ACCCAGAGGAGAAGGCCACCCCGCCTGGAGGCACAGGCCATGAGGGGCTCTCAGGAGGTG
CTGCTGATGTGGCTTCTGGTGTGGCAGTGGGCGGCACAGAGCAGCCTACCGGCCCGGC
CGTTAGGGTGTGTGCTGTCCCGGGCTCACGGGGACCCTGTCTCCGAGTCGTTCTGTGCAGC
GTGTGTACCAGCCCTTCTCACCACCTGCGACGGGCACCGGGCCTGCAGCACCTACCGAA
CCATTATAGGACCGCCTACCGCCGACGCCCTGGGCTGGCCCCCTGCCAGGCCTCGCTACG
CGTCTGCCCGGGCTGGAAGAGGACAGCGGGCTTCTTGGGGCCTGTGGAGCAGCAATAT
GCCAGCCGCATGCCGGAACGGAGGGAGCTGTGTCCAGCCTGGCCGCTGCCGCTGCCCTG
CAGGATGGCGGGGTGACACTTGCCAGTCAGATGTGGATGAATGCAGTGCTAGGAGGGGCG
GCTGTCCCCAGCGCTGCATCAACACCGCCGGCAGTTACTGGTGCCAGTGTTGGGAGGGGC
ACAGCCTGTCTGCAGACGGTACACTCTGTGTGCCCAAGGGAGGGCCCCCAGGGTGGCCC
CCAACCCGACAGGAGTGGACAGTGCAATGAAGGAAGAAGTGCAGAGGCTGCAGTCCAGGG
TGGACCTGCTGGAGGAGAAGCTGCAGCTGGTGCTGGCCCCACTGCACAGCCTGGCCTCGC
AGGCACTGGAGCATGGGCTCCCGGACCCCGGCAGCCTCCTGGTGCACTCCTTCCAGCAGC
TCGGCCGCATCGACTCCCTGAGCGAGCAGATTTCTTCTTGGAGGAGCAGCTGGGGTCTCT
GCTCCTGCAAGAAAGACTCGTGACTGCCCAGCGCCCCAGGCTGGACTGAGCCCCCTCACGC
CGCCCTGCAGCCCCCATGCCCCCTGCCCAACATGCTGGGGGTCCAGAAGCCACCTCGGGGT
GACTGAGCGGAAGGCCAGGCAGGGCCTTCTCCTTTCTCCTCCCTTCCCTCGGGAGG
GTCCCCAGACCCTGGCATGGGATGGGCTGGGATTTTTTTTGTGAATCCACCCCTGGCTAC
CCCCACCCTGGTTACCCCAACGGCATCCCAAGGCCAGGTGGGCCCTCAGCTGAGGGAAGG
TACGAGTTCCTCTGCTGGAGCCTGGGACCCATGGCACAGGCCAGGCAGCCCGGAGGCTGG
GTGGGGCCTCAGTGGGGGCTGCTGCCTGACCCCCAGCACAATAAAAATGAAACGTGAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAGGGCGGCCGCGACTCTAGAGTC
GACCTGCAGAAGCTTGGCCGCCATGGCCCAACTTGTTTATTGCAGCTTATAATGGTTACAAAT

389/392


FIGURE 374

MTDSPPPGHPEEKATPPGGTGHEGLSGGAADVASGVGSGRHRARLPARPLGCVLSRAHGD
PVSESFVQRVYQPFLTTCDGHRACSTYRTIYRTAYRRSPGLAPARPRYACCPGWKRTSGL
PGACGAATICQPPCRNGGSCVQPGRCRCPAGWRGDTQCSDVDECSARRGGCPQRCINTAGS
YWCQCWEGHSLSADGTLCPKGGPPRVAPNPTGVDSAMKEEVQRLQSRVDLLEEKQLQLVL
APLHSLASQALEHGLPDPGSLLVHSFQQLGRIDSLSEQISFLEEQLGSCSCKKDS

390/392

FIGURE 375


Wholemount In Situ with Prol449 Orthologue




PRO1449 is expressed in vasculature of many inflamed and diseased tissues

FIGURE 376

Human tumor tissue (BF)



Human tumor tissue (DF)




DNA64907 in situ

392/392

FIGURE 377

Mouse orthologue of PRO1449 induces ectopic vessels in the eyes of chicken embryos



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 January 2002 (03.01.2002)

PCT

(10) International Publication Number
WO 02/000690 A3

(51) International Patent Classification ⁷ : C07K 14/47, C12N 15/12, A61K 38/17, C07K 16/18		09/808,689	14 March 2001 (14.03.2001)	US
		09/816,744	22 March 2001 (22.03.2001)	US
		09/828,366	5 April 2001 (05.04.2001)	US
(21) International Application Number: PCT/US01/19692		09/854,208	10 May 2001 (10.05.2001)	US
		09/854,280	10 May 2001 (10.05.2001)	US
(22) International Filing Date: 20 June 2001 (20.06.2001)		09/866,028	25 May 2001 (25.05.2001)	US
		09/866,034	25 May 2001 (25.05.2001)	US
(25) Filing Language: English		PCT/US01/17092	25 May 2001 (25.05.2001)	US
		09/870,574	30 May 2001 (30.05.2001)	US
(26) Publication Language: English		PCT/US01/17443	30 May 2001 (30.05.2001)	US
		PCT/US01/17800	1 June 2001 (01.06.2001)	US
(30) Priority Data:				
60/213,637	23 June 2000 (23.06.2000)	US	(71) Applicant (for all designated States except US): GENEN- TECH, INC. [US/US]; 1 DNA Way, South San Francisco, CA 94080-4990 (US).	
60/219,556	20 July 2000 (20.07.2000)	US		
60/220,624	25 July 2000 (25.07.2000)	US	(72) Inventors; and	
60/220,664	25 July 2000 (25.07.2000)	US		
PCT/US00/20710	28 July 2000 (28.07.2000)	US	(75) Inventors/Applicants (for US only): BAKER, Kevin, P. [GB/US]; 14006 Indian Run Drive, Darnestown, MD 20878 (US). FERRARA, Napoleone [US/US]; #704, 2090 Pacific Avenue, San Francisco, CA 94109 (US). GERBER, Hanspeter [CH/US]; #5, 1121 Tennessee Street, San Francisco, CA 94107 (US). GERRITSEN, Mary, E. [CA/US]; 541 Parrott Drive, San Mateo, CA 94402 (US). GODDARD, Audrey [CA/US]; 110 Congo Street, San Francisco, CA 94131 (US). GODOWSKI, Paul, J. [US/US]; 25 Orange Court, Hillsborough, CA 94010 (US). GURNEY, Austin, L. [US/US]; 1 Debbie Lane, Belmont, CA 94002 (US). HILLAN, Kenneth, J. [GB/US]; 64 Seward Street, San Francisco, CA 94114 (US). MARSTERS, Scot, A. [US/US]; 990 Cherry Street, San Carlos, CA 94070 (US). PAN, James [CA/US]; 2705 Coronet Boulevard, Belmont, CA 94002 (US). PAONI, Nicholas, F. [US/US]; 1756 Terrace Drive, Belmont, CA 94002 (US). STEPHAN, Jean-Philippe, F. [FR/US]; 320 C Lansdale Avenue, Millbrae, CA 94030 (US). WATAN- ABE, Colin, K. [US/US]; 128 Corliss Drive, Moraga, CA 94556 (US). WILLIAMS, P., Mickey [US/US]; 509 Alto Avenue, Half Moon Bay, CA 94019 (US). WOOD, William, I. [US/US]; 35 Southdown Court, Hillsborough, CA 94010 (US). YE, Weilan [CN/US]; 119 Barkentine Street, Foster City, CA 94404 (US).	
PCT/US00/20710	28 July 2000 (28.07.2000)	US		
60/222,695	2 August 2000 (02.08.2000)	US		
09/643,657	17 August 2000 (17.08.2000)	US		
PCT/US00/23522	23 August 2000 (23.08.2000)	US		
PCT/US00/23328	24 August 2000 (24.08.2000)	US		
60/230,978	7 September 2000 (07.09.2000)	US		
60/000,000	15 September 2000 (15.09.2000)	US		
09/664,610	18 September 2000 (18.09.2000)	US		
09/665,350	18 September 2000 (18.09.2000)	US		
60/242,922	24 October 2000 (24.10.2000)	US		
09/709,238	8 November 2000 (08.11.2000)	US		
PCT/US00/30952	8 November 2000 (08.11.2000)	US		
PCT/US00/30873	10 November 2000 (10.11.2000)	US		
PCT/US00/32678	1 December 2000 (01.12.2000)	US		
09/747,259	20 December 2000 (20.12.2000)	US		
PCT/US00/34956	20 December 2000 (20.12.2000)	US		
09/767,609	22 January 2001 (22.01.2001)	US		
09/796,498	28 February 2001 (28.02.2001)	US		
PCT/US01/06520	28 February 2001 (28.02.2001)	US		
PCT/US01/06666	1 March 2001 (01.03.2001)	US		
09/802,706	9 March 2001 (09.03.2001)	US		

[Continued on next page]

(54) Title: COMPOSITIONS AND METHODS FOR THE DIAGNOSIS AND TREATMENT OF DISORDERS INVOLVING ANGIOGENESIS

(57) Abstract: Compositions and methods are disclosed for stimulating or inhibiting angiogenesis and/or cardiovascularization in mammals, including humans. Pharmaceutical compositions are based on polypeptides or antagonists thereto that have been identified for one or more of these uses. Disorders that can be diagnosed, prevented, or treated by the compositions herein include trauma such as wounds, various cancers, and disorders of the vessels including atherosclerosis and cardiac hypertrophy. In addition, the present invention is directed to novel polypeptides and to nucleic acid molecules encoding those polypeptides. Also provided herein are vectors and host cells comprising those nucleic acid sequences, chimeric polypeptide molecules comprising the polypeptides of the present invention fused to heterologous polypeptide sequences, antibodies which bind to the polypeptides of the present invention and to methods for producing the polypeptides of the present invention.

WO 02/000690 A3



(74) **Agents:** AGARWAL, Atulya, R. et al.; c/o GENENTECH, INC., MS49, 1 DNA Way, South San Francisco, CA 94080-4990 (US).

patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(81) **Designated States (national):** AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

Published:

— with international search report

(88) **Date of publication of the international search report:**
13 March 2003

(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/19692

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C07K14/47 A61K38/17 C07K16/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EP0-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,L	WO 99 46281 A (BAKER KEVIN P ;CHEN JIAN (US); GENENTECH INC (US); GURNEY AUSTIN () 16 September 1999 (1999-09-16) see PR0181	1-19
X,L	US 5 968 744 A (CORLEY-NEIL C ET AL) 19 October 1999 (1999-10-19) L: priority. the whole document	1-19
X	WO 99 50405 A (GENETICS INST) 7 October 1999 (1999-10-07) see pk65_4 and seq.ID's 11 and 12	1-19
X	WO 99 43802 A (KATO SEISHI ;KIMURA TOMOKO (JP); PROTEGENE INC (JP); SEKINE SHINGO) 2 September 1999-(1999-09-02)- see clone HP02239.	1-19
-/--		

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

21 August 2002

Date of mailing of the international search report

29. 11. 2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Smalt, R

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/19692

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 10363 A (WEI YING FEI ;HUMAN GENOME SCIENCES INC (US); ROSEN CRAIG A (US);) 4 March 1999 (1999-03-04) see secreted protein 24 ---	1-19
X	WO 99 33979 A (CHIRON CORP) 8 July 1999 (1999-07-08) the whole document ---	1-19
A	WO 95 29242 A (CHILDRENS MEDICAL CENTER) 2 November 1995 (1995-11-02) the whole document ---	
A	EP 0 335 243 A (HARVARD COLLEGE) 4 October 1989 (1989-10-04) the whole document ---	
A	FOLKMAN J: "CLINICAL APPLICATIONS OF RESEARCH ON ANGIOGENESIS" NEW ENGLAND JOURNAL OF MEDICINE, THE, MASSACHUSETTS MEDICAL SOCIETY, WALTHAM, MA, US, vol. 333, no. 26, 28 December 1995 (1995-12-28); pages 1757-1763, XP000654557 ISSN: 0028-4793 the whole document ---	
A	KLASBRUN M: "REGULATORS OF ANGIOGENESIS: STIMULATORS, INHIBITORS, AND EXTRACELLULAR MATRIX" JOURNAL OF CELLULAR BIOCHEMISTRY, LISS, NEW YORK, NY, US, vol. 47, no. 3, 1 November 1991 (1991-11-01), pages 199-200, XP000616588 ISSN: 0730-2312 the whole document ---	
P,X, L	WO 00 53756 A (BOTSTEIN DAVID;BAKER KEVIN P ; GENENTECH INC (US); ASHKENAZI AVI J) 14 September 2000 (2000-09-14) see seq.ID.322; L: priority. ---	1-19
P,X, L	WO 00 53751 A (BAKER KEVIN P ;GENENTECH INC (US); NAPIER MARY A (US); ASHKENAZI A) 14 September 2000 (2000-09-14) the whole document ---	1-21,23, 27-29, 31,32, 34,35
P,X	WO 01 00828 A (CORIXA CORP ;FANGER GARY R (US); RETTER MARC W (US); WANG TONGTONG) 4 January 2001 (2001-01-04) see seq.ID.327 ---	1-21,23, 27-29, 31,32, 34,35

-/--

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/19692

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E,L	WO 02 08284 A (FERRARA NAPOLEONE ;STEPHAN JEAN PHILIPPE F (US); WILLIAMS P MICKEY) 31 January 2002 (2002-01-31) See seq.ID's 1 and 120. L: priority. -----	1-35,37, 41

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 01/19692

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 20-34, 37 and 41 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-35, 37, 41 all partially

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Present claims 20-28,31,34,35,37, and 41 relate to products defined by reference to a desirable characteristic or property, namely having (ant)agonistic activity towards the protein(s) of claim 1.

The application provides no support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for any such products. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the product by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search impossible. Consequently, the search of said claims, in as far as the (ant)agonists are concerned, has been carried out for those aspects which appear to be clear, supported and disclosed, namely those parts relating to antibodies directed against the protein(s) of claim 1.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

Invention 1: 1-35, 37, 41 all partially

Nucleic acid with at least 80% identity to seq.ID.1 and protein encoded thereby, vector, host cell, method for producing the protein, chimeric protein, antibody, and pharmaceutical compositions.

Inventions 2-187: 1-42, all partially,
and as far as applicable

Subject matter as defined for invention 1. above, but limited to the respective nucleic acid sequences 3-373 (odd numbers) and the polypeptides encoded thereby (seq.ID's 4-374, even numbers).

For the sake of conciseness, the first subject matter is explicitly defined, the other subject matters are defined by analogy thereto.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/19692

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9946281	A	16-09-1999	US 6391311 B1	21-05-2002
			AU 3072199 A	27-09-1999
			CA 2321677 A1	16-09-1999
			US 2002127584 A1	12-09-2002
			US 2002094416 A1	18-07-2002
			WO 9946281 A2	16-09-1999
			US 2002156006 A1	24-10-2002
			US 2002169284 A1	14-11-2002
			ZA 9901937 A	17-04-2000
			EP 1241187 A2	18-09-2002
			AU 3075099 A	11-10-1999
			CA 2322792 A1	23-09-1999
			EP 1064382 A2	03-01-2001
			WO 9947677 A2	23-09-1999
			US 6455283 B1	24-09-2002
			EP 1241183 A2	18-09-2002
			US 2002072496 A1	13-06-2002
			US 2002142961 A1	03-10-2002
			US 2002072067 A1	13-06-2002
			US 2002072092 A1	13-06-2002
			US 2002072497 A1	13-06-2002
			US 2002103125 A1	01-08-2002
			US 2002123463 A1	05-09-2002
			US 2002132252 A1	19-09-2002
			US 2002127576 A1	12-09-2002
			US 2002132253 A1	19-09-2002
			US 2002160384 A1	31-10-2002
			AU 735081 B2	28-06-2001
			AU 1532499 A	15-06-1999
			CA 2309358 A1	03-06-1999
			EP 1241180 A2	18-09-2002
			EP 1032667 A2	06-09-2000
			WO 9927098 A2	03-06-1999
			US 6410708 B1	25-06-2002
			EP 1241185 A2	18-09-2002
			EP 1249491 A2	16-10-2002
			EP 1251139 A2	23-10-2002
			US 2002161199 A1	31-10-2002
			EP 1241179 A2	18-09-2002
			EP 1241186 A2	18-09-2002
			AU 741133 B2	22-11-2001
			AU 3757099 A	08-11-1999
			CA 2324297 A1	28-10-1999
			EP 1071773 A1	31-01-2001
			JP 2002512032 T	23-04-2002
			WO 9954467 A1	28-10-1999
			US 2002147322 A1	10-10-2002
			EP 1241251 A2	18-09-2002
			US 2002119118 A1	29-08-2002
US 5968744	A	19-10-1999	US 2002103342 A1	01-08-2002
			US 6348576 B1	19-02-2002
WO 9950405	A	07-10-1999	AU 3457299 A	18-10-1999
			CA 2324492 A1	07-10-1999
			EP 1068307 A1	17-01-2001
			JP 2002509722 T	02-04-2002
			WO 9950405 A1	07-10-1999

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/19692

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9943802	A	02-09-1999	AU	2549399 A	15-09-1999
			CA	2322711 A1	02-09-1999
			EP	1056847 A2	06-12-2000
			WO	9943802 A2	02-09-1999
			JP	2002506615 T	05-03-2002
WO 9910363	A	04-03-1999	AU	8921598 A	16-03-1999
			CA	2302387 A1	04-03-1999
			EP	1017707 A1	12-07-2000
			JP	2002532054 T	02-10-2002
			WO	9910363 A1	04-03-1999
			US	6410709 B1	25-06-2002
			US	2001016647 A1	23-08-2001
WO 9933979	A	08-07-1999	AU	1929599 A	19-07-1999
			EP	1042470 A2	11-10-2000
			JP	2002511231 T	16-04-2002
			WO	9933979 A2	08-07-1999
			US	2002034800 A1	21-03-2002
WO 9529242	A	02-11-1995	US	5639725 A	17-06-1997
			US	5792845 A	11-08-1998
			AU	692865 B2	18-06-1998
			AU	2461795 A	16-11-1995
			BR	9507479 A	16-09-1997
			CA	2188813 A1	02-11-1995
			CN	1149319 A	07-05-1997
			CZ	9603122 A3	14-05-1997
			EP	0758390 A1	19-02-1997
			HU	76095 A2	30-06-1997
			JP	9512173 T	09-12-1997
			NZ	285501 A	26-02-1998
			WO	9529242 A1	02-11-1995
			US	5885795 A	23-03-1999
			US	5733876 A	31-03-1998
			US	5776704 A	07-07-1998
			ZA	9503419 A	11-01-1996
EP 0335243	A	04-10-1989	US	4900673 A	13-02-1990
			AT	96844 T	15-11-1993
			AU	621358 B2	12-03-1992
			AU	3433189 A	16-10-1989
			CA	1331356 A1	09-08-1994
			DE	68910354 D1	09-12-1993
			DE	68910354 T2	03-03-1994
			DK	233390 A	26-11-1990
			EP	0335243 A2	04-10-1989
			IE	890954 L	28-09-1989
			JP	3503641 T	15-08-1991
			PT	90129 A	10-11-1989
			WO	8909277 A1	05-10-1989
			ZA	8902270 A	24-12-1991
WO 0053756	A	14-09-2000	AU	1748200 A	19-06-2000
			AU	1749900 A	12-07-2000
			AU	2192800 A	12-07-2000
			AU	2224800 A	28-09-2000
			AU	2399300 A	28-09-2000

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/19692

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0053756	A		AU 2495200 A	28-09-2000
		AU 2596700 A	28-09-2000	
		AU 2600800 A	28-09-2000	
		AU 2879400 A	31-07-2001	
		AU 2883600 A	28-09-2000	
		AU 3072199 A	27-09-1999	
		AU 3107700 A	28-09-2000	
		AU 3381600 A	28-09-2000	
		AU 3514400 A	28-09-2000	
		CA 2321677 A1	16-09-1999	
		EP 1173563 A1	23-01-2002	
		EP 1159419 A1	05-12-2001	
		EP 1220905 A2	10-07-2002	
		EP 1135485 A2	26-09-2001	
		EP 1141284 A2	10-10-2001	
		EP 1159422 A1	05-12-2001	
		EP 1141289 A2	10-10-2001	
		US 2002119130 A1	29-08-2002	
		WO 0053753 A2	14-09-2000	
		WO 0053754 A1	14-09-2000	
		WO 0053755 A2	14-09-2000	
		WO 0153486 A1	26-07-2001	
		WO 0053756 A2	14-09-2000	
		WO 0053757 A2	14-09-2000	
		WO 0053758 A2	14-09-2000	
		US 2002127584 A1	12-09-2002	
		US 2002094416 A1	18-07-2002	
		WO 0032221 A2	08-06-2000	
		WO 0053750 A1	14-09-2000	
		WO 0037638 A2	29-06-2000	
		WO 0037640 A2	29-06-2000	
		WO 0053751 A1	14-09-2000	
		WO 0053752 A2	14-09-2000	
		US 2002119118 A1	29-08-2002	
		US 2002161199 A1	31-10-2002	
		US 2002156006 A1	24-10-2002	
		US 2002169284 A1	14-11-2002	
		US 2002072496 A1	13-06-2002	
		US 2002142961 A1	03-10-2002	
		US 2002072067 A1	13-06-2002	
		US 2002072092 A1	13-06-2002	
		US 2002072497 A1	13-06-2002	
		US 2002103125 A1	01-08-2002	
<hr/>				
WO 0053751	A	14-09-2000	AU 1748200 A	19-06-2000
		AU 1749900 A	12-07-2000	
		AU 2192800 A	12-07-2000	
		AU 2224800 A	28-09-2000	
		AU 2399300 A	28-09-2000	
		AU 2495200 A	28-09-2000	
		AU 2596700 A	28-09-2000	
		AU 2600800 A	28-09-2000	
		AU 2879400 A	31-07-2001	
		AU 2883600 A	28-09-2000	
		AU 3072199 A	27-09-1999	
		AU 3107700 A	28-09-2000	
		AU 3381600 A	28-09-2000	
		AU 3514400 A	28-09-2000	

INTERNATIONAL SEARCH REPORT
Information on patent family members

International Application No
PCT/US 01/19692

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0053751	A		CA 2321677 A1	16-09-1999
			EP 1173563 A1	23-01-2002
			EP 1159419 A1	05-12-2001
			EP 1220905 A2	10-07-2002
			EP 1135485 A2	26-09-2001
			EP 1141284 A2	10-10-2001
			EP 1159422 A1	05-12-2001
			EP 1141289 A2	10-10-2001
			US 2002119130 A1	29-08-2002
			WO 0053753 A2	14-09-2000
			WO 0053754 A1	14-09-2000
			WO 0053755 A2	14-09-2000
			WO 0153486 A1	26-07-2001
			WO 0053756 A2	14-09-2000
			WO 0053757 A2	14-09-2000
			WO 0053758 A2	14-09-2000
			US 2002127584 A1	12-09-2002
			US 2002094416 A1	18-07-2002
			WO 0032221 A2	08-06-2000
			WO 0053750 A1	14-09-2000
			WO 0037638 A2	29-06-2000
			WO 0037640 A2	29-06-2000
			WO 0053751 A1	14-09-2000
			WO 0053752 A2	14-09-2000
			US 2002119118 A1	29-08-2002
			US 2002161199 A1	31-10-2002
			US 2002156006 A1	24-10-2002
			US 2002169284 A1	14-11-2002
			US 2002072496 A1	13-06-2002
			US 2002142961 A1	03-10-2002
			US 2002072067 A1	13-06-2002
			US 2002072092 A1	13-06-2002
			US 2002072497 A1	13-06-2002
			US 2002103125 A1	01-08-2002

WO 0100828	A	04-01-2001	AU 5902700 A	31-01-2001
			BR 0012069 A	27-08-2002
			CN 1367829 T	04-09-2002
			EP 1196565 A2	17-04-2002
			HU 3201757 A2	28-09-2002
			NO 20016427 A	25-02-2002
			WO 0100828 A2	04-01-2001
			US 2002168637 A1	14-11-2002
			TR 200200691 T2	21-10-2002

WO 0208284	A	31-01-2002	US 2002042367 A1	11-04-2002
			US 2002010137 A1	24-01-2002
			US 2002106743 A1	08-08-2002
			US 2002052027 A1	02-05-2002
			US 2002058309 A1	16-05-2002
			US 2002102723 A1	01-08-2002
			AU 6531101 A	17-12-2001
			AU 7197301 A	05-02-2002
			AU 7315001 A	30-01-2002
			AU 7885201 A	08-01-2002
			WO 0193983 A1	13-12-2001
			WO 0200690 A2	03-01-2002
			WO 0208288 A2	31-01-2002

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/19692

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0208284	A	WO 0208284 A2	31-01-2002
		AU 1916701 A	16-07-2001
		AU 2055401 A	12-06-2001
		AU 6391000 A	19-02-2001
		AU 6802801 A	24-09-2001
		AU 6901800 A	12-06-2001
		EP 1244784 A2	02-10-2002
		EP 1250426 A2	23-10-2002
		WO 0140464 A1	07-06-2001
		WO 0149715 A2	12-07-2001
		WO 0140466 A2	07-06-2001
		WO 0168848 A2	20-09-2001
		US 2002127584 A1	12-09-2002
		US 2002094416 A1	18-07-2002
		US 2002160374 A1	31-10-2002
		US 2002146709 A1	10-10-2002
		US 2002132240 A1	19-09-2002
		US 2002142958 A1	03-10-2002
		US 2002150976 A1	17-10-2002
		US 2002132981 A1	19-09-2002
		US 2002165143 A1	07-11-2002
		US 2002156004 A1	24-10-2002
		US 2002142419 A1	03-10-2002
		US 2002102647 A1	01-08-2002
		US 2002110859 A1	15-08-2002
		US 2002142959 A1	03-10-2002
		US 2002115145 A1	22-08-2002
		US 2002132768 A1	19-09-2002
		US 2002127643 A1	12-09-2002
		US 2002156006 A1	24-10-2002
		US 2002169284 A1	14-11-2002
		US 2002072496 A1	13-06-2002
		US 2002142961 A1	03-10-2002
		US 2002072067 A1	13-06-2002
		US 2002072092 A1	13-06-2002
		US 2002072497 A1	13-06-2002
		US 2002103125 A1	01-08-2002